

HANDBOOK OF

---

**LINEAR PARTIAL  
DIFFERENTIAL  
EQUATIONS for  
ENGINEERS  
and SCIENTISTS**

---

Andrei D. Polyanin



CHAPMAN & HALL/CRC

---

A CRC Press Company

Boca Raton London New York Washington, D.C.

## Library of Congress Cataloging-in-Publication Data

---

Polianin, A. D. (Andrei Dmitrievich)

Handbook of linear partial differential equations for engineers and scientists / by Andrei D. Polyanin

p. cm.

Includes bibliographical references and index.

ISBN 1-58488-299-9

1. Differential equations, Linear--Numerical solution--Handbooks, manuals, etc. I. Title.

QA377 .P568 2001

515'.354—dc21

2001052427

CIP

This book contains information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. A wide variety of references are listed. Reasonable efforts have been made to publish reliable data and information, but the author and the publisher cannot assume responsibility for the validity of all materials or for the consequences of their use.

Apart from any fair dealing for the purpose of research or private study, or criticism or review, as permitted under the UK Copyright Designs and Patents Act, 1988, this publication may not be reproduced, stored or transmitted, in any form or by any means, electronic or mechanical, including photocopying, micro-filming, and recording, or by any information storage or retrieval system, without the prior permission in writing of the publishers, or in the case of reprographic reproduction only in accordance with the terms of the licenses issued by the Copyright Licensing Agency in the UK, or in accordance with the terms of the license issued by the appropriate Reproduction Rights Organization outside the UK.

All rights reserved. Authorization to photocopy items for internal or personal use, or the personal or internal use of specific clients, may be granted by CRC Press LLC, provided that \$1.50 per page photocopied is paid directly to Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923 USA. The fee code for users of the Transactional Reporting Service is ISBN 1-58488-299-9/02/\$0.00+\$1.50. The fee is subject to change without notice. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

The consent of CRC Press LLC does not extend to copying for general distribution, for promotion, for creating new works, or for resale. Specific permission must be obtained in writing from CRC Press LLC for such copying.

Direct all inquiries to CRC Press LLC, 2000 N.W. Corporate Blvd., Boca Raton, Florida 33431.

**Trademark Notice:** Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation, without intent to infringe.

Visit the CRC Press Web site at [www.crcpress.com](http://www.crcpress.com)

---

© 2002 by Chapman & Hall/CRC

No claim to original U.S. Government works

International Standard Book Number 1-58488-299-9

Library of Congress Card Number 2001052427

Printed in the United States of America 1 2 3 4 5 6 7 8 9 0

# FOREWORD

Linear partial differential equations arise in various fields of science and numerous applications, e.g., heat and mass transfer theory, wave theory, hydrodynamics, aerodynamics, elasticity, acoustics, electrostatics, electrodynamics, electrical engineering, diffraction theory, quantum mechanics, control theory, chemical engineering sciences, and biomechanics.

This book presents brief statements and exact solutions of more than 2000 linear equations and problems of mathematical physics. Nonstationary and stationary equations with constant and variable coefficients of parabolic, hyperbolic, and elliptic types are considered. A number of new solutions to linear equations and boundary value problems are described. Special attention is paid to equations and problems of general form that depend on arbitrary functions. Formulas for the effective construction of solutions to nonhomogeneous boundary value problems of various types are given. We consider second-order and higher-order equations as well as the corresponding boundary value problems. All in all, the handbook presents more equations and problems of mathematical physics than any other book currently available.

For the reader's convenience, the introduction outlines some definitions and basic equations, problems, and methods of mathematical physics. It also gives useful formulas that enable one to express solutions to stationary and nonstationary boundary value problems of general form in terms of the Green's function.

Two supplements are given at the end of the book. Supplement A lists properties of the most common special functions (the gamma function, Bessel functions, degenerate hypergeometric functions, Mathieu functions, etc.). Supplement B describes the methods of generalized and functional separation of variables for nonlinear partial differential equations. We give specific examples and an overview application of these methods to construct exact solutions for various classes of second-, third-, fourth-, and higher-order equations (in total, about 150 nonlinear equations with solutions are described). Special attention is paid to equations of heat and mass transfer theory, wave theory, and hydrodynamics as well as to mathematical physics equations of general form that involve arbitrary functions.

The equations in all chapters are in ascending order of complexity. Many sections can be read independently, which facilitates working with the material. An extended table of contents will help the reader find the desired equations and boundary value problems. We refer to specific equations using notation like "1.8.5.2," which means "Equation 2 in Subsection 1.8.5."

To extend the range of potential readers with diverse mathematical backgrounds, the author strove to avoid the use of special terminology wherever possible. For this reason, some results are presented schematically, in a simplified manner (without details), which is however quite sufficient in most applications.

Separate sections of the book can serve as a basis for practical courses and lectures on equations of mathematical physics.

The author thanks Alexei Zhurov for useful remarks on the manuscript.

The author hopes that the handbook will be useful for a wide range of scientists, university teachers, engineers, and students in various areas of mathematics, physics, mechanics, control, and engineering sciences.

*Andrei D. Polyinin*

# BASIC NOTATION

## Latin Characters

|                      |                                                                                                                                             |
|----------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| $\mathcal{E}$        | fundamental solution                                                                                                                        |
| $\text{Im}[A]$       | imaginary part of a complex quantity $A$                                                                                                    |
| $G$                  | Green's function                                                                                                                            |
| $\mathbb{R}^n$       | $n$ -dimensional Euclidean space, $\mathbb{R}^n = \{-\infty < x_k < \infty; k = 1, \dots, n\}$                                              |
| $\text{Re}[A]$       | real part of a complex quantity $A$                                                                                                         |
| $r, \varphi, z$      | cylindrical coordinates, $r = \sqrt{x^2 + y^2}$ and $x = r \cos \varphi, y = r \sin \varphi$                                                |
| $r, \theta, \varphi$ | spherical coordinates, $r = \sqrt{x^2 + y^2 + z^2}$ and $x = r \sin \theta \cos \varphi, y = r \sin \theta \sin \varphi, z = r \cos \theta$ |
| $t$                  | time ( $t \geq 0$ )                                                                                                                         |
| $w$                  | unknown function (dependent variable)                                                                                                       |
| $x, y, z$            | space (Cartesian) coordinates                                                                                                               |
| $x_1, \dots, x_n$    | Cartesian coordinates in $n$ -dimensional space                                                                                             |
| $\mathbf{x}$         | $n$ -dimensional vector, $\mathbf{x} = \{x_1, \dots, x_n\}$                                                                                 |
| $ \mathbf{x} $       | magnitude (length) of $n$ -dimensional vector, $ \mathbf{x}  = \sqrt{x_1^2 + x_2^2 + \dots + x_n^2}$                                        |
| $\mathbf{y}$         | $n$ -dimensional vector, $\mathbf{y} = \{y_1, \dots, y_n\}$                                                                                 |

## Greek Characters

|                |                                                                                                                                                      |
|----------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| $\Delta$       | Laplace operator                                                                                                                                     |
| $\Delta_2$     | two-dimensional Laplace operator, $\Delta_2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$                                     |
| $\Delta_3$     | three-dimensional Laplace operator, $\Delta_3 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ |
| $\Delta_n$     | $n$ -dimensional Laplace operator, $\Delta_n = \sum_{k=1}^n \frac{\partial^2}{\partial x_k^2}$                                                       |
| $\delta(x)$    | Dirac delta function; $\int_{-a}^a f(y)\delta(x-y) dy = f(x)$ , where $f(x)$ is any continuous function, $a > 0$                                     |
| $\delta_{nm}$  | Kronecker delta, $\delta_{nm} = \begin{cases} 1 & \text{if } n = m, \\ 0 & \text{if } n \neq m \end{cases}$                                          |
| $\vartheta(x)$ | Heaviside unit step function, $\vartheta(x) = \begin{cases} 1 & \text{if } x \geq 0, \\ 0 & \text{if } x < 0 \end{cases}$                            |

## Brief Notation for Derivatives

$$\partial_t w = \frac{\partial w}{\partial t}, \quad \partial_x w = \frac{\partial w}{\partial x}, \quad \partial_{tt} w = \frac{\partial^2 w}{\partial t^2}, \quad \partial_{xx} w = \frac{\partial^2 w}{\partial x^2} \quad (\text{partial derivatives})$$

$$f'_x = \frac{df}{dx}, \quad f''_{xx} = \frac{d^2 f}{dx^2}, \quad f'''_{xxx} = \frac{d^3 f}{dx^3}, \quad f^{(n)}_x = \frac{d^n f}{dx^n} \quad (\text{derivatives for } f = f(x))$$

## Special Functions (See Also Supplement A)

$$\text{Ai}(x) = \frac{1}{\pi} \int_0^\infty \cos\left(\frac{1}{3}t^3 + xt\right) dt \quad \text{Airy function; } \text{Ai}(x) = \frac{1}{\pi} \sqrt{\frac{1}{3}x} K_{1/3}\left(\frac{2}{3}x^{3/2}\right)$$

$$\text{Ce}_{2n+p}(x, q) = \sum_{k=0}^\infty A_{2k+p}^{2n+p} \cosh[(2k+p)x] \quad \text{even modified Mathieu functions, where } p = 0, 1;$$

$$\text{Ce}_{2n+p}(x, q) = \text{ce}_{2n+p}(ix, q)$$



$$\operatorname{ce}_{2n}(x, q) = \sum_{k=0}^{\infty} A_{2k}^{2n} \cos 2kx$$

$$\operatorname{ce}_{2n+1}(x, q) = \sum_{k=0}^{\infty} A_{2k+1}^{2n+1} \cos[(2k+1)x]$$

$$D_{\nu} = D_{\nu}(x)$$

$$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_0^x \exp(-\xi^2) d\xi$$

$$\operatorname{erfc} x = \frac{2}{\sqrt{\pi}} \int_x^{\infty} \exp(-\xi^2) d\xi$$

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2})$$

$$H_{\nu}^{(1)}(x) = J_{\nu}(x) + iY_{\nu}(x)$$

$$H_{\nu}^{(2)}(x) = J_{\nu}(x) - iY_{\nu}(x)$$

$$F(a, b, c; x) = 1 + \sum_{n=1}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{x^n}{n!}$$

$$I_{\nu}(x) = \sum_{n=0}^{\infty} \frac{(x/2)^{\nu+2n}}{n! \Gamma(\nu+n+1)}$$

$$J_{\nu}(x) = \sum_{n=0}^{\infty} \frac{(-1)^n (x/2)^{\nu+2n}}{n! \Gamma(\nu+n+1)}$$

$$K_{\nu}(x) = \frac{\pi}{2} \frac{I_{-\nu}(x) - I_{\nu}(x)}{\sin(\pi\nu)}$$

$$L_n^s(x) = \frac{1}{n!} x^{-s} e^x \frac{d^n}{dx^n} (x^{n+s} e^{-x})$$

$$P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n$$

$$P_n^m(x) = (1 - x^2)^{m/2} \frac{d^m}{dx^m} P_n(x)$$

$$\operatorname{Se}_{2n+p}(x, q) = \sum_{k=0}^{\infty} B_{2k+p}^{2n+p} \sinh[(2k+p)x]$$

$$\operatorname{se}_{2n}(x, q) = \sum_{k=0}^{\infty} B_{2k}^{2n} \sin 2kx$$

$$\operatorname{se}_{2n+1}(x, q) = \sum_{k=0}^{\infty} B_{2k+1}^{2n+1} \sin[(2k+1)x]$$

$$Y_{\nu}(x) = \frac{J_{\nu}(x) \cos(\pi\nu) - J_{-\nu}(x)}{\sin(\pi\nu)}$$

$$\gamma(\alpha, x) = \int_0^x e^{-\xi} \xi^{\alpha-1} d\xi$$

$$\Gamma(\alpha) = \int_0^{\infty} e^{-\xi} \xi^{\alpha-1} d\xi$$

$$\Phi(a, b; x) = 1 + \sum_{n=1}^{\infty} \frac{(a)_n}{(b)_n} \frac{x^n}{n!}$$

even  $\pi$ -periodic Mathieu functions; these satisfy the equation  $y'' + (a - 2q \cos 2x)y = 0$ , where  $a = a_{2n}(q)$  are eigenvalues

even  $2\pi$ -periodic Mathieu functions; these satisfy the equation  $y'' + (a - 2q \cos 2x)y = 0$ , where  $a = a_{2n+1}(q)$  are eigenvalues

parabolic cylinder function (see Paragraph 7.3.4-1); it satisfies the equation  $y'' + (\nu + \frac{1}{2} - \frac{1}{4}x^2)y = 0$

error function

complementary error function

Hermite polynomial

Hankel function of first kind,  $i^2 = -1$

Hankel function of second kind,  $i^2 = -1$

hypergeometric function,  $(a)_n = a(a+1) \dots (a+n-1)$

modified Bessel function of first kind

Bessel function of first kind

modified Bessel function of second kind

generalized Laguerre polynomial

Legendre polynomial

associated Legendre functions

odd modified Mathieu functions, where  $p = 0, 1$ ;

$$\operatorname{Se}_{2n+p}(x, q) = -i \operatorname{se}_{2n+p}(ix, q)$$

odd  $\pi$ -periodic Mathieu functions; these satisfy the equation  $y'' + (a - 2q \cos 2x)y = 0$ , where  $a = b_{2n}(q)$  are eigenvalues

odd  $2\pi$ -periodic Mathieu functions; these satisfy the equation  $y'' + (a - 2q \cos 2x)y = 0$ , where  $a = b_{2n+1}(q)$  are eigenvalues


Bessel function of second kind

incomplete gamma function

gamma function

degenerate hypergeometric function,  
 $(a)_n = a(a+1) \dots (a+n-1)$

## AUTHOR



**Andrei D. Polyanin, D.Sc., Ph.D.**, is a noted scientist of broad interests, who works in various areas of mathematics, mechanics, and chemical engineering sciences.

A. D. Polyanin graduated from the Department of Mechanics and Mathematics of the Moscow State University in 1974. He received his Ph.D. degree in 1981 and D.Sc. degree in 1986 at the Institute for Problems in Mechanics of the Russian (former USSR) Academy of Sciences. Since 1975, A. D. Polyanin has been a member of the staff of the Institute for Problems in Mechanics of the Russian Academy of Sciences.

Professor Polyanin has made important contributions to developing new exact and approximate analytical methods of the theory of differential equations, mathematical physics, integral equations, engineering mathematics, nonlinear mechanics, theory of heat and mass transfer, and chemical hydrodynamics. He ob-

tained exact solutions for several thousand ordinary differential, partial differential, mathematical physics, and integral equations.

Professor Polyanin is an author of 27 books in English, Russian, German, and Bulgarian, as well as over 120 research papers and three patents. He has written a number of fundamental handbooks, including A. D. Polyanin and V. F. Zaitsev, *Handbook of Exact Solutions for Ordinary Differential Equations*, CRC Press, 1995; A. D. Polyanin and A. V. Manzhirov, *Handbook of Integral Equations*, CRC Press, 1998; and A. D. Polyanin, V. F. Zaitsev, and A. Moussiaux, *Handbook of First Order Partial Differential Equations*, Gordon and Breach, 2001.

In 1991, A. D. Polyanin was awarded a Chaplygin Prize of the USSR Academy of Sciences for his research in mechanics.

**Address:** Institute for Problems in Mechanics, RAS, 101 Vernadsky Avenue, Building 1, 117526 Moscow, Russia

**E-mail:** polyanin@ipmnet.ru

# CONTENTS

## Foreword

## Basic Notation and Remarks

## Author

## Introduction. Some Definitions, Formulas, Methods, and Solutions

- 0.1. Classification of Second-Order Partial Differential Equations
  - 0.1.1. Equations with Two Independent Variables
  - 0.1.2. Equations with Many Independent Variables
- 0.2. Basic Problems of Mathematical Physics
  - 0.2.1. Initial and Boundary Conditions. Cauchy Problem. Boundary Value Problems
  - 0.2.2. First, Second, Third, and Mixed Boundary Value Problems
- 0.3. Properties and Particular Solutions of Linear Equations
  - 0.3.1. Homogeneous Linear Equations
  - 0.3.2. Nonhomogeneous Linear Equations
- 0.4. Separation of Variables Method
  - 0.4.1. General Description of the Separation of Variables Method
  - 0.4.2. Solution of Boundary Value Problems for Parabolic and Hyperbolic Equations
- 0.5. Integral Transforms Method
  - 0.5.1. Main Integral Transforms
  - 0.5.2. Laplace Transform and Its Application in Mathematical Physics
  - 0.5.3. Fourier Transform and Its Application in Mathematical Physics
- 0.6. Representation of the Solution of the Cauchy Problem via the Fundamental Solution
  - 0.6.1. Cauchy Problem for Parabolic Equations
  - 0.6.2. Cauchy Problem for Hyperbolic Equations
- 0.7. Nonhomogeneous Boundary Value Problems with One Space Variable. Representation of Solutions via the Green's Function
  - 0.7.1. Problems for Parabolic Equations
  - 0.7.2. Problems for Hyperbolic Equations
- 0.8. Nonhomogeneous Boundary Value Problems with Many Space Variables. Representation of Solutions via the Green's Function
  - 0.8.1. Problems for Parabolic Equations
  - 0.8.2. Problems for Hyperbolic Equations
  - 0.8.3. Problems for Elliptic Equations
  - 0.8.4. Comparison of the Solution Structures for Boundary Value Problems for Equations of Various Types
- 0.9. Construction of the Green's Functions. General Formulas and Relations
  - 0.9.1. Green's Functions of Boundary Value Problems for Equations of Various Types in Bounded Domains
  - 0.9.2. Green's Functions Admitting Incomplete Separation of Variables
  - 0.9.3. Construction of Green's Functions via Fundamental Solutions

- 0.10. Duhamel's Principles in Nonstationary Problems
  - 0.10.1. Problems for Homogeneous Linear Equations
  - 0.10.2. Problems for Nonhomogeneous Linear Equations
- 0.11. Transformations Simplifying Initial and Boundary Conditions
  - 0.11.1. Transformations That Lead to Homogeneous Boundary Conditions
  - 0.11.2. Transformations That Lead to Homogeneous Initial and Boundary Conditions

## 1. Parabolic Equations with One Space Variable

- 1.1. Constant Coefficient Equations
  - 1.1.1. Heat Equation  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2}$
  - 1.1.2. Equation of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + \Phi(x, t)$
  - 1.1.3. Equation of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + bw + \Phi(x, t)$
  - 1.1.4. Equation of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + \Phi(x, t)$
  - 1.1.5. Equation of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw + \Phi(x, t)$
- 1.2. Heat Equation with Axial or Central Symmetry and Related Equations
  - 1.2.1. Equation of the Form  $\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right)$
  - 1.2.2. Equation of the Form  $\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) + \Phi(r, t)$
  - 1.2.3. Equation of the Form  $\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right)$
  - 1.2.4. Equation of the Form  $\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right) + \Phi(r, t)$
  - 1.2.5. Equation of the Form  $\frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \frac{1-2\beta}{x} \frac{\partial w}{\partial x}$
  - 1.2.6. Equation of the Form  $\frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \frac{1-2\beta}{x} \frac{\partial w}{\partial x} + \Phi(x, t)$
- 1.3. Equations Containing Power Functions and Arbitrary Parameters
  - 1.3.1. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t)w$
  - 1.3.2. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x}$
  - 1.3.3. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w + h(x, t)$
  - 1.3.4. Equations of the Form  $\frac{\partial w}{\partial t} = (ax + b) \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.3.5. Equations of the Form  $\frac{\partial w}{\partial t} = (ax^2 + bx + c) \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.3.6. Equations of the Form  $\frac{\partial w}{\partial t} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x, t) \frac{\partial w}{\partial x} + h(x, t)w$
  - 1.3.7. Equations of the Form  $\frac{\partial w}{\partial t} = f(x, t) \frac{\partial^2 w}{\partial x^2} + g(x, t) \frac{\partial w}{\partial x} + h(x, t)w$
  - 1.3.8. Liquid-Film Mass Transfer Equation  $(1 - y^2) \frac{\partial w}{\partial x} = a \frac{\partial^2 w}{\partial y^2}$
  - 1.3.9. Equations of the Form  $f(x, y) \frac{\partial w}{\partial x} + g(x, y) \frac{\partial w}{\partial y} = \frac{\partial^2 w}{\partial y^2} + h(x, y)$
- 1.4. Equations Containing Exponential Functions and Arbitrary Parameters
  - 1.4.1. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t)w$
  - 1.4.2. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x}$
  - 1.4.3. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.4.4. Equations of the Form  $\frac{\partial w}{\partial t} = ax^n \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.4.5. Equations of the Form  $\frac{\partial w}{\partial t} = ae^{\beta x} \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.4.6. Other Equations
- 1.5. Equations Containing Hyperbolic Functions and Arbitrary Parameters
  - 1.5.1. Equations Containing a Hyperbolic Cosine
  - 1.5.2. Equations Containing a Hyperbolic Sine
  - 1.5.3. Equations Containing a Hyperbolic Tangent
  - 1.5.4. Equations Containing a Hyperbolic Cotangent

- 1.6. Equations Containing Logarithmic Functions and Arbitrary Parameters
  - 1.6.1. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.6.2. Equations of the Form  $\frac{\partial w}{\partial t} = ax^k \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
- 1.7. Equations Containing Trigonometric Functions and Arbitrary Parameters
  - 1.7.1. Equations Containing a Cosine
  - 1.7.2. Equations Containing a Sine
  - 1.7.3. Equations Containing a Tangent
  - 1.7.4. Equations Containing a Cotangent
- 1.8. Equations Containing Arbitrary Functions
  - 1.8.1. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t)w$
  - 1.8.2. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x}$
  - 1.8.3. Equations of the Form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.8.4. Equations of the Form  $\frac{\partial w}{\partial t} = ax^n \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.8.5. Equations of the Form  $\frac{\partial w}{\partial t} = ae^{\beta x} \frac{\partial^2 w}{\partial x^2} + f(x, t) \frac{\partial w}{\partial x} + g(x, t)w$
  - 1.8.6. Equations of the Form  $\frac{\partial w}{\partial t} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x, t) \frac{\partial w}{\partial x} + h(x, t)w$
  - 1.8.7. Equations of the Form  $\frac{\partial w}{\partial t} = f(t) \frac{\partial^2 w}{\partial x^2} + g(x, t) \frac{\partial w}{\partial x} + h(x, t)w$
  - 1.8.8. Equations of the Form  $\frac{\partial w}{\partial t} = f(x, t) \frac{\partial^2 w}{\partial x^2} + g(x, t) \frac{\partial w}{\partial x} + h(x, t)w$
  - 1.8.9. Equations of the Form  $s(x) \frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ p(x) \frac{\partial w}{\partial x} \right] - q(x)w + \Phi(x, t)$
- 1.9. Equations of Special Form
  - 1.9.1. Equations of the Diffusion (Thermal) Boundary Layer
  - 1.9.2. One-Dimensional Schrödinger Equation  $i\hbar \frac{\partial w}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 w}{\partial x^2} + U(x)w$

## 2. Parabolic Equations with Two Space Variables

- 2.1. Heat Equation  $\frac{\partial w}{\partial t} = a\Delta_2 w$ 
  - 2.1.1. Boundary Value Problems in Cartesian Coordinates
  - 2.1.2. Problems in Polar Coordinates
  - 2.1.3. Axisymmetric Problems
- 2.2. Heat Equation with a Source  $\frac{\partial w}{\partial t} = a\Delta_2 w + \Phi(x, y, t)$ 
  - 2.2.1. Problems in Cartesian Coordinates
  - 2.2.2. Problems in Polar Coordinates
  - 2.2.3. Axisymmetric Problems
- 2.3. Other Equations
  - 2.3.1. Equations Containing Arbitrary Parameters
  - 2.3.2. Equations Containing Arbitrary Functions

## 3. Parabolic Equations with Three or More Space Variables

- 3.1. Heat Equation  $\frac{\partial w}{\partial t} = a\Delta_3 w$ 
  - 3.1.1. Problems in Cartesian Coordinates
  - 3.1.2. Problems in Cylindrical Coordinates
  - 3.1.3. Problems in Spherical Coordinates
- 3.2. Heat Equation with Source  $\frac{\partial w}{\partial t} = a\Delta_3 w + \Phi(x, y, z, t)$ 
  - 3.2.1. Problems in Cartesian Coordinates
  - 3.2.2. Problems in Cylindrical Coordinates
  - 3.2.3. Problems in Spherical Coordinates
- 3.3. Other Equations with Three Space Variables
  - 3.3.1. Equations Containing Arbitrary Parameters
  - 3.3.2. Equations Containing Arbitrary Functions
  - 3.3.3. Equations of the Form  $\rho(x, y, z) \frac{\partial w}{\partial t} = \text{div}[a(x, y, z)\nabla w] - q(x, y, z)w + \Phi(x, y, z, t)$

- 3.4. Equations with  $n$  Space Variables
  - 3.4.1. Equations of the Form  $\frac{\partial w}{\partial t} = a\Delta_n w + \Phi(x_1, \dots, x_n, t)$
  - 3.4.2. Other Equations Containing Arbitrary Parameters
  - 3.4.3. Equations Containing Arbitrary Functions

#### 4. Hyperbolic Equations with One Space Variable

- 4.1. Constant Coefficient Equations
  - 4.1.1. Wave Equation  $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2}$
  - 4.1.2. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} + \Phi(x, t)$
  - 4.1.3. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} - bw + \Phi(x, t)$
  - 4.1.4. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} - b \frac{\partial w}{\partial x} + \Phi(x, t)$
  - 4.1.5. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw + \Phi(x, t)$
- 4.2. Wave Equation with Axial or Central Symmetry
  - 4.2.1. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right)$
  - 4.2.2. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) + \Phi(r, t)$
  - 4.2.3. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right)$
  - 4.2.4. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right) + \Phi(r, t)$
  - 4.2.5. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) - bw + \Phi(r, t)$
  - 4.2.6. Equation of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right) - bw + \Phi(r, t)$
- 4.3. Equations Containing Power Functions and Arbitrary Parameters
  - 4.3.1. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} = (ax + b) \frac{\partial^2 w}{\partial x^2} + c \frac{\partial w}{\partial x} + kw + \Phi(x, t)$
  - 4.3.2. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} = (ax^2 + b) \frac{\partial^2 w}{\partial x^2} + cx \frac{\partial w}{\partial x} + kw + \Phi(x, t)$
  - 4.3.3. Other Equations
- 4.4. Equations Containing the First Time Derivative
  - 4.4.1. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw + \Phi(x, t)$
  - 4.4.2. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x) \frac{\partial w}{\partial x} + h(x)w + \Phi(x, t)$
  - 4.4.3. Other Equations
- 4.5. Equations Containing Arbitrary Functions
  - 4.5.1. Equations of the Form  $s(x) \frac{\partial^2 w}{\partial t^2} = \frac{\partial}{\partial x} \left[ p(x) \frac{\partial w}{\partial x} \right] - q(x)w + \Phi(x, t)$
  - 4.5.2. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a(t) \frac{\partial w}{\partial t} = b(t) \left\{ \frac{\partial}{\partial x} \left[ p(x) \frac{\partial w}{\partial x} \right] - q(x)w \right\} + \Phi(x, t)$
  - 4.5.3. Other Equations

#### 5. Hyperbolic Equations with Two Space Variables

- 5.1. Wave Equation  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_2 w$ 
  - 5.1.1. Problems in Cartesian Coordinates
  - 5.1.2. Problems in Polar Coordinates
  - 5.1.3. Axisymmetric Problems
- 5.2. Nonhomogeneous Wave Equation  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_2 w + \Phi(x, y, t)$ 
  - 5.2.1. Problems in Cartesian Coordinates
  - 5.2.2. Problems in Polar Coordinates
  - 5.2.3. Axisymmetric Problems
- 5.3. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_2 w - bw + \Phi(x, y, t)$ 
  - 5.3.1. Problems in Cartesian Coordinates
  - 5.3.2. Problems in Polar Coordinates
  - 5.3.3. Axisymmetric Problems

5.4. Telegraph Equation  $\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \Delta_2 w - bw + \Phi(x, y, t)$

5.4.1. Problems in Cartesian Coordinates

5.4.2. Problems in Polar Coordinates

5.4.3. Axisymmetric Problems

5.5. Other Equations with Two Space Variables

## 6. Hyperbolic Equations with Three or More Space Variables

6.1. Wave Equation  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_3 w$

6.1.1. Problems in Cartesian Coordinates

6.1.2. Problems in Cylindrical Coordinates

6.1.3. Problems in Spherical Coordinates

6.2. Nonhomogeneous Wave Equation  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_3 w + \Phi(x, y, z, t)$

6.2.1. Problems in Cartesian Coordinates

6.2.2. Problems in Cylindrical Coordinates

6.2.3. Problems in Spherical Coordinates

6.3. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_3 w - bw + \Phi(x, y, z, t)$

6.3.1. Problems in Cartesian Coordinates

6.3.2. Problems in Cylindrical Coordinates

6.3.3. Problems in Spherical Coordinates

6.4. Telegraph Equation  $\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \Delta_3 w - bw + \Phi(x, y, z, t)$

6.4.1. Problems in Cartesian Coordinates

6.4.2. Problems in Cylindrical Coordinates

6.4.3. Problems in Spherical Coordinates

6.5. Other Equations with Three Space Variables

6.5.1. Equations Containing Arbitrary Parameters

6.5.2. Equation of the Form  $\rho(x, y, z) \frac{\partial^2 w}{\partial t^2} = \operatorname{div} [a(x, y, z) \nabla w] - q(x, y, z)w + \Phi(x, y, z, t)$

6.6. Equations with  $n$  Space Variables

6.6.1. Wave Equation  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_n w$

6.6.2. Nonhomogeneous Wave Equation  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_n w + \Phi(x_1, \dots, x_n, t)$

6.6.3. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_n w - bw + \Phi(x_1, \dots, x_n, t)$

6.6.4. Equations Containing the First Time Derivative

## 7. Elliptic Equations with Two Space Variables

7.1. Laplace Equation  $\Delta_2 w = 0$

7.1.1. Problems in Cartesian Coordinate System

7.1.2. Problems in Polar Coordinate System

7.1.3. Other Coordinate Systems. Conformal Mappings Method

7.2. Poisson Equation  $\Delta_2 w = -\Phi(x)$

7.2.1. Preliminary Remarks. Solution Structure

7.2.2. Problems in Cartesian Coordinate System

7.2.3. Problems in Polar Coordinate System

7.2.4. Arbitrary Shape Domain. Conformal Mappings Method

7.3. Helmholtz Equation  $\Delta_2 w + \lambda w = -\Phi(x)$

7.3.1. General Remarks, Results, and Formulas

7.3.2. Problems in Cartesian Coordinate System

7.3.3. Problems in Polar Coordinate System

7.3.4. Other Orthogonal Coordinate Systems. Elliptic Domain

- 7.4. Other Equations
  - 7.4.1. Stationary Schrödinger Equation  $\Delta_2 w = f(x, y)w$
  - 7.4.2. Convective Heat and Mass Transfer Equations
  - 7.4.3. Equations of Heat and Mass Transfer in Anisotropic Media
  - 7.4.4. Other Equations Arising in Applications
  - 7.4.5. Equations of the Form  $a(x)\frac{\partial^2 w}{\partial x^2} + b(x)\frac{\partial^2 w}{\partial y^2} + c(x)\frac{\partial w}{\partial x} + d(x)y = -\Phi(x, y)$

## 8. Elliptic Equations with Three or More Space Variables

- 8.1. Laplace Equation  $\Delta_3 w = 0$ 
  - 8.1.1. Problems in Cartesian Coordinates
  - 8.1.2. Problems in Cylindrical Coordinates
  - 8.1.3. Problems in Spherical Coordinates
  - 8.1.4. Other Orthogonal Curvilinear Systems of Coordinates
- 8.2. Poisson Equation  $\Delta_3 w + \Phi(x) = 0$ 
  - 8.2.1. Preliminary Remarks. Solution Structure
  - 8.2.2. Problems in Cartesian Coordinates
  - 8.2.3. Problems in Cylindrical Coordinates
  - 8.2.4. Problems in Spherical Coordinates
- 8.3. Helmholtz Equation  $\Delta_3 w + \lambda w = -\Phi(x)$ 
  - 8.3.1. General Remarks, Results, and Formulas
  - 8.3.2. Problems in Cartesian Coordinates
  - 8.3.3. Problems in Cylindrical Coordinates
  - 8.3.4. Problems in Spherical Coordinates
  - 8.3.5. Other Orthogonal Curvilinear Coordinates
- 8.4. Other Equations with Three Space Variables
  - 8.4.1. Equations Containing Arbitrary Functions
  - 8.4.2. Equations of the Form  $\operatorname{div}[a(x, y, z)\nabla w] - q(x, y, z)w = -\Phi(x, y, z)$
- 8.5. Equations with  $n$  Space Variables
  - 8.5.1. Laplace Equation  $\Delta_n w = 0$
  - 8.5.2. Other Equations

## 9. Higher-Order Partial Differential Equations

- 9.1. Third-Order Partial Differential Equations
- 9.2. Fourth-Order One-Dimensional Nonstationary Equations
  - 9.2.1. Equations of the Form  $\frac{\partial w}{\partial t} + a^2 \frac{\partial^4 w}{\partial x^4} = \Phi(x, t)$
  - 9.2.2. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \frac{\partial^4 w}{\partial x^4} = 0$
  - 9.2.3. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \frac{\partial^4 w}{\partial x^4} = \Phi(x, t)$
  - 9.2.4. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \frac{\partial^4 w}{\partial x^4} + kw = \Phi(x, t)$
  - 9.2.5. Other Equations
- 9.3. Two-Dimensional Nonstationary Fourth-Order Equations
  - 9.3.1. Equations of the Form  $\frac{\partial w}{\partial t} + a^2 \left( \frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} \right) = \Phi(x, y, t)$
  - 9.3.2. Two-Dimensional Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \Delta \Delta w = 0$
  - 9.3.3. Three- and  $n$ -Dimensional Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \Delta \Delta w = 0$
  - 9.3.4. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \Delta \Delta w + kw = \Phi(x, y, t)$
  - 9.3.5. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \left( \frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} \right) + kw = \Phi(x, y, t)$
- 9.4. Fourth-Order Stationary Equations
  - 9.4.1. Biharmonic Equation  $\Delta \Delta w = 0$
  - 9.4.2. Equations of the Form  $\Delta \Delta w = \Phi(x, y)$



- 9.4.3. Equations of the Form  $\Delta\Delta w - \lambda w = \Phi(x, y)$
- 9.4.4. Equations of the Form  $\frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} = \Phi(x, y)$
- 9.4.5. Equations of the Form  $\frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} + kw = \Phi(x, y)$
- 9.4.6. Stokes Equation (Axisymmetric Flows of Viscous Fluids)
- 9.5. Higher-Order Linear Equations with Constant Coefficients
  - 9.5.1. Fundamental Solutions. Cauchy Problem
  - 9.5.2. Elliptic Equations
  - 9.5.3. Hyperbolic Equations
  - 9.5.4. Regular Equations. Number of Initial Conditions in the Cauchy Problem
  - 9.5.5. Some Special-Type Equations
- 9.6. Higher-Order Linear Equations with Variable Coefficients
  - 9.6.1. Equations Containing the First Time Derivative
  - 9.6.2. Equations Containing the Second Time Derivative
  - 9.6.3. Nonstationary Problems with Many Space Variables
  - 9.6.4. Some Special-Type Equations

## **Supplement A. Special Functions and Their Properties**

- A.1. Some Symbols and Coefficients
  - A.1.1. Factorials
  - A.1.2. Binomial Coefficients
  - A.1.3. Pochhammer Symbol
  - A.1.4. Bernoulli Numbers
- A.2. Error Functions and Exponential Integral
  - A.2.1. Error Function and Complementary Error Function
  - A.2.2. Exponential Integral
  - A.2.3. Logarithmic Integral
- A.3. Sine Integral and Cosine Integral. Fresnel Integrals
  - A.3.1. Sine Integral
  - A.3.2. Cosine Integral
  - A.3.3. Fresnel Integrals
- A.4. Gamma and Beta Functions
  - A.4.1. Gamma Function
  - A.4.2. Beta Function
- A.5. Incomplete Gamma and Beta Functions
  - A.5.1. Incomplete Gamma Function
  - A.5.2. Incomplete Beta Function
- A.6. Bessel Functions
  - A.6.1. Definitions and Basic Formulas
  - A.6.2. Integral Representations and Asymptotic Expansions
  - A.6.3. Zeros and Orthogonality Properties of Bessel Functions
  - A.6.4. Hankel Functions (Bessel Functions of the Third Kind)
- A.7. Modified Bessel Functions
  - A.7.1. Definitions. Basic Formulas
  - A.7.2. Integral Representations and Asymptotic Expansions
- A.8. Airy Functions
  - A.8.1. Definition and Basic Formulas
  - A.8.2. Power Series and Asymptotic Expansions

- A.9. **Degenerate Hypergeometric Functions**
  - A.9.1. **Definitions and Basic Formulas**
  - A.9.2. **Integral Representations and Asymptotic Expansions**
- A.10. **Hypergeometric Functions**
  - A.10.1. **Definition and Some Formulas**
  - A.10.2. **Basic Properties and Integral Representations**
- A.11. **Whittaker Functions**
- A.12. **Legendre Polynomials and Legendre Functions**
  - A.12.1. **Definitions. Basic Formulas**
  - A.12.2. **Zeros of Legendre Polynomials and the Generating Function**
  - A.12.3. **Associated Legendre Functions**
- A.13. **Parabolic Cylinder Functions**
  - A.13.1. **Definitions. Basic Formulas**
  - A.13.2. **Integral Representations and Asymptotic Expansions**
- A.14. **Mathieu Functions**
  - A.14.1. **Definitions and Basic Formulas**
- A.15. **Modified Mathieu Functions**
- A.16. **Orthogonal Polynomials**
  - A.16.1. **Laguerre Polynomials and Generalized Laguerre Polynomials**
  - A.16.2. **Chebyshev Polynomials and Functions**
  - A.16.3. **Hermite Polynomial**
  - A.16.4. **Jacobi Polynomials**

## **Supplement B. Methods of Generalized and Functional Separation of Variables in Nonlinear Equations of Mathematical Physics**

- B.1. **Introduction**
  - B.1.1. **Preliminary Remarks**
  - B.1.2. **Simple Cases of Variable Separation in Nonlinear Equations**
  - B.1.3. **Examples of Nontrivial Variable Separation in Nonlinear Equations**
- B.2. **Methods of Generalized Separation of Variables**
  - B.2.1. **Structure of Generalized Separable Solutions**
  - B.2.2. **Solution of Functional Differential Equations by Differentiation**
  - B.2.3. **Solution of Functional Differential Equations by Splitting**
  - B.2.4. **Simplified Scheme for Constructing Exact Solutions of Equations with Quadratic Nonlinearities**
- B.3. **Methods of Functional Separation of Variables**
  - B.3.1. **Structure of Functional Separable Solutions**
  - B.3.2. **Special Functional Separable Solutions**
  - B.3.3. **Differentiation Method**
  - B.3.4. **Splitting Method. Reduction to a Functional Equation with Two Variables**
  - B.3.5. **Some Functional Equations and Their Solutions. Exact Solutions of Heat and Wave Equations**
- B.4. **First-Order Nonlinear Equations**
  - B.4.1. **Preliminary Remarks**
  - B.4.2. **Individual Equations**
- B.5. **Second-Order Nonlinear Equations**
  - B.5.1. **Parabolic Equations**
  - B.5.2. **Hyperbolic Equations**
  - B.5.3. **Elliptic Equations**
  - B.5.4. **Equations Containing Mixed Derivatives**

- B.5.5. [General Form Equations](#)
- B.6. [Third-Order Nonlinear Equations](#)
  - B.6.1. [Stationary Hydrodynamic Boundary Layer Equations](#)
  - B.6.2. [Nonstationary Hydrodynamic Boundary Layer Equations](#)
- B.7. [Fourth-Order Nonlinear Equations](#)
  - B.7.1. [Stationary Hydrodynamic Equations \(Navier–Stokes Equations\)](#)
  - B.7.2. [Nonstationary Hydrodynamic Equations](#)
- B.8. [Higher-Order Nonlinear Equations](#)
  - B.8.1. [Equations of the Form  \$\frac{\partial w}{\partial t} = F\left\(x, t, w, \frac{\partial w}{\partial x}, \dots, \frac{\partial^n w}{\partial x^n}\right\)\$](#)
  - B.8.2. [Equations of the Form  \$\frac{\partial^2 w}{\partial t^2} = F\left\(x, t, w, \frac{\partial w}{\partial x}, \dots, \frac{\partial^n w}{\partial x^n}\right\)\$](#)
  - B.8.3. [Other Equations](#)

## [References](#)

## Chapter 2

# Parabolic Equations with Two Space Variables

---

### 2.1. Heat Equation $\frac{\partial w}{\partial t} = a \Delta_2 w$

#### 2.1.1. Boundary Value Problems in Cartesian Coordinates

In rectangular Cartesian coordinates, the two-dimensional sourceless heat equation has the form

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right).$$

It governs two-dimensional unsteady heat transfer processes in quiescent media or solid bodies with constant thermal diffusivity  $a$ . A similar equation is used to study analogous two-dimensional unsteady mass transfer phenomena with constant diffusivity; in this case the equation is called a diffusion equation.

##### 2.1.1-1. Particular solutions:

$$\begin{aligned} w(x, y) &= Axy + C_1x + C_2y + C_3, \\ w(x, y, t) &= Ax^2 + By^2 + 2a(A + B)t, \\ w(x, y, t) &= A(x^2 + 2at)(y^2 + 2at) + B, \\ w(x, y, t) &= A \exp[k_1x + k_2y + (k_1^2 + k_2^2)at] + B, \\ w(x, y, t) &= A \cos(k_1x + C_1) \cos(k_2y + C_2) \exp[-(k_1^2 + k_2^2)at], \\ w(x, y, t) &= A \cos(k_1x + C_1) \sinh(k_2y + C_2) \exp[-(k_1^2 - k_2^2)at], \\ w(x, y, t) &= A \cos(k_1x + C_1) \cosh(k_2y + C_2) \exp[-(k_1^2 - k_2^2)at], \\ w(x, y, t) &= A \exp(-\mu x - \lambda y) \cos(\mu x - 2a\mu^2t + C_1) \cos(\lambda y - 2a\lambda^2t + C_2), \\ w(x, y, t) &= \frac{A}{t - t_0} \exp \left[ -\frac{(x - x_0)^2 + (y - y_0)^2}{4a(t - t_0)} \right], \\ w(x, y, t) &= A \operatorname{erf} \left( \frac{x - x_0}{2\sqrt{at}} \right) \operatorname{erf} \left( \frac{y - y_0}{2\sqrt{at}} \right) + B, \end{aligned}$$

where  $A, B, C_1, C_2, C_3, k_1, k_2, x_0, y_0$ , and  $t_0$  are arbitrary constants.

Fundamental solution:

$$\mathcal{E}(x, y, t) = \frac{1}{4\pi at} \exp \left( -\frac{x^2 + y^2}{4at} \right).$$

##### 2.1.1-2. Formulas to construct particular solutions. Remarks on the Green's functions.

1°. Apart from usual separable solutions  $w(x, y, t) = f_1(x)f_2(y)f_3(t)$ , the equation in question has more sophisticated solutions in the product form

$$w(x, y, t) = u(x, t)v(y, t),$$

where  $u = u(x, t)$  and  $v = v(y, t)$  are solutions of the one-dimensional heat equations

$$\frac{\partial u}{\partial t} = a \frac{\partial^2 u}{\partial x^2}, \quad \frac{\partial v}{\partial t} = a \frac{\partial^2 v}{\partial y^2},$$

considered in Subsection 1.1.1.

2°. Suppose  $w = w(x, y, t)$  is a solution of the heat equation. Then the functions

$$\begin{aligned} w_1 &= Aw(\pm\lambda x + C_1, \pm\lambda y + C_2, \lambda^2 t + C_3), \\ w_2 &= Aw(x \cos \beta - y \sin \beta + C_1, x \sin \beta + y \cos \beta + C_2, t + C_3), \\ w_3 &= A \exp[\lambda_1 x + \lambda_2 y + a(\lambda_1^2 + \lambda_2^2)t] w(x + 2a\lambda_1 t + C_1, y + 2a\lambda_2 t + C_2, t + C_3), \\ w_4 &= \frac{A}{\delta + \beta t} \exp\left[-\frac{\beta(x^2 + y^2)}{4a(\delta + \beta t)}\right] w\left(\frac{x}{\delta + \beta t}, \frac{y}{\delta + \beta t}, \frac{\gamma + \lambda t}{\delta + \beta t}\right), \quad \lambda\delta - \beta\gamma = 1, \end{aligned}$$

where  $A, C_1, C_2, C_3, \beta, \delta, \lambda, \lambda_1$  and  $\lambda_2$  are arbitrary constants, are also solutions of this equation. The signs at  $\lambda$ 's in the formula for  $w_1$  are taken arbitrarily, independently of each other.

⊙ *Reference:* W. Miller, Jr. (1977).

3°. For all two-dimensional boundary value problems discussed in Subsection 2.1.1, the Green's function can be represented in the product form

$$G(x, y, \xi, \eta, t) = G_1(x, \xi, t)G_2(y, \eta, t),$$

where  $G_1(x, \xi, t)$  and  $G_2(y, \eta, t)$  are the Green's functions of the corresponding one-dimensional boundary value problems (these functions are specified in Subsections 1.1.1 and 1.1.2).

**Example 1.** The Green's function of the first boundary value problem for a semiinfinite strip ( $0 \leq x \leq l, 0 \leq y < \infty$ ), considered in Subsection 2.1.1-12, is the product of two one-dimensional Green's functions. The first Green's function is that of the first boundary value problem on a closed interval ( $0 \leq x \leq l$ ) presented in Subsection 1.1.2-5. The second Green's function is that of the first boundary value problem on a semiinfinite interval ( $0 \leq y < \infty$ ) presented in Subsection 1.1.2-2, where  $x$  and  $\xi$  must be renamed  $y$  and  $\eta$ , respectively.

### 2.1.1-3. Transformations that allow separation of variables.

**Table 18** lists possible transformations that allow reduction of the two-dimensional heat equation to a separable equation. All transformations of the independent variables have the form  $(x, y, t) \mapsto (\xi, \eta, t)$ . The transformations that can be obtained by interchange of independent variables,  $x \rightleftharpoons y$ , are omitted.

The anharmonic oscillator functions are solutions of the second-order ordinary differential equation  $F''_{zz} + (az^4 + bz^2 + c)F = 0$ . The Ince polynomials are the  $2\pi$ -periodic solutions of the Whittaker–Hill equation  $F''_{zz} + k \sin 2z F'_z + (a - bk \cos 2z)F = 0$ ; see Arscott (1964, 1967).

⊙ *Reference:* W. Miller, Jr. (1977).

### 2.1.1-4. Domain: $-\infty < x < \infty, -\infty < y < \infty$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(x, y) \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, y, t) = \frac{1}{4\pi at} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta) \exp\left[-\frac{(x - \xi)^2 + (y - \eta)^2}{4at}\right] d\xi d\eta.$$

**Example 2.** The initial temperature is piecewise-constant and equal to  $w_1$  in the domain  $|x| < x_0, |y| < y_0$  and  $w_2$  in the domain  $|x| > x_0, |y| > y_0$ , specifically,

$$f(x, y) = \begin{cases} w_1 & \text{for } |x| < x_0, |y| < y_0, \\ w_2 & \text{for } |x| > x_0, |y| > y_0. \end{cases}$$

TABLE 18

Transformations  $(x, y, t) \mapsto (\xi, \eta, t)$  that allow solutions with  $\mathcal{R}$ -separated variables,  $w = \exp[\mathcal{R}(\xi, \eta, t)] f(\xi) g(\eta) h(t)$ , for the two-dimensional heat equation  $\partial_t w = \partial_{xx} w + \partial_{yy} w$ . Everywhere, the function  $h(t)$  is exponential

| No | Transformations                                                               | Factor $\exp \mathcal{R}$                                                                       | Function $f(\xi)$           | Function $g(\eta)$          |
|----|-------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------|-----------------------------|-----------------------------|
| 1  | $x = \xi,$<br>$y = \eta$                                                      | $\mathcal{R} = 0$                                                                               | Exponential function        | Exponential function        |
| 2  | $x = \xi,$<br>$y = \eta\sqrt{ t }$                                            | $\mathcal{R} = 0$                                                                               | Exponential function        | Hermite function            |
| 3  | $x = \xi\sqrt{ t },$<br>$y = \eta\sqrt{ t }$                                  | $\mathcal{R} = 0$                                                                               | Hermite function            | Hermite function            |
| 4  | $x = \frac{1}{2}(\xi^2 - \eta^2),$<br>$y = \xi\eta$                           | $\mathcal{R} = 0$                                                                               | Parabolic cylinder function | Parabolic cylinder function |
| 5  | $x = \xi \cos \eta,$<br>$y = \xi \sin \eta$                                   | $\mathcal{R} = 0$                                                                               | Bessel function             | Exponential function        |
| 6  | $x = \cosh \xi \cos \eta,$<br>$y = \sinh \xi \sin \eta$                       | $\mathcal{R} = 0$                                                                               | Modified Mathieu function   | Mathieu function            |
| 7  | $x = \sqrt{ t } \xi \cos \eta,$<br>$y = \sqrt{ t } \xi \sin \eta$             | $\mathcal{R} = 0$                                                                               | Laguerre function           | Exponential function        |
| 8  | $x = \sqrt{ t } \cosh \xi \cos \eta,$<br>$y = \sqrt{ t } \sinh \xi \sin \eta$ | $\mathcal{R} = 0$                                                                               | Ince polynomial             | Ince polynomial             |
| 9  | $x = \xi,$<br>$y = \eta + at^2$                                               | $\mathcal{R} = -a\eta t$                                                                        | Exponential function        | Airy function               |
| 10 | $x = \xi,$<br>$y = \eta t + b/t$                                              | $\mathcal{R} = -\frac{1}{4}\eta^2 t + \frac{1}{2}b\eta/t$                                       | Exponential function        | Airy function               |
| 11 | $x = \xi,$<br>$y = \eta\sqrt{1+t^2}$                                          | $\mathcal{R} = -\frac{1}{4}\eta^2 t$                                                            | Exponential function        | Parabolic cylinder function |
| 12 | $x = \xi,$<br>$y = \eta\sqrt{ 1-t^2 }$                                        | $\mathcal{R} = -\frac{1}{4}\varepsilon\eta^2 t,$<br>$\varepsilon = \text{sign}(1-t^2)$          | Exponential function        | Hermite function            |
| 13 | $x = \xi t,$<br>$y = \eta t$                                                  | $\mathcal{R} = -\frac{1}{4}(\xi^2 + \eta^2)t$                                                   | Exponential function        | Exponential function        |
| 14 | $x = \xi + at^2,$<br>$y = \eta + bt^2$                                        | $\mathcal{R} = -(a\xi + b\eta)t$                                                                | Airy function               | Airy function               |
| 15 | $x = \xi t + a/t,$<br>$y = \eta t + b/t$                                      | $\mathcal{R} = -\frac{1}{4}(\xi^2 + \eta^2)t$<br>$+ \frac{1}{2}(a\xi + b\eta)/t$                | Airy function               | Airy function               |
| 16 | $x = \frac{1}{2}(\xi^2 - \eta^2)t,$<br>$y = \xi\eta t$                        | $\mathcal{R} = -\frac{1}{16}(\xi^2 + \eta^2)^2 t$                                               | Parabolic cylinder function | Parabolic cylinder function |
| 17 | $x = \xi\sqrt{1+t^2},$<br>$y = \eta\sqrt{1+t^2}$                              | $\mathcal{R} = -\frac{1}{4}(\xi^2 + \eta^2)t$                                                   | Parabolic cylinder function | Parabolic cylinder function |
| 18 | $x = \xi\sqrt{ 1-t^2 },$<br>$y = \eta\sqrt{ 1-t^2 }$                          | $\mathcal{R} = -\frac{1}{4}\varepsilon(\xi^2 + \eta^2)t,$<br>$\varepsilon = \text{sign}(1-t^2)$ | Hermite function            | Hermite function            |

TABLE 18  
(continued)

| No | Transformations                                                                       | Factor $\exp \mathcal{R}$                                                                                  | Function $f(\xi)$              | Function $g(\eta)$             |
|----|---------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|--------------------------------|--------------------------------|
| 19 | $x = \frac{1}{2}(\xi^2 - \eta^2) + at^2,$<br>$y = \xi\eta$                            | $\mathcal{R} = -\frac{1}{2}a(\xi^2 - \eta^2)t$                                                             | Anharmonic oscillator function | Anharmonic oscillator function |
| 20 | $x = \frac{1}{2}(\xi^2 - \eta^2)t + a/t,$<br>$y = \xi\eta t$                          | $\mathcal{R} = -\frac{1}{16}(\xi^2 + \eta^2)^2t$<br>$+\frac{1}{4}a(\xi^2 - \eta^2)/t$                      | Anharmonic oscillator function | Anharmonic oscillator function |
| 21 | $x = \xi t \cos \eta,$<br>$y = \xi t \sin \eta$                                       | $\mathcal{R} = -\frac{1}{4}\xi^2 t$                                                                        | Bessel function                | Exponential function           |
| 22 | $x = t \cosh \xi \cos \eta,$<br>$y = t \sinh \xi \sin \eta$                           | $\mathcal{R} = -\frac{1}{4}(\sinh^2 \xi + \cos^2 \eta)t$                                                   | Modified Mathieu function      | Mathieu function               |
| 23 | $x = \sqrt{1+t^2} \xi \cos \eta,$<br>$y = \sqrt{1+t^2} \xi \sin \eta$                 | $\mathcal{R} = -\frac{1}{4}\xi^2 t$                                                                        | Whittaker function             | Exponential function           |
| 24 | $x = \sqrt{ 1-t^2 } \xi \cos \eta,$<br>$y = \sqrt{ 1-t^2 } \xi \sin \eta$             | $\mathcal{R} = -\frac{1}{4}\varepsilon \xi^2 t,$<br>$\varepsilon = \text{sign}(1-t^2)$                     | Laguerre function              | Exponential function           |
| 25 | $x = \sqrt{1+t^2} \cosh \xi \cos \eta,$<br>$y = \sqrt{1+t^2} \sinh \xi \sin \eta$     | $\mathcal{R} = -\frac{1}{4}(\sinh^2 \xi + \cos^2 \eta)t$                                                   | Ince polynomial                | Ince polynomial                |
| 26 | $x = \sqrt{ 1-t^2 } \cosh \xi \cos \eta,$<br>$y = \sqrt{ 1-t^2 } \sinh \xi \sin \eta$ | $\mathcal{R} = -\frac{1}{4}\varepsilon(\sinh^2 \xi + \cos^2 \eta)t,$<br>$\varepsilon = \text{sign}(1-t^2)$ | Ince polynomial                | Ince polynomial                |

Solution:

$$w = \frac{1}{4}(w_1 - w_2) \left[ \text{erf} \left( \frac{x_0 - x}{2\sqrt{at}} \right) + \text{erf} \left( \frac{x_0 + x}{2\sqrt{at}} \right) \right] \left[ \text{erf} \left( \frac{y_0 - y}{2\sqrt{at}} \right) + \text{erf} \left( \frac{y_0 + y}{2\sqrt{at}} \right) \right] + w_2.$$

If the initial temperature distribution  $f(x, y)$  is an infinitely differentiable function in both arguments, then the solution can be represented in the series form

$$w(x, y, t) = f(x, y) + \sum_{n=1}^{\infty} \frac{(at)^n}{n!} \mathbf{L}^n [f(x, y)], \quad \mathbf{L} \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}.$$

Such a representation is useful for small  $t$ .

© *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

2.1.1-5. Domain:  $0 \leq x < \infty, -\infty < y < \infty$ . First boundary value problem.

A half-plane is considered. The following conditions are prescribed:

$$w = f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$w = g(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}).$$

Solution:

$$w(x, y, t) = \int_0^\infty \int_{-\infty}^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ + a \int_0^t \int_{-\infty}^\infty g(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau,$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] - \exp \left[ -\frac{(x + \xi)^2 + (y - \eta)^2}{4at} \right] \right\}.$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**2.1.1-6. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ . Second boundary value problem.**

A half-plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(x, y, t) = \int_0^\infty \int_{-\infty}^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi - a \int_0^t \int_{-\infty}^\infty g(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau,$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] + \exp \left[ -\frac{(x + \xi)^2 + (y - \eta)^2}{4at} \right] \right\}.$$

**2.1.1-7. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ . Third boundary value problem.**

A half-plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - kw &= g(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.1.1-6 where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{4\pi at} \exp \left[ -\frac{(y - \eta)^2}{4at} \right] \left\{ \exp \left[ -\frac{(x - \xi)^2}{4at} \right] + \exp \left[ -\frac{(x + \xi)^2}{4at} \right] \right. \\ &\quad \left. - 2k \int_0^\infty \exp \left[ -\frac{(x + \xi + s)^2}{4at} - ks \right] ds \right\}. \end{aligned}$$

**2.1.1-8. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.**

A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad + a \int_0^t \int_0^\infty g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &\quad + a \int_0^t \int_0^\infty g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau, \end{aligned}$$



where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] - \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right\} \left\{ \exp \left[ -\frac{(y-\eta)^2}{4at} \right] - \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right\}.$$

**Example 3.** The initial temperature is uniform,  $f(x, y) = w_0$ . The boundary is maintained at zero temperature,  $g_1(y, t) = g_2(x, t) = 0$ .

Solution:

$$w = w_0 \operatorname{erf} \left( \frac{x}{2\sqrt{at}} \right) \operatorname{erf} \left( \frac{y}{2\sqrt{at}} \right).$$

⊙ *References:* A. G. Butkovskiy (1979), H. S. Carslaw and J. C. Jaeger (1984).

#### 2.1.1-9. Domain: $0 \leq x < \infty$ , $0 \leq y < \infty$ . Second boundary value problem.

A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_y w &= g_2(x, t) & \text{at } y &= 0 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad - a \int_0^t \int_0^\infty g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau - a \int_0^t \int_0^\infty g_2(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] + \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right\} \left\{ \exp \left[ -\frac{(y-\eta)^2}{4at} \right] + \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right\}.$$

#### 2.1.1-10. Domain: $0 \leq x < \infty$ , $0 \leq y < \infty$ . Third boundary value problem.

A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - k_1 w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_y w - k_2 w &= g_2(x, t) & \text{at } y &= 0 & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.1.1-9 where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] + \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right. \\ &\quad \left. - 2k_1 \sqrt{\pi at} \exp[ak_1^2 t + k_1(x+\xi)] \operatorname{erfc} \left( \frac{x+\xi}{2\sqrt{at}} + k_1 \sqrt{at} \right) \right\} \\ &\quad \times \left\{ \exp \left[ -\frac{(y-\eta)^2}{4at} \right] + \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right. \\ &\quad \left. - 2k_2 \sqrt{\pi at} \exp[ak_2^2 t + k_2(y+\eta)] \operatorname{erfc} \left( \frac{y+\eta}{2\sqrt{at}} + k_2 \sqrt{at} \right) \right\}. \end{aligned}$$

**Example 4.** The initial temperature is constant,  $f(x, y) = w_0$ . The temperature of the environment is zero,  $g_1(y, t) = g_2(x, t) = 0$ .

Solution:

$$\begin{aligned} w &= w_0 \left[ \operatorname{erf} \left( \frac{x}{2\sqrt{at}} \right) + \exp(k_1 x + ak_1^2 t) \operatorname{erfc} \left( \frac{x}{2\sqrt{at}} + k_1 \sqrt{at} \right) \right] \\ &\quad \times \left[ \operatorname{erf} \left( \frac{y}{2\sqrt{at}} \right) + \exp(k_2 y + ak_2^2 t) \operatorname{erfc} \left( \frac{y}{2\sqrt{at}} + k_2 \sqrt{at} \right) \right]. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

1°. A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_2(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad + a \int_0^t \int_0^\infty g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &\quad - a \int_0^t \int_0^\infty g_2(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] - \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right\} \left\{ \exp \left[ -\frac{(y-\eta)^2}{4at} \right] + \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right\}.$$

2°. A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - kw &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad - a \int_0^t \int_0^\infty g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\ &\quad + a \int_0^t \int_0^\infty g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{4\pi at} \left\{ \exp \left[ \frac{(y-\eta)^2}{4at} \right] - \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right\} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] + \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right. \\ &\quad \left. - 2k\sqrt{\pi at} \exp[ak^2t + k(x+\xi)] \operatorname{erfc} \left( \frac{x+\xi}{2\sqrt{at}} + k\sqrt{at} \right) \right\}. \end{aligned}$$

**Example 5.** The initial temperature is uniform,  $f(x, y) = w_0$ . Heat exchange with the environment of zero temperature occurs at one side and the other side is maintained at zero temperature:  $g_1(y, t) = g_2(x, t) = 0$ .

Solution:

$$w = w_0 \left[ \operatorname{erf} \left( \frac{x}{2\sqrt{at}} \right) + \exp(kx + ak^2t) \operatorname{erfc} \left( \frac{x}{2\sqrt{at}} + k\sqrt{at} \right) \right] \operatorname{erf} \left( \frac{y}{2\sqrt{at}} \right).$$

**2.1.1-12. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . First boundary value problem.**

A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x = l & \quad (\text{boundary condition}), \\ w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^l f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad + a \int_0^t \int_0^\infty g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &\quad - a \int_0^t \int_0^\infty g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l} d\eta d\tau \\ &\quad + a \int_0^t \int_0^l g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= G_1(x, \xi, t) G_2(y, \eta, t), \\ G_1(x, \xi, t) &= \frac{2}{l} \sum_{n=1}^\infty \sin\left(\frac{n\pi x}{l}\right) \sin\left(\frac{n\pi \xi}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right), \\ G_2(y, \eta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(y-\eta)^2}{4at}\right] - \exp\left[-\frac{(y+\eta)^2}{4at}\right] \right\}. \end{aligned}$$

**Example 6.** The initial temperature is uniform,  $f(x, y) = w_0$ . The boundary is maintained at zero temperature,  $g_1(y, t) = g_2(y, t) = g_3(x, t) = 0$ .

Solution:

$$w = \frac{4w_0}{\pi} \operatorname{erf}\left(\frac{y}{2\sqrt{at}}\right) \sum_{n=0}^\infty \frac{1}{2n+1} \sin\left[\frac{(2n+1)\pi x}{l}\right] \exp\left[-\frac{\pi^2(2n+1)^2 at}{l^2}\right].$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**2.1.1-13. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . Second boundary value problem.**

A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) & \text{at } x = l & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^l f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad - a \int_0^t \int_0^\infty g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\ &\quad + a \int_0^t \int_0^\infty g_2(\eta, \tau) G(x, y, l, \eta, t - \tau) d\eta d\tau \\ &\quad - a \int_0^t \int_0^l g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = G_1(x, \xi, t)G_2(y, \eta, t),$$

$$G_1(x, \xi, t) = \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi \xi}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right),$$

$$G_2(y, \eta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(y-\eta)^2}{4at}\right] + \exp\left[-\frac{(y+\eta)^2}{4at}\right] \right\}.$$

**2.1.1-14. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . Third boundary value problem.**

A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - k_1 w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w + k_2 w &= g_2(y, t) & \text{at } x &= l & \text{(boundary condition),} \\ \partial_y w - k_3 w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.1.1-13 where the Green's function  $G(x, y, \xi, \eta, t)$  is the product of the Green's function of Subsection 1.1.1-11 and that of Subsection 1.1.1-8; one should replace  $x$ ,  $\xi$ , and  $k$  by  $y$ ,  $\eta$  and  $k_3$ , respectively, in the last Green's function.

**2.1.1-15. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . Mixed boundary value problems.**

1°. A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(y, t) & \text{at } x &= l & \text{(boundary condition),} \\ \partial_y w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^{\infty} \int_0^l f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &+ a \int_0^t \int_0^{\infty} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &- a \int_0^t \int_0^{\infty} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l} d\eta d\tau \\ &- a \int_0^t \int_0^l g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = G_1(x, \xi, t)G_2(y, \eta, t),$$

$$G_1(x, \xi, t) = \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{l}\right) \sin\left(\frac{n\pi \xi}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right),$$

$$G_2(y, \eta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(y-\eta)^2}{4at}\right] + \exp\left[-\frac{(y+\eta)^2}{4at}\right] \right\}.$$

2°. A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) & \text{at } x = l & \quad (\text{boundary condition}), \\ w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^l f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad - a \int_0^t \int_0^\infty g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\ &\quad + a \int_0^t \int_0^\infty g_2(\eta, \tau) G(x, y, l, \eta, t - \tau) d\eta d\tau \\ &\quad + a \int_0^t \int_0^l g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= G_1(x, \xi, t) G_2(y, \eta, t), \\ G_1(x, \xi, t) &= \frac{1}{l} + \frac{2}{l} \sum_{n=1}^\infty \cos\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi \xi}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right), \\ G_2(y, \eta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(y - \eta)^2}{4at}\right] - \exp\left[-\frac{(y + \eta)^2}{4at}\right] \right\}. \end{aligned}$$

2.1.1-16. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . First boundary value problem.

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &\quad + a \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &\quad - a \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\ &\quad + a \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad - a \int_0^t \int_0^{l_1} g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=l_2} d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin \frac{n\pi x}{l_1} \sin \frac{n\pi \xi}{l_1} \sin \frac{m\pi y}{l_2} \sin \frac{m\pi \eta}{l_2} \exp \left[ -\pi^2 \left( \frac{n^2}{l_1^2} + \frac{m^2}{l_2^2} \right) at \right].$$

**Example 7.** The initial temperature is uniform,  $f(x, y) = w_0$ . The boundary is maintained at zero temperature,  $g_1(y, t) = g_2(y, t) = g_3(x, t) = g_4(x, t) = 0$ .

Solution:

$$w = \frac{16w_0}{\pi^2} \left\{ \sum_{n=0}^{\infty} \frac{1}{2n+1} \sin \left[ \frac{(2n+1)\pi x}{l_1} \right] \exp \left[ -\frac{\pi^2(2n+1)^2 at}{l_1^2} \right] \right\} \\ \times \left\{ \sum_{m=0}^{\infty} \frac{1}{2m+1} \sin \left[ \frac{(2m+1)\pi y}{l_2} \right] \exp \left[ -\frac{\pi^2(2m+1)^2 at}{l_2^2} \right] \right\}.$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

2.1.1-17. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Second boundary value problem.

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w &= g_2(y, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ \partial_y w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ \partial_y w &= g_4(x, t) & \text{at } y &= l_2 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &\quad - a \int_0^t \int_0^{l_2} g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\ &\quad + a \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\ &\quad - a \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + a \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{l_1 l_2} \left[ 1 + 2 \sum_{n=1}^{\infty} \exp \left( -\frac{\pi^2 n^2 at}{l_1^2} \right) \cos \frac{n\pi x}{l_1} \cos \frac{n\pi \xi}{l_1} \right] \\ &\quad \times \left[ 1 + 2 \sum_{m=1}^{\infty} \exp \left( -\frac{\pi^2 m^2 at}{l_2^2} \right) \cos \frac{m\pi y}{l_2} \cos \frac{m\pi \eta}{l_2} \right]. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

2.1.1-18. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Third boundary value problem.

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - k_1 w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w + k_2 w &= g_2(y, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ \partial_y w - k_3 w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ \partial_y w + k_4 w &= g_4(x, t) & \text{at } y &= l_2 & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.1.1-17 where

$$G(x, y, \xi, \eta, t) = \left\{ \sum_{n=1}^{\infty} \frac{\varphi_n(x)\varphi_n(\xi)}{\|\varphi_n\|^2} \exp(-a\mu_n^2 t) \right\} \left\{ \sum_{m=1}^{\infty} \frac{\psi_m(y)\psi_m(\eta)}{\|\psi_m\|^2} \exp(-a\lambda_m^2 t) \right\},$$

$$\varphi_n(x) = \cos(\mu_n x) + \frac{k_1}{\mu_n} \sin(\mu_n x), \quad \|\varphi_n\|^2 = \frac{k_2}{2\mu_n^2} \frac{\mu_n^2 + k_1^2}{\mu_n^2 + k_2^2} + \frac{k_1}{2\mu_n^2} + \frac{l_1}{2} \left( 1 + \frac{k_1^2}{\mu_n^2} \right),$$

$$\psi_m(y) = \cos(\lambda_m y) + \frac{k_3}{\lambda_m} \sin(\lambda_m y), \quad \|\psi_m\|^2 = \frac{k_4}{2\lambda_m^2} \frac{\lambda_m^2 + k_3^2}{\lambda_m^2 + k_4^2} + \frac{k_3}{2\lambda_m^2} + \frac{l_2}{2} \left( 1 + \frac{k_3^2}{\lambda_m^2} \right).$$

Here, the  $\mu_n$  and  $\lambda_m$  are positive roots of the transcendental equations

$$\frac{\tan(\mu l_1)}{\mu} = \frac{k_1 + k_2}{\mu^2 - k_1 k_2}, \quad \frac{\tan(\lambda l_2)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}.$$

2.1.1-19. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Mixed boundary value problems.

1°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &+ a \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &- a \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\ &- a \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\ &+ a \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{4}{l_1 l_2} \left[ \sum_{n=1}^{\infty} \sin \frac{n\pi x}{l_1} \sin \frac{n\pi \xi}{l_1} \exp \left( -\frac{\pi^2 n^2 a t}{l_1^2} \right) \right] \\ &\times \left[ \frac{1}{2} + \sum_{m=1}^{\infty} \cos \frac{m\pi y}{l_2} \cos \frac{m\pi \eta}{l_2} \exp \left( -\frac{\pi^2 m^2 a t}{l_2^2} \right) \right]. \end{aligned}$$

2°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + a \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 & + a \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\
 & + a \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & + a \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & \frac{4}{l_1 l_2} \left\{ \sum_{n=0}^{\infty} \sin \left[ \frac{\pi(2n+1)x}{2l_1} \right] \sin \left[ \frac{\pi(2n+1)\xi}{2l_1} \right] \exp \left[ -\frac{a\pi^2(2n+1)^2 t}{4l_1^2} \right] \right\} \\
 & \times \left\{ \sum_{m=0}^{\infty} \sin \left[ \frac{\pi(2m+1)y}{2l_2} \right] \sin \left[ \frac{\pi(2m+1)\eta}{2l_2} \right] \exp \left[ -\frac{a\pi^2(2m+1)^2 t}{4l_2^2} \right] \right\}.
 \end{aligned}$$

### 2.1.2. Problems in Polar Coordinates

The sourceless heat equation with two space variables in the polar coordinate system  $r, \varphi$  has the form

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} \right), \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional problems with axial symmetry that have solutions of the form  $w = w(r, t)$  are considered in Subsection 1.2.1.

2.1.2-1. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq 2\pi$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(r, \varphi) \quad \text{at} \quad t = 0.$$

Solution:

$$w(r, \varphi, t) = \frac{1}{4\pi a t} \int_0^{2\pi} \int_0^{\infty} \xi \exp \left[ -\frac{r^2 + \xi^2 - 2r\xi \cos(\varphi - \eta)}{4at} \right] f(\xi, \eta) d\xi d\eta.$$

2.1.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A circle is considered. The following conditions are prescribed:

$$w = f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$w = g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}).$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \int_0^{2\pi} \int_0^R f(\xi, \eta) G(r, \varphi, \xi, \eta, t) d\xi d\eta \\
 & - aR \int_0^t \int_0^{2\pi} g(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau.
 \end{aligned}$$



Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t),$$

$$A_0 = 1, \quad A_n = 2 \quad (n = 1, 2, \dots),$$

where  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument), and  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**2.1.2-3. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

A circle is considered. The following conditions are prescribed:

$$w = f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\partial_r w = g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}).$$

Solution:

$$w(r, \varphi, t) = \int_0^{2\pi} \int_0^R f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + aR \int_0^t \int_0^{2\pi} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau.$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{\pi R^2} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J'_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t),$$

$$A_0 = 1, \quad A_n = 2 \quad (n = 1, 2, \dots),$$

where  $J_n(\xi)$  are the Bessel functions, and  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

**2.1.2-4. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi$ . Third boundary value problem.**

A circle is considered. The following conditions are prescribed:

$$w = f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\partial_r w + kw = g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}).$$

The solution  $w(r, \varphi, t)$  is determined by the formula in Paragraph 2.1.2-3 where

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 + k^2 R^2 - n^2) [J'_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t),$$

$$A_0 = 1, \quad A_n = 2 \quad (n = 1, 2, \dots).$$

Here,  $J_n(\xi)$  are the Bessel functions, and  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k J_n(\mu R) = 0.$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**2.1.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad + aR_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\ &\quad - aR_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{\pi}{2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ A_n &= \begin{cases} 1/2 & \text{if } n = 0, \\ 1 & \text{if } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)}, \\ Z_n(\mu_{nm} r) &= J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r), \end{aligned}$$

where  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**2.1.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad - aR_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) G(r, \varphi, R_1, \eta, t - \tau) d\eta d\tau \\ &\quad + aR_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) G(r, \varphi, R_2, \eta, t - \tau) d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi(R_2^2 - R_1^2)} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t)}{(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\ Z_n(\mu_{nm} r) &= J_n'(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n'(\mu_{nm} R_1) J_n(\mu_{nm} r), \end{aligned}$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ;  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n'(\mu R_1) Y_n'(\mu R_2) - Y_n'(\mu R_1) J_n'(\mu R_2) = 0.$$

**2.1.2-7. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad - a R_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) G(r, \varphi, R_1, \eta, t - \tau) d\eta d\tau \\ &\quad + a R_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) G(r, \varphi, R_2, \eta, t - \tau) d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t)}{(k_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\ Z_n(\mu_{nm} r) &= [\mu_{nm} J_n'(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) \\ &\quad - [\mu_{nm} Y_n'(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r), \end{aligned}$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ;  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned} [\mu J_n'(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y_n'(\mu R_2) + k_2 Y_n(\mu R_2)] \\ = [\mu Y_n'(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J_n'(\mu R_2) + k_2 J_n(\mu R_2)]. \end{aligned}$$

☉ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**2.1.2-8. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ . First boundary value problem.**

A wedge domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(r, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\ w &= g_2(r, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{\varphi_0} \int_0^{\infty} f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad + a \int_0^t \int_0^{\infty} g_1(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad - a \int_0^t \int_0^{\infty} g_2(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{a \varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \sum_{n=1}^{\infty} I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right),$$

where  $I_\nu(r)$  are the modified Bessel functions.

☉ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**2.1.2-9. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ . Second boundary value problem.**

A wedge domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ r^{-1} \partial_\varphi w &= g_1(r, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_2(r, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad - a \int_0^t \int_0^\infty g_1(\xi, \tau) G(r, \varphi, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + a \int_0^t \int_0^\infty g_2(\xi, \tau) G(r, \varphi, \xi, \varphi_0, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{a\varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \left[ \frac{1}{2} I_0\left(\frac{r\xi}{2at}\right) + \sum_{n=1}^{\infty} I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \right],$$

where  $I_\nu(r)$  are the modified Bessel functions.

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**2.1.2-10. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . First boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ w &= g_2(r, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\ w &= g_3(r, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{\varphi_0} \int_0^R f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad - aR \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad + a \int_0^t \int_0^R g_2(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad - a \int_0^t \int_0^R g_3(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{4}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \exp(-\mu_{nm}^2 at),$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

**Example.** The initial temperature is uniform,  $f(r, \varphi) = w_0$ . The boundary is maintained at zero temperature,  $g_1(\varphi, t) = g_2(r, t) = g_3(r, t) = 0$ .

Solution:

$$w = \frac{8w_0}{\pi R^2} \sum_{n=0}^{\infty} \frac{1}{2n+1} \sin(s_n \varphi) \sum_{m=1}^{\infty} \exp(-\mu_{nm}^2 at) \frac{J_{s_n}(\mu_{nm} r)}{[J'_{s_n}(\mu_{nm} R)]^2} \int_0^R J_{s_n}(\mu_{nm} \xi) \xi d\xi, \quad s_n = \frac{(2n+1)\pi}{\varphi_0},$$

where the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{s_n}(\mu R) = 0$ .

☉ *References:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), H. S. Carslaw and J. C. Jaeger (1984).

**2.1.2-11. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Second boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, t) & \text{at } r &= R & (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_2(r, t) & \text{at } \varphi &= 0 & (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_3(r, t) & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{\varphi_0} \int_0^R f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad + aR \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\ &\quad - a \int_0^t \int_0^R g_2(\xi, \tau) G(r, \varphi, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + a \int_0^t \int_0^R g_3(\xi, \tau) G(r, \varphi, \xi, \varphi_0, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{2}{R^2 \varphi_0} + 4\varphi_0 \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_{nm}^2 J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{(R^2 \varphi_0^2 \mu_{nm}^2 - n^2 \pi^2) [J_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \\ &\quad \times \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \exp(-\mu_{nm}^2 at), \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_{n\pi/\varphi_0}(\mu R) = 0$ .

☉ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**2.1.2-12. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Mixed boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_r w - kw &= g(\varphi, t) & \text{at } r &= R & (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi &= 0 & (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{\varphi_0} \int_0^R f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad + aR \int_0^t \int_0^{\varphi_0} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_{nm} J_{s_n}(\mu_{nm} r) J_{s_n}(\mu_{nm} \xi) \cos(s_n \varphi) \cos(s_n \eta) \exp(-\mu_{nm}^2 a t),$$

$$s_n = \frac{n\pi}{\varphi_0}, \quad A_{nm} = \frac{4\mu_{nm}^2}{\varphi_0(\mu_{nm}^2 R^2 + k^2 R^2 - s_n^2) [J_{s_n}(\mu_{nm} R)]^2},$$

where  $J_{s_n}(r)$  are the Bessel functions, and  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_{s_n}(\mu R) + k J_{s_n}(\mu R) = 0.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

2.1.2-13. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq \varphi_0$ . Different boundary value problems.

Some problems for this domain were studied in Budak, Samarskii, and Tikhonov (1980).

### 2.1.3. Axisymmetric Problems

In the case of angular symmetry, the two-dimensional sourceless heat equation in the cylindrical coordinate system has the form

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} \right), \quad r = \sqrt{x^2 + y^2}.$$

This equation governs two-dimensional unsteady thermal processes in quiescent media or solid bodies (bounded by coordinate surfaces of the cylindrical system) in the case where the initial and boundary conditions are independent of the angular coordinate. A similar equation is used to study analogous two-dimensional unsteady mass transfer phenomena.

2.1.3-1. Particular solutions. Remarks on the Green's functions.

1°. Apart from usual separable solutions  $w(r, z, t) = f_1(r)f_2(z)f_3(t)$ , the equation in question has more sophisticated solutions in the product form

$$w(r, z, t) = u(r, t)v(z, t),$$

where  $u = u(r, t)$  and  $v = v(z, t)$  are solutions of the simpler one-dimensional equations

$$\frac{\partial u}{\partial t} = a \left( \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} \right) \quad (\text{see Subsection 1.2.1-1 for particular solutions of this equation}),$$

$$\frac{\partial v}{\partial t} = a \frac{\partial^2 v}{\partial z^2} \quad (\text{see Subsection 1.1.1-1 for particular solutions of this equation}).$$

2°. For all two-dimensional boundary value problems considered in Subsection 2.1.3, the Green's function can be represented in the product form

$$G(r, z, \xi, \eta, t) = G_1(r, \xi, t)G_2(z, \eta, t),$$

where  $G_1(r, \xi, t)$  and  $G_2(z, \eta, t)$  are the Green's functions of appropriate one-dimensional boundary value problems.

### 2.1.3-2. Domain: $0 \leq r \leq R, 0 \leq z < \infty$ . First boundary value problem.

A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= 2\pi \int_0^\infty \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad - 2\pi a R \int_0^t \int_0^\infty g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad + 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, z, \xi, \eta, t) &= G_1(r, \xi, t) G_2(z, \eta, t), \\ G_1(r, \xi, t) &= \frac{1}{\pi R^2} \sum_{n=1}^\infty \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a \mu_n^2 t}{R^2}\right), \\ G_2(z, \eta, t) &= \frac{1}{2\sqrt{\pi a t}} \left\{ \exp\left[-\frac{(z - \eta)^2}{4at}\right] - \exp\left[-\frac{(z + \eta)^2}{4at}\right] \right\}, \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu_n) = 0$ .

**Example 1.** The initial temperature is the same at every point of the cylinder,  $f(r, z) = w_0$ . The lateral surface and the end face are maintained at zero temperature,  $g_1(r, t) = g_2(z, t) = 0$ .

Solution:

$$w(r, z, t) = \frac{2w_0}{R} \operatorname{erf}\left(\frac{z}{2\sqrt{at}}\right) \sum_{n=1}^\infty \frac{J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n R)} \exp(-\lambda_n^2 at), \quad \lambda_n = \frac{\mu_n}{R}.$$

**Example 2.** The initial temperature of the cylinder is everywhere zero,  $f(r, z) = 0$ . The lateral surface  $r = R$  is maintained at a constant temperature  $w_0$ , and the end face  $z = 0$  at zero temperature.

Solution:

$$\begin{aligned} w(r, z, t) &= w_0 - \frac{w_0}{R} \sum_{n=1}^\infty \frac{J_0(\lambda_n r)}{\lambda_n J_1(\lambda_n R)} \left[ 2 \exp(-\lambda_n^2 at) \operatorname{erf}\left(\frac{z}{2\sqrt{at}}\right) \right. \\ &\quad \left. + \exp(\lambda_n z) \operatorname{erfc}\left(\frac{z}{2\sqrt{at}} + \lambda_n \sqrt{at}\right) + \exp(-\lambda_n z) \operatorname{erfc}\left(\frac{z}{2\sqrt{at}} - \lambda_n \sqrt{at}\right) \right], \end{aligned}$$

where the  $\lambda_n$  are positive zeros of the Bessel function,  $J_0(\lambda R) = 0$ .

☉ *References:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), H. S. Carslaw and J. C. Jaeger (1984).

### 2.1.3-3. Domain: $0 \leq r \leq R, 0 \leq z < \infty$ . Second boundary value problem.

A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & 2\pi \int_0^\infty \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\
 & + 2\pi a R \int_0^t \int_0^\infty g_1(\eta, \tau) G(r, z, R, \eta, t - \tau) d\eta d\tau \\
 & - 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) G(r, z, \xi, 0, t - \tau) d\xi d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, z, \xi, \eta, t) &= G_1(r, \xi, t) G_2(z, \eta, t), \\
 G_1(r, \xi, t) &= \frac{1}{\pi R^2} + \frac{1}{\pi R^2} \sum_{n=1}^\infty \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a\mu_n^2 t}{R^2}\right), \\
 G_2(z, \eta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(z-\eta)^2}{4at}\right] + \exp\left[-\frac{(z+\eta)^2}{4at}\right] \right\},
 \end{aligned}$$

where the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu_n) = 0$ .

2.1.3-4. Domain:  $0 \leq r \leq R$ ,  $0 \leq z < \infty$ . Third boundary value problem.

A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_r w + k_1 w &= g_1(z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\
 \partial_z w - k_2 w &= g_2(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(r, z, t)$  is determined by the formula in Paragraph 2.1.3-3 where

$$\begin{aligned}
 G(r, z, \xi, \eta, t) &= G_1(r, \xi, t) G_2(z, \eta, t), \\
 G_1(r, \xi, t) &= \frac{1}{\pi R^2} \sum_{n=1}^\infty \frac{\mu_n^2}{(k_1^2 R^2 + \mu_n^2) J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a\mu_n^2 t}{R^2}\right), \\
 G_2(z, \eta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(z-\eta)^2}{4at}\right] + \exp\left[-\frac{(z+\eta)^2}{4at}\right] - 2k_2 \int_0^\infty \exp\left[-\frac{(z+\eta+s)^2}{4at}\right] - k_2 s \, ds \right\}.
 \end{aligned}$$

Here,  $J_0(\mu)$  is the zeroth Bessel function and the  $\mu_n$  are positive roots of the transcendental equation

$$\mu J_1(\mu) - k_1 R J_0(\mu) = 0.$$

**Example 3.** The initial temperature is the same at every point of the cylinder,  $f(r, z) = w_0$ . At the lateral surface and the end face, heat exchange of the cylinder with the zero temperature environment occurs,  $g_1(z, t) = g_2(r, t) = 0$ .

Solution:

$$w(r, z, t) = \frac{2w_0 k_1}{R} \left[ \operatorname{erf}\left(\frac{z}{2\sqrt{at}}\right) + \exp(k_2 z + k_2^2 at) \operatorname{erfc}\left(\frac{z}{2\sqrt{at}} + k_2 \sqrt{at}\right) \right] \sum_{n=1}^\infty \frac{J_0(\nu_n r) \exp(-\nu_n^2 at)}{(k_1^2 + \nu_n^2) J_0(\nu_n R)},$$

where the  $\nu_n$  are positive roots of the transcendental equation  $\nu J_1(\nu R) - k_1 J_0(\nu R) = 0$ .

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).



**2.1.3-5. Domain:  $0 \leq r \leq R, 0 \leq z < \infty$ . Mixed boundary value problems.**

1°. A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= 2\pi \int_0^\infty \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad - 2\pi a R \int_0^t \int_0^\infty g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad - 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) G(r, z, \xi, 0, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, z, \xi, \eta, t) &= G_1(r, \xi, t) G_2(z, \eta, t), \\ G_1(r, \xi, t) &= \frac{1}{\pi R^2} \sum_{n=1}^\infty \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a\mu_n^2 t}{R^2}\right), \\ G_2(z, \eta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(z - \eta)^2}{4at}\right] + \exp\left[-\frac{(z + \eta)^2}{4at}\right] \right\}, \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu_n) = 0$ .

2°. A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= 2\pi \int_0^\infty \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad + 2\pi a R \int_0^t \int_0^\infty g_1(\eta, \tau) G(r, z, R, \eta, t - \tau) d\eta d\tau \\ &\quad + 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, z, \xi, \eta, t) &= G_1(r, \xi, t) G_2(z, \eta, t), \\ G_1(r, \xi, t) &= \frac{1}{\pi R^2} + \frac{1}{\pi R^2} \sum_{n=1}^\infty \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a\mu_n^2 t}{R^2}\right), \\ G_2(z, \eta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(z - \eta)^2}{4at}\right] - \exp\left[-\frac{(z + \eta)^2}{4at}\right] \right\}, \end{aligned}$$

where the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu_n) = 0$ .

3°. A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g_1(z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ w &= g_2(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, z, t)$  is determined by the formula in Paragraph 2.1.3-5, Item 2° where

$$\begin{aligned} G(r, z, \xi, \eta, t) &= G_1(r, \xi, t)G_2(z, \eta, t), \\ G_1(r, \xi, t) &= \frac{1}{\pi R^2} \sum_{n=1}^{\infty} \frac{\mu_n^2}{(k^2 R^2 + \mu_n^2) J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a \mu_n^2 t}{R^2}\right), \\ G_2(z, \eta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(z-\eta)^2}{4at}\right] - \exp\left[-\frac{(z+\eta)^2}{4at}\right] \right\}, \end{aligned}$$

where the  $\mu_n$  are positive roots of the transcendental equation

$$\mu J_1(\mu) - kR J_0(\mu) = 0.$$

**Example 4.** The initial temperature is the same at every point of the cylinder,  $f(r, z) = w_0$ . Heat exchange of the cylinder with the zero temperature environment occurs at the lateral surface,  $g_1(z, t) = 0$ . The end face is maintained at zero temperature,  $g_2(r, t) = 0$ .

Solution:

$$w(r, z, t) = \frac{2w_0 k}{R} \operatorname{erf}\left(\frac{z}{2\sqrt{at}}\right) \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{(k^2 + \lambda_n^2) J_0(\lambda_n R)} \exp(-\lambda_n^2 at), \quad \lambda_n = \frac{\mu_n}{R}.$$

**Example 5.** The initial temperature of the cylinder is everywhere zero,  $f(r, z) = 0$ . Heat exchange of the cylinder with the zero temperature environment occurs at the lateral surface,  $g_1(z, t) = 0$ . The end face is maintained at a constant temperature,  $g_2(r, t) = w_0$ .

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{k w_0}{R} \sum_{n=1}^{\infty} \frac{J_0(\lambda_n r)}{(\lambda_n^2 + k^2) J_0(\lambda_n R)} \left[ 2 \exp(-\lambda_n z) \right. \\ &\quad \left. + \exp(\lambda_n z) \operatorname{erfc}\left(\lambda_n \sqrt{at} + \frac{z}{2\sqrt{at}}\right) - \exp(-\lambda_n z) \operatorname{erfc}\left(\lambda_n \sqrt{at} - \frac{z}{2\sqrt{at}}\right) \right], \quad \lambda_n = \frac{\mu_n}{R}. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

### 2.1.3-6. Domain: $0 \leq r \leq R$ , $0 \leq z \leq l$ . First boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ w &= g_2(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_3(r, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= 2\pi \int_0^l \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad - 2\pi a R \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad + 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad - 2\pi a \int_0^t \int_0^R \xi g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau. \end{aligned}$$

Here,

$$G(r, z, \xi, \eta, t) = G_1(r, \xi, t)G_2(z, \eta, t),$$

$$G_1(r, \xi, t) = \frac{1}{\pi R^2} \sum_{n=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a\mu_n^2 t}{R^2}\right),$$

$$G_2(z, \eta, t) = \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi \eta}{l}\right) \exp\left(-\frac{an^2 \pi^2 t}{l^2}\right),$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu_n) = 0$ .

**Example 6.** The initial temperature is the same at every point of the cylinder,  $f(r, z) = w_0$ . The lateral surface and the end faces are maintained at zero temperature,  $g_1(z, t) = g_2(r, t) = g_3(r, t) = 0$ .

Solution:

$$w = \frac{8w_0}{\pi} \left\{ \sum_{n=0}^{\infty} \frac{1}{2n+1} \sin\left[\frac{(2n+1)\pi z}{l}\right] \exp\left[-\frac{a(2n+1)^2 \pi^2 t}{l^2}\right] \right\} \left\{ \sum_{n=1}^{\infty} \frac{1}{\mu_n J_1(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) \exp\left(-\frac{\mu_n^2 a t}{R^2}\right) \right\},$$

where  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu_n) = 0$ .

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

### 2.1.3-7. Domain: $0 \leq r \leq R$ , $0 \leq z \leq l$ . Second boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(z, t) & \text{at } r &= R & \text{(boundary condition),} \\ \partial_z w &= g_2(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= 2\pi \int_0^l \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &+ 2\pi a R \int_0^t \int_0^l g_1(\eta, \tau) G(r, z, R, \eta, t - \tau) d\eta d\tau \\ &- 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) G(r, z, \xi, 0, t - \tau) d\xi d\tau \\ &+ 2\pi a \int_0^t \int_0^R \xi g_3(\xi, \tau) G(r, z, \xi, l, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, z, \xi, \eta, t) = G_1(r, \xi, t)G_2(z, \eta, t),$$

$$G_1(r, \xi, t) = \frac{1}{\pi R^2} + \frac{1}{\pi R^2} \sum_{n=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a\mu_n^2 t}{R^2}\right),$$

$$G_2(z, \eta, t) = \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi \eta}{l}\right) \exp\left(-\frac{an^2 \pi^2 t}{l^2}\right),$$

where the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu_n) = 0$ .

**2.1.3-8. Domain:  $0 \leq r \leq R, 0 \leq z \leq l$ . Third boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + k_1 w &= g_1(z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ \partial_z w - k_2 w &= g_2(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w + k_3 w &= g_3(r, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, z, t)$  is determined by the formula in Paragraph 2.1.3-7 where

$$\begin{aligned} G(r, z, \xi, \eta, t) &= G_1(r, \xi, t)G_2(z, \eta, t), \\ G_1(r, \xi, t) &= \frac{1}{\pi R^2} \sum_{n=1}^{\infty} \frac{\mu_n^2}{(k_1^2 R^2 + \mu_n^2) J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a \mu_n^2 t}{R^2}\right), \\ G_2(z, \eta, t) &= \sum_{m=1}^{\infty} \frac{\varphi_m(z) \varphi_m(\eta)}{\|\varphi_m\|^2} \exp(-a \lambda_m^2 t), \quad \varphi_m(z) = \cos(\lambda_m z) + \frac{k_2}{\lambda_m} \sin(\lambda_m z), \\ \|\varphi_m\|^2 &= \frac{k_3}{2 \lambda_m^2} \frac{\lambda_m^2 + k_2^2}{\lambda_m^2 + k_3^2} + \frac{k_2}{2 \lambda_m^2} + \frac{l}{2} \left(1 + \frac{k_2^2}{\lambda_m^2}\right), \end{aligned}$$

and the  $\mu_n$  and  $\lambda_m$  are positive roots of the transcendental equations

$$\mu J_1(\mu) - k_1 R J_0(\mu) = 0, \quad \frac{\tan(\lambda l)}{\lambda} = \frac{k_2 + k_3}{\lambda^2 - k_2 k_3}.$$

**2.1.3-9. Domain:  $0 \leq r \leq R, 0 \leq z \leq l$ . Mixed boundary value problems.**

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= 2\pi \int_0^l \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad - 2\pi a R \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad - 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) G(r, z, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + 2\pi a \int_0^t \int_0^R \xi g_3(\xi, \tau) G(r, z, \xi, l, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, z, \xi, \eta, t) &= G_1(r, \xi, t)G_2(z, \eta, t), \\ G_1(r, \xi, t) &= \frac{1}{\pi R^2} \sum_{n=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a \mu_n^2 t}{R^2}\right), \\ G_2(z, \eta, t) &= \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi \eta}{l}\right) \exp\left(-\frac{a n^2 \pi^2 t}{l^2}\right), \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu_n) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ w &= g_2(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_3(r, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}).\end{aligned}$$

Solution:

$$\begin{aligned}w(r, z, t) &= 2\pi \int_0^l \int_0^R \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &+ 2\pi a R \int_0^t \int_0^l g_1(\eta, \tau) G(r, z, R, \eta, t - \tau) d\eta d\tau \\ &+ 2\pi a \int_0^t \int_0^R \xi g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- 2\pi a \int_0^t \int_0^R \xi g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau.\end{aligned}$$

Here,

$$\begin{aligned}G(r, z, \xi, \eta, t) &= G_1(r, \xi, t) G_2(z, \eta, t), \\ G_1(r, \xi, t) &= \frac{1}{\pi R^2} + \frac{1}{\pi R^2} \sum_{n=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \exp\left(-\frac{a \mu_n^2 t}{R^2}\right), \\ G_2(z, \eta, t) &= \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n \pi z}{l}\right) \sin\left(\frac{n \pi \eta}{l}\right) \exp\left(-\frac{a n^2 \pi^2 t}{l^2}\right),\end{aligned}$$

where the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu_n) = 0$ .

2.1.3-10. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . First boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(z, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(z, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}), \\ w &= g_3(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_4(r, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}).\end{aligned}$$

Solution:

$$\begin{aligned}w(r, z, t) &= 2\pi \int_0^l \int_{R_1}^{R_2} \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &+ 2\pi a R_1 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\ &- 2\pi a R_2 \int_0^t \int_0^l g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau \\ &+ 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau.\end{aligned}$$

Here,

$$G(r, z, \xi, \eta, t) = G_1(z, \eta, t)G_2(r, \xi, t),$$

$$G_1(z, \eta, t) = \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi \eta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right),$$

$$G_2(r, \xi, t) = \frac{\pi}{4R_1^2} \sum_{n=1}^{\infty} \frac{\mu_n^2 J_0^2(s\mu_n)}{J_0^2(\mu_n) - J_0^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \exp\left(-\frac{a\mu_n^2 t}{R_1^2}\right),$$

$$\Psi_n(r) = Y_0(\mu_n)J_0\left(\frac{\mu_n r}{R_1}\right) - J_0(\mu_n)Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1},$$

where  $J_0(\mu)$  and  $Y_0(\mu)$  are the Bessel functions, the  $\mu_n$  are positive roots of the transcendental equation

$$J_0(\mu)Y_0(s\mu) - J_0(s\mu)Y_0(\mu) = 0.$$

2.1.3-11. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(z, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ \partial_r w &= g_2(z, t) & \text{at } r &= R_2 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_4(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= 2\pi \int_0^l \int_{R_1}^{R_2} \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad - 2\pi a R_1 \int_0^t \int_0^l g_1(\eta, \tau) G(r, z, R_1, \eta, t - \tau) d\eta d\tau \\ &\quad + 2\pi a R_2 \int_0^t \int_0^l g_2(\eta, \tau) G(r, z, R_2, \eta, t - \tau) d\eta d\tau \\ &\quad - 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_3(\xi, \tau) G(r, z, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_4(\xi, \tau) G(r, z, \xi, l, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, z, \xi, \eta, t) = G_1(z, \eta, t)G_2(r, \xi, t),$$

$$G_1(z, \eta, t) = \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi \eta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right),$$

$$G_2(r, \xi, t) = \frac{1}{\pi(R_2^2 - R_1^2)} + \frac{\pi}{4R_1^2} \sum_{n=1}^{\infty} \frac{\mu_n^2 J_1^2(s\mu_n)}{J_1^2(\mu_n) - J_1^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \exp\left(-\frac{a\mu_n^2 t}{R_1^2}\right),$$

$$\Psi_n(r) = Y_1(\mu_n)J_0\left(\frac{\mu_n r}{R_1}\right) - J_1(\mu_n)Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1},$$

where  $J_k(\mu)$  and  $Y_k(\mu)$  are the Bessel functions of order  $k = 0, 1$  and the  $\mu_n$  are positive roots of the transcendental equation

$$J_1(\mu)Y_1(s\mu) - J_1(s\mu)Y_1(\mu) = 0.$$

**2.1.3-12. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . Third boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(z, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(z, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}), \\ \partial_z w - k_3 w &= g_3(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w + k_4 w &= g_4(r, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

For the solution of this problem, see Subsection 2.2.3-4 with  $\Phi \equiv 0$ .

## 2.2. Heat Equation with a Source $\frac{\partial w}{\partial t} = a\Delta_2 w + \Phi(x, y, t)$

### 2.2.1. Problems in Cartesian Coordinates

In the rectangular Cartesian coordinate system, the heat equation has the form

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + \Phi(x, y, t).$$

It governs two-dimensional unsteady thermal processes in quiescent media or solids with constant thermal diffusivity in the cases where there are volume thermal sources or sinks.

**2.2.1-1. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ . Cauchy problem.**

An initial condition is prescribed:

$$w = f(x, y) \quad \text{at} \quad t = 0.$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &+ \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right].$$

☉ *Reference:* A. G. Butkovskiy (1979).

**2.2.1-2. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ . First boundary value problem.**

A half-plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &+ a \int_0^t \int_{-\infty}^{\infty} g(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &+ \int_0^t \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] - \exp \left[ -\frac{(x + \xi)^2 + (y - \eta)^2}{4at} \right] \right\}.$$

⊙ *References:* A. G. Butkovskiy (1979), H. S. Carslaw and J. C. Jaeger (1984).

2.2.1-3. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ . Second boundary value problem.

A half-plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_{-\infty}^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi - a \int_0^t \int_{-\infty}^\infty g(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\ &\quad + \int_0^t \int_0^\infty \int_{-\infty}^\infty \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] + \exp \left[ -\frac{(x + \xi)^2 + (y - \eta)^2}{4at} \right] \right\}.$$

2.2.1-4. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ . Third boundary value problem.

A half-plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - kw &= g(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.2.1-3 where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{4\pi at} \exp \left[ -\frac{(y - \eta)^2}{4at} \right] \left\{ \exp \left[ -\frac{(x - \xi)^2}{4at} \right] + \exp \left[ -\frac{(x + \xi)^2}{4at} \right] \right. \\ &\quad \left. - 2k \int_0^\infty \exp \left[ -\frac{(x + \xi + s)^2}{4at} - ks \right] ds \right\}. \end{aligned}$$

2.2.1-5. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.

A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad + a \int_0^t \int_0^\infty g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &\quad + a \int_0^t \int_0^\infty g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad + \int_0^t \int_0^\infty \int_0^\infty \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\xi d\eta d\tau, \end{aligned}$$



where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] - \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right\} \left\{ \exp \left[ -\frac{(y-\eta)^2}{4at} \right] - \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right\}.$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**2.2.1-6. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . Second boundary value problem.**

A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_2(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad - a \int_0^t \int_0^\infty g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\ &\quad - a \int_0^t \int_0^\infty g_2(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + \int_0^t \int_0^\infty \int_0^\infty \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] + \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right\} \left\{ \exp \left[ -\frac{(y-\eta)^2}{4at} \right] + \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right\}.$$

**2.2.1-7. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . Third boundary value problem.**

A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_y w - k_2 w &= g_2(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.2.1-6 where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] + \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right. \\ &\quad \left. - 2k_1 \sqrt{\pi at} \exp[ak_1^2 t + k_1(x+\xi)] \operatorname{erfc} \left( \frac{x+\xi}{2\sqrt{at}} + k_1 \sqrt{at} \right) \right\} \\ &\quad \times \left\{ \exp \left[ -\frac{(y-\eta)^2}{4at} \right] + \exp \left[ -\frac{(y+\eta)^2}{4at} \right] \right. \\ &\quad \left. - 2k_2 \sqrt{\pi at} \exp[ak_2^2 t + k_2(y+\eta)] \operatorname{erfc} \left( \frac{y+\eta}{2\sqrt{at}} + k_2 \sqrt{at} \right) \right\}. \end{aligned}$$

**2.2.1-8. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . Mixed boundary value problem.**

A quadrant of the plane is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_2(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^\infty f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta \\ &\quad + a \int_0^t \int_0^\infty g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &\quad - a \int_0^t \int_0^\infty g_2(\xi, \tau) G(x, y, 0, t - \tau) d\xi d\tau \\ &\quad + \int_0^t \int_0^\infty \int_0^\infty \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi at} \left\{ \exp \left[ -\frac{(x - \xi)^2}{4at} \right] - \exp \left[ -\frac{(x + \xi)^2}{4at} \right] \right\} \left\{ \exp \left[ -\frac{(y - \eta)^2}{4at} \right] + \exp \left[ -\frac{(y + \eta)^2}{4at} \right] \right\}.$$

**2.2.1-9. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . First boundary value problem.**

A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x = l & \quad (\text{boundary condition}), \\ w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution is given by the formula of Subsection 2.1.1-12 with the additional term

$$\int_0^t \int_0^l \int_0^\infty \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,$$

which takes into account the equation's nonhomogeneity.

**2.2.1-10. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . Second boundary value problem.**

A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) & \text{at } x = l & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^\infty \int_0^l f(\xi, \eta) G(x, y, \xi, \eta, t) d\xi d\eta + \int_0^t \int_0^\infty \int_0^l \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\xi d\eta d\tau \\ &\quad - a \int_0^t \int_0^\infty g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau + a \int_0^t \int_0^\infty g_2(\eta, \tau) G(x, y, l, \eta, t - \tau) d\eta d\tau \\ &\quad - a \int_0^t \int_0^l g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = G_1(x, \xi, t)G_2(y, \eta, t),$$

$$G_1(x, \xi, t) = \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi \xi}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right),$$

$$G_2(y, \eta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp\left[-\frac{(y-\eta)^2}{4at}\right] + \exp\left[-\frac{(y+\eta)^2}{4at}\right] \right\}.$$

**2.2.1-11. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . Third boundary value problem.**

A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(y, t) & \text{at } x = l & \quad (\text{boundary condition}), \\ \partial_y w - k_3 w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.2.1-10 where the Green's function  $G(x, y, \xi, \eta, t)$  is the product of the Green's function of Subsection 1.1.1-11 and that of Subsection 1.1.1-8;  $x$ ,  $\xi$ , and  $k$  in the last Green's function must be replaced by  $y$ ,  $\eta$ , and  $k_3$ , respectively.

**2.2.1-12. Domain:  $0 \leq x \leq l$ ,  $0 \leq y < \infty$ . Mixed boundary value problem.**

A semiinfinite strip is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x = l & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution is given by the formula of Subsection 2.1.1-15 (Item 1°) with the additional term

$$\int_0^t \int_0^l \int_0^\infty \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,$$

which takes into account the equation's nonhomogeneity.

**2.2.1-13. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . First boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution is given by the formula of Subsection 2.1.1-16 with the additional term

$$\int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,$$

which takes into account the equation's nonhomogeneity.

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**2.2.1-14. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Second boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

**Solution:**

$$\begin{aligned} w(x, y, t) &= \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t-\tau) d\eta d\xi d\tau \\ &\quad - a \int_0^t \int_0^{l_2} g_1(\eta, \tau) G(x, y, 0, \eta, t-\tau) d\eta d\tau + a \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t-\tau) d\eta d\tau \\ &\quad - a \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t-\tau) d\xi d\tau + a \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t-\tau) d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{l_1 l_2} \left[ 1 + 2 \sum_{n=1}^{\infty} \exp\left(-\frac{\pi^2 n^2 a t}{l_1^2}\right) \cos \frac{n\pi x}{l_1} \cos \frac{n\pi \xi}{l_1} \right] \\ &\quad \times \left[ 1 + 2 \sum_{m=1}^{\infty} \exp\left(-\frac{\pi^2 m^2 a t}{l_2^2}\right) \cos \frac{m\pi y}{l_2} \cos \frac{m\pi \eta}{l_2} \right]. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**2.2.1-15. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Third boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w - k_3 w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w + k_4 w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 2.2.1-14 where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \left\{ \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\|\varphi_n\|^2} \exp(-a\mu_n^2 t) \right\} \left\{ \sum_{m=1}^{\infty} \frac{\psi_m(y) \psi_m(\eta)}{\|\psi_m\|^2} \exp(-a\lambda_m^2 t) \right\}, \\ \varphi_n(x) &= \cos(\mu_n x) + \frac{k_1}{\mu_n} \sin(\mu_n x), \quad \|\varphi_n\|^2 = \frac{k_2}{2\mu_n^2} \frac{\mu_n^2 + k_1^2}{\mu_n^2 + k_2^2} + \frac{k_1}{2\mu_n^2} + \frac{l_1}{2} \left( 1 + \frac{k_1^2}{\mu_n^2} \right), \\ \psi_m(y) &= \cos(\lambda_m y) + \frac{k_3}{\lambda_m} \sin(\lambda_m y), \quad \|\psi_m\|^2 = \frac{k_4}{2\lambda_m^2} \frac{\lambda_m^2 + k_3^2}{\lambda_m^2 + k_4^2} + \frac{k_3}{2\lambda_m^2} + \frac{l_2}{2} \left( 1 + \frac{k_3^2}{\lambda_m^2} \right). \end{aligned}$$

Here, the  $\mu_n$  and  $\lambda_m$  are positive roots of the transcendental equations

$$\frac{\tan(\mu l_1)}{\mu} = \frac{k_1 + k_2}{\mu^2 - k_1 k_2}, \quad \frac{\tan(\lambda l_2)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}.$$

**2.2.1-16. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Mixed boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(y, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution is given by the formula of Subsection 2.1.1-19 (Item 1°) with the additional term

$$\int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,$$

which takes into account the equation's nonhomogeneity.

## 2.2.2. Problems in Polar Coordinates

The heat equation with a volume source in the polar coordinate system  $r, \varphi$  is written as

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} \right) + \Phi(r, \varphi, t).$$

Solutions of the form  $w = w(r, t)$  that are independent of the angular coordinate  $\varphi$  and govern plane thermal processes with central symmetry, are presented in Subsection 1.2.2.

**2.2.2-1. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq 2\pi$ . Cauchy problem.**

An initial condition is prescribed:

$$w = f(r, \varphi) \quad \text{at} \quad t = 0.$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \int_0^{2\pi} \int_0^\infty f(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &+ \int_0^t \int_0^{2\pi} \int_0^\infty \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau, \end{aligned}$$

where

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{4\pi a t} \exp \left[ -\frac{r^2 + \xi^2 - 2r\xi \cos(\varphi - \eta)}{4at} \right].$$

**2.2.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a circle of radius  $R$  is given by the formula from Subsection 2.1.2-2 with the additional term

$$\int_0^t \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a circle is given by the formula in Paragraph 2.1.2-3 with the additional term (1).

3°. The solution of the third boundary value problem for a circle is given by the formula in Paragraph 2.1.2-4 with the additional term (1).

**2.2.2-3. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for an annular domain is given by the formula in Paragraph 2.1.2-5 with the additional term

$$\int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the third boundary value problem for an annular domain is given by the formula in Paragraph 2.1.2-7 with the additional term (2).

**2.2.2-4. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a wedge domain is given by the formula in Paragraph 2.1.2-8 with the additional term

$$\int_0^t \int_0^{\varphi_0} \int_0^\infty \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau, \quad (3)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a wedge domain is given by the formula in Paragraph 2.1.2-9 with the additional term (3).

**2.2.2-5. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a sector of a circle is given by the formula of Paragraph 2.1.2-10 with the additional term

$$\int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau, \quad (4)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the mixed boundary value problem for a sector of a circle is given by the formula of Paragraph 2.1.2-11 with the additional term (4).

## 2.2.3. Axisymmetric Problems

In the case of axial symmetry, the heat equation in the cylindrical coordinate system is written as

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} \right) + \Phi(r, z, t),$$

provided there are heat sources or sinks.

One-dimensional axisymmetric problems that have solutions of the form  $w = w(r, t)$  can be found in Subsection 1.2.2.

**2.2.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq z < \infty$ . Different boundary value problems.**

1°. The solution to the first boundary value problem for a semiinfinite circular cylinder of radius  $R$  is given by the formula of Subsection 2.1.3-2 with the term

$$2\pi \int_0^t \int_0^\infty \int_0^R \xi \Phi(\xi, \eta, \tau) G(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau \quad (1)$$

added; this term takes into account the nonhomogeneity of the equation.

2°. The solution to the second boundary value problem for a semiinfinite circular cylinder is given by the formula of Subsection 2.1.3-3 with the additional term (1).

3°. The solution to the third boundary value problem for a semiinfinite circular cylinder is given by the formula of Subsection 2.1.3-4 with the additional term (1).

4°. The solutions to various mixed boundary value problems for a semiinfinite circular cylinder are defined by formulas of Subsection 2.1.3-5 with additional terms of the form (1).

**2.2.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution to the first boundary value problem for a circular cylinder of radius  $R$  and length  $l$  is given by the formula of Subsection 2.1.3-6 with the term

$$2\pi \int_0^t \int_0^l \int_0^R \xi \Phi(\xi, \eta, \tau) G(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau, \quad (2)$$

added; this term takes into account the nonhomogeneity of the equation.

2°. The solution to the second boundary value problem for a finite circular cylinder is given by the formula of Subsection 2.1.3-7 with the additional term (2).

3°. The solution to the third boundary value problem for a finite circular cylinder is given by the formula of Subsection 2.1.3-8 with the additional term (2).

4°. The solutions to various mixed boundary value problems for a finite circular cylinder are defined by formulas of Subsection 2.1.3-9 with additional terms of the form (2).

**2.2.3-3. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . First and second boundary value problems.**

1°. The solution to the first boundary value problem for a hollow circular cylinder of interior radius  $R_1$ , exterior radius  $R_2$ , and length  $l$  is given by the formula of Subsection 2.1.3-10 with the term

$$2\pi \int_0^t \int_0^l \int_{R_1}^{R_2} \xi \Phi(\xi, \eta, \tau) G(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau \quad (3)$$

added; this term takes into account the equation's nonhomogeneity.

2°. The solution to the second boundary value problem for a finite hollow circular cylinder is given by the formula of Subsection 2.1.3-11 with the additional term (3).

**2.2.3-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . Third boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ \partial_z w - k_3 w &= g_3(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w + k_4 w &= g_4(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & 2\pi \int_0^l \int_{R_1}^{R_2} \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\
 & - 2\pi a R_1 \int_0^t \int_0^l g_1(\eta, \tau) G(r, z, R_1, \eta, t - \tau) d\eta d\tau \\
 & + 2\pi a R_2 \int_0^t \int_0^l g_2(\eta, \tau) G(r, z, R_2, \eta, t - \tau) d\eta d\tau \\
 & - 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_3(\xi, \tau) G(r, z, \xi, 0, t - \tau) d\xi d\tau \\
 & + 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_4(\xi, \tau) G(r, z, \xi, l, t - \tau) d\xi d\tau \\
 & + 2\pi \int_0^t \int_0^l \int_{R_1}^{R_2} \xi \Phi(\xi, \eta, \tau) G(r, \xi, z, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here, the Green's function is given by

$$\begin{aligned}
 G(r, z, \xi, \eta, t) &= G_1(r, \xi, t) G_2(z, \eta, t), \\
 G_1(r, \xi, t) &= \frac{\pi}{4} \sum_{n=1}^{\infty} \frac{\lambda_n^2}{B_n} [k_2 J_0(\lambda_n R_2) - \lambda_n J_1(\lambda_n R_2)]^2 H_n(r) H_n(\xi) \exp(-\lambda_n^2 a t), \\
 G_2(z, \eta, t) &= \sum_{m=1}^{\infty} \frac{\varphi_m(z) \varphi_m(\eta)}{\|\varphi_m\|^2} \exp(-\mu_m^2 a t),
 \end{aligned}$$

where

$$\begin{aligned}
 B_n &= (\lambda_n^2 + k_2^2) [k_1 J_0(\lambda_n R_1) + \lambda_n J_1(\lambda_n R_1)]^2 - (\lambda_n^2 + k_1^2) [k_2 J_0(\lambda_n R_2) - \lambda_n J_1(\lambda_n R_2)]^2, \\
 H_n(r) &= [k_1 Y_0(\lambda_n R_1) + \lambda_n Y_1(\lambda_n R_1)] J_0(\lambda_n r) - [k_1 J_0(\lambda_n R_1) + \lambda_n J_1(\lambda_n R_1)] Y_0(\lambda_n r), \\
 \varphi_m(z) &= \mu_m \cos(\mu_m z) + k_3 \sin(\mu_m z), \quad \|\varphi_m\|^2 = \frac{k_4}{2} \frac{\mu_m^2 + k_3^2}{\mu_m^2 + k_4^2} + \frac{k_3}{2} + \frac{l}{2} (\mu_m^2 + k_3^2),
 \end{aligned}$$

$J_0(\lambda)$ ,  $J_1(\lambda)$ ,  $Y_0(\lambda)$ , and  $Y_1(\lambda)$  are the Bessel functions, the  $\lambda_n$  are positive roots of the transcendental equation

$$\begin{aligned}
 & [k_1 J_0(\lambda R_1) + \lambda J_1(\lambda R_1)] [k_2 Y_0(\lambda R_2) - \lambda Y_1(\lambda R_2)] \\
 & - [k_2 J_0(\lambda R_2) - \lambda J_1(\lambda R_2)] [k_1 Y_0(\lambda R_1) + \lambda Y_1(\lambda R_1)] = 0,
 \end{aligned}$$

and the  $\mu_m$  are positive roots of the transcendental equation

$$\frac{\tan \mu l}{\mu} = \frac{k_3 + k_4}{\mu^2 - k_3 k_4}.$$

⊙ Reference: A. G. Butkovskiy (1979).

2.2.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . Mixed boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(r, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_r w - k_1 w &= g_1(z, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\
 \partial_r w + k_2 w &= g_2(z, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}), \\
 w &= g_3(r, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
 w &= g_4(r, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}).
 \end{aligned}$$



Solution:

$$\begin{aligned}
 w(r, z, t) = & 2\pi \int_0^l \int_{R_1}^{R_2} \xi f(\xi, \eta) G(r, z, \xi, \eta, t) d\xi d\eta \\
 & - 2\pi a R_1 \int_0^t \int_0^l g_1(\eta, \tau) G(r, z, R_1, \eta, t - \tau) d\eta d\tau \\
 & + 2\pi a R_2 \int_0^t \int_0^l g_2(\eta, \tau) G(r, z, R_2, \eta, t - \tau) d\eta d\tau \\
 & + 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & - 2\pi a \int_0^t \int_{R_1}^{R_2} \xi g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau \\
 & + 2\pi \int_0^t \int_0^l \int_{R_1}^{R_2} \xi \Phi(\xi, \eta, \tau) G(r, \xi, z, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$G(r, \xi, z, \eta, t) = G_1(r, \xi, t) \frac{2}{l} \sum_{m=1}^{\infty} \sin\left(\frac{\pi m z}{l}\right) \sin\left(\frac{\pi m \eta}{l}\right) \exp\left(-\frac{\pi^2 m^2 a t}{l^2}\right),$$

where the expression of  $G_1(r, \xi, t)$  is specified in Subsection 2.2.3-4.

► Subsection 3.2.2 presents solutions of other boundary value problems; a more general, three-dimensional equation is discussed there.

## 2.3. Other Equations

### 2.3.1. Equations Containing Arbitrary Parameters

$$1. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + (bx + cy + k)w.$$

The transformation

$$w(x, y, t) = u(\xi, \eta, t) \exp\left[(bx + cy + k)t + \frac{1}{3}a(b^2 + c^2)t^3\right], \quad \xi = x + abt^2, \quad \eta = y + act^2$$

leads to the two-dimensional heat equation  $\partial_t u = a(\partial_{\xi\xi} u + \partial_{\eta\eta} u)$ .

See also Niederer (1973) and Boyer (1974).

$$2. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) - (bx^2 + by^2 + k)w, \quad b > 0.$$

The transformation ( $C$  is an arbitrary constant)

$$\begin{aligned}
 w(x, y, t) = & u(\xi, \eta, \tau) \exp\left[\frac{1}{2}\sqrt{\frac{b}{a}}(x^2 + y^2) + (2\sqrt{ab} - k)t\right], \\
 \xi = & x \exp(2\sqrt{ab}t), \quad \eta = y \exp(2\sqrt{ab}t), \quad \tau = \frac{1}{4\sqrt{ab}} \exp(4\sqrt{ab}t) + C
 \end{aligned}$$

leads to the two-dimensional heat equation  $\partial_{\tau} u = a(\partial_{\xi\xi} u + \partial_{\eta\eta} u)$ .

See also Niederer (1973) and Boyer (1974).

$$3. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + (bx^2 + by^2 - k)w, \quad b > 0.$$

The transformation

$$w(x, y, t) = \frac{1}{\cos(2\sqrt{ab}t)} \exp \left[ \frac{\sqrt{b}}{2\sqrt{a}} \tan(2\sqrt{ab}t) (x^2 + y^2) - kt \right] u(\xi, \eta, \tau),$$

$$\xi = \frac{x}{\cos(2\sqrt{ab}t)}, \quad \eta = \frac{y}{\cos(2\sqrt{ab}t)}, \quad \tau = \frac{\sqrt{a}}{2\sqrt{b}} \tan(2\sqrt{ab}t)$$

leads to the two-dimensional heat equation  $\partial_\tau u = \partial_{\xi\xi} u + \partial_{\eta\eta} u$ .

See also Niederer (1973) and Boyer (1974).

$$4. \quad \frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + (ax^{-2} + by^{-2})w.$$

This is a special case of equation 2.3.2.7. Boyer (1976) showed that this equation admits the separation of variables into 25 systems of coordinates for  $ab = 0$  and 15 systems of coordinates for  $ab \neq 0$ .

$$5. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + (bt^n x + ct^m y + st^k)w.$$

This is a special case of equation 2.3.2.2 with  $f(t) = bt^n$ ,  $g(t) = ct^m$ , and  $h(t) = st^k$ .

$$6. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + [-b(x^2 + y^2) + c_1 t^{n_1} x + c_2 t^{n_2} y + st^k]w.$$

This is a special case of equation 2.3.2.3 with  $f(t) = c_1 t^{n_1}$ ,  $g(t) = c_2 t^{n_2}$ , and  $h(t) = st^k$ .

$$7. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + b_1 \frac{\partial w}{\partial x} + b_2 \frac{\partial w}{\partial y} + cw.$$

This equation describes an unsteady temperature (concentration) field in a medium moving with a constant velocity, provided there is volume release (absorption) of heat proportional to temperature.

The substitution

$$w(x, y, t) = \exp(A_1 x + A_2 y + Bt) U(x, y, t),$$

$$A_1 = -\frac{b_1}{2a}, \quad A_2 = -\frac{b_2}{2a}, \quad B = c - \frac{b_1^2 + b_2^2}{4a},$$

leads to the two-dimensional heat equation  $\partial_t U = a\Delta_2 U$  that is considered in Subsection 2.1.1.

$$8. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + b_1 \frac{\partial w}{\partial x} + b_2 \frac{\partial w}{\partial y} + (c_1 x + c_2 y + k)w.$$

The transformation

$$w(x, y, t) = \exp[(c_1 x + c_2 y)t + \frac{1}{3}a(c_1^2 + c_2^2)t^3 + \frac{1}{2}(b_1 c_1 + b_2 c_2)t^2 + kt] U(\xi, \eta, t),$$

$$\xi = x + ac_1 t^2 + b_1 t, \quad \eta = y + ac_2 t^2 + b_2 t$$

leads to the two-dimensional heat equation  $\partial_t U = a(\partial_{\xi\xi} U + \partial_{\eta\eta} U)$  that is considered in Subsection 2.1.1.

$$9. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + b_1 t^{n_1} \frac{\partial w}{\partial x} + b_2 t^{n_2} \frac{\partial w}{\partial y} + (c_1 t^{m_1} x + c_2 t^{m_2} y + st^k)w.$$

This is a special case of equation 2.3.2.5. The equation can be reduced to the two-dimensional heat equation treated in Subsection 2.1.1.

$$10. \quad i\hbar \frac{\partial w}{\partial t} + \frac{\hbar^2}{2m} \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) = 0.$$

Two-dimensional Schrödinger equation,  $i^2 = -1$ .

Fundamental solution:

$$\mathcal{E}(x, y, t) = -\frac{im}{2\pi\hbar^2 t} \exp \left[ \frac{im}{2\hbar t} (x^2 + y^2) - i\frac{\pi}{2} \right].$$

⊙ Reference: V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

### 2.3.2. Equations Containing Arbitrary Functions

$$1. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + f(t)w.$$

This equation describes two-dimensional thermal phenomena in quiescent media or solids with constant thermal diffusivities in the case of unsteady volume heat release proportional to temperature.

The substitution  $w(x, y, t) = \exp \left[ \int f(t) dt \right] U(x, y, t)$  leads to the two-dimensional heat equation  $\partial_t U = a(\partial_{xx} U + \partial_{yy} U)$  treated in Subsection 2.1.1.

$$2. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + [xf(t) + yg(t) + h(t)]w.$$

The transformation

$$w(x, y, t) = u(\xi, \eta, t) \exp \left[ xF(t) + yG(t) + H(t) + a \int F^2(t) dt + a \int G^2(t) dt \right],$$

$$\xi = x + 2a \int F(t) dt, \quad \eta = y + 2a \int G(t) dt,$$

where

$$F(t) = \int f(t) dt, \quad G(t) = \int g(t) dt, \quad H(t) = \int h(t) dt,$$

leads to the two-dimensional heat equation  $\partial_t u = a(\partial_{\xi\xi} u + \partial_{\eta\eta} u)$ .

$$3. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + [-b(x^2 + y^2) + f(t)x + g(t)y + h(t)]w.$$

1°. Case  $b > 0$ . The transformation

$$w(x, y, t) = u(\xi, \eta, \tau) \exp \left[ \frac{1}{2} \sqrt{\frac{b}{a}} (x^2 + y^2) \right],$$

$$\xi = x \exp(2\sqrt{ab}t), \quad \eta = y \exp(2\sqrt{ab}t), \quad \tau = \frac{1}{4\sqrt{ab}} \exp(4\sqrt{ab}t)$$

leads to an equation of the form 2.3.2.2:

$$\frac{\partial u}{\partial \tau} = a \left( \frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \eta^2} \right) + [F(\tau)\xi + G(\tau)\eta + H(\tau)]u,$$

$$F(\tau) = \frac{1}{(c\tau)^{3/2}} f \left( \frac{\ln(c\tau)}{c} \right), \quad G(\tau) = \frac{1}{(c\tau)^{3/2}} g \left( \frac{\ln(c\tau)}{c} \right), \quad H(\tau) = \frac{1}{c\tau} h \left( \frac{\ln(c\tau)}{c} \right) + \frac{1}{2\tau}, \quad c = 4\sqrt{ab}.$$

2°. Case  $b < 0$ . The transformation

$$w(x, y, t) = v(\bar{\xi}, \bar{\eta}, \bar{\tau}) \exp \left[ \frac{\sqrt{-b}}{2\sqrt{a}} \tan(2\sqrt{-ab}t) (x^2 + y^2) \right],$$

$$\bar{\xi} = \frac{x}{\cos(2\sqrt{-ab}t)}, \quad \bar{\eta} = \frac{y}{\cos(2\sqrt{-ab}t)}, \quad \bar{\tau} = \frac{1}{2\sqrt{-ab}} \tan(2\sqrt{-ab}t)$$

also leads to an equation of the form 2.3.2.2 (the transformed equation is not written out here).

$$4. \quad \frac{\partial w}{\partial t} = a_1(t) \frac{\partial^2 w}{\partial x^2} + a_2(t) \frac{\partial^2 w}{\partial y^2} + \Phi(x, y, t).$$

This is a special case of equation 2.3.2.8. Let  $0 < a_1(t) < \infty$  and  $0 < a_2(t) < \infty$ .

For the first, second, third, and mixed boundary value problems treated in rectangular, finite, or infinite domains ( $x_1 \leq x \leq x_2$ ,  $y_1 \leq y \leq y_2$ ), the Green's function can be represented in the product form

$$G(x, y, \xi, \eta, t, \tau) = G_1(x, \xi, T_1)G_2(y, \eta, T_2),$$

$$T_1 = \int_{\tau}^t a_1(\eta) d\eta, \quad T_2 = \int_{\tau}^t a_2(\eta) d\eta.$$

Here,  $G_1 = G_1(x, \xi, t)$  is the auxiliary Green's function that corresponds to the one-dimensional heat equation for  $a_1(t) = 1$ ,  $a_2(t) = 0$ , and  $\Phi(x, y, t) = 0$  with homogeneous boundary conditions at  $x = x_1$  and  $x = x_2$  (the  $G_1$ 's for various boundary value problems can be found in Subsections 1.1.1 and 1.1.2). Similarly,  $G_2 = G_2(y, \eta, t)$  is the auxiliary Green's function that corresponds to the one-dimensional heat equation for  $a_1(t) = 0$ ,  $a_2(t) = 1$ , and  $\Phi(x, y, t) = 0$  with homogeneous boundary conditions at  $y = y_1$  and  $y = y_2$ . Note that the Green's functions  $G_1$  and  $G_2$  are introduced for  $\tau = 0$ .

See Subsection 0.8.1 for solution of various boundary value problems with the help of the Green's function.

**Example 1.** Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(x, y) \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, y, t) = \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t, \tau) d\xi d\eta d\tau + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta) G(x, y, \xi, \eta, t, 0) d\xi d\eta,$$

where

$$G(x, y, \xi, \eta, t, \tau) = \frac{1}{4\pi\sqrt{T_1 T_2}} \exp\left[-\frac{(x-\xi)^2}{4T_1} - \frac{(y-\eta)^2}{4T_2}\right], \quad T_1 = \int_{\tau}^t a_1(\eta) d\eta, \quad T_2 = \int_{\tau}^t a_2(\eta) d\eta.$$

**Example 2.** Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_2(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^t \int_0^{\infty} \int_0^{\infty} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t, \tau) d\xi d\eta d\tau + \int_0^{\infty} \int_0^{\infty} f(\xi, \eta) G(x, y, \xi, \eta, t, 0) d\xi d\eta \\ &\quad - \int_0^t \int_0^{\infty} a_1(\tau) g_1(\eta, \tau) G(x, y, 0, \eta, t, \tau) d\eta d\tau - \int_0^t \int_0^{\infty} a_2(\tau) g_2(\xi, \tau) G(x, y, \xi, 0, t, \tau) d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t, \tau) &= G_1(x, \xi, T_1)G_2(y, \eta, T_2), \\ G_1(x, \xi, T_1) &= \frac{1}{2\sqrt{\pi T_1}} \left\{ \exp\left[-\frac{(x-\xi)^2}{4T_1}\right] + \exp\left[-\frac{(x+\xi)^2}{4T_1}\right] \right\}, \quad T_1 = \int_{\tau}^t a_1(\eta) d\eta, \\ G_2(y, \eta, T_2) &= \frac{1}{2\sqrt{\pi T_2}} \left\{ \exp\left[-\frac{(y-\eta)^2}{4T_2}\right] + \exp\left[-\frac{(y+\eta)^2}{4T_2}\right] \right\}, \quad T_2 = \int_{\tau}^t a_2(\eta) d\eta. \end{aligned}$$

**Example 3.** Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x &= 0 & (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x &= l_1 & (\text{boundary condition}), \\ w &= h_1(x, t) & \text{at } y &= 0 & (\text{boundary condition}), \\ w &= h_2(x, t) & \text{at } y &= l_2 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t, \tau) d\eta d\xi d\tau + \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t, 0) d\eta d\xi \\ &+ \int_0^t \int_0^{l_2} a_1(\tau) g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t, \tau) \right]_{\xi=0} d\eta d\tau - \int_0^t \int_0^{l_2} a_1(\tau) g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t, \tau) \right]_{\xi=l_1} d\eta d\tau \\ &+ \int_0^t \int_0^{l_1} a_2(\tau) h_1(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t, \tau) \right]_{\eta=0} d\xi d\tau - \int_0^t \int_0^{l_1} a_2(\tau) h_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t, \tau) \right]_{\eta=l_2} d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t, \tau) &= G_1(x, \xi, T_1) G_2(y, \eta, T_2), \\ G_1(x, \xi, T_1) &= \frac{2}{l_1} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi x}{l_1}\right) \sin\left(\frac{n\pi \xi}{l_1}\right) \exp\left(-\frac{n^2 \pi^2 T_1}{l_1^2}\right), \quad T_1 = \int_{\tau}^t a_1(\eta) d\eta, \\ G_2(y, \eta, T_2) &= \frac{2}{l_2} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi y}{l_2}\right) \sin\left(\frac{n\pi \eta}{l_2}\right) \exp\left(-\frac{n^2 \pi^2 T_2}{l_2^2}\right), \quad T_2 = \int_{\tau}^t a_2(\eta) d\eta. \end{aligned}$$

$$\begin{aligned} 5. \quad \frac{\partial w}{\partial t} &= a_1(t) \frac{\partial^2 w}{\partial x^2} + a_2(t) \frac{\partial^2 w}{\partial y^2} + [b_1(t)x + c_1(t)] \frac{\partial w}{\partial x} \\ &+ [b_2(t)y + c_2(t)] \frac{\partial w}{\partial y} + [s_1(t)x + s_2(t)y + p(t)]w. \end{aligned}$$

The transformation

$$w(x, y, t) = \exp[f_1(t)x + f_2(t)y + g(t)] u(\xi, \eta, t), \quad \xi = h_1(t)x + r_1(t), \quad \eta = h_2(t)y + r_2(t),$$

where

$$\begin{aligned} h_k(t) &= A_k \exp\left[\int b_k(t) dt\right], \\ f_k(t) &= h_k(t) \int \frac{s_k(t)}{h_k(t)} dt + B_k h_k(t), \\ r_k(t) &= \int [2a_k(t)f_k(t) + c_k(t)] h_k(t) dt + C_k, \\ g(t) &= \int [a_1(t)f_1^2(t) + a_2(t)f_2^2(t) + c_1(t)f_1(t) + c_2(t)f_2(t) + p(t)] dt + D, \end{aligned}$$

( $k = 1, 2$ ;  $A_k, B_k, C_k$ , and  $D$  are arbitrary constants), leads to an equation of the form 2.3.2.4:

$$\frac{\partial u}{\partial t} = a_1(t) h_1^2(t) \frac{\partial^2 u}{\partial \xi^2} + a_2(t) h_2^2(t) \frac{\partial^2 u}{\partial \eta^2}.$$

$$\begin{aligned} 6. \quad \frac{\partial w}{\partial t} &= a_1(t) \frac{\partial^2 w}{\partial x^2} + a_2(t) \frac{\partial^2 w}{\partial y^2} + [b_1(t)x + c_1(t)] \frac{\partial w}{\partial x} + [b_2(t)y + c_2(t)] \frac{\partial w}{\partial y} \\ &+ [s_1(t)x^2 + s_2(t)y^2 + p_1(t)x + p_2(t)y + q(t)]w. \end{aligned}$$

The substitution

$$w(x, y, t) = \exp[f_1(t)x^2 + f_2(t)y^2] u(x, y, t),$$

where the functions  $f_1 = f_1(t)$  and  $f_2 = f_2(t)$  are solutions of the Riccati equations

$$\begin{aligned} f_1' &= 4a_1(t)f_1^2 + 2b_1(t)f_1 + s_1(t), \\ f_2' &= 4a_2(t)f_2^2 + 2b_2(t)f_2 + s_2(t), \end{aligned}$$

leads to an equation of the form 2.3.2.5 for  $u = u(x, y, t)$ .

$$7. \quad \frac{\partial w}{\partial t} = a_1(x) \frac{\partial^2 w}{\partial x^2} + a_2(y) \frac{\partial^2 w}{\partial y^2} + b_1(x) \frac{\partial w}{\partial x} + b_2(y) \frac{\partial w}{\partial y} + [c_1(x) + c_2(y)]w + \Phi(x, y, t).$$

Domain:  $x_1 \leq x \leq x_2$ ,  $y_1 \leq y \leq y_2$ . Different boundary value problems:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ s_1 \partial_x w - k_1 w &= g_1(y, t) & \text{at } x &= x_1 & \text{(boundary condition),} \\ s_2 \partial_x w + k_2 w &= g_2(y, t) & \text{at } x &= x_2 & \text{(boundary condition),} \\ s_3 \partial_y w - k_3 w &= g_3(x, t) & \text{at } y &= y_1 & \text{(boundary condition),} \\ s_4 \partial_y w + k_4 w &= g_4(x, t) & \text{at } y &= y_2 & \text{(boundary condition).} \end{aligned}$$

By choosing appropriate parameters  $s_n$ ,  $k_n$  ( $n = 1, 2, 3, 4$ ), one obtains the first, second, third, or mixed boundary value problem. If the domain is infinite, say,  $x_2 = \infty$ , the corresponding boundary condition should be omitted; this is also valid for  $x_1 = -\infty$ ,  $y_1 = -\infty$ , or  $y_2 = \infty$ .

The Green's function admits incomplete separation of variables; specifically, it can be represented in the product form

$$G(x, y, \xi, \eta, t) = G_1(x, \xi, t)G_2(y, \eta, t).$$

Here,  $G_1 = G_1(x, \xi, t)$  and  $G_2 = G_2(y, \eta, t)$  are auxiliary Green's functions that are determined by solving the following simpler one-dimensional problems with homogeneous boundary conditions:

$$\begin{aligned} \frac{\partial G_1}{\partial t} &= a_1(x) \frac{\partial^2 G_1}{\partial x^2} + b_1(x) \frac{\partial G_1}{\partial x} + c_1(x)G_1, & \frac{\partial G_2}{\partial t} &= a_2(y) \frac{\partial^2 G_2}{\partial y^2} + b_2(y) \frac{\partial G_2}{\partial y} + c_2(y)G_2, \\ G_1 &= \delta(x - \xi) & \text{at } t &= 0, & G_2 &= \delta(y - \eta) & \text{at } t &= 0, \\ s_1 \partial_x G_1 - k_1 G_1 &= 0 & \text{at } x &= x_1, & s_3 \partial_y G_2 - k_3 G_2 &= 0 & \text{at } y &= y_1, \\ s_2 \partial_x G_1 + k_2 G_1 &= 0 & \text{at } x &= x_2, & s_4 \partial_y G_2 + k_4 G_2 &= 0 & \text{at } y &= y_2, \end{aligned}$$

where  $\xi$  and  $\eta$  are free parameters, and  $\delta(x)$  is the Dirac delta function.

The equation for  $G_1$  coincides with equation 1.8.6.5, which is reduced to the equation of Subsection 1.8.9 (where the expression of the Green's function can also be found). In the general case, the equation for  $G_2$  differs from the equation for  $G_1$  in only notation.

$$8. \quad \frac{\partial w}{\partial t} = a_1(x, t) \frac{\partial^2 w}{\partial x^2} + a_2(y, t) \frac{\partial^2 w}{\partial y^2} + b_1(x, t) \frac{\partial w}{\partial x} + b_2(y, t) \frac{\partial w}{\partial y} + [c_1(x, t) + c_2(y, t)]w + \Phi(x, y, t).$$

Suppose this equation is subject to the same initial and boundary conditions as equation 2.3.2.7. Then the Green's function for this problem can be represented in the product form

$$G(x, y, \xi, \eta, t, \tau) = G_1(x, \xi, t, \tau)G_2(y, \eta, t, \tau).$$

Here,  $G_1 = G_1(x, \xi, t, \tau)$  and  $G_2 = G_2(y, \eta, t, \tau)$  are auxiliary Green's functions that are determined by solving the following simpler boundary value problems with homogeneous boundary conditions:

$$\begin{aligned} \frac{\partial G_1}{\partial t} &= a_1(x, t) \frac{\partial^2 G_1}{\partial x^2} + b_1(x, t) \frac{\partial G_1}{\partial x} + c_1(x, t)G_1, & \frac{\partial G_2}{\partial t} &= a_2(y, t) \frac{\partial^2 G_2}{\partial y^2} + b_2(y, t) \frac{\partial G_2}{\partial y} + c_2(y, t)G_2, \\ G_1 &= \delta(x - \xi) & \text{at } t &= \tau, & G_2 &= \delta(y - \eta) & \text{at } t &= \tau, \\ s_1 \partial_x G_1 - k_1 G_1 &= 0 & \text{at } x &= x_1, & s_3 \partial_y G_2 - k_3 G_2 &= 0 & \text{at } y &= y_1, \\ s_2 \partial_x G_1 + k_2 G_1 &= 0 & \text{at } x &= x_2, & s_4 \partial_y G_2 + k_4 G_2 &= 0 & \text{at } y &= y_2, \end{aligned}$$

where  $\xi$ ,  $\eta$ , and  $\tau$  are free parameters, and  $\delta(x)$  is the Dirac delta function,  $t \geq \tau$ .

See Subsection 0.8.1 for the solution of boundary value problems with the help of the Green's function.

## Chapter 3

# Parabolic Equations with Three or More Space Variables

---

### 3.1. Heat Equation $\frac{\partial w}{\partial t} = a \Delta_3 w$

#### 3.1.1. Problems in Cartesian Coordinates

The three-dimensional sourceless heat equation in the rectangular Cartesian system of coordinates has the form

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right).$$

It governs three-dimensional thermal phenomena in quiescent media or solids with constant thermal diffusivity. A similar equation is used to study the corresponding three-dimensional unsteady mass-exchange processes with constant diffusivity.

##### 3.1.1-1. Particular solutions:

$$w(x, y, z, t) = Ax^2 + By^2 + Cz^2 + 2a(A + B + C)t,$$

$$w(x, y, z, t) = A(x^2 + 2at)(y^2 + 2at)(z^2 + 2at) + B,$$

$$w(x, y, z, t) = A \exp[k_1 x + k_2 y + k_3 z + (k_1^2 + k_2^2 + k_3^2)at] + B,$$

$$w(x, y, z, t) = A \cos(k_1 x + C_1) \cos(k_2 y + C_2) \cos(k_3 z + C_3) \exp[-(k_1^2 + k_2^2 + k_3^2)at],$$

$$w(x, y, z, t) = A \cos(k_1 x + C_1) \cos(k_2 y + C_2) \sinh(k_3 z + C_3) \exp[-(k_1^2 + k_2^2 - k_3^2)at],$$

$$w(x, y, z, t) = A \cos(k_1 x + C_1) \cos(k_2 y + C_2) \cosh(k_3 z + C_3) \exp[-(k_1^2 + k_2^2 - k_3^2)at],$$

$$w(x, y, z, t) = A \exp(-k_1 x - k_2 y - k_3 z) \cos(k_1 x - 2ak_1^2 t) \cos(k_2 y - 2ak_2^2 t) \cos(k_3 z - 2ak_3^2 t),$$

$$w(x, y, z, t) = \frac{A}{(t - t_0)^{3/2}} \exp \left[ -\frac{(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2}{4a(t - t_0)} \right],$$

$$w(x, y, z, t) = A \operatorname{erf} \left( \frac{x - x_0}{2\sqrt{at}} \right) \operatorname{erf} \left( \frac{y - y_0}{2\sqrt{at}} \right) \operatorname{erf} \left( \frac{z - z_0}{2\sqrt{at}} \right) + B,$$

where  $A, B, C, C_1, C_2, C_3, k_1, k_2, k_3, x_0, y_0, z_0$ , and  $t_0$  are arbitrary constants.

Fundamental solution:

$$\mathcal{E}(x, y, z, t) = \frac{1}{8(\pi at)^{3/2}} \exp \left( -\frac{x^2 + y^2 + z^2}{4at} \right).$$

##### 3.1.1-2. Formulas to construct particular solutions. Remarks on the Green's functions.

1°. Apart from usual solutions with separated variables,

$$w(x, y, z, t) = f_1(x)f_2(y)f_3(z)f_4(t),$$

the equation in question admits more sophisticated solutions in the product form

$$w(x, y, z, t) = u_1(x, t)u_2(y, t)u_3(z, t),$$

where the functions  $u_1 = u_1(x, t)$ ,  $u_2 = u_2(y, t)$ , and  $u_3 = u_3(z, t)$  are solutions of the one-dimensional heat equations

$$\frac{\partial u_1}{\partial t} = a \frac{\partial^2 u_1}{\partial x^2}, \quad \frac{\partial u_2}{\partial t} = a \frac{\partial^2 u_2}{\partial y^2}, \quad \frac{\partial u_3}{\partial t} = a \frac{\partial^2 u_3}{\partial z^2},$$

treated in Subsection 1.1.1.

2°. Suppose  $w = w(x, y, z, t)$  is a solution of the three-dimensional heat equation. Then the functions

$$\begin{aligned} w_1 &= Aw(\pm\lambda x + C_1, \pm\lambda y + C_2, \pm\lambda z + C_3, \lambda^2 t + C_4), \\ w_2 &= A \exp[\lambda_1 x + \lambda_2 y + \lambda_3 z + (\lambda_1^2 + \lambda_2^2 + \lambda_3^2)at] w(x + 2a\lambda_1 t, y + 2a\lambda_2 t, z + 2a\lambda_3 t, t), \\ w_3 &= \frac{A}{|\delta + \beta t|^{3/2}} \exp\left[-\frac{\beta(x^2 + y^2 + z^2)}{4a(\delta + \beta t)}\right] w\left(\frac{x}{\delta + \beta t}, \frac{y}{\delta + \beta t}, \frac{z}{\delta + \beta t}, \frac{\gamma + \lambda t}{\delta + \beta t}\right), \quad \lambda\delta - \beta\gamma = 1, \end{aligned}$$

where  $A$ ,  $C_1$ ,  $C_2$ ,  $C_3$ ,  $C_4$ ,  $\lambda$ ,  $\lambda_1$ ,  $\lambda_2$ ,  $\lambda_3$ ,  $\beta$ , and  $\delta$  are arbitrary constants, are also solutions of this equation. The signs at  $\lambda$  in the formula for  $w_1$  can be taken independently of one another.

3°. For the three-dimensional boundary value problems considered in Subsection 3.1.1, the Green's function can be represented in the product form

$$G(x, y, z, \xi, \eta, \zeta, t) = G_1(x, \xi, t)G_2(y, \eta, t)G_3(z, \zeta, t),$$

where  $G_1(x, \xi, t)$ ,  $G_2(y, \eta, t)$ ,  $G_3(z, \zeta, t)$  are the Green's functions of the corresponding one-dimensional boundary value problems; these functions can be found in Subsections 1.1.1 and 1.1.2.

**Example 1.** The Green's function of the mixed boundary value problem for a semiinfinite layer ( $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < l$ ) presented in Paragraph 3.1.1-14 is the product of three one-dimensional Green's functions from Paragraph 1.1.2-1 (Cauchy problem for  $-\infty < x < \infty$ ), Paragraph 1.1.2-2 (first boundary value problem for  $0 \leq y < \infty$ ), and Paragraph 1.1.2-6 (second boundary value problem for  $0 \leq z < l$ ), in which one needs to carry out obvious renaming of variables.

3.1.1-3. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(x, y, z) \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, y, z, t) = \frac{1}{8(\pi at)^{3/2}} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) \exp\left[-\frac{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}{4at}\right] d\xi d\eta d\zeta.$$

**Example 2.** The initial temperature is constant and is equal to  $w_1$  in the domain  $|x| < x_0$ ,  $|y| < y_0$ ,  $|z| < z_0$  and is equal to  $w_2$  in the domain  $|x| > x_0$ ,  $|y| > y_0$ ,  $|z| > z_0$ ; specifically,

$$f(x, y, z) = \begin{cases} w_1 & \text{for } |x| < x_0, |y| < y_0, |z| < z_0, \\ w_2 & \text{for } |x| > x_0, |y| > y_0, |z| > z_0. \end{cases}$$

Solution:

$$\begin{aligned} w &= \frac{1}{8}(w_1 - w_2) \left[ \operatorname{erf}\left(\frac{x_0 - x}{2\sqrt{at}}\right) + \operatorname{erf}\left(\frac{x_0 + x}{2\sqrt{at}}\right) \right] \\ &\quad \times \left[ \operatorname{erf}\left(\frac{y_0 - y}{2\sqrt{at}}\right) + \operatorname{erf}\left(\frac{y_0 + y}{2\sqrt{at}}\right) \right] \left[ \operatorname{erf}\left(\frac{z_0 - z}{2\sqrt{at}}\right) + \operatorname{erf}\left(\frac{z_0 + z}{2\sqrt{at}}\right) \right] + w_2. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).



**3.1.1-4. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . First boundary value problem.**

A half-space is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_0^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{8(\pi at)^{3/2}} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] - \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right\} \exp \left[ -\frac{(y-\eta)^2 + (z-\zeta)^2}{4at} \right].$$

⊙ *References:* A. G. Butkovskiy (1979), H. S. Carslaw and J. C. Jaeger (1984).

**3.1.1-5. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Second boundary value problem.**

A half-space is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_0^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{8(\pi at)^{3/2}} \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] + \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right\} \exp \left[ -\frac{(y-\eta)^2 + (z-\zeta)^2}{4at} \right].$$

⊙ *Reference:* A. G. Butkovskiy (1979).

**3.1.1-6. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Third boundary value problem.**

A half-space is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - kw &= g(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 3.1.1-5 where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{8(\pi at)^{3/2}} \exp \left[ -\frac{(y-\eta)^2 + (z-\zeta)^2}{4at} \right] \left\{ \exp \left[ -\frac{(x-\xi)^2}{4at} \right] + \exp \left[ -\frac{(x+\xi)^2}{4at} \right] \right. \\ &\quad \left. - 2k\sqrt{\pi at} \exp[k^2 at + k(x+\xi)] \operatorname{erfc} \left( \frac{x+\xi}{2\sqrt{at}} + k\sqrt{at} \right) \right\}. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**3.1.1-7. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq l$ . First boundary value problem.**

An infinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_2(x, y, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^l \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g_1(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{2\pi a l t} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] \sum_{n=1}^{\infty} \sin \frac{n\pi z}{l} \sin \frac{n\pi \zeta}{l} \exp \left( -\frac{n^2 \pi^2 a t}{l^2} \right),$$

or

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{8(\pi a t)^{3/2}} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] \\ &\times \sum_{n=-\infty}^{\infty} \left\{ \exp \left[ -\frac{(2nl + z - \zeta)^2}{4at} \right] - \exp \left[ -\frac{(2nl + z + \zeta)^2}{4at} \right] \right\}. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**3.1.1-8. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

An infinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_z w &= g_1(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_2(x, y, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^l \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g_1(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g_2(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{4\pi a l t} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] \\ &\times \left[ 1 + 2 \sum_{n=1}^{\infty} \cos \frac{n\pi z}{l} \cos \frac{n\pi \zeta}{l} \exp \left( -\frac{n^2 \pi^2 a t}{l^2} \right) \right], \end{aligned}$$

or

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{(2\sqrt{\pi at})^3} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] \\ \times \sum_{n=-\infty}^{\infty} \left\{ \exp \left[ -\frac{(z - \zeta + 2nl)^2}{4at} \right] + \exp \left[ -\frac{(z + \zeta + 2nl)^2}{4at} \right] \right\}.$$

⊙ Reference: H. S. Carslaw and J. C. Jaeger (1984).

3.1.1-9. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq l$ . Third boundary value problem.

An infinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_z w - k_1 w &= g_1(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w + k_2 w &= g_2(x, y, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 3.1.1-8 where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{4\pi at} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] \sum_{n=1}^{\infty} \frac{\varphi_n(z)\varphi_n(\zeta)}{\|\varphi_n\|^2} \exp(-a\mu_n^2 t), \\ \varphi_n(z) = \cos(\mu_n z) + \frac{k_1}{\mu_n} \sin(\mu_n z), \quad \|\varphi_n\|^2 = \frac{k_2}{2\mu_n^2} \frac{\mu_n^2 + k_1^2}{\mu_n^2 + k_2^2} + \frac{k_1}{2\mu_n^2} + \frac{l}{2} \left( 1 + \frac{k_1^2}{\mu_n^2} \right).$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\frac{\tan(\mu l)}{\mu} = \frac{k_1 + k_2}{\mu^2 - k_1 k_2}$ .

3.1.1-10. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq l$ . Mixed boundary value problem.

An infinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_2(x, y, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^l \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g_1(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g_2(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{2\pi alt} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] \\ \times \sum_{n=0}^{\infty} \sin \left[ \frac{(2n+1)\pi z}{2l} \right] \sin \left[ \frac{(2n+1)\pi \zeta}{2l} \right] \exp \left[ -\frac{(2n+1)^2 \pi^2 at}{4l^2} \right],$$

or

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{(2\sqrt{\pi at})^3} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2}{4at} \right] \\ \times \sum_{n=-\infty}^{\infty} (-1)^n \left\{ \exp \left[ -\frac{(z - \zeta + 2nl)^2}{4at} \right] - \exp \left[ -\frac{(z + \zeta + 2nl)^2}{4at} \right] \right\}.$$

⊙ Reference: A. G. Butkovskiy (1979).

3.1.1-11. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z \leq l$ . First boundary value problem.

A semiinfinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \text{(initial condition),} \\ w &= g_1(x, z, t) & \text{at } y = 0 & \text{(boundary condition),} \\ w &= g_2(x, y, t) & \text{at } z = 0 & \text{(boundary condition),} \\ w &= g_3(x, y, t) & \text{at } z = l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^l \int_0^\infty \int_{-\infty}^\infty f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^l \int_{-\infty}^\infty g_1(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &+ a \int_0^t \int_0^\infty \int_{-\infty}^\infty g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a \int_0^t \int_0^\infty \int_{-\infty}^\infty g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{2\pi a l t} \exp \left[ -\frac{(x - \xi)^2}{4at} \right] \left\{ \exp \left[ -\frac{(y - \eta)^2}{4at} \right] - \exp \left[ -\frac{(y + \eta)^2}{4at} \right] \right\} \\ &\times \sum_{n=1}^\infty \sin \frac{n\pi z}{l} \sin \frac{n\pi \zeta}{l} \exp \left( -\frac{n^2 \pi^2 a t}{l^2} \right). \end{aligned}$$

3.1.1-12. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A semiinfinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \text{(initial condition),} \\ \partial_y w &= g_1(x, z, t) & \text{at } y = 0 & \text{(boundary condition),} \\ \partial_z w &= g_2(x, y, t) & \text{at } z = 0 & \text{(boundary condition),} \\ \partial_z w &= g_3(x, y, t) & \text{at } z = l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^l \int_0^\infty \int_{-\infty}^\infty f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a \int_0^t \int_0^l \int_{-\infty}^\infty g_1(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^\infty \int_{-\infty}^\infty g_2(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a \int_0^t \int_0^\infty \int_{-\infty}^\infty g_3(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{4\pi a l t} \exp \left[ -\frac{(x - \xi)^2}{4at} \right] \left\{ \exp \left[ -\frac{(y - \eta)^2}{4at} \right] + \exp \left[ -\frac{(y + \eta)^2}{4at} \right] \right\} \\ &\times \left[ 1 + 2 \sum_{n=1}^\infty \cos \frac{n\pi z}{l} \cos \frac{n\pi \zeta}{l} \exp \left( -\frac{n^2 \pi^2 a t}{l^2} \right) \right]. \end{aligned}$$

**3.1.1-13. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z \leq l$ . Third boundary value problem.**

A semiinfinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_y w - k_1 w &= g_1(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_z w - k_2 w &= g_2(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w + k_3 w &= g_3(x, y, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 3.1.1-12 where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{4\pi at} \exp\left[-\frac{(x-\xi)^2}{4at}\right] H(y, \eta, t) \sum_{n=1}^{\infty} \frac{\varphi_n(z)\varphi_n(\zeta)}{\|\varphi_n\|^2} \exp(-a\mu_n^2 t), \\ H(y, \eta, t) &= \exp\left[-\frac{(y-\eta)^2}{4at}\right] + \exp\left[-\frac{(y+\eta)^2}{4at}\right] - 2k_1 \int_0^{\infty} \exp\left[-\frac{(y+\eta+s)^2}{4at} - k_1 s\right] ds. \end{aligned}$$

Here,

$$\varphi_n(z) = \cos(\mu_n z) + \frac{k_2}{\mu_n} \sin(\mu_n z), \quad \|\varphi_n\|^2 = \frac{k_3}{2\mu_n^2} \frac{\mu_n^2 + k_2^2}{\mu_n^2 + k_3^2} + \frac{k_2}{2\mu_n^2} + \frac{l}{2} \left(1 + \frac{k_2^2}{\mu_n^2}\right);$$

the  $\mu_n$  are positive roots of the transcendental equation  $\frac{\tan(\mu l)}{\mu} = \frac{k_2 + k_3}{\mu^2 - k_2 k_3}$ .

**3.1.1-14. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.**

1°. A semiinfinite layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_2(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(x, y, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^l \int_0^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^l \int_{-\infty}^{\infty} g_1(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t-\tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^{\infty} \int_{-\infty}^{\infty} g_2(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t-\tau) d\xi d\eta d\tau \\ &+ a \int_0^t \int_0^{\infty} \int_{-\infty}^{\infty} g_3(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l, t-\tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{4\pi a l t} \exp\left[-\frac{(x-\xi)^2}{4at}\right] \left\{ \exp\left[-\frac{(y-\eta)^2}{4at}\right] - \exp\left[-\frac{(y+\eta)^2}{4at}\right] \right\} \\ &\times \left[ 1 + 2 \sum_{n=1}^{\infty} \cos \frac{n\pi z}{l} \cos \frac{n\pi \zeta}{l} \exp\left(-\frac{n^2 \pi^2 a t}{l^2}\right) \right]. \end{aligned}$$

2°. A semiinfinite layer is considered. The following conditions are prescribed:

$$\begin{aligned}w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_y w &= g_1(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ w &= g_2(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_3(x, y, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}).\end{aligned}$$

Solution:

$$\begin{aligned}w(x, y, z, t) &= \int_0^l \int_0^\infty \int_{-\infty}^\infty f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a \int_0^t \int_0^l \int_{-\infty}^\infty g_1(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a \int_0^t \int_0^\infty \int_{-\infty}^\infty g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a \int_0^t \int_0^\infty \int_{-\infty}^\infty g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau,\end{aligned}$$

where

$$\begin{aligned}G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{2\pi a l t} \exp\left[-\frac{(x - \xi)^2}{4at}\right] \left\{ \exp\left[-\frac{(y - \eta)^2}{4at}\right] + \exp\left[-\frac{(y + \eta)^2}{4at}\right] \right\} \\ &\times \sum_{n=1}^\infty \sin \frac{n\pi z}{l} \sin \frac{n\pi \zeta}{l} \exp\left(-\frac{n^2 \pi^2 a t}{l^2}\right).\end{aligned}$$

3.1.1-15. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . First boundary value problem.

An octant is considered. The following conditions are prescribed:

$$\begin{aligned}w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ w &= g_3(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}).\end{aligned}$$

Solution:

$$\begin{aligned}w(x, y, z, t) &= \int_0^\infty \int_0^\infty \int_0^\infty G(x, y, z, \xi, \eta, \zeta, t) f(\xi, \eta, \zeta) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^\infty \int_0^\infty g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^\infty \int_0^\infty g_2(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &+ a \int_0^t \int_0^\infty \int_0^\infty g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau,\end{aligned}$$

where

$$\begin{aligned}G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{(2\sqrt{\pi a t})^3} H(x, \xi, t) H(y, \eta, t) H(z, \zeta, t), \\ H(x, \xi, t) &= \exp\left[-\frac{(x - \xi)^2}{4at}\right] - \exp\left[-\frac{(x + \xi)^2}{4at}\right].\end{aligned}$$

**Example 3.** The initial temperature is uniform,  $f(x, y, z) = w_0$ . The faces are maintained at zero temperature,  $g_1 = g_2 = g_3 = 0$ .  
Solution:

$$w = w_0 \operatorname{erf}\left(\frac{x}{2\sqrt{at}}\right) \operatorname{erf}\left(\frac{y}{2\sqrt{at}}\right) \operatorname{erf}\left(\frac{z}{2\sqrt{at}}\right).$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

3.1.1-16. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . Second boundary value problem.

An octant is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \text{(initial condition),} \\ \partial_x w &= g_1(y, z, t) & \text{at } x = 0 & \text{(boundary condition),} \\ \partial_y w &= g_2(x, z, t) & \text{at } y = 0 & \text{(boundary condition),} \\ \partial_z w &= g_3(x, y, t) & \text{at } z = 0 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^\infty \int_0^\infty \int_0^\infty G(x, y, z, \xi, \eta, \zeta, t) f(\xi, \eta, \zeta) d\xi d\eta d\zeta \\ &\quad - a \int_0^t \int_0^\infty \int_0^\infty g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &\quad - a \int_0^t \int_0^\infty \int_0^\infty g_2(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &\quad - a \int_0^t \int_0^\infty \int_0^\infty g_3(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{(2\sqrt{\pi at})^3} H(x, \xi, t) H(y, \eta, t) H(z, \zeta, t), \\ H(x, \xi, t) &= \exp\left[-\frac{(x - \xi)^2}{4at}\right] + \exp\left[-\frac{(x + \xi)^2}{4at}\right]. \end{aligned}$$

3.1.1-17. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . Third boundary value problem.

An octant is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \text{(initial condition),} \\ \partial_x w - k_1 w &= g_1(y, z, t) & \text{at } x = 0 & \text{(boundary condition),} \\ \partial_y w - k_2 w &= g_2(x, z, t) & \text{at } y = 0 & \text{(boundary condition),} \\ \partial_z w - k_3 w &= g_3(x, y, t) & \text{at } z = 0 & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 3.1.1-16 where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{(2\sqrt{\pi at})^3} H(x, \xi, t; k_1) H(y, \eta, t; k_2) H(z, \zeta, t; k_3), \\ H(x, \xi, t; k) &= \exp\left[-\frac{(x - \xi)^2}{4at}\right] + \exp\left[-\frac{(x + \xi)^2}{4at}\right] \\ &\quad - 2k\sqrt{\pi at} \exp[ak^2 t + k(x + \xi)] \operatorname{erfc}\left(\frac{x + \xi}{2\sqrt{at}} + k\sqrt{at}\right). \end{aligned}$$

**Example 4.** The initial temperature is uniform,  $f(x, y, z) = w_0$ . The temperature of the contacting media is zero,  $g_1 = g_2 = g_3 = 0$ .  
Solution:

$$w = w_0 \left[ \operatorname{erf} \left( \frac{x}{2\sqrt{at}} \right) + \exp(k_1 x + k_1^2 at) \operatorname{erfc} \left( \frac{x}{2\sqrt{at}} + k_1 \sqrt{at} \right) \right] \\ \times \left[ \operatorname{erf} \left( \frac{y}{2\sqrt{at}} \right) + \exp(k_2 y + k_2^2 at) \operatorname{erfc} \left( \frac{y}{2\sqrt{at}} + k_2 \sqrt{at} \right) \right] \\ \times \left[ \operatorname{erf} \left( \frac{z}{2\sqrt{at}} \right) + \exp(k_3 z + k_3^2 at) \operatorname{erfc} \left( \frac{z}{2\sqrt{at}} + k_3 \sqrt{at} \right) \right].$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

3.1.1-18. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . Mixed boundary value problems.

1°. An octant is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_2(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^\infty \int_0^\infty \int_0^\infty G(x, y, z, \xi, \eta, \zeta, t) f(\xi, \eta, \zeta) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^\infty \int_0^\infty g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a \int_0^t \int_0^\infty \int_0^\infty g_2(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^\infty \int_0^\infty g_3(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{(2\sqrt{\pi at})^3} \left\{ \exp \left[ -\frac{(x - \xi)^2}{4at} \right] - \exp \left[ -\frac{(x + \xi)^2}{4at} \right] \right\} H(y, \eta, t) H(z, \zeta, t), \\ H(y, \eta, t) = \exp \left[ -\frac{(y - \eta)^2}{4at} \right] + \exp \left[ -\frac{(y + \eta)^2}{4at} \right].$$

2°. An octant is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^\infty \int_0^\infty \int_0^\infty G(x, y, z, \xi, \eta, \zeta, t) f(\xi, \eta, \zeta) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^\infty \int_0^\infty g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^\infty \int_0^\infty g_2(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^\infty \int_0^\infty g_3(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau, \end{aligned}$$



where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{(2\sqrt{\pi at})^3} H(x, \xi, t) H(y, \eta, t) \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] + \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\},$$

$$H(x, \xi, t) = \exp \left[ -\frac{(x - \xi)^2}{4at} \right] - \exp \left[ -\frac{(x + \xi)^2}{4at} \right].$$

3.1.1-19. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $-\infty < z < \infty$ . First boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_{-\infty}^{\infty} \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] H_1(x, \xi, t) H_2(y, \eta, t), \\ H_1(x, \xi, t) &= \frac{2}{l_1} \sum_{n=1}^{\infty} \sin \left( \frac{\pi n x}{l_1} \right) \sin \left( \frac{\pi n \xi}{l_1} \right) \exp \left( -\frac{\pi^2 n^2 at}{l_1^2} \right), \\ H_2(y, \eta, t) &= \frac{2}{l_2} \sum_{n=1}^{\infty} \sin \left( \frac{\pi n y}{l_2} \right) \sin \left( \frac{\pi n \eta}{l_2} \right) \exp \left( -\frac{\pi^2 n^2 at}{l_2^2} \right). \end{aligned}$$

3.1.1-20. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $-\infty < z < \infty$ . Second boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) = & \int_{-\infty}^{\infty} \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & - a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_2} g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & - a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \exp\left[-\frac{(z - \zeta)^2}{4at}\right] H_1(x, \xi, t) H_2(y, \eta, t), \\
 H_1(x, \xi, t) &= \frac{1}{l_1} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos\left(\frac{\pi n x}{l_1}\right) \cos\left(\frac{\pi n \xi}{l_1}\right) \exp\left(-\frac{\pi^2 n^2 at}{l_1^2}\right) \right], \\
 H_2(y, \eta, t) &= \frac{1}{l_2} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos\left(\frac{\pi n y}{l_2}\right) \cos\left(\frac{\pi n \eta}{l_2}\right) \exp\left(-\frac{\pi^2 n^2 at}{l_2^2}\right) \right].
 \end{aligned}$$

3.1.1-21. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $-\infty < z < \infty$ . Third boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_x w - k_1 w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\
 \partial_x w + k_2 w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\
 \partial_y w - k_3 w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\
 \partial_y w + k_4 w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 3.1.1-20 where

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \exp\left[-\frac{(z - \zeta)^2}{4at}\right] H_1(x, \xi, t) H_2(y, \eta, t), \\
 H_1(x, \xi, t) &= \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\|\varphi_n\|^2} \exp(-a\mu_n^2 t), \quad H_2(y, \eta, t) = \sum_{m=1}^{\infty} \frac{\psi_m(y) \psi_m(\eta)}{\|\psi_m\|^2} \exp(-a\lambda_m^2 t).
 \end{aligned}$$

Here,

$$\begin{aligned}
 \varphi_n(x) &= \cos(\mu_n x) + \frac{k_1}{\mu_n} \sin(\mu_n x), \quad \|\varphi_n\|^2 = \frac{k_2}{2\mu_n^2} \frac{\mu_n^2 + k_1^2}{\mu_n^2 + k_2^2} + \frac{k_1}{2\mu_n^2} + \frac{l_1}{2} \left( 1 + \frac{k_1^2}{\mu_n^2} \right), \\
 \psi_m(y) &= \cos(\lambda_m y) + \frac{k_3}{\lambda_m} \sin(\lambda_m y), \quad \|\psi_m\|^2 = \frac{k_4}{2\lambda_m^2} \frac{\lambda_m^2 + k_3^2}{\lambda_m^2 + k_4^2} + \frac{k_3}{2\lambda_m^2} + \frac{l_2}{2} \left( 1 + \frac{k_3^2}{\lambda_m^2} \right);
 \end{aligned}$$

the  $\mu_n$  and  $\lambda_m$  are positive roots of the transcendental equations

$$\frac{\tan(\mu l_1)}{\mu} = \frac{k_1 + k_2}{\mu^2 - k_1 k_2}, \quad \frac{\tan(\lambda l_2)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}.$$

3.1.1-22. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $-\infty < z < \infty$ . Mixed boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_{-\infty}^{\infty} \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ &- a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a \int_0^t \int_{-\infty}^{\infty} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{2}{l_1 l_2 \sqrt{\pi a t}} \exp \left[ -\frac{(z - \zeta)^2}{4 a t} \right] \left[ \sum_{n=1}^{\infty} \sin \left( \frac{\pi n x}{l_1} \right) \sin \left( \frac{\pi n \xi}{l_1} \right) \exp \left( -\frac{\pi^2 n^2 a t}{l_1^2} \right) \right] \\ &\times \left[ \frac{1}{2} + \sum_{m=1}^{\infty} \cos \left( \frac{\pi m x}{l_2} \right) \cos \left( \frac{\pi m \eta}{l_2} \right) \exp \left( -\frac{\pi^2 m^2 a t}{l_2^2} \right) \right]. \end{aligned}$$

3.1.1-23. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z < \infty$ . First boundary value problem.

A semiinfinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\ w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) = & \int_0^\infty \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + a \int_0^t \int_0^\infty \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\
 & - a \int_0^t \int_0^\infty \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^\infty \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^\infty \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta, t) &= G_1(x, \xi, t; l_1) G_1(y, \eta, t; l_2) G_2(z, \zeta, t), \\
 G_1(x, \xi, t; l) &= \frac{2}{l} \sum_{n=1}^\infty \sin\left(\frac{\pi n x}{l}\right) \sin\left(\frac{\pi n \xi}{l}\right) \exp\left(-\frac{\pi^2 n^2 a t}{l^2}\right), \\
 G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \left\{ \exp\left[-\frac{(z - \zeta)^2}{4 a t}\right] - \exp\left[-\frac{(z + \zeta)^2}{4 a t}\right] \right\}.
 \end{aligned}$$

3.1.1-24. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z < \infty$ . Second boundary value problem.

A semiinfinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_x w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\
 \partial_x w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\
 \partial_y w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\
 \partial_y w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\
 \partial_z w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) = & \int_0^\infty \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & - a \int_0^t \int_0^\infty \int_0^{l_2} g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^\infty \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & - a \int_0^t \int_0^\infty \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^\infty \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau,
 \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] + \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} G_1(x, \xi, t) G_2(y, \eta, t),$$

$$G_1(x, \xi, t) = \frac{1}{l_1} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos \left( \frac{\pi n x}{l_1} \right) \cos \left( \frac{\pi n \xi}{l_1} \right) \exp \left( -\frac{\pi^2 n^2 at}{l_1^2} \right) \right],$$

$$G_2(y, \eta, t) = \frac{1}{l_2} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos \left( \frac{\pi n y}{l_2} \right) \cos \left( \frac{\pi n \eta}{l_2} \right) \exp \left( -\frac{\pi^2 n^2 at}{l_2^2} \right) \right].$$

3.1.1-25. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z < \infty$ . Third boundary value problem.

A semiinfinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w - k_3 w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w + k_4 w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ \partial_z w - k_5 w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 3.1.1-24 where

$$G(x, y, z, \xi, \eta, \zeta, t) = H_1(x, \xi, t) H_2(y, \eta, t) H_3(z, \zeta, t),$$

$$H_3(z, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] + \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} \\ - k_5 \exp[k_5^2 at + k_5(z + \zeta)] \operatorname{erfc} \left( \frac{z + \zeta}{2\sqrt{at}} + k_5 \sqrt{at} \right),$$

and the functions  $H_1(x, \xi, t)$  and  $H_2(y, \eta, t)$  can be found in Paragraph 3.1.1-21.

3.1.1-26. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z < \infty$ . Mixed boundary value problems.

1°. A semiinfinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) = & \int_0^\infty \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + a \int_0^t \int_0^\infty \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\
 & - a \int_0^t \int_0^\infty \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^\infty \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^\infty \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] + \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} H(x, \xi, t; l_1) H(y, \eta, t; l_2), \\
 H(x, \xi, t; l) &= \frac{2}{l} \sum_{n=1}^\infty \sin \left( \frac{\pi n x}{l} \right) \sin \left( \frac{\pi n \xi}{l} \right) \exp \left( -\frac{\pi^2 n^2 at}{l^2} \right).
 \end{aligned}$$

2°. A semiinfinite cylindrical domain of a rectangular cross-section is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_x w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\
 \partial_x w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\
 \partial_y w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\
 \partial_y w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\
 w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) = & \int_0^\infty \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & - a \int_0^t \int_0^\infty \int_0^{l_2} g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^\infty \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & - a \int_0^t \int_0^\infty \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^\infty \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau,
 \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] - \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} H(x, \xi, t; l_1) H(y, \eta, t; l_2),$$

$$H(x, \xi, t; l) = \frac{1}{l} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos \left( \frac{\pi n x}{l} \right) \cos \left( \frac{\pi n \xi}{l} \right) \exp \left( -\frac{\pi^2 n^2 at}{l^2} \right) \right].$$

3.1.1-27. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . First boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(y, z, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(y, z, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ w &= g_3(x, z, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ w &= g_4(x, z, t) & \text{at } y &= l_2 & \text{(boundary condition),} \\ w &= g_5(x, y, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_6(x, y, t) & \text{at } z &= l_3 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau \\ &+ a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l_3} d\xi d\eta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= G_1(x, \xi, t) G_2(y, \eta, t) G_3(z, \zeta, t), \\ G_1(x, \xi, t) &= \frac{2}{l_1} \sum_{n=1}^{\infty} \sin \left( \frac{\pi n x}{l_1} \right) \sin \left( \frac{\pi n \xi}{l_1} \right) \exp \left( -\frac{\pi^2 n^2 at}{l_1^2} \right), \\ G_2(y, \eta, t) &= \frac{2}{l_2} \sum_{n=1}^{\infty} \sin \left( \frac{\pi n y}{l_2} \right) \sin \left( \frac{\pi n \eta}{l_2} \right) \exp \left( -\frac{\pi^2 n^2 at}{l_2^2} \right), \\ G_3(z, \zeta, t) &= \frac{2}{l_3} \sum_{n=1}^{\infty} \sin \left( \frac{\pi n z}{l_3} \right) \sin \left( \frac{\pi n \zeta}{l_3} \right) \exp \left( -\frac{\pi^2 n^2 at}{l_3^2} \right). \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**3.1.1-28. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Second boundary value problem.**

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &\quad - a \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &\quad + a \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &\quad + a \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &\quad + a \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= G_1(x, \xi, t) G_2(y, \eta, t) G_3(z, \zeta, t), \\ G_1(x, \xi, t) &= \frac{1}{l_1} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos\left(\frac{\pi n x}{l_1}\right) \cos\left(\frac{\pi n \xi}{l_1}\right) \exp\left(-\frac{\pi^2 n^2 a t}{l_1^2}\right) \right], \\ G_2(y, \eta, t) &= \frac{1}{l_2} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos\left(\frac{\pi n y}{l_2}\right) \cos\left(\frac{\pi n \eta}{l_2}\right) \exp\left(-\frac{\pi^2 n^2 a t}{l_2^2}\right) \right], \\ G_3(z, \zeta, t) &= \frac{1}{l_3} \left[ 1 + 2 \sum_{n=1}^{\infty} \cos\left(\frac{\pi n z}{l_3}\right) \cos\left(\frac{\pi n \zeta}{l_3}\right) \exp\left(-\frac{\pi^2 n^2 a t}{l_3^2}\right) \right]. \end{aligned}$$

**3.1.1-29. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Third boundary value problem.**

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w - k_3 w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w + k_4 w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ \partial_z w - k_5 w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w + k_6 w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}). \end{aligned}$$



The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 3.1.1-28 where

$$G(x, y, z, \xi, \eta, \zeta, t) = H_1(x, \xi, t) \overline{H_2(y, \eta, t)} \overline{H_3(z, \zeta, t)}.$$

The functions  $H_1(x, \xi, t)$  and  $H_2(y, \eta, t)$  can be found in Paragraph 3.1.1-21, and the function  $H_3(z, \zeta, t)$  is given by

$$H_3(z, \zeta, t) = \sum_{n=1}^{\infty} \frac{\rho_n(z) \rho_n(\zeta)}{\|\rho_n\|^2} \exp(-a\nu_n^2 t),$$

$$\rho_n(x) = \cos(\nu_n x) + \frac{k_5}{\nu_n} \sin(\nu_n x), \quad \|\rho_n\|^2 = \frac{k_6}{2\nu_n^2} \frac{\nu_n^2 + k_5^2}{\nu_n^2 + k_6^2} + \frac{k_5}{2\nu_n^2} + \frac{l_3}{2} \left( 1 + \frac{k_5^2}{\nu_n^2} \right),$$

where the  $\nu_n$  are positive roots of the transcendental equation  $\frac{\tan(\nu l_3)}{\nu} = \frac{k_5 + k_6}{\nu^2 - k_5 k_6}$ .

3.1.1-30. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Mixed boundary value problems.

1°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = G_1(x, \xi, t)G_2(y, \eta, t)G_3(z, \zeta, t),$$

$$G_1(x, \xi, t) = \frac{2}{l_1} \sum_{n=1}^{\infty} \sin\left(\frac{\pi n x}{l_1}\right) \sin\left(\frac{\pi n \xi}{l_1}\right) \exp\left(-\frac{\pi^2 n^2 a t}{l_1^2}\right),$$

$$G_2(y, \eta, t) = \frac{2}{l_2} \sum_{k=1}^{\infty} \sin\left(\frac{\pi k y}{l_2}\right) \sin\left(\frac{\pi k \eta}{l_2}\right) \exp\left(-\frac{\pi^2 k^2 a t}{l_2^2}\right),$$

$$G_3(z, \zeta, t) = \frac{1}{l_3} + \frac{2}{l_3} \sum_{m=1}^{\infty} \cos\left(\frac{\pi m z}{l_3}\right) \cos\left(\frac{\pi m \zeta}{l_3}\right) \exp\left(-\frac{\pi^2 m^2 a t}{l_3^2}\right).$$

2°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$w = f(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$w = g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}),$$

$$w = g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}),$$

$$\partial_y w = g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}),$$

$$\partial_y w = g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}),$$

$$\partial_z w = g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}),$$

$$\partial_z w = g_6(x, y, t) \quad \text{at} \quad z = l_3 \quad (\text{boundary condition}).$$

Solution:

$$\begin{aligned} w(x, y, z, t) = & \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ & + a \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ & - a \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ & - a \int_0^t \int_0^{l_3} \int_0^{l_2} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ & + a \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\ & - a \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ & + a \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = G_1(x, \xi, t)G_2(y, \eta, t)G_3(z, \zeta, t),$$

$$G_1(x, \xi, t) = \frac{2}{l_1} \sum_{n=1}^{\infty} \sin\left(\frac{\pi n x}{l_1}\right) \sin\left(\frac{\pi n \xi}{l_1}\right) \exp\left(-\frac{\pi^2 n^2 a t}{l_1^2}\right),$$

$$G_2(y, \eta, t) = \frac{1}{l_2} + \frac{2}{l_2} \sum_{k=1}^{\infty} \cos\left(\frac{\pi k y}{l_2}\right) \cos\left(\frac{\pi k \eta}{l_2}\right) \exp\left(-\frac{\pi^2 k^2 a t}{l_2^2}\right),$$

$$G_3(z, \zeta, t) = \frac{1}{l_3} + \frac{2}{l_3} \sum_{m=1}^{\infty} \cos\left(\frac{\pi m z}{l_3}\right) \cos\left(\frac{\pi m \zeta}{l_3}\right) \exp\left(-\frac{\pi^2 m^2 a t}{l_3^2}\right).$$

### 3.1.2. Problems in Cylindrical Coordinates

The three-dimensional sourceless heat equation in the cylindrical coordinate system has the form

$$\frac{\partial w}{\partial t} = a \left[ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} \right], \quad r = \sqrt{x^2 + y^2}.$$

It is used to describe nonsymmetric unsteady processes in moving media or solids with cylindrical or plane boundaries. A similar equation is used to study the corresponding three-dimensional unsteady mass-exchange processes with constant diffusivity.

One-dimensional problems with axial symmetry that have solutions of the form  $w = w(r, t)$  are discussed in Subsection 1.2.1. Two-dimensional problems whose solutions have the form  $w = w(r, \varphi, t)$  or  $w = w(r, z, t)$  are considered in Subsections 2.1.2 and 2.1.3.

#### 3.1.2-1. Remarks on the Green's functions.

For the three-dimensional problems dealt with in Subsection 3.1.2, the Green's function can be represented in the product form

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t),$$

where  $G_1(r, \varphi, \xi, \eta, t)$  is the Green's function of the two-dimensional boundary value problem (such functions are presented in Subsection 2.1.2), and  $G_2(z, \zeta, t)$  is the Green's function of the corresponding one-dimensional boundary value problem (such functions can be found in Subsections 1.1.1 and 1.1.2).

**Example.** The Green's function of the first boundary value problem for a semiinfinite circular cylinder ( $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ ) of Paragraph 3.1.2-5 is the product of the two-dimensional Green's function of the first boundary value problem of Paragraph 2.1.2-2 ( $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ) and the one-dimensional Green's function of the first boundary value problem of Paragraph 1.1.2-2 ( $0 \leq z < \infty$ ), in which one should perform obvious renaming of variables.

General formulas that enable one to obtain solutions of basic boundary value problems with the help of the Green's function can be found in Subsection 0.8.1.

#### 3.1.2-2. Domain: $0 \leq r \leq R$ , $0 \leq \varphi \leq 2\pi$ , $-\infty < z < \infty$ . First boundary value problem.

An infinite circular cylinder is considered. The following conditions are prescribed:

$$w = f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$w = g(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}).$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) = & \int_{-\infty}^{\infty} \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ & - aR \int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} g(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \exp\left[-\frac{(z - \zeta)^2}{4 a t}\right], \quad A_n = \begin{cases} 1 & \text{for } n=0, \\ 2 & \text{for } n=1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**3.1.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Second boundary value problem.**

An infinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_{-\infty}^{\infty} \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ aR \int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} g(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \exp\left[-\frac{(z - \zeta)^2}{4at}\right], \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**3.1.2-4. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Third boundary value problem.**

An infinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 3.1.2-3 where

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 + k^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \exp\left[-\frac{(z - \zeta)^2}{4at}\right], \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots \end{cases} \end{aligned}$$

Here, the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k J_n(\mu R) = 0.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**3.1.2-5. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . First boundary value problem.**

A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^\infty \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &\quad - aR \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &\quad + a \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} \sum_{n=0}^\infty \sum_{m=1}^\infty \frac{A_n}{[J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \left\{ \exp\left[-\frac{(z - \zeta)^2}{4at}\right] - \exp\left[-\frac{(z + \zeta)^2}{4at}\right] \right\}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**3.1.2-6. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Second boundary value problem.**

A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^\infty \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &\quad + aR \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} + \frac{1}{\pi} \sum_{n=0}^\infty \sum_{m=1}^\infty \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \left\{ \exp\left[-\frac{(z - \zeta)^2}{4at}\right] + \exp\left[-\frac{(z + \zeta)^2}{4at}\right] \right\}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**3.1.2-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Third boundary value problem.**

A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + k_1 w &= g(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ \partial_z w - k_2 w &= g_2(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 3.1.2-6 where

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 + k_1^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \left\{ \exp\left[-\frac{(z - \zeta)^2}{4at}\right] + \exp\left[-\frac{(z + \zeta)^2}{4at}\right] - 2k_2 \int_0^{\infty} \exp\left[-\frac{(z + \zeta + s)^2}{4at}\right] - k_2 s \, ds \right\}. \end{aligned}$$

Here,  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k_1 J_n(\mu R) = 0.$$

**3.1.2-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Mixed boundary value problems.**

1°. A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^{\infty} \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &\quad - aR \int_0^t \int_0^{2\pi} \int_0^R g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \left\{ \exp\left[-\frac{(z - \zeta)^2}{4at}\right] + \exp\left[-\frac{(z + \zeta)^2}{4at}\right] \right\}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

2°. A semiinfinite circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^\infty \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ aR \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^\infty \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} + \frac{1}{\pi} \sum_{n=0}^\infty \sum_{m=1}^\infty \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{2\sqrt{\pi a t}} \left\{ \exp\left[-\frac{(z - \zeta)^2}{4at}\right] - \exp\left[-\frac{(z + \zeta)^2}{4at}\right] \right\}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

3.1.2-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- aR \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^\infty \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a \int_0^t \int_0^\infty \int_0^{2\pi} \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} \sum_{n=0}^\infty \sum_{m=1}^\infty \frac{A_n}{[J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{2}{l} \sum_{n=1}^\infty \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{an^2 \pi^2 t}{l^2}\right), \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

3.1.2-10. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ aR \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &- a \int_0^t \int_0^l \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a \int_0^t \int_0^l \int_0^{2\pi} \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi \xi}{l}\right) \exp\left(-\frac{an^2 \pi^2 t}{l^2}\right), \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

3.1.2-11. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + k_1 w &= g(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ \partial_z w - k_2 w &= g_2(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w + k_3 w &= g_3(r, \varphi, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 3.1.2-10 where

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) \sum_{s=1}^{\infty} \frac{h_s(z) h_s(\zeta)}{\|h_s\|^2} \exp(-a \lambda_s^2 t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 + k_1^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ h_s(z) &= \cos(\lambda_s z) + \frac{k_2}{\lambda_s} \sin(\lambda_s z), \quad \|h_s\|^2 = \frac{k_3}{2\lambda_s^2} \frac{\lambda_s^2 + k_2^2}{\lambda_s^2 + k_3^2} + \frac{k_2}{2\lambda_s^2} + \frac{l}{2} \left(1 + \frac{k_2^2}{\lambda_s^2}\right). \end{aligned}$$



Here,  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(\xi)$  are the Bessel functions; and the  $\mu_{nm}$  and  $\lambda_s$  are positive roots of the transcendental equations

$$\mu J'_n(\mu R) + k_1 J_n(\mu R) = 0, \quad \frac{\tan(\lambda l)}{\lambda} = \frac{k_2 + k_3}{\lambda^2 - k_2 k_3}.$$

**3.1.2-12. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.**

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \text{(initial condition),} \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \text{(boundary condition),} \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &\quad - aR \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &\quad + a \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ G_2(z, \zeta, t) &= \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{a n^2 \pi^2 t}{l^2}\right), \quad A_n = \begin{cases} 1 & \text{for } n=0, \\ 2 & \text{for } n=1, 2, \dots, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \text{(initial condition),} \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \text{(boundary condition),} \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \text{(boundary condition),} \\ w &= g_3(r, \varphi, t) & \text{at } z = l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{2\pi} \int_0^R \xi f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &\quad + aR \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &\quad + a \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &\quad - a \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t)G_2(z, \zeta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{\pi R^2} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t),$$

$$G_2(z, \zeta, t) = \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{an^2 \pi^2 t}{l^2}\right), \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n = 1, 2, \dots, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

3.1.2-13. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . First boundary value problem.

An infinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_{-\infty}^{\infty} \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &\quad + aR_1 \int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\ &\quad - aR_2 \int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \exp\left[-\frac{(z - \zeta)^2}{4at}\right] G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{\pi}{2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t),$$

$$A_n = \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)},$$

$$Z_n(\mu_{nm} r) = J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

3.1.2-14. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Second boundary value problem.

An infinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_{-\infty}^{\infty} \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - aR_1 \int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \xi, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + aR_2 \int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{1}{2\sqrt{\pi at}} \exp\left[-\frac{(z-\zeta)^2}{4at}\right] G_1(r, \varphi, \xi, \eta, t), \\
 G_1(r, \varphi, \xi, \eta, t) = & \frac{1}{\pi(R_2^2 - R_1^2)} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at)}{(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\
 & Z_n(\mu_{nm} r) = J_n'(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n'(\mu_{nm} R_1) J_n(\mu_{nm} r),
 \end{aligned}$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions (the prime denotes the derivative with respect to the argument); and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n'(\mu R_1) Y_n'(\mu R_2) - Y_n'(\mu R_1) J_n'(\mu R_2) = 0.$$

3.1.2-15. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Third boundary value problem.

An infinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_r w - k_1 w = g_1(\varphi, z, t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
 \partial_r w + k_2 w = g_2(\varphi, z, t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition).}
 \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is given by relations in Paragraph 3.1.2-14 in which

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{1}{2\sqrt{\pi at}} \exp\left[-\frac{(z-\zeta)^2}{4at}\right] G_1(r, \varphi, \xi, \eta, t), \\
 G_1(r, \varphi, \xi, \eta, t) = & \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at)}{(k_1^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\
 & Z_n(\mu_{nm} r) = [\mu_{nm} J_n'(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) \\
 & - [\mu_{nm} Y_n'(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r).
 \end{aligned}$$

Here,  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ;  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned}
 & [\mu J_n'(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y_n'(\mu R_2) + k_2 Y_n(\mu R_2)] \\
 & = [\mu Y_n'(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J_n'(\mu R_2) + k_2 J_n(\mu R_2)].
 \end{aligned}$$

3.1.2-16. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . First boundary value problem.

A semiinfinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^\infty \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ aR_1 \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\ &- aR_2 \int_0^t \int_0^\infty \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] - \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{\pi}{2} \sum_{n=0}^\infty \sum_{m=1}^\infty A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at), \\ A_n &= \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)}, \\ Z_n(\mu_{nm} r) &= J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r), \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

3.1.2-17. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Second boundary value problem.

A semiinfinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^\infty \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- aR_1 \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ aR_2 \int_0^t \int_0^\infty \int_0^{2\pi} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &- a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z-\zeta)^2}{4at} \right] + \exp \left[ -\frac{(z+\zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{\pi(R_2^2 - R_1^2)} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at)}{(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)},$$

$$Z_n(\mu_{nm} r) = J_n'(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n'(\mu_{nm} R_1) J_n(\mu_{nm} r),$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n'(\mu R_1) Y_n'(\mu R_2) - Y_n'(\mu R_1) J_n'(\mu R_2) = 0.$$

3.1.2-18. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Third boundary value problem.

A semiinfinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(\varphi, z, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(\varphi, z, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}), \\ \partial_z w - k_3 w &= g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 3.1.2-17 where

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(z, \zeta, t) G_2(r, \varphi, \xi, \eta, t),$$

$$G_1(z, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z-\zeta)^2}{4at} \right] + \exp \left[ -\frac{(z+\zeta)^2}{4at} \right] - 2k_3 \int_0^{\infty} \exp \left[ -\frac{(z+\zeta+s)^2}{4at} - k_3 s \right] ds \right\},$$

$$G_2(r, \varphi, \xi, \eta, t) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at)}{(k_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)},$$

$$Z_n(\mu_{nm} r) = [\mu_{nm} J_n'(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) - [\mu_{nm} Y_n'(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r).$$

Here,  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned} &[\mu J_n'(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y_n'(\mu R_2) + k_2 Y_n(\mu R_2)] \\ &= [\mu Y_n'(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J_n'(\mu R_2) + k_2 J_n(\mu R_2)]. \end{aligned}$$

3.1.2-19. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Mixed boundary value problems.

1°. A semiinfinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^\infty \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & + aR_1 \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\
 & - aR_2 \int_0^t \int_0^\infty \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\
 & - a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] + \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t), \\
 G_1(r, \varphi, \xi, \eta, t) = & \frac{\pi}{2} \sum_{n=0}^\infty \sum_{m=1}^\infty A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at), \\
 A_n = & \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)}, \\
 Z_n(\mu_{nm} r) = & J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),
 \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

2°. A semiinfinite hollow circular cylinder is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_r w = g_1(\varphi, z, t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
 \partial_r w = g_2(\varphi, z, t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition),} \\
 w = g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^\infty \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - aR_1 \int_0^t \int_0^\infty \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + aR_2 \int_0^t \int_0^\infty \int_0^{2\pi} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] - \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t), \\
 G_1(r, \varphi, \xi, \eta, t) = & \frac{1}{\pi(R_2^2 - R_1^2)} + \frac{1}{\pi} \sum_{n=0}^\infty \sum_{m=1}^\infty \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at)}{(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\
 Z_n(\mu_{nm} r) = & J_n'(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n'(\mu_{nm} R_1) J_n(\mu_{nm} r),
 \end{aligned}$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions (the prime denotes the derivative with respect to the argument); and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1)Y'_n(\mu R_2) - Y'_n(\mu R_1)J'_n(\mu R_2) = 0.$$

3.1.2-20. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_4(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ aR_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\ &- aR_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\ &+ a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t) \left[ \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right) \right],$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{\pi}{2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 at),$$

$$A_n = \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)},$$

$$Z_n(\mu_{nm} r) = J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1)Y_n(\mu R_2) - Y_n(\mu R_1)J_n(\mu R_2) = 0.$$

3.1.2-21. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - a R_1 \int_0^t \int_0^{2\pi} \int_0^{R_1} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + a R_2 \int_0^t \int_0^{2\pi} \int_0^{R_2} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & - a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
 & + a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & G_1(r, \varphi, \xi, \eta, t) \left[ \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right) \right], \\
 G_1(r, \varphi, \xi, \eta, t) = & \frac{1}{\pi(R_2^2 - R_1^2)} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t)}{(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\
 Z_n(\mu_{nm} r) = & J_n'(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n'(\mu_{nm} R_1) J_n(\mu_{nm} r),
 \end{aligned}$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions (the prime denotes the derivative with respect to the argument); and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n'(\mu R_1) Y_n'(\mu R_2) - Y_n'(\mu R_1) J_n'(\mu R_2) = 0.$$

3.1.2-22. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f(r, \varphi, z) \quad & \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_r w - k_1 w = g_1(\varphi, z, t) \quad & \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\
 \partial_r w + k_2 w = g_2(\varphi, z, t) \quad & \text{at} \quad r = R_2 \quad (\text{boundary condition}), \\
 \partial_z w - k_3 w = g_3(r, \varphi, t) \quad & \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
 \partial_z w + k_4 w = g_4(r, \varphi, t) \quad & \text{at} \quad z = l \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 3.1.2-21 where

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t).$$

Here, the first factor has the form

$$\begin{aligned}
 G_1(r, \varphi, \xi, \eta, t) = & \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t)}{(k_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\
 Z_n(\mu_{nm} r) = & [\mu_{nm} J_n'(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) \\
 & - [\mu_{nm} Y_n'(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r),
 \end{aligned}$$



where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned} & [\mu J'_n(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y'_n(\mu R_2) + k_2 Y_n(\mu R_2)] \\ &= [\mu Y'_n(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J'_n(\mu R_2) + k_2 J_n(\mu R_2)]. \end{aligned}$$

The second factor is given by

$$\begin{aligned} G_2(z, \zeta, t) &= \sum_{s=1}^{\infty} \frac{h_s(z)h_s(\zeta)}{\|h_s\|^2} \exp(-a\lambda_s^2 t), \\ h_s(z) &= \cos(\lambda_s z) + \frac{k_3}{\lambda_s} \sin(\lambda_s z), \quad \|h_s\|^2 = \frac{k_4}{2\lambda_s^2} \frac{\lambda_s^2 + k_3^2}{\lambda_s^2 + k_4^2} + \frac{k_3}{2\lambda_s^2} + \frac{l}{2} \left(1 + \frac{k_3^2}{\lambda_s^2}\right), \end{aligned}$$

where the  $\lambda_s$  are positive roots of the transcendental equation  $\frac{\tan(\lambda l)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}$ .

3.1.2-23. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_4(r, \varphi, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ a R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\ &- a R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\ &- a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\ &+ a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) \left[ \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{a n^2 \pi^2 t}{l^2}\right) \right], \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{\pi}{2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t), \\ A_n &= \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)}, \\ Z_n(\mu_{nm} r) &= J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r), \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1)Y_n(\mu R_2) - Y_n(\mu R_1)J_n(\mu R_2) = 0.$$

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_4(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &\quad - a R_1 \int_0^t \int_0^{2\pi} \int_0^l g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &\quad + a R_2 \int_0^t \int_0^{2\pi} \int_0^l g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &\quad + a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &\quad - a \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t) \left[ \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{an^2 \pi^2 t}{l^2}\right) \right],$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{\pi(R_2^2 - R_1^2)} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \exp(-\mu_{nm}^2 a t)}{(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)},$$

$$Z_n(\mu_{nm} r) = J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r),$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions (the prime denotes the derivative with respect to the argument); and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

3.1.2-24. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $-\infty < z < \infty$ . First boundary value problem.

A dihedral angle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(r, z, t) & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ w &= g_2(r, z, t) & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_{-\infty}^{\infty} \int_0^{\varphi_0} \int_0^{\infty} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &\quad + a \int_0^t \int_{-\infty}^{\infty} \int_0^{\infty} g_1(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &\quad - a \int_0^t \int_{-\infty}^{\infty} \int_0^{\infty} g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \exp\left[-\frac{(z-\zeta)^2}{4at}\right] G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{a\varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \sum_{n=1}^{\infty} I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right),$$

where the  $I_\nu(r)$  are the modified Bessel functions.

3.1.2-25. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $-\infty < z < \infty$ . Second boundary value problem.

A dihedral angle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ r^{-1} \partial_\varphi w &= g_1(r, z, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_2(r, z, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_{-\infty}^{\infty} \int_0^{\varphi_0} \int_0^{\infty} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &\quad - a \int_0^t \int_{-\infty}^{\infty} \int_0^{\infty} g_1(\xi, \zeta, \tau) G(r, \varphi, z, \xi, 0, \zeta, t-\tau) d\xi d\zeta d\tau \\ &\quad + a \int_0^t \int_{-\infty}^{\infty} \int_0^{\infty} g_2(\xi, \zeta, \tau) G(r, \varphi, z, \xi, \varphi_0, \zeta, t-\tau) d\xi d\zeta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \exp\left[-\frac{(z-\zeta)^2}{4at}\right] G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{a\varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \left[ \frac{1}{2} I_0\left(\frac{r\xi}{2at}\right) + \sum_{n=1}^{\infty} I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \right],$$

where the  $I_\nu(r)$  are the modified Bessel functions.

3.1.2-26. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ . First boundary value problem.

The upper half of a dihedral angle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(r, z, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\ w &= g_2(r, z, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^{\infty} \int_0^{\varphi_0} \int_0^{\infty} f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &\quad + a \int_0^t \int_0^{\infty} \int_0^{\infty} g_1(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t-\tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{\infty} \int_0^{\infty} g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t-\tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &\quad + a \int_0^t \int_0^{\varphi_0} \int_0^{\infty} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t-\tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z-\zeta)^2}{4at} \right] - \exp \left[ -\frac{(z+\zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{a\varphi_0 t} \exp \left( -\frac{r^2 + \xi^2}{4at} \right) \sum_{n=1}^{\infty} I_{n\pi/\varphi_0} \left( \frac{r\xi}{2at} \right) \sin \left( \frac{n\pi\varphi}{\varphi_0} \right) \sin \left( \frac{n\pi\eta}{\varphi_0} \right),$$

where the  $I_\nu(r)$  are the modified Bessel functions.

3.1.2-27. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ . Second boundary value problem.

The upper half of a dihedral angle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ r^{-1} \partial_\varphi w &= g_1(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ r^{-1} \partial_\varphi w &= g_2(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^\infty \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &\quad - a \int_0^t \int_0^\infty \int_0^\infty g_1(\xi, \zeta, \tau) G(r, \varphi, z, \xi, 0, \zeta, t-\tau) d\xi d\zeta d\tau \\ &\quad + a \int_0^t \int_0^\infty \int_0^\infty g_2(\xi, \zeta, \tau) G(r, \varphi, z, \xi, \varphi_0, \zeta, t-\tau) d\xi d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t-\tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z-\zeta)^2}{4at} \right] + \exp \left[ -\frac{(z+\zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{a\varphi_0 t} \exp \left( -\frac{r^2 + \xi^2}{4at} \right) \left[ \frac{1}{2} I_0 \left( \frac{r\xi}{2at} \right) + \sum_{n=1}^{\infty} I_{n\pi/\varphi_0} \left( \frac{r\xi}{2at} \right) \cos \left( \frac{n\pi\varphi}{\varphi_0} \right) \cos \left( \frac{n\pi\eta}{\varphi_0} \right) \right],$$

where the  $I_\nu(r)$  are the modified Bessel functions.

3.1.2-28. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ . Mixed boundary value problems.

1°. The upper half of a dihedral angle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ w &= g_2(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^\infty \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & + a \int_0^t \int_0^\infty \int_0^\infty g_1(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^\infty \int_0^\infty g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] + \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t), \\
 G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{a\varphi_0 t} \exp \left( -\frac{r^2 + \xi^2}{4at} \right) \sum_{n=1}^\infty I_{n\pi/\varphi_0} \left( \frac{r\xi}{2at} \right) \sin \left( \frac{n\pi\varphi}{\varphi_0} \right) \sin \left( \frac{n\pi\eta}{\varphi_0} \right),
 \end{aligned}$$

where the  $I_\nu(r)$  are the modified Bessel functions.

2°. The upper half of a dihedral angle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 r^{-1} \partial_\varphi w &= g_1(r, z, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\
 r^{-1} \partial_\varphi w &= g_2(r, z, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}), \\
 w &= g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^\infty \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - a \int_0^t \int_0^\infty \int_0^\infty g_1(\xi, \zeta, \tau) G(r, \varphi, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^\infty \int_0^\infty g_2(\xi, \zeta, \tau) G(r, \varphi, z, \xi, \varphi_0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z - \zeta)^2}{4at} \right] - \exp \left[ -\frac{(z + \zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t), \\
 G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{a\varphi_0 t} \exp \left( -\frac{r^2 + \xi^2}{4at} \right) \left[ \frac{1}{2} I_0 \left( \frac{r\xi}{2at} \right) + \sum_{n=1}^\infty I_{n\pi/\varphi_0} \left( \frac{r\xi}{2at} \right) \cos \left( \frac{n\pi\varphi}{\varphi_0} \right) \cos \left( \frac{n\pi\eta}{\varphi_0} \right) \right],
 \end{aligned}$$

where the  $I_\nu(r)$  are the modified Bessel functions.

**3.1.2-29. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . First boundary value problem.**

A wedge domain of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ w &= g_2(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\ w &= g_3(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_4(r, \varphi, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ a \int_0^t \int_0^l \int_0^\infty g_1(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a \int_0^t \int_0^l \int_0^\infty g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &+ a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) \left[ \frac{2}{l} \sum_{n=1}^\infty \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right) \right], \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{a\varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \sum_{n=1}^\infty I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right), \end{aligned}$$

where the  $I_\nu(r)$  are the modified Bessel functions.

**3.1.2-30. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

A wedge domain of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ r^{-1} \partial_\varphi w &= g_1(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ r^{-1} \partial_\varphi w &= g_2(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^l \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - a \int_0^t \int_0^l \int_0^\infty g_1(\xi, \zeta, \tau) G(r, \varphi, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^l \int_0^\infty g_2(\xi, \zeta, \tau) G(r, \varphi, z, \xi, \varphi_0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
 & + a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\
 G_1(r, \varphi, \xi, \eta, t) &= \frac{1}{a\varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \left[ \frac{1}{2} I_0\left(\frac{r\xi}{2at}\right) + \sum_{n=1}^{\infty} I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \right], \\
 G_2(z, \zeta, t) &= \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi\zeta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right),
 \end{aligned}$$

where the  $I_\nu(r)$  are the modified Bessel functions.

3.1.2-31. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.

1°. A wedge domain of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 w &= g_1(r, z, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\
 w &= g_2(r, z, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}), \\
 \partial_z w &= g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
 \partial_z w &= g_4(r, \varphi, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^l \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & + a \int_0^t \int_0^l \int_0^\infty g_1(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^l \int_0^\infty g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
 & + a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{a\varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \sum_{n=1}^{\infty} I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right),$$

$$G_2(z, \zeta, t) = \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi\zeta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right),$$

where the  $I_\nu(r)$  are the modified Bessel functions.

2°. A wedge domain of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ r^{-1} \partial_\varphi w &= g_1(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ r^{-1} \partial_\varphi w &= g_2(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\ w &= g_3(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_4(r, \varphi, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^l \int_0^{\varphi_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a \int_0^t \int_0^l \int_0^\infty g_1(\xi, \zeta, \tau) G(r, \varphi, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a \int_0^t \int_0^l \int_0^\infty g_2(\xi, \zeta, \tau) G(r, \varphi, z, \xi, \varphi_0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a \int_0^t \int_0^{\varphi_0} \int_0^\infty g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t) \left[ \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi\zeta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right) \right],$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{1}{a\varphi_0 t} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) \left[ \frac{1}{2} I_0\left(\frac{r\xi}{2at}\right) + \sum_{n=1}^{\infty} I_{n\pi/\varphi_0}\left(\frac{r\xi}{2at}\right) \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \right],$$

where the  $I_\nu(r)$  are the modified Bessel functions.

3.1.2-32. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $-\infty < z < \infty$ . First boundary value problem.

An infinite cylindrical sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(\varphi, z, t) & \text{at } r &= R & \text{(boundary condition),} \\ w &= g_2(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition).} \end{aligned}$$



Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_{-\infty}^{\infty} \int_0^{\varphi_0} \int_0^R f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - aR \int_0^t \int_{-\infty}^{\infty} \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\
 & + a \int_0^t \int_{-\infty}^{\infty} \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_{-\infty}^{\infty} \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau.
 \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \exp \left[ -\frac{(z-\zeta)^2}{4at} \right] G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{4}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin \left( \frac{n\pi \varphi}{\varphi_0} \right) \sin \left( \frac{n\pi \eta}{\varphi_0} \right) \exp(-\mu_{nm}^2 at),$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

3.1.2-33. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ . First boundary value problem.

A semiinfinite cylindrical sector is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\
 w &= g_1(\varphi, z, t) & \text{at } r &= R & \text{(boundary condition),} \\
 w &= g_2(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\
 w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\
 w &= g_4(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^{\infty} \int_0^{\varphi_0} \int_0^R f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - aR \int_0^t \int_0^{\varphi_0} \int_0^R g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^{\varphi_0} \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^{\varphi_0} \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z-\zeta)^2}{4at} \right] - \exp \left[ -\frac{(z+\zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t),$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{4}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin \left( \frac{n\pi \varphi}{\varphi_0} \right) \sin \left( \frac{n\pi \eta}{\varphi_0} \right) \exp(-\mu_{nm}^2 at),$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

**3.1.2-34. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ . Mixed boundary value problem.**

A semiinfinite cylindrical sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r &= R & (\text{boundary condition}), \\ w &= g_2(r, z, t) & \text{at } \varphi &= 0 & (\text{boundary condition}), \\ w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}), \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z &= 0 & (\text{boundary condition}). \end{aligned}$$

**Solution:**

$$\begin{aligned} w(r, \varphi, z, t) &= \int_0^\infty \int_0^{\varphi_0} \int_0^R f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &\quad - aR \int_0^t \int_0^\infty \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &\quad + a \int_0^t \int_0^\infty \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &\quad - a \int_0^t \int_0^\infty \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &\quad - a \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{2\sqrt{\pi at}} \left\{ \exp \left[ -\frac{(z-\zeta)^2}{4at} \right] + \exp \left[ -\frac{(z+\zeta)^2}{4at} \right] \right\} G_1(r, \varphi, \xi, \eta, t), \\ G_1(r, \varphi, \xi, \eta, t) &= \frac{4}{R^2 \varphi_0} \sum_{n=1}^\infty \sum_{m=1}^\infty \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin \left( \frac{n\pi \varphi}{\varphi_0} \right) \sin \left( \frac{n\pi \eta}{\varphi_0} \right) \exp(-\mu_{nm}^2 at), \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

**3.1.2-35. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . First boundary value problem.**

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r &= R & (\text{boundary condition}), \\ w &= g_2(r, z, t) & \text{at } \varphi &= 0 & (\text{boundary condition}), \\ w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}), \\ w &= g_4(r, \varphi, t) & \text{at } z &= 0 & (\text{boundary condition}), \\ w &= g_5(r, \varphi, t) & \text{at } z &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^l \int_0^{\varphi_0} \int_0^R f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - aR \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\
 & + a \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\
 & - a \int_0^t \int_0^{\varphi_0} \int_0^R g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) &= G_1(r, \varphi, \xi, \eta, t) G_2(z, \zeta, t), \\
 G_1(r, \varphi, \xi, \eta, t) &= \frac{4}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \exp(-\mu_{nm}^2 a t), \\
 G_2(z, \zeta, t) &= \frac{2}{l} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi z}{l}\right) \sin\left(\frac{n\pi\zeta}{l}\right) \exp\left(-\frac{a n^2 \pi^2 t}{l^2}\right),
 \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

3.1.2-36. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . Mixed boundary value problem.

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\
 w &= g_1(\varphi, z, t) & \text{at } r &= R & \text{(boundary condition),} \\
 w &= g_2(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\
 w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\
 \partial_z w &= g_4(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition),} \\
 \partial_z w &= g_5(r, \varphi, t) & \text{at } z &= l & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \int_0^l \int_0^{\varphi_0} \int_0^R f(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & - aR \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\
 & + a \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\
 & - a \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
 & + a \int_0^t \int_0^{\varphi_0} \int_0^R g_5(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = G_1(r, \varphi, \xi, \eta, t) \left[ \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos\left(\frac{n\pi z}{l}\right) \cos\left(\frac{n\pi \zeta}{l}\right) \exp\left(-\frac{an^2\pi^2 t}{l^2}\right) \right],$$

$$G_1(r, \varphi, \xi, \eta, t) = \frac{4}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi) \sin\left(\frac{n\pi \varphi}{\varphi_0}\right) \sin\left(\frac{n\pi \eta}{\varphi_0}\right) \exp(-\mu_{nm}^2 at),$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

### 3.1.3. Problems in Spherical Coordinates

The heat equation in the spherical coordinate system has the form

$$\frac{\partial w}{\partial t} = a \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} \right], \quad r = \sqrt{x^2 + y^2 + z^2}.$$

This representation is convenient to describe three-dimensional heat and mass exchange phenomena in domains bounded by coordinate surfaces of the spherical coordinate system.

One-dimensional problems with central symmetry that have solutions of the form  $w = w(r, t)$  are discussed in Subsection 1.2.3.

3.1.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \int_0^{2\pi} \int_0^{\pi} \int_0^R f(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &\quad - aR^2 \int_0^t \int_0^{2\pi} \int_0^{\pi} g(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} \sin \eta \, d\eta \, d\zeta \, d\tau, \end{aligned}$$

where

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= \frac{1}{2\pi R^2 \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\ &\quad \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \exp(-\lambda_{nm}^2 at), \\ A_k &= \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{(2n+1)(n-k)!}{(n+k)! [J'_{n+1/2}(\lambda_{nm} R)]^2}. \end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as follows:

$$P_n^k(\mu) = (1 - \mu^2)^{k/2} \frac{d^k}{d\mu^k} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n;$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $J_{n+1/2}(\lambda R) = 0$ .

⊙ *References:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), H. S. Carslaw and J. C. Jaeger (1984).

**3.1.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \int_0^{2\pi} \int_0^\pi \int_0^R f(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &\quad + aR^2 \int_0^t \int_0^{2\pi} \int_0^\pi g(\eta, \zeta, \tau) G(r, \theta, \varphi, R, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau, \end{aligned}$$

where

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= \frac{3}{4\pi R^3} + \frac{1}{2\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\ &\quad \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \exp(-\lambda_{nm}^2 a t), \\ A_k &= \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{\lambda_{nm}^2 (2n+1)(n-k)!}{(n+k)! [R^2 \lambda_{nm}^2 - n(n+1)] [J_{n+1/2}(\lambda_{nm} R)]^2}. \end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions (see Paragraph 3.1.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$2\lambda R J'_{n+1/2}(\lambda R) - J_{n+1/2}(\lambda R) = 0.$$

**3.1.3-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.**

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 3.1.3-2 where

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= \frac{1}{2\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n A_s B_{nms} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\ &\quad \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \exp(-\lambda_{nm}^2 a t), \\ A_s &= \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{\lambda_{nm}^2 (2n+1)(n-s)!}{(n+s)! [R^2 \lambda_{nm}^2 + (kR+n)(kR-n-1)] [J_{n+1/2}(\lambda_{nm} R)]^2}. \end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 3.1.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda R J'_{n+1/2}(\lambda R) + \left(kR - \frac{1}{2}\right) J_{n+1/2}(\lambda R) = 0.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

3.1.3-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\theta, \varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(\theta, \varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &+ a R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} \sin \eta \, d\eta \, d\zeta \, d\tau \\ &- a R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} \sin \eta \, d\eta \, d\zeta \, d\tau, \end{aligned}$$

where

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= \frac{\pi}{8\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} Z_{n+1/2}(\lambda_{nm} r) Z_{n+1/2}(\lambda_{nm} \xi) \\ &\times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \exp(-\lambda_{nm}^2 a t). \end{aligned}$$

Here,

$$\begin{aligned} Z_{n+1/2}(\lambda_{nm} r) &= J_{n+1/2}(\lambda_{nm} R_1) Y_{n+1/2}(\lambda_{nm} r) - Y_{n+1/2}(\lambda_{nm} R_1) J_{n+1/2}(\lambda_{nm} r), \\ A_k &= \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{\lambda_{nm}^2 (2n+1)(n-k)! J_{n+1/2}^2(\lambda_{nm} R_2)}{(n+k)! [J_{n+1/2}^2(\lambda_{nm} R_1) - J_{n+1/2}^2(\lambda_{nm} R_2)]}, \end{aligned}$$

where the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as follows:

$$P_n^k(\mu) = (1 - \mu^2)^{k/2} \frac{d^k}{d\mu^k} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n;$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$Z_{n+1/2}(\lambda R_2) = 0.$$

⊙ Reference: B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

3.1.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g_1(\theta, \varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\theta, \varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &- a R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) G(r, \theta, \varphi, R_1, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\ &+ a R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) G(r, \theta, \varphi, R_2, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau, \end{aligned}$$

where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{3}{4\pi(R_2^3 - R_1^3)} + \frac{1}{4\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n \frac{A_k}{B_{nmk}} Z_{n+1/2}(\lambda_{nm}r) Z_{n+1/2}(\lambda_{nm}\xi) \\ \times P_n^k(\cos\theta) P_n^k(\cos\eta) \cos[k(\varphi - \zeta)] \exp(-\lambda_{nm}^2 at).$$

Here,

$$A_k = \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{(n+k)!}{(2n+1)(n-k)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm}r) dr, \\ Z_{n+1/2}(\lambda r) = \left[ \lambda J'_{n+1/2}(\lambda R_1) - \frac{1}{2R_1} J_{n+1/2}(\lambda R_1) \right] Y_{n+1/2}(\lambda r) \\ - \left[ \lambda Y'_{n+1/2}(\lambda R_1) - \frac{1}{2R_1} Y_{n+1/2}(\lambda R_1) \right] J_{n+1/2}(\lambda r),$$

where the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions (see Paragraph 3.1.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) - \frac{1}{2R_2} Z_{n+1/2}(\lambda R_2) = 0.$$

The integrals that determine the coefficients  $B_{nmk}$  can be expressed in terms of the Bessel functions and their derivatives; see Budak, Samarskii, and Tikhonov (1980).

3.1.3-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$w = f(r, \theta, \varphi) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w = g_1(\theta, \varphi, t) \quad \text{at } r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w = g_2(\theta, \varphi, t) \quad \text{at } r = R_2 \quad (\text{boundary condition}).$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 3.1.3-5 where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{1}{4\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n \frac{A_s}{B_{nms}} Z_{n+1/2}(\lambda_{nm}r) Z_{n+1/2}(\lambda_{nm}\xi) \\ \times P_n^s(\cos\theta) P_n^s(\cos\eta) \cos[s(\varphi - \zeta)] \exp(-\lambda_{nm}^2 at).$$

Here,

$$A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{(n+s)!}{(2n+1)(n-s)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm}r) dr, \\ Z_{n+1/2}(\lambda r) = \left[ \lambda J'_{n+1/2}(\lambda R_1) - \left( k_1 + \frac{1}{2R_1} \right) J_{n+1/2}(\lambda R_1) \right] Y_{n+1/2}(\lambda r) \\ - \left[ \lambda Y'_{n+1/2}(\lambda R_1) - \left( k_1 + \frac{1}{2R_1} \right) Y_{n+1/2}(\lambda R_1) \right] J_{n+1/2}(\lambda r),$$

where the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 3.1.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) + \left( k_2 - \frac{1}{2R_2} \right) Z_{n+1/2}(\lambda R_2) = 0.$$

The integrals that determine the coefficients  $B_{nms}$  can be expressed in terms of the Bessel functions and their derivatives.

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**3.1.3-7. Domain:  $0 \leq r < \infty$ ,  $0 \leq \theta \leq \theta_0$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

A cone is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(r, \varphi, t) \quad \text{at} \quad \theta = \theta_0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \int_0^{2\pi} \int_0^{\theta_0} \int_0^\infty f(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &\quad - a \int_0^t \int_0^{2\pi} \int_0^\infty g(\xi, \zeta, \tau) \left[ \sin \eta \frac{\partial}{\partial \eta} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\theta_0} d\xi \, d\zeta \, d\tau, \end{aligned}$$

where

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= -\frac{1}{4\pi a t \sqrt{r\xi}} \sum_{m=0}^\infty \sum_{\nu} \frac{A_m(2\nu+1)}{B_{m\nu}} \exp\left(-\frac{r^2 + \xi^2}{4at}\right) I_{\nu+1/2}\left(\frac{r\xi}{2at}\right) \\ &\quad \times P_\nu^{-m}(\cos \theta) P_\nu^{-m}(\cos \eta) \cos[m(\varphi - \zeta)], \\ A_m &= \begin{cases} 1 & \text{for } m=0, \\ 2 & \text{for } m \neq 0, \end{cases} \quad B_{m\nu} = \left[ (1-\mu)^2 \frac{d}{d\mu} P_\nu^{-m}(\mu) \frac{d}{d\nu} P_\nu^{-m}(\mu) \right]_{\mu=\cos \theta_0}. \end{aligned}$$

Here,  $P_\nu^{-m}(\mu)$  is the modified Legendre function expressed as

$$P_\nu^{-m}(\mu) = \frac{1}{\Gamma(1+m)} \left( \frac{1-\mu}{1+\mu} \right)^{m/2} F(-\nu, \nu+1, 1+m; \frac{1}{2} - \frac{1}{2}\mu),$$

where  $F(a, b, c; \mu)$  is the Gaussian hypergeometric function and  $\Gamma(z)$  is the gamma function. The summation with respect to  $\nu$  is performed over all roots of the equation  $P_\nu^{-m}(\cos \theta_0) = 0$  that are greater than  $-1/2$ .

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

## 3.2. Heat Equation with Source $\frac{\partial w}{\partial t} = a \Delta_3 w + \Phi(x, y, z, t)$

### 3.2.1. Problems in Cartesian Coordinates

In the Cartesian coordinate system, the three-dimensional heat equation with a volume source has the form

$$\frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + \Phi(x, y, z, t).$$

It describes three-dimensional unsteady thermal phenomena in quiescent media or solids with constant thermal diffusivity. A similar equation is used to study the corresponding three-dimensional mass transfer processes with constant diffusivity.

**3.2.1-1. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Cauchy problem.**

An initial condition is prescribed:

$$w = f(x, y, z) \quad \text{at} \quad t = 0.$$



Solution:

$$w(x, y, z, t) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ + \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau,$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{8(\pi at)^{3/2}} \exp \left[ -\frac{(x - \xi)^2 + (y - \eta)^2 + (z - \zeta)^2}{4at} \right].$$

⊙ *References:* A. G. Butkovskiy (1979).

**3.2.1-2. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a half-space is given by the formula in Paragraph 3.1.1-4 with the additional term

$$\int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_0^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a half-space is given by the formula in Paragraph 3.1.1-5 with the additional term (1).

3°. The solution of the third boundary value problem for a half-space is given by the formula in Paragraph 3.1.1-6 with the additional term (1).

⊙ *References:* A. G. Butkovskiy (1979), H. S. Carslaw and J. C. Jaeger (1984).

**3.2.1-3. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for an infinite layer is given by the formula in Paragraph 3.1.1-7 with the additional term

$$\int_0^t \int_0^l \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for an infinite layer is given by the formula in Paragraph 3.1.1-8 with the additional term (2).

3°. The solution of the third boundary value problem for an infinite layer is given by the formula in Paragraph 3.1.1-9 with the additional term (2).

4°. The solution of a mixed boundary value problem for an infinite layer is given by the formula in Paragraph 3.1.1-10 with the additional term (2).

**3.2.1-4. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a semiinfinite layer is given by the formula in Paragraph 3.1.1-11 with the additional term

$$\int_0^t \int_0^l \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau, \quad (3)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a semiinfinite layer is given by the formula in Paragraph 3.1.1-12 with the additional term (3).

3°. The solution of the third boundary value problem for a semiinfinite layer is given by the formula in Paragraph 3.1.1-13 with the additional term (3).

4°. The solutions of mixed boundary value problems for a semiinfinite layer are given by the formulas in Paragraph 3.1.1-14 with additional terms of the form (3).

⊙ *References:* A. G. Butkovskiy (1979), H. S. Carslaw and J. C. Jaeger (1984).

3.2.1-5. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . Different boundary value problems.

1°. The solution of the first boundary value problem for the first octant is given by the formula in Paragraph 3.1.1-15 with the additional term

$$\int_0^t \int_0^\infty \int_0^\infty \int_0^\infty \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau, \quad (4)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for the first octant is given by the formula in Paragraph 3.1.1-16 with the additional term (4).

3°. The solution of the third boundary value problem for the first octant is given by the formula in Paragraph 3.1.1-17 with the additional term (4).

4°. The solutions of mixed boundary value problems for the first octant are given by the formulas in Paragraph 3.1.1-18 with additional terms of the form (4).

3.2.1-6. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $-\infty < z < \infty$ . Different boundary value problems.

1°. The solution of the first boundary value problem in an infinite rectangular domain is given by the formula in Paragraph 3.1.1-19 with the additional term

$$\int_0^t \int_{-\infty}^\infty \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau, \quad (5)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem in an infinite rectangular domain is given by the formula in Paragraph 3.1.1-20 with the additional term (5).

3°. The solution of the third boundary value problem in an infinite rectangular domain is given by the formula in Paragraph 3.1.1-21 with the additional term (5).

4°. The solution of a mixed boundary value problem in an infinite rectangular domain is given by the formula in Paragraph 3.1.1-22 with the additional term (5).

3.2.1-7. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z < \infty$ . Different boundary value problems.

1°. The solution of the first boundary value problem in a semiinfinite rectangular domain is given by the formula of Paragraph 3.1.1-23 with the additional term

$$\int_0^t \int_0^\infty \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau, \quad (6)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem in a semiinfinite rectangular domain is given by the formula in Paragraph 3.1.1-24 with the additional term (6).

3°. The solution of the third boundary value problem in a semiinfinite rectangular domain is given by the formula in Paragraph 3.1.1-25 with the additional term (6).

4°. The solutions of mixed boundary value problems in a semiinfinite rectangular domain are given by the formulas in Paragraph 3.1.1-26 with additional terms of the form (6).

**3.2.1-8. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a rectangular parallelepiped is given by the formula in Paragraph 3.1.1-27 with the additional term

$$\int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau, \quad (7)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a rectangular parallelepiped is given by the formula in Paragraph 3.1.1-28 with the additional term (7).

3°. The solution of the third boundary value problem for a rectangular parallelepiped is given by the formula in Paragraph 3.1.1-29 with the additional term (7).

4°. The solutions of mixed boundary value problems for a rectangular parallelepiped are given by the formulas in Paragraph 3.1.1-30 with additional terms of the form (7).

⊙ *References:* A. G. Butkovskiy (1979), H. S. Carslaw and J. C. Jaeger (1984).

## 3.2.2. Problems in Cylindrical Coordinates

In the cylindrical coordinate system, the heat equation with a volume source is written as

$$\frac{\partial w}{\partial t} = a \left[ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} \right] + \Phi(r, \varphi, z, t).$$

This representation is used to describe nonsymmetric unsteady thermal (diffusion) processes in quiescent media or solids bounded by cylindrical surfaces and planes.

**3.2.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for an infinite circular cylinder is given by the formula in Paragraph 3.1.2-2 with the additional term

$$\int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for an infinite circular cylinder is given by the formula in Paragraph 3.1.2-3 with the additional term (1).

3°. The solution of the third boundary value problem for an infinite circular cylinder is the sum of the solution presented in Paragraph 3.1.2-4 and expression (1).

**3.2.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a semiinfinite circular cylinder is given by the formula in Paragraph 3.1.2-5 with the additional term

$$\int_0^t \int_0^{\infty} \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a semiinfinite circular cylinder is given by the formula in Paragraph 3.1.2-6 with the additional term (2).

3°. The solution of the third boundary value problem for a semiinfinite circular cylinder is the sum of the solution presented in Paragraph 3.1.2-7 and expression (2).

4°. The solutions of mixed boundary value problems for a semiinfinite circular cylinder are given by the formulas in Paragraph 3.1.2-8 with additional terms of the form (2).

**3.2.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a circular cylinder of finite length is given by the formula in Paragraph 3.1.2-9 with the additional term

$$\int_0^t \int_0^l \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (3)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a circular cylinder of finite length is given by the formula in Paragraph 3.1.2-10 with the additional term (3).

3°. The solution of the third boundary value problem for a circular cylinder of finite length is the sum of the solution presented in Paragraph 3.1.2-11 and expression (3).

4°. The solutions of mixed boundary value problems for a circular cylinder of finite length are given by the formulas in Paragraph 3.1.2-12 with additional terms of the form (3).

**3.2.2-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for an infinite hollow cylinder is given by the formula in Paragraph 3.1.2-13 with the additional term

$$\int_0^t \int_{-\infty}^{\infty} \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (4)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for an infinite hollow cylinder is given by the formula in Paragraph 3.1.2-14 with the additional term (4).

3°. The solution of the third boundary value problem for an infinite hollow cylinder is the sum of the solution presented in Paragraph 3.1.2-15 and expression (4).

**3.2.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a semiinfinite hollow cylinder is given by the formula in Paragraph 3.1.2-16 with the additional term

$$\int_0^t \int_0^{\infty} \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (5)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a semiinfinite hollow cylinder is given by the formula in Paragraph 3.1.2-17 with the additional term (5).

3°. The solution of the third boundary value problem for a semiinfinite hollow cylinder is the sum of the solution presented in Paragraph 3.1.2-18 and expression (5).

4°. The solutions of mixed boundary value problems for a semiinfinite hollow cylinder are given by the formulas in Paragraph 3.1.2-19 with additional terms of the form (5).

**3.2.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a hollow cylinder of finite length is given by the formula in Paragraph 3.1.2-20 with the additional term

$$\int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (6)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a hollow cylinder of finite length is given by the formula in Paragraph 3.1.2-21 with the additional term (6).

3°. The solution of the third boundary value problem for a hollow cylinder of finite length is the sum of the solution specified in Paragraph 3.1.2-22 and expression (6).

4°. The solutions of mixed boundary value problems for a hollow cylinder of finite length are given by the formulas in Paragraph 3.1.2-23 with additional terms of the form (6).

**3.2.2-7. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $-\infty < z < \infty$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for an infinite wedge domain is given by the formula in Paragraph 3.1.2-24 with the additional term

$$\int_0^t \int_{-\infty}^{\infty} \int_0^{\varphi_0} \int_0^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (7)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for an infinite wedge domain is given by the formula in Paragraph 3.1.2-25 with the additional term (7).

**3.2.2-8. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a semiinfinite wedge domain is given by the formula in Paragraph 3.1.2-26 with the additional term

$$\int_0^t \int_0^{\infty} \int_0^{\varphi_0} \int_0^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (8)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a semiinfinite wedge domain is given by the formula in Paragraph 3.1.2-27 with the additional term (8).

3°. The solutions of mixed boundary value problems for a semiinfinite wedge domain are given by the formulas in Paragraph 3.1.2-28 with additional terms of the form (8).

**3.2.2-9. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a wedge domain of finite height is given by the formula in Paragraph 3.1.2-29 with the additional term

$$\int_0^t \int_0^l \int_0^{\varphi_0} \int_0^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (9)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a wedge domain of finite height is given by the formula in Paragraph 3.1.2-30 with the additional term (9).

3°. The solutions of mixed boundary value problems for a wedge domain of finite height are given by the formulas in Paragraph 3.1.2-31 with additional terms of the form (9).

### 3.2.2-10. Different boundary value problems for a cylindrical sector.

1°. The solution of the first boundary value problem for an unbounded cylindrical sector ( $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $-\infty < z < \infty$ ) is given by the formula in Paragraph 3.1.2-32 with the additional term

$$\int_0^t \int_{-\infty}^{\infty} \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau,$$

which allows for the equation's nonhomogeneity.

2°. The solution of the first boundary value problem for a semibounded cylindrical sector ( $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ ) is given by the formula in Paragraph 3.1.2-33 with the additional term

$$\int_0^t \int_0^{\infty} \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (10)$$

which allows for the equation's nonhomogeneity.

3°. The solution of the mixed boundary value problem for a semibounded cylindrical sector ( $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z < \infty$ ) is given by the formula in Paragraph 3.1.2-34 with the additional term (10).

4°. The solution of the first boundary value problem for a cylindrical sector of finite height ( $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ ) is given by the formula in Paragraph 3.1.2-35 with the additional term

$$\int_0^t \int_0^l \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (11)$$

which allows for the equation's nonhomogeneity.

3°. The solution of a mixed boundary value problem for a cylindrical sector of finite height is given by the formula in Paragraph 3.1.2-36 with the additional term (11).

## 3.2.3. Problems in Spherical Coordinates

In the spherical coordinate system, the heat equation with a volume source has the form

$$\frac{\partial w}{\partial t} = a \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} \right] + \Phi(r, \theta, \varphi, t).$$

One-dimensional problems with central symmetry that have solutions of the form  $w = w(r, t)$  are discussed in Subsection 1.2.4.

### 3.2.3-1. Domain: $0 \leq r \leq R$ , $0 \leq \theta \leq \pi$ , $0 \leq \varphi \leq 2\pi$ . Different boundary value problems.

1°. The solution of the first boundary value problem for a spherical domain is given by the formula in Paragraph 3.1.3-1 with the additional term

$$\int_0^t \int_0^{2\pi} \int_0^{\pi} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta d\xi d\eta d\zeta d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a spherical domain is given by the formula in Paragraph 3.1.3-2 with the additional term (1).

3°. The solution of the third boundary value problem for a spherical domain is the sum of the solution specified in Paragraph 3.1.3-3 and expression (1).

**3.2.3-2. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a spherical layer is given by the formula in Paragraph 3.1.3-4 with the additional term

$$\int_0^t \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a spherical layer is given by the formula in Paragraph 3.1.3-5 with the additional term (2).

3°. The solution of the third boundary value problem for a spherical layer is the sum of the solution specified in Paragraph 3.1.3-6, and expression (2).

**3.2.3-3. Domain:  $0 \leq r < \infty$ ,  $0 \leq \theta \leq \theta_0$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

The solution of the first boundary value problem for an infinite cone is given by the formula in Paragraph 3.1.3-7 with the additional term

$$\int_0^t \int_0^{2\pi} \int_0^{\theta_0} \int_0^\infty \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau,$$

which allows for the equation's nonhomogeneity.

## 3.3. Other Equations with Three Space Variables

### 3.3.1. Equations Containing Arbitrary Parameters

$$1. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + (b_1 x + b_2 y + b_3 z + c)w.$$

The transformation

$$w(x, y, z, t) = \exp \left[ (b_1 x + b_2 y + b_3 z)t + \frac{1}{3}a(b_1^2 + b_2^2 + b_3^2)t^3 + ct \right] u(\xi, \eta, \zeta, t),$$

$$\xi = x + ab_1 t^2, \quad \eta = y + ab_2 t^2, \quad \zeta = z + ab_3 t^2$$

leads to the three-dimensional heat equation  $\partial_t u = a(\partial_{\xi\xi} u + \partial_{\eta\eta} u + \partial_{\zeta\zeta} u)$  that is dealt with in Subsection 3.1.1.

$$2. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) - [b(x^2 + y^2 + z^2) + c]w, \quad b > 0.$$

The transformation ( $A$  is any number)

$$w(x, y, z, t) = \exp \left[ \frac{\sqrt{ab}}{2a} (x^2 + y^2 + z^2) + (3\sqrt{ab} - c) t \right] u(\xi, \eta, \zeta, \tau),$$

$$\xi = x \exp(2\sqrt{ab}t), \quad \eta = y \exp(2\sqrt{ab}t), \quad \zeta = z \exp(2\sqrt{ab}t), \quad \tau = \frac{1}{4\sqrt{ab}} \exp(4\sqrt{ab}t) + A$$

leads to the three-dimensional heat equation  $\partial_\tau u = a(\partial_{\xi\xi} u + \partial_{\eta\eta} u + \partial_{\zeta\zeta} u)$  that is dealt with in Subsection 3.1.1.

$$3. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + [-b(x^2 + y^2 + z^2) + c_1 x + c_2 y + c_3 z + s]w.$$

This is a special case of equation 3.3.2.3 with  $f_k(t) = c_k$ ,  $g(t) = s$ .

$$4. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + b_1 \frac{\partial w}{\partial x} + b_2 \frac{\partial w}{\partial y} + b_3 \frac{\partial w}{\partial z} + cw.$$

This equation governs the nonstationary temperature (concentration) field in a medium moving with a constant velocity, provided there is volume release (absorption) of heat proportional to temperature (concentration).

The substitution

$$w(x, y, z, t) = \exp(A_1 x + A_2 y + A_3 z + Bt) U(x, y, z, t),$$

where

$$A_1 = -\frac{b_1}{2a}, \quad A_2 = -\frac{b_2}{2a}, \quad A_3 = -\frac{b_3}{2a}, \quad B = c - \frac{1}{4a} (b_1^2 + b_2^2 + b_3^2),$$

leads to the three-dimensional heat equation  $\partial_t U = a \Delta_3 U$  that is considered in Subsection 3.1.1.

$$5. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) - by \frac{\partial w}{\partial x}.$$

This equation is encountered in problems of convective heat and mass transfer in a simple shear flow.

Fundamental solution:

$$\mathcal{E}(x, y, z, \xi, \eta, \zeta, t) = \frac{1}{(4\pi at)^{3/2} \left(1 + \frac{1}{12} b^2 t^2\right)^{1/2}} \exp \left\{ -\frac{\left[x - \xi - \frac{1}{2} bt(y + \eta)\right]^2}{4at \left(1 + \frac{1}{12} b^2 t^2\right)} - \frac{(y - \eta)^2 + (z - \zeta)^2}{4at} \right\}.$$

⊙ *Reference:* E. A. Novikov (1958).

$$6. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + b_1 x \frac{\partial w}{\partial x} + b_2 y \frac{\partial w}{\partial x} + b_3 z \frac{\partial w}{\partial z} + \Phi(x, y, z, t).$$

This equation is encountered in problems of convective heat and mass transfer in a linear shear flow.

Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(x, y, z) \quad \text{at} \quad t = 0.$$

Solution:

$$\begin{aligned} w(x, y, z, t) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) dx dy dz \\ & + \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) dx dy dz d\tau, \end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = H(x, \xi, t; b_1) H(y, \eta, t; b_2) H(z, \zeta, t; b_3),$$

$$H(x, \xi, t; b) = \left[ \frac{2\pi a}{b} (e^{2bt} - 1) \right]^{-1/2} \exp \left[ -\frac{b(xe^{bt} - \xi)^2}{2a(e^{2bt} - 1)} \right].$$

$$\begin{aligned} 7. \quad \frac{\partial w}{\partial t} = & a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + (b_1 x + c_1) \frac{\partial w}{\partial x} + (b_2 y + c_2) \frac{\partial w}{\partial y} \\ & + (b_3 z + c_3) \frac{\partial w}{\partial z} + (s_1 x + s_2 y + s_3 z + p)w. \end{aligned}$$

This is a special case of equation 3.3.2.5.



$$8. \quad i\hbar \frac{\partial w}{\partial t} + \frac{\hbar^2}{2m} \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) = 0.$$

Three-dimensional Schrödinger equation,  $i^2 = -1$ .

Fundamental solution:

$$\mathcal{E}(x, y, z, t) = -\frac{i}{\hbar} \left( \frac{m}{2\pi\hbar t} \right)^{3/2} \exp \left[ i \frac{m}{2\hbar t} (x^2 + y^2 + z^2) - i \frac{3\pi}{4} \right].$$

⊙ *Reference:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

$$9. \quad \frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( by^m \frac{\partial w}{\partial y} \right) + \frac{\partial}{\partial z} \left( cz^k \frac{\partial w}{\partial z} \right).$$

This equation describes unsteady heat and mass transfer processes in inhomogeneous (anisotropic) media. It admits separable solutions, as well as solutions with incomplete separation of variables (see Subsection 0.9.2-1). In addition, for  $n \neq 2$ ,  $m \neq 2$ ,  $k \neq 2$  there are particular solutions of the form

$$w = w(\xi, t), \quad \xi^2 = 4 \left[ \frac{x^{2-n}}{a(2-n)^2} + \frac{y^{2-m}}{b(2-m)^2} + \frac{z^{2-k}}{c(2-k)^2} \right],$$

where the function  $w(\xi, t)$  is determined by the one-dimensional nonstationary equation

$$\frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial \xi^2} + \frac{A}{\xi} \frac{\partial w}{\partial \xi}, \quad A = 2 \left( \frac{1}{2-n} + \frac{1}{2-m} + \frac{1}{2-k} \right) - 1.$$

For solutions of this equation, see Subsections 1.2.1, 1.2.3, and 1.2.5.

### 3.3.2. Equations Containing Arbitrary Functions

$$1. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + f(t)w.$$

This equation describes three-dimensional unsteady thermal phenomena in quiescent media or solids with constant thermal diffusivity, provided there is unsteady volume heat release proportional to temperature.

The substitution  $w(x, y, z, t) = \exp \left[ \int f(t) dt \right] U(x, y, z, t)$  leads to the usual heat equation  $\partial_t U = a \Delta_3 U$  that is dealt with in Subsection 3.1.1.

$$2. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + [x f_1(t) + y f_2(t) + z f_3(t) + g(t)] w.$$

The transformation

$$w(x, y, z, t) = \exp \left\{ x F_1(t) + y F_2(t) + z F_3(t) + a \int [F_1^2(t) + F_2^2(t) + F_3^2(t)] dt + \int g(t) dt \right\} u(\xi, \eta, \zeta, t),$$

$$\xi = x + 2a \int F_1(t) dt, \quad \eta = y + 2a \int F_2(t) dt, \quad \zeta = z + 2a \int F_3(t) dt, \quad F_k(t) = \int f_k(t) dt,$$

leads to the three-dimensional heat equation  $\partial_t u = a(\partial_{\xi\xi} u + \partial_{\eta\eta} u + \partial_{\zeta\zeta} u)$  that is dealt with in Subsection 3.1.1.

$$3. \quad \frac{\partial w}{\partial t} = a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + [-b(x^2 + y^2 + z^2) + x f_1(t) + y f_2(t) + z f_3(t) + g(t)] w.$$

1°. Case  $b > 0$ . The transformation

$$w(x, y, z, t) = u(\xi, \eta, \zeta, \tau) \exp \left[ \frac{\sqrt{ab}}{2a} (x^2 + y^2 + z^2) \right],$$

$$\xi = x \exp(2\sqrt{ab}t), \quad \eta = y \exp(2\sqrt{ab}t), \quad \zeta = z \exp(2\sqrt{ab}t), \quad \tau = \frac{1}{4\sqrt{ab}} \exp(4\sqrt{ab}t) + A,$$

where  $A$  is an arbitrary constant, leads to an equation of the form 3.3.2.2:

$$\frac{\partial u}{\partial \tau} = a \left( \frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \eta^2} + \frac{\partial^2 u}{\partial \zeta^2} \right) + [\xi F_1(\tau) + \eta F_2(\tau) + \zeta F_3(\tau) + G(\tau)] u,$$

$$F_k(\tau) = \frac{1}{(c\tau)^{3/2}} f_k \left( \frac{\ln(c\tau)}{c} \right), \quad G(\tau) = \frac{1}{c\tau} g \left( \frac{\ln(c\tau)}{c} \right) + \frac{3}{4\tau}, \quad c = 4\sqrt{ab}, \quad k = 1, 2, 3.$$

2°. Case  $b < 0$ . The transformation

$$w(x, y, z, t) = v(\xi_1, \eta_1, \zeta_1, \tau_1) \exp \left[ \frac{\sqrt{-ab}}{2a} (x^2 + y^2 + z^2) \tan(2\sqrt{-ab}t) \right],$$

$$\xi_1 = \frac{x}{\cos(2\sqrt{-ab}t)}, \quad \eta_1 = \frac{y}{\cos(2\sqrt{-ab}t)}, \quad \zeta_1 = \frac{z}{\cos(2\sqrt{-ab}t)}, \quad \tau_1 = \frac{1}{2\sqrt{-ab}} \tan(2\sqrt{-ab}t)$$

also leads to an equation of the form 3.3.2.2 for  $v = v(\xi_1, \eta_1, \zeta_1, \tau_1)$  (the equation for  $v$  is not written out here).

$$4. \quad \frac{\partial w}{\partial t} = a_1(t) \frac{\partial^2 w}{\partial x^2} + a_2(t) \frac{\partial^2 w}{\partial y^2} + a_3(t) \frac{\partial^2 w}{\partial z^2} + \Phi(x, y, z, t).$$

Here,  $0 < a_k(t) < \infty$ ;  $k = 1, 2, 3$ .

Domain:  $-\infty < x < \infty, -\infty < y < \infty, -\infty < z < \infty$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(x, y, z) \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, y, z, t) = \int_0^t \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t, \tau) d\xi d\eta d\zeta d\tau$$

$$+ \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t, 0) d\xi d\eta d\zeta,$$

where

$$G(x, y, z, \xi, \eta, \zeta, t, \tau) = \frac{1}{8\pi^{3/2}\sqrt{T_1 T_2 T_3}} \exp \left[ -\frac{(x-\xi)^2}{4T_1} - \frac{(y-\eta)^2}{4T_2} - \frac{(z-\zeta)^2}{4T_3} \right],$$

$$T_1 = \int_{\tau}^t a_1(\sigma) d\sigma, \quad T_2 = \int_{\tau}^t a_2(\sigma) d\sigma, \quad T_3 = \int_{\tau}^t a_3(\sigma) d\sigma.$$

See also the more general equation 3.4.3.3, where other boundary value problems are considered.

$$5. \quad \frac{\partial w}{\partial t} = a_1(t) \frac{\partial^2 w}{\partial x^2} + a_2(t) \frac{\partial^2 w}{\partial y^2} + a_3(t) \frac{\partial^2 w}{\partial z^2} + [b_1(t)x + c_1(t)] \frac{\partial w}{\partial x} \\ + [b_2(t)y + c_2(t)] \frac{\partial w}{\partial y} + [b_3(t)z + c_3(t)] \frac{\partial w}{\partial z} + [s_1(t)x + s_2(t)y + s_3(t)z + p(t)] w.$$

The transformation

$$w(x, y, z, t) = \exp[f_1(t)x + f_2(t)y + f_3(t)z + g(t)] u(\xi, \eta, \zeta, t), \\ \xi = h_1(t)x + r_1(t), \quad \eta = h_2(t)y + r_2(t), \quad \zeta = h_3(t)z + r_3(t),$$

where

$$h_k(t) = A_k \exp \left[ \int b_k(t) dt \right], \\ f_k(t) = h_k(t) \int \frac{s_k(t)}{h_k(t)} dt + B_k h_k(t), \\ r_k(t) = \int [2a_k(t)f_k(t) + c_k(t)] h_k(t) dt + C_k, \\ g(t) = \int \sum_{k=1}^3 [a_k(t)f_k^2(t) + c_k(t)f_k(t)] dt + \int p(t) dt + D,$$

( $k = 1, 2, 3$ ;  $A_k, B_k, C_k, D$  are arbitrary constants) leads to an equation of the form 3.3.2.4:

$$\frac{\partial u}{\partial t} = a_1(t)h_1^2(t) \frac{\partial^2 u}{\partial \xi^2} + a_2(t)h_2^2(t) \frac{\partial^2 u}{\partial \eta^2} + a_3(t)h_3^2(t) \frac{\partial^2 u}{\partial \zeta^2}.$$

$$6. \quad \frac{\partial w}{\partial t} = a_1(t) \frac{\partial^2 w}{\partial x^2} + a_2(t) \frac{\partial^2 w}{\partial y^2} + a_3(t) \frac{\partial^2 w}{\partial z^2} + [b_1(t)x + c_1(t)] \frac{\partial w}{\partial x} + [b_2(t)y + c_2(t)] \frac{\partial w}{\partial y} \\ + [b_3(t)z + c_3(t)] \frac{\partial w}{\partial z} + [s_1(t)x^2 + s_2(t)y^2 + s_3(t)z^2 + p_1(t)x + p_2(t)y + p_3(t)z + q(t)] w.$$

The substitution

$$w(x, y, z, t) = \exp[f_1(t)x^2 + f_2(t)y^2 + f_3(t)z^2] u(x, y, z, t),$$

where the functions  $f_k = f_k(t)$  are solutions of the respective Riccati equations

$$f'_k = 4a_k(t)f_k^2 + 2b_k(t)f_k + s_k(t) \quad (k = 1, 2, 3),$$

leads to an equation of the form 3.3.2.5 for  $u = u(x, y, z, t)$ .

$$7. \quad \frac{\partial w}{\partial t} + \sum_{n=1}^2 [f_n(t) + g_n(t)x_3] \frac{\partial w}{\partial x_n} - a \frac{\partial w}{\partial x_3} = \sum_{n,m=1}^2 K_{nm}(t) \frac{\partial^2 w}{\partial x_n \partial x_m} + K_{33}(t) \frac{\partial^2 w}{\partial x_3^2}.$$

*Equation of turbulent diffusion.* It describes the diffusion of an admixture in a horizontal stream whose velocity components are linear functions of the height.

Fundamental solution:

$$\mathcal{C}(x_1, x_2, x_3, \xi_1, \xi_2, \xi_3) = \frac{1}{(4\pi)^{3/2} \sqrt{\det |T|}} \exp \left[ -\frac{1}{4} \sum_{i,j=1}^3 T_{ij}^{-1}(t) y_i y_j \right], \quad T_{ij}(t) = \int_0^t S_{ij}(\tau) d\tau.$$

Here, the following notation is used ( $n, m = 1, 2$ ):

$$y_n = x_n - \xi_n - F_n(t) - x_3 G_n(t) - a \int_0^t (t - \tau) g_n(\tau) d\tau, \quad y_3 = x_3 - \xi_3 + at, \\ S_{nm}(t) = K_{nm}(t) + K_{33}(t) G_n(t) G_m(t), \quad S_{n3}(t) = S_{3n}(t) = -K_{33}(t) G_n(t), \\ S_{33}(t) = K_{33}(t), \quad F_n(t) = \int_0^t f_n(t) dt, \quad G_n(t) = \int_0^t g_n(t) dt,$$

$\det |T|$  is the determinant of the matrix  $\mathbf{T}$  with entries  $T_{ij}(t)$ ,  $T_{ij}^{-1}(t)$  are the entries of the inverse of  $\mathbf{T}$ . The inequalities  $T_{11}(t) > 0$ ,  $T_{11}(t)T_{22}(t) - T_{12}^2(t) > 0$ , and  $\det |T| > 0$  are assumed to hold.

⊙ *Reference:* E. A. Novikov (1958).

### 3.3.3. Equations of the Form

$$\rho(x, y, z) \frac{\partial w}{\partial t} = \operatorname{div}[a(x, y, z) \nabla w] - q(x, y, z)w + \Phi(x, y, z, t)$$

Equations of this form are often encountered in the theory of heat and mass transfer. For brevity, the following notation is used:

$$\operatorname{div}[a(\mathbf{r}) \nabla w] = \frac{\partial}{\partial x} \left[ a(\mathbf{r}) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ a(\mathbf{r}) \frac{\partial w}{\partial y} \right] + \frac{\partial}{\partial z} \left[ a(\mathbf{r}) \frac{\partial w}{\partial z} \right], \quad \mathbf{r} = \{x, y, z\}.$$

The problems presented in this subsection are assumed to refer to a simply connected bounded domain  $V$  with smooth boundary  $S$ . It is also assumed that  $\rho(\mathbf{r}) > 0$ ,  $a(\mathbf{r}) > 0$ , and  $q(\mathbf{r}) \geq 0$ .

#### 3.3.3-1. First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{r}) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(\mathbf{r}, t) \quad \text{for} \quad \mathbf{r} \in S \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{r}, t) &= \int_0^t \int_V \Phi(\boldsymbol{\xi}, \tau) \mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t - \tau) dV_{\boldsymbol{\xi}} d\tau + \int_V f(\boldsymbol{\xi}) \rho(\boldsymbol{\xi}) \mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t) dV_{\boldsymbol{\xi}} \\ &\quad - \int_0^t \int_S g(\boldsymbol{\xi}, \tau) a(\boldsymbol{\xi}) \left[ \frac{\partial}{\partial N_{\boldsymbol{\xi}}} \mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t - \tau) \right] dS_{\boldsymbol{\xi}} d\tau. \end{aligned} \quad (1)$$

Here, the modified Green's function is given by

$$\mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t) = \sum_{n=1}^{\infty} \frac{u_n(\mathbf{r}) u_n(\boldsymbol{\xi})}{\|u_n\|^2} \exp(-\lambda_n t), \quad \|u_n\|^2 = \int_V \rho(\mathbf{r}) u_n^2(\mathbf{r}) dV, \quad \boldsymbol{\xi} = \{\xi_1, \xi_2, \xi_3\}, \quad (2)$$

where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the following elliptic second-order equation with a homogeneous boundary condition of the first kind:

$$\operatorname{div}[a(\mathbf{r}) \nabla u] - q(\mathbf{r})u + \lambda \rho(\mathbf{r})u = 0, \quad (3)$$

$$u = 0 \quad \text{for} \quad \mathbf{r} \in S. \quad (4)$$

The integration in solution (1) is carried out with respect to  $\xi_1, \xi_2, \xi_3$ ;  $\frac{\partial}{\partial N_{\boldsymbol{\xi}}}$  denotes the derivative along the outward normal to the surface  $S$  with respect to  $\xi_1, \xi_2$ , and  $\xi_3$ .

General properties of the Sturm–Liouville problem (3)–(4):

1°. There are countably many eigenvalues. All eigenvalues are real and can be ordered so that  $\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$ , with  $\lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ ; consequently, there can exist only finitely many negative eigenvalues.

2°. For  $\rho(\mathbf{r}) > 0$ ,  $a(\mathbf{r}) > 0$ , and  $q(\mathbf{r}) \geq 0$ , all eigenvalues are positive:  $\lambda_n > 0$ .

3°. The eigenfunctions are defined up to a constant multiplier. Any two eigenfunctions  $u_n(\mathbf{r})$  and  $u_m(\mathbf{r})$  corresponding to different eigenvalues  $\lambda_n$  and  $\lambda_m$  are orthogonal with weight  $\rho(\mathbf{r})$  in the domain  $V$ :

$$\int_V \rho(\mathbf{r}) u_n(\mathbf{r}) u_m(\mathbf{r}) dV = 0 \quad \text{for} \quad n \neq m.$$

4°. An arbitrary function  $F(\mathbf{r})$  that is twice continuously differentiable and satisfies the boundary condition of the Sturm–Liouville problem ( $F = 0$  for  $\mathbf{r} \in S$ ) can be expanded into an absolutely and uniformly convergent series in the eigenvalues:

$$F(\mathbf{r}) = \sum_{n=1}^{\infty} F_n u_n(\mathbf{r}), \quad F_n = \frac{1}{\|u_n\|^2} \int_V F(\mathbf{r}) \rho(\mathbf{r}) u_n(\mathbf{r}) dV,$$

where the formula for  $\|u_n\|^2$  is given in (2).

*Remark.* In a three-dimensional problem, to each eigenvalue  $\lambda_n$  there generally correspond finitely many linearly independent eigenfunctions  $u_n^{(1)}, u_n^{(2)}, \dots, u_n^{(m)}$ . These function can always be replaced by their linear combinations

$$\bar{u}_n^{(k)} = A_{k,1} u_n^{(1)} + \dots + A_{k,k-1} u_n^{(k-1)} + u_n^{(k)}, \quad k = 1, 2, \dots, m,$$

such that  $\bar{u}_n^{(1)}, \bar{u}_n^{(2)}, \dots, \bar{u}_n^{(m)}$  are now pairwise orthogonal. Thus, without loss of generality, we assume that all eigenfunctions are orthogonal.

### 3.3.3-2. Second boundary value problem.

The following conditions are prescribed:

$$w = f(\mathbf{r}) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\frac{\partial w}{\partial N} = g(\mathbf{r}, t) \quad \text{for} \quad \mathbf{r} \in S \quad (\text{boundary condition}).$$

Solution:

$$\begin{aligned} w(\mathbf{r}, t) = & \int_0^t \int_V \Phi(\xi, \tau) \mathcal{G}(\mathbf{r}, \xi, t - \tau) dV_\xi d\tau + \int_V f(\xi) \rho(\xi) \mathcal{G}(\mathbf{r}, \xi, t) dV_\xi \\ & + \int_0^t \int_S g(\xi, \tau) a(\xi) \mathcal{G}(\mathbf{r}, \xi, t - \tau) dS_\xi d\tau. \end{aligned} \quad (5)$$

Here, the modified Green's function is given by (2), where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the elliptic second-order equation (3) with a homogeneous boundary condition of the second kind,

$$\frac{\partial u}{\partial N} = 0 \quad \text{for} \quad \mathbf{r} \in S. \quad (6)$$

For  $q(\mathbf{r}) > 0$  the general properties of the eigenvalue problem (3), (6) are the same as for the first boundary value problem (with all  $\lambda_n > 0$ ).

### 3.3.3-3. Third boundary value problem.

The following conditions are prescribed:

$$w = f(\mathbf{r}) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\frac{\partial w}{\partial N} + k(\mathbf{r})w = g(\mathbf{r}, t) \quad \text{for} \quad \mathbf{r} \in S \quad (\text{boundary condition}).$$

The solution of the third boundary value problem is given by formulas (5) and (2), where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the second-order elliptic equation (3) with a homogeneous boundary condition of the third kind,

$$\frac{\partial u}{\partial N} + k(\mathbf{r})u = 0 \quad \text{for} \quad \mathbf{r} \in S. \quad (7)$$

For  $q(\mathbf{r}) \geq 0$  and  $k(\mathbf{r}) > 0$  the general properties of the eigenvalue problem (3), (7) are the same as for the first boundary value problem (see Paragraph 3.3.3-1).

Let  $k(\mathbf{r}) = k = \text{const}$ . Denote the Green's functions of the second and third boundary value problems by  $G_2(\mathbf{r}, \xi, t)$  and  $G_3(\mathbf{r}, \xi, t, k)$ , respectively. If  $q(\mathbf{r}) > 0$ , then the following limiting relation holds:  $G_2(\mathbf{r}, \xi, t) = \lim_{k \rightarrow 0} G_3(\mathbf{r}, \xi, t, k)$ .

© *References for Subsection 3.3.3:* V. S. Vladimirov (1988), A. D. Polyanin (2000a, 2000c).

## 3.4. Equations with $n$ Space Variables

### 3.4.1. Equations of the Form $\frac{\partial w}{\partial t} = a\Delta_n w + \Phi(x_1, \dots, x_n, t)$

This is an  $n$ -dimensional nonhomogeneous heat equation. In the Cartesian system of coordinates, it is represented as

$$\frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} + \Phi(\mathbf{x}, t), \quad \mathbf{x} = \{x_1, \dots, x_n\}.$$

The solutions of various problems for this equation can be constructed on the basis of incomplete separation of variables (see Paragraphs 0.6.1-2 and 0.9.2-1) taking into account the results of Subsections 1.1.1 and 1.1.2. Some examples of solving such problems can be found below in Paragraphs 3.4.1-2 through 3.4.1-4.

#### 3.4.1-1. Homogeneous equation ( $\Phi \equiv 0$ ).

1°. Particular solutions:

$$\begin{aligned} w(\mathbf{x}, t) &= A \exp\left(\sum_{m=1}^n k_m x_m + at \sum_{m=1}^n k_m^2\right), \\ w(\mathbf{x}, t) &= A \exp\left(-at \sum_{m=1}^n k_m^2\right) \prod_{m=1}^n \cos(k_m x_m + C_m), \\ w(\mathbf{x}, t) &= A \exp\left(-\sum_{m=1}^n k_m x_m\right) \prod_{m=1}^n \cos(k_m x_m - 2ak_m^2 t + C_m), \\ w(\mathbf{x}, t) &= \frac{A}{(t-t_0)^{n/2}} \exp\left[-\frac{1}{4a(t-t_0)} \sum_{m=1}^n (x_m - C_m)^2\right], \\ w(\mathbf{x}, t) &= A \prod_{m=1}^n \operatorname{erf}\left(\frac{x_m - C_m}{2\sqrt{at}}\right), \end{aligned}$$

where  $\mathbf{x} = \{x_1, \dots, x_n\}$ ;  $A$ ,  $k_m$ ,  $C_m$ , and  $t_0$  are arbitrary constants.

2°. Fundamental solution:

$$\mathcal{E}(\mathbf{x}, t) = \frac{1}{(2\sqrt{\pi at})^n} \exp\left(-\frac{|\mathbf{x}|^2}{4at}\right), \quad |\mathbf{x}|^2 = \sum_{k=1}^n x_k^2.$$

3°. Suppose  $w = w(x_1, \dots, x_n, t)$  is a solution of the homogeneous equation. Then the functions

$$\begin{aligned} w_1 &= Aw(\pm\lambda x_1 + C_1, \dots, \pm\lambda x_n + C_n, \lambda^2 t + C_{n+1}), \\ w_2 &= A \exp\left(\sum_{k=1}^n \lambda_k x_k + at \sum_{k=1}^n \lambda_k^2\right) w(x_1 + 2a\lambda_1 t + C_1, \dots, x_n + 2a\lambda_n t + C_n, t + C_{n+1}), \\ w_3 &= \frac{A}{|\delta + \beta t|^{n/2}} \exp\left[-\frac{\beta}{4a(\delta + \beta t)} \sum_{k=1}^n x_k^2\right] w\left(\frac{x_1}{\delta + \beta t}, \dots, \frac{x_n}{\delta + \beta t}, \frac{\gamma + \lambda t}{\delta + \beta t}\right), \quad \lambda\delta - \beta\gamma = 1, \end{aligned}$$

where  $A$ ,  $C_1, \dots, C_{n+1}$ ,  $\lambda$ ,  $\lambda_1, \dots, \lambda_n$ ,  $\beta$ , and  $\delta$  are arbitrary constants, are also solutions of the equation. The signs at  $\lambda$  in the formula for  $w_1$  can be taken independently of one another.

3.4.1-2. Domain:  $\mathbb{R}^n = \{-\infty < x_k < \infty; k = 1, \dots, n\}$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(\mathbf{x}) \quad \text{at} \quad t = 0.$$

Solution:

$$w(\mathbf{x}, t) = \frac{1}{(2\sqrt{\pi at})^n} \int_{\mathbb{R}^n} f(\mathbf{y}) \exp\left(-\frac{|\mathbf{x}-\mathbf{y}|^2}{4at}\right) d\mathbf{y} + \int_0^t \int_{\mathbb{R}^n} \frac{\Phi(\mathbf{y}, \tau)}{(2\sqrt{\pi a(t-\tau)})^n} \exp\left(-\frac{|\mathbf{x}-\mathbf{y}|^2}{4a(t-\tau)}\right) d\mathbf{y} d\tau,$$

where  $\mathbf{y} = \{y_1, \dots, y_n\}$ ,  $|\mathbf{x}-\mathbf{y}| = \sqrt{(x_1 - y_1)^2 + \dots + (x_n - y_n)^2}$ ,  $d\mathbf{y} = dy_1 dy_2 \dots dy_n$ .

⊙ Reference: V. S. Vladimirov (1988).

3.4.1-3. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) & \text{at} \quad t &= 0 & \text{(initial condition),} \\ w &= g_k(\mathbf{x}, t) & \text{at} \quad x_k &= 0 & \text{(boundary conditions),} \\ w &= h_k(\mathbf{x}, t) & \text{at} \quad x_k &= l_k & \text{(boundary conditions).} \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ &+ a \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\ &- a \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ h_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau, \end{aligned}$$

where the following notation is used:

$$dS_y^{(k)} = dy_1 \dots dy_{k-1} dy_{k+1} \dots dy_n, \quad S^{(k)} = \{0 \leq y_m \leq l_m \text{ for } m = 1, \dots, k-1, k+1, \dots, n\}.$$

The Green's function can be represented in the product form

$$G(\mathbf{x}, \mathbf{y}, t) = \prod_{k=1}^n G_k(x_k, y_k, t), \quad (1)$$

where the  $G_k(x_k, y_k, t)$  are the Green's functions of the respective one-dimensional boundary value problems (see Paragraph 1.1.2-5):

$$G_k(x_k, y_k, t) = \frac{2}{l_k} \sum_{m=1}^{\infty} \sin\left(\frac{m\pi x}{l_k}\right) \sin\left(\frac{m\pi y}{l_k}\right) \exp\left(-\frac{am^2\pi^2 t}{l_k^2}\right).$$

3.4.1-4. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) & \text{at} \quad t &= 0 & \text{(initial condition),} \\ \partial_{x_k} w &= g_k(\mathbf{x}, t) & \text{at} \quad x_k &= 0 & \text{(boundary conditions),} \\ \partial_{x_k} w &= h_k(\mathbf{x}, t) & \text{at} \quad x_k &= l_k & \text{(boundary conditions).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(\mathbf{x}, t) = & \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\
 & - a \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [g_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=0} dS_y^{(k)} d\tau \\
 & + a \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=l_k} dS_y^{(k)} d\tau.
 \end{aligned} \tag{2}$$

The Green's function can be represented as the product (1) of the corresponding one-dimensional Green's functions of the form (see Paragraph 1.1.2-6)

$$G_k(x_k, y_k, t) = \frac{1}{l_k} + \frac{2}{l_k} \sum_{m=1}^{\infty} \cos\left(\frac{m\pi x}{l_k}\right) \cos\left(\frac{m\pi y}{l_k}\right) \exp\left(-\frac{am^2\pi^2 t}{l_k^2}\right).$$

### 3.4.2. Other Equations Containing Arbitrary Parameters

$$1. \quad \frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} + \left(c + \sum_{k=1}^n b_k x_k\right) w.$$

This is a special case of equation 3.4.3.1. The transformation

$$w(x_1, \dots, x_n, t) = \exp\left(t \sum_{k=1}^n b_k x_k + \frac{1}{3} a t^3 \sum_{k=1}^n b_k^2 + c t\right) u(\xi_1, \dots, \xi_n, t), \quad \xi_k = x_k + a b_k t^2$$

leads to the  $n$ -dimensional heat equation  $\partial_t u = a \sum_{k=1}^n \partial_{\xi_k \xi_k} u$  that is dealt with in Subsection 3.4.1.

$$2. \quad \frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} - \left(c + b \sum_{k=1}^n x_k^2\right) w, \quad b > 0.$$

The transformation ( $A$  is any number)

$$\begin{aligned}
 w(x_1, \dots, x_n, t) = & u(\xi_1, \dots, \xi_n, \tau) \exp\left[\frac{1}{2} \sqrt{\frac{b}{a}} \sum_{k=1}^n x_k^2 + (n\sqrt{ab} - c) t\right], \\
 \xi_1 = & x_1 \exp(2\sqrt{ab} t), \quad \dots, \quad \xi_n = x_n \exp(2\sqrt{ab} t), \quad \tau = \frac{1}{4\sqrt{ab}} \exp(4\sqrt{ab} t) + A
 \end{aligned}$$

leads to the  $n$ -dimensional heat equation  $\partial_\tau u = a \sum_{k=1}^n \partial_{\xi_k \xi_k} u$  that is dealt with in Subsection 3.4.1.

$$3. \quad \frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} + \left(-b \sum_{k=1}^n x_k^2 + \sum_{k=1}^n c_k x_k + s\right) w.$$

This is a special case of equation 3.4.3.2 with  $f_k(t) = c_k$  and  $g(t) = s$ .

$$4. \quad \frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} + \sum_{k=1}^n b_k \frac{\partial w}{\partial x_k} + c w.$$

The substitution

$$w(x_1, \dots, x_n, t) = \exp\left(At - \frac{1}{2a} \sum_{k=1}^n b_k x_k\right) U(x_1, \dots, x_n, t), \quad \text{where} \quad A = c - \frac{1}{4a} \sum_{k=1}^n b_k^2,$$

leads to the  $n$ -dimensional heat equation  $\partial_t U = a \Delta_n U$  that is dealt with in Subsection 3.4.1.



$$5. \quad \frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} + \sum_{k=1}^n (b_k x_k + c_k) \frac{\partial w}{\partial x_k} + \left( \sum_{k=1}^n s_k x_k + p \right) w.$$

This is a special case of equation 3.4.3.4.

$$6. \quad i\hbar \frac{\partial w}{\partial t} + \frac{\hbar^2}{2m} \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} = 0.$$

This is the  $n$ -dimensional Schrödinger equation,  $i^2 = -1$ .

Fundamental solution:

$$\mathcal{E}(\mathbf{x}, t) = -\frac{i}{\hbar} \left( \frac{m}{2\pi\hbar t} \right)^{n/2} \exp \left( i \frac{m}{2\hbar t} |\mathbf{x}|^2 - i \frac{\pi n}{4} \right), \quad |\mathbf{x}|^2 = x_1^2 + \cdots + x_n^2.$$

⊙ Reference: V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

### 3.4.3. Equations Containing Arbitrary Functions

$$1. \quad \frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} + \left[ \sum_{k=1}^n x_k f_k(t) + g(t) \right] w.$$

The transformation

$$w(x_1, \dots, x_n, t) = \exp \left[ \sum_{k=1}^n x_k F_k(t) + a \sum_{k=1}^n \int F_k^2(t) dt + G(t) \right] u(\xi_1, \dots, \xi_n, t),$$

$$\xi_k = x_k + 2a \int F_k(t) dt, \quad F_k(t) = \int f_k(t) dt, \quad G(t) = \int g(t) dt,$$

leads to the  $n$ -dimensional heat equation  $\partial_t u = a \sum_{k=1}^n \partial_{\xi_k \xi_k} u$  that is discussed in Subsection 3.4.1.

$$2. \quad \frac{\partial w}{\partial t} = a \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2} + \left[ -b \sum_{k=1}^n x_k^2 + \sum_{k=1}^n x_k f_k(t) + g(t) \right] w.$$

1°. Case  $b > 0$ . The transformation

$$w(x_1, \dots, x_n, t) = u(\xi_1, \dots, \xi_n, \tau) \exp \left( \frac{1}{2} \sqrt{\frac{b}{a}} \sum_{k=1}^n x_k^2 \right),$$

$$\xi_1 = x_1 \exp(2\sqrt{ab}t), \quad \dots, \quad \xi_n = x_n \exp(2\sqrt{ab}t), \quad \tau = \frac{1}{4\sqrt{ab}} \exp(4\sqrt{ab}t) + C,$$

where  $C$  is an arbitrary constant, leads to an equation of the form 3.4.3.1:

$$\frac{\partial u}{\partial \tau} = a \sum_{k=1}^n \frac{\partial^2 u}{\partial \xi_k^2} + \left[ \sum_{k=1}^n \xi_k F_k(\tau) + G(\tau) \right] u,$$

$$F_k(\tau) = \frac{1}{(s\tau)^{3/2}} f_k \left( \frac{\ln(s\tau)}{s} \right), \quad G(\tau) = \frac{1}{s\tau} g \left( \frac{\ln(s\tau)}{s} \right) + \frac{n}{4\tau}, \quad s = 4\sqrt{ab}.$$

2°. Case  $b < 0$ . The transformation

$$w(x_1, \dots, x_n, t) = v(z_1, \dots, z_n, \tau) \exp \left[ \frac{\sqrt{-b}}{2\sqrt{a}} \tan(2\sqrt{-ab}t) \sum_{k=1}^n x_k^2 \right],$$

$$z_1 = \frac{x_1}{\cos(2\sqrt{-ab}t)}, \quad \dots, \quad z_n = \frac{x_n}{\cos(2\sqrt{-ab}t)}, \quad \tau = \frac{1}{2\sqrt{-ab}} \tan(2\sqrt{-ab}t)$$

also leads to an equation of the form 3.4.3.1 (this equation is not specified here).

$$3. \quad \frac{\partial w}{\partial t} = \sum_{k=1}^n a_k(t) \frac{\partial^2 w}{\partial x_k^2} + \Phi(x_1, \dots, x_n, t).$$

The solutions of various problems for this equation can be constructed on the basis of incomplete separation of variables (see Paragraphs 0.6.1-2 and 0.9.2-1) taking into account the results of Subsections 1.1.1 and 1.1.2. Some examples of solving such problems are given below. It is assumed that  $0 < a_k(t) < \infty$ ,  $k = 1, \dots, n$ .

1°. Domain:  $\mathbb{R}^n = \{-\infty < x_k < \infty; k = 1, \dots, n\}$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(\mathbf{x}) \quad \text{at} \quad t = 0.$$

Solution:

$$w(\mathbf{x}, t) = \int_0^t \int_{\mathbb{R}^n} \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau + \int_{\mathbb{R}^n} f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t, 0) d\mathbf{y},$$

where

$$G(\mathbf{x}, \mathbf{y}, t, \tau) = \frac{1}{2^{n/2} \pi^{n/2} \sqrt{T_1 T_2 \dots T_n}} \exp \left[ - \sum_{k=1}^n \frac{(x_k - y_k)^2}{4T_k} \right], \quad T_k = \int_{\tau}^t a_k(\eta) d\eta,$$

$$\mathbf{x} = \{x_1, \dots, x_n\}, \quad \mathbf{y} = \{y_1, \dots, y_n\}, \quad d\mathbf{y} = dy_1 dy_2 \dots dy_n.$$

2°. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) & \text{at} \quad t &= 0 & \text{(initial condition),} \\ w &= g_k(\mathbf{x}, t) & \text{at} \quad x_k &= 0 & \text{(boundary conditions),} \\ w &= h_k(\mathbf{x}, t) & \text{at} \quad x_k &= l_k & \text{(boundary conditions).} \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t, 0) d\mathbf{y} \\ &+ \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\tau) \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t, \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\ &- \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\tau) \left[ h_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t, \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau, \end{aligned}$$

where the following notation is used:

$$dS_y^{(k)} = dy_1 \dots dy_{k-1} dy_{k+1} \dots dy_n, \quad S^{(k)} = \{0 \leq y_m \leq l_m \text{ for } m = 1, \dots, k-1, k+1, \dots, n\}.$$

The Green's function can be represented in the product form

$$G(\mathbf{x}, \mathbf{y}, t, \tau) = \prod_{k=1}^n G_k(x_k, y_k, t, \tau), \quad (1)$$

where the  $G_k(x_k, y_k, t, \tau)$  are the Green's functions of the respective boundary value problems,

$$G_k(x_k, y_k, t, \tau) = \frac{2}{l_k} \sum_{m=1}^{\infty} \sin \left( \frac{m\pi x_k}{l_k} \right) \sin \left( \frac{m\pi y_k}{l_k} \right) \exp \left( - \frac{m^2 \pi^2 T_k}{l_k^2} \right), \quad T_k = \int_{\tau}^t a_k(\sigma) d\sigma. \quad (2)$$

3°. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_{x_k} w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ &\quad - \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\tau) [g_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau)]_{y_k=0} dS_y^{(k)} d\tau \\ &\quad + \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\tau) [h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau)]_{y_k=l_k} dS_y^{(k)} d\tau. \end{aligned}$$

The Green's function can be represented as the product (1) of the corresponding one-dimensional Green's functions

$$G_k(x_k, y_k, t, \tau) = \frac{1}{l_k} + \frac{2}{l_k} \sum_{m=1}^{\infty} \cos\left(\frac{m\pi x_k}{l_k}\right) \cos\left(\frac{m\pi y_k}{l_k}\right) \exp\left(-\frac{m^2 \pi^2 T_k}{l_k^2}\right), \quad T_k = \int_{\tau}^t a_k(\sigma) d\sigma.$$

⊙ Reference: A. D. Polyanin (2000a, 2000b).

$$4. \quad \frac{\partial w}{\partial t} = \sum_{k=1}^n a_k(t) \frac{\partial^2 w}{\partial x_k^2} + \sum_{k=1}^n [b_k(t)x_k + c_k(t)] \frac{\partial w}{\partial x_k} + \left[ \sum_{k=1}^n s_k(t)x_k + p(t) \right] w.$$

Let us perform the transformation

$$w(x_1, \dots, x_n, t) = \exp \left[ \sum_{k=1}^n f_k(t)x_k + g(t) \right] u(z_1, \dots, z_n, t), \quad z_k = h_k(t)x_k + r_k(t),$$

where the functions  $f_k(t)$ ,  $g(t)$ ,  $h_k(t)$ , and  $r_k(t)$  are given by ( $A_k$ ,  $B_k$ ,  $C_k$ , and  $D$  are arbitrary constants):

$$\begin{aligned} h_k(t) &= A_k \exp \left[ \int b_k(t) dt \right], \\ f_k(t) &= h_k(t) \int \frac{s_k(t)}{h_k(t)} dt + B_k h_k(t), \\ r_k(t) &= \int [2a_k(t)f_k(t) + c_k(t)] h_k(t) dt + C_k, \\ g(t) &= \int \left[ p(t) + \sum_{k=1}^n a_k(t)f_k^2(t) + \sum_{k=1}^n c_k(t)f_k(t) \right] dt + D. \end{aligned}$$

As a result, we arrive at an equation of the form 3.4.3.3 for the new dependent variable  $u = u(z_1, \dots, z_n, t)$ :

$$\frac{\partial u}{\partial t} = \sum_{k=1}^n a_k(t) h_k^2(t) \frac{\partial^2 u}{\partial z_k^2}.$$

$$5. \quad \frac{\partial w}{\partial t} = \sum_{k=1}^n a_k(t) \frac{\partial^2 w}{\partial x_k^2} + \sum_{k=1}^n [b_k(t)x_k + c_k(t)] \frac{\partial w}{\partial x_k} + \left[ \sum_{k=1}^n s_k(t)x_k^2 + \sum_{k=1}^n p_k(t)x_k + q(t) \right] w.$$

The substitution

$$w(x_1, \dots, x_n, t) = \exp \left[ \sum_{k=1}^n f_k(t)x_k^2 \right] u(x_1, \dots, x_n, t),$$

where the functions  $f_k = f_k(t)$  are solutions of the Riccati equation

$$f'_k = 4a_k(t)f_k^2 + 2b_k(t)f_k + s_k(t) \quad (k = 1, \dots, n),$$

leads to an equation of the form 3.4.3.4 for  $u = u(x_1, \dots, x_n, t)$ .

$$6. \quad \frac{\partial w}{\partial t} - \sum_{k=1}^n \left[ a_k(x_k, t) \frac{\partial^2 w}{\partial x_k^2} + b_k(x_k, t) \frac{\partial w}{\partial x_k} + c_k(x_k, t) w \right] = \Phi(x_1, \dots, x_n, t).$$

Here,  $0 < a_k(x_k, t) < \infty$  for all  $k$ . We introduce the notation  $\mathbf{x} = \{x_1, \dots, x_n\}$ ,  $\mathbf{y} = \{y_1, \dots, y_n\}$  and consider the domain  $V = \{\alpha_k \leq x_k \leq \beta_k, k = 1, \dots, n\}$ , which is an  $n$ -dimensional parallelepiped.

1°. First boundary value problem. The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_k(\mathbf{x}, t) & \text{at } x_k &= \alpha_k & (\text{boundary conditions}), \\ w &= h_k(\mathbf{x}, t) & \text{at } x_k &= \beta_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t, 0) d\mathbf{y} \\ &+ \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\alpha_k, \tau) \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t, \tau) \right]_{y_k=\alpha_k} dS_y^{(k)} d\tau \\ &- \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\beta_k, \tau) \left[ h_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t, \tau) \right]_{y_k=\beta_k} dS_y^{(k)} d\tau, \end{aligned}$$

where

$$\begin{aligned} d\mathbf{y} &= dy_1 dy_2 \dots dy_n, \quad dS_y^{(k)} = dy_1 \dots dy_{k-1} dy_{k+1} \dots dy_n, \\ S^{(k)} &= \{\alpha_m \leq y_m \leq \beta_m \text{ for } m = 1, \dots, k-1, k+1, \dots, n\}. \end{aligned}$$

The Green's function can be represented in the product form

$$G(\mathbf{x}, \mathbf{y}, t, \tau) = \prod_{k=1}^n G_k(x_k, y_k, t, \tau). \quad (1)$$

Here, the  $G_k = G_k(x_k, y_k, t, \tau)$  are auxiliary Green's functions that, for  $t > \tau \geq 0$ , satisfy the one-dimensional linear homogeneous equations

$$\frac{\partial G_k}{\partial t} - a_k(x_k, t) \frac{\partial^2 G_k}{\partial x_k^2} - b_k(x_k, t) \frac{\partial G_k}{\partial x_k} - c_k(x_k, t) G_k = 0 \quad (k = 1, \dots, n) \quad (2)$$

with nonhomogeneous initial conditions of a special form,

$$G_k = \delta(x_k - y_k) \quad \text{at } t = \tau, \quad (3)$$

and homogeneous boundary conditions of the first kind,

$$\begin{aligned} G_k &= 0 & \text{at } x_k &= \alpha_k, \\ G_k &= 0 & \text{at } x_k &= \beta_k. \end{aligned}$$

In determining the function  $G_k$ , the quantities  $y_k$  and  $\tau$  play the role of parameters;  $\delta(x)$  is the Dirac delta function.

2°. The second and third boundary value problems. The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_{x_k} w - s_k w &= g_k(\mathbf{x}, t) & \text{at } x_k &= \alpha_k & (\text{boundary conditions}), \\ \partial_{x_k} w + p_k w &= h_k(\mathbf{x}, t) & \text{at } x_k &= \beta_k & (\text{boundary conditions}). \end{aligned}$$

The second boundary value problem corresponds to  $s_k = p_k = 0$ .

Solution:

$$\begin{aligned}
 w(\mathbf{x}, t) = & \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t, 0) d\mathbf{y} \\
 & - \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\alpha_k, \tau) [g_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau)]_{y_k=\alpha_k} dS_y^{(k)} d\tau \\
 & + \sum_{k=1}^n \int_0^t \int_{S^{(k)}} a_k(\beta_k, \tau) [h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t, \tau)]_{y_k=\beta_k} dS_y^{(k)} d\tau.
 \end{aligned}$$

The Green's function can be represented as the product (1) of the corresponding one-dimensional Green's functions satisfying the linear equations (2) with the initial conditions (3) and the homogeneous boundary conditions

$$\begin{aligned}
 \partial_{x_k} G_k - s_k G_k &= 0 \quad \text{at} \quad x_k = \alpha_k, \\
 \partial_{x_k} G_k + p_k G_k &= 0 \quad \text{at} \quad x_k = \beta_k.
 \end{aligned}$$

⊙ *Reference:* A. D. Polyanin (2000a, 2000b).

$$7. \quad \frac{\partial w}{\partial t} = \sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left[ a_{ij}(x_1, \dots, x_n) \frac{\partial w}{\partial x_j} \right] - q(x_1, \dots, x_n) w + \Phi(x_1, \dots, x_n, t).$$

The problems considered below are assumed to refer to a bounded domain  $V$  with smooth surface  $S$ . We introduce the brief notation  $\mathbf{x} = \{x_1, \dots, x_n\}$  and assume that the condition

$$\sum_{i,j=1}^n a_{ij}(\mathbf{x}) \lambda_i \lambda_j \geq c \sum_{i=1}^n \lambda_i^2, \quad c > 0,$$

is satisfied; this condition imposes the requirement that the differential operator on the right-hand side of the equation is elliptic.

1°. First boundary value problem. The following conditions are prescribed:

$$\begin{aligned}
 w &= f(\mathbf{x}) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 w &= g(\mathbf{x}, t) \quad \text{for} \quad \mathbf{x} \in S \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(\mathbf{x}, t) = & \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t-\tau) dV_y d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) dV_y \\
 & - \int_0^t \int_S g(\mathbf{y}, \tau) \left[ \frac{\partial}{\partial M_y} G(\mathbf{x}, \mathbf{y}, t-\tau) \right] dS_y d\tau.
 \end{aligned} \tag{1}$$

Here, the Green's function is given by

$$G(\mathbf{x}, \mathbf{y}, t) = \sum_{n=1}^{\infty} \frac{u_n(\mathbf{x}) u_n(\mathbf{y})}{\|u_n\|^2} \exp(-\lambda_n t), \quad \|u_n\|^2 = \int_V u_n^2(\mathbf{x}) dV, \quad \mathbf{y} = \{y_1, \dots, y_n\}, \tag{2}$$

where the  $\lambda_n$  and  $u_n(\mathbf{x})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the following elliptic second-order equation with homogeneous boundary condition of the first kind:

$$\sum_{i,j=1}^n \frac{\partial}{\partial x_i} \left[ a_{ij}(\mathbf{x}) \frac{\partial u}{\partial x_j} \right] - q(\mathbf{x}) u + \lambda u = 0, \tag{3}$$

$$u = 0 \quad \text{for} \quad \mathbf{x} \in S. \tag{4}$$

The integration in solution (1) is carried out with respect to  $y_1, \dots, y_n$ ;  $\frac{\partial}{\partial M_y}$  is the differential operator defined as

$$\frac{\partial G}{\partial M_y} \equiv \sum_{i,j=1}^n a_{ij}(\mathbf{y}) N_j \frac{\partial G}{\partial y_i}, \quad (5)$$

where  $\mathbf{N} = \{N_1, \dots, N_n\}$  is the unit outward normal to the surface  $S$ . In the special case where  $a_{ii}(\mathbf{x}) = 1$  and  $a_{ij}(\mathbf{x}) = 0$  for  $i \neq j$ , the operator of (5) coincides with the usual operator of differentiation along the direction of the outward normal to the surface  $S$ .

General properties of the Sturm–Liouville problem (3)–(4):

1. There are countably many eigenvalues. All eigenvalues are real and can be ordered so that  $\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$ , with  $\lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ ; consequently, there can exist only finitely many negative eigenvalues.

2. For  $q(\mathbf{x}) \geq 0$  all eigenvalues are positive:  $\lambda_n > 0$ .

3. The eigenfunctions are defined up to a constant multiplier. Any two eigenfunctions  $u_n(\mathbf{x})$  and  $u_m(\mathbf{x})$  corresponding to different eigenvalues  $\lambda_n$  and  $\lambda_m$  are orthogonal in the domain  $V$ :

$$\int_V u_n(\mathbf{x}) u_m(\mathbf{x}) dV = 0 \quad \text{for } n \neq m.$$

*Remark.* To each eigenvalue  $\lambda_n$  there generally correspond finitely many linearly independent eigenfunctions  $u_n^{(1)}, u_n^{(2)}, \dots, u_n^{(m)}$ . These functions can always be replaced by their linear combinations

$$\bar{u}_n^{(k)} = A_{k,1} u_n^{(1)} + \dots + A_{k,k-1} u_n^{(k-1)} + u_n^{(k)}, \quad k = 1, 2, \dots, m,$$

such that  $\bar{u}_n^{(1)}, \bar{u}_n^{(2)}, \dots, \bar{u}_n^{(m)}$  are now pairwise orthogonal. Thus, without loss of generality, we assume that all eigenfunctions are orthogonal.

2°. Second boundary value problem. The following conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \frac{\partial w}{\partial M_x} &= g(\mathbf{x}, t) \quad \text{for } \mathbf{x} \in S \quad (\text{boundary condition}). \end{aligned}$$

Here, the left-hand side of the boundary condition is determined with the help of (5), where  $G, y, \mathbf{y}$ , and  $y_k$  must be replaced by  $w, x, \mathbf{x}$ , and  $x_k$ , respectively.

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) dV_y d\tau + \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) dV_y \\ &+ \int_0^t \int_S g(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) dS_y d\tau. \end{aligned} \quad (6)$$

Here, the Green's function is defined by (2), where the  $\lambda_n$  and  $u_n(\mathbf{x})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the elliptic second-order equation (3) with a homogeneous boundary condition of the second kind:

$$\frac{\partial u}{\partial M_x} = 0 \quad \text{for } \mathbf{x} \in S. \quad (7)$$

For  $q(\mathbf{x}) > 0$  the general properties of the eigenvalue problem (3), (7) are the same as for the first boundary value problem (see Item 1°). For  $q(\mathbf{x}) \equiv 0$  the zero eigenvalue  $\lambda_0 = 0$  arises which corresponds to the eigenfunction  $u_0 = \text{const}$ .

It should be noted that the Green's function of the second boundary value problem can be expressed in terms of the Green's function of the third boundary value problem (see Item 3°).

3°. Third boundary value problem. The following conditions are prescribed:

$$w = f(\mathbf{x}) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\frac{\partial w}{\partial M_x} + k(\mathbf{x})w = g(\mathbf{x}, t) \quad \text{for} \quad \mathbf{x} \in S \quad (\text{boundary condition}).$$

The solution of the third boundary value problem is given by relations (6) and (2), where the  $\lambda_n$  and  $u_n(\mathbf{x})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the second-order elliptic equation (3) with a homogeneous boundary condition of the third kind:

$$\frac{\partial u}{\partial M_x} + k(\mathbf{x})u = 0 \quad \text{for} \quad \mathbf{x} \in S. \quad (8)$$

For  $q(\mathbf{x}) \geq 0$  and  $k(\mathbf{x}) > 0$ , the general properties of the eigenvalue problem (3), (8) are the same as for the first boundary value problem (see Item 1°).

Let  $k(\mathbf{x}) = k = \text{const}$ . Denote the Green's functions of the second and third boundary value problems by  $G_2(\mathbf{x}, \mathbf{y}, t)$  and  $G_3(\mathbf{x}, \mathbf{y}, t, k)$ , respectively. Then the following relations hold:

$$G_2(\mathbf{x}, \mathbf{y}, t) = \begin{cases} \lim_{k \rightarrow 0} G_3(\mathbf{x}, \mathbf{y}, t, k), & \text{if } q(\mathbf{x}) > 0; \\ \frac{1}{V_0} + \lim_{k \rightarrow 0} G_3(\mathbf{x}, \mathbf{y}, t, k), & \text{if } q(\mathbf{x}) \equiv 0; \end{cases}$$

where  $V_0 = \int_V dV$  is the volume of the domain in question.

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), A. D. Polyinin (2000a, 2000b).

## Chapter 4

# Hyperbolic Equations with One Space Variable

---

### 4.1. Constant Coefficient Equations

#### 4.1.1. Wave Equation $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2}$

This equation is also known as the *equation of vibration of a string*. It is often encountered in elasticity, aerodynamics, acoustics, and electrodynamics.

4.1.1-1. General solution. Some formulas.

1°. General solution:

$$w(x, t) = \varphi(x + at) + \psi(x - at),$$

where  $\varphi(x)$  and  $\psi(x)$  are arbitrary functions. *Physical interpretation*: The solution represents two traveling waves that propagate, respectively, to the left and right along the  $x$ -axis at a constant speed  $a$ .

2°. Fundamental solution:

$$\mathcal{E}(x, t) = \frac{1}{2a} \vartheta(at - |x|), \quad \vartheta(z) = \begin{cases} 0 & \text{for } z < 0, \\ 1 & \text{for } z > 0. \end{cases}$$

3°. Infinite series solutions containing arbitrary functions of the space variable:

$$w(x, t) = f(x) + \sum_{n=1}^{\infty} \frac{(at)^{2n}}{(2n)!} f_x^{(2n)}(x), \quad f_x^{(m)}(x) = \frac{d^m}{dx^m} f(x),$$
$$w(x, t) = tg(x) + t \sum_{n=1}^{\infty} \frac{(at)^{2n}}{(2n+1)!} g_x^{(2n)}(x),$$

where  $f(x)$  and  $g(x)$  are any infinitely differentiable functions. The first solution satisfies the initial conditions  $w(x, 0) = f(x)$  and  $\partial_t w(x, 0) = 0$ , and the second  $w(x, 0) = 0$  and  $\partial_t w(x, 0) = g(x)$ . The sums are finite if  $f(x)$  and  $g(x)$  are polynomials.

4°. Infinite series solutions containing arbitrary functions of time:

$$w(x, t) = f(t) + \sum_{n=1}^{\infty} \frac{1}{a^{2n}(2n)!} x^{2n} f_t^{(2n)}(t), \quad f_t^{(m)}(t) = \frac{d^m}{dt^m} f(t),$$
$$w(x, t) = xg(t) + x \sum_{n=1}^{\infty} \frac{1}{a^{2n}(2n+1)!} x^{2n} g_t^{(2n)}(t),$$

where  $f(t)$  and  $g(t)$  are any infinitely differentiable functions. The sums are finite if  $f(t)$  and  $g(t)$  are polynomials. The first solution satisfies the boundary condition of the first kind  $w(0, t) = f(t)$ , and the second solution to the boundary condition of the second kind  $\partial_x w(0, t) = g(t)$ .



5°. If  $w(x, t)$  is a solution of the wave equation, then the functions

$$\begin{aligned}w_1 &= Aw(\pm\lambda x + C_1, \pm\lambda t + C_2), \\w_2 &= Aw\left(\frac{x - vt}{\sqrt{1 - (v/a)^2}}, \frac{t - va^{-2}x}{\sqrt{1 - (v/a)^2}}\right), \\w_3 &= Aw\left(\frac{x}{x^2 - a^2t^2}, \frac{t}{x^2 - a^2t^2}\right),\end{aligned}$$

are also solutions of the equation everywhere these functions are defined ( $A$ ,  $C_1$ ,  $C_2$ ,  $v$ , and  $\lambda$  are arbitrary constants). The signs at  $\lambda$ 's in the formula for  $w_1$  are taken arbitrarily, independently of each other. The function  $w_2$  results from the invariance of the wave equation under the Lorentz transformations.

⊙ *References:* G. N. Polozhii (1964), A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 4.1.1-2. Domain: $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned}w &= f(x) & \text{at } t &= 0, \\ \partial_t w &= g(x) & \text{at } t &= 0.\end{aligned}$$

Solution (D'Alembert's formula):

$$w(x, t) = \frac{1}{2}[f(x + at) + f(x - at)] + \frac{1}{2a} \int_{x-at}^{x+at} g(\xi) d\xi.$$

#### 4.1.1-3. Domain: $0 \leq x < \infty$ . First boundary value problem.

1°. Problem with a homogeneous boundary condition:

$$\begin{aligned}w &= f(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= g(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= 0 & \text{at } x &= 0 & \text{(boundary condition).}\end{aligned}$$

Solution:

$$w(x, t) = \begin{cases} \frac{1}{2}[f(x + at) + f(x - at)] + \frac{1}{2a} \int_{x-at}^{x+at} g(\xi) d\xi & \text{for } t < \frac{x}{a}, \\ \frac{1}{2}[f(x + at) - f(at - x)] + \frac{1}{2a} \int_{at-x}^{x+at} g(\xi) d\xi & \text{for } t > \frac{x}{a}. \end{cases}$$

2°. Problem with a nonhomogeneous boundary condition:

$$\begin{aligned}w &= f(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= g(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= h(t) & \text{at } x &= 0 & \text{(boundary condition).}\end{aligned}$$

Solution:

$$w(x, t) = \begin{cases} \frac{1}{2}[f(x + at) + f(x - at)] + \frac{1}{2a} \int_{x-at}^{x+at} g(\xi) d\xi & \text{for } t < \frac{x}{a}, \\ \frac{1}{2}[f(x + at) - f(at - x)] + \frac{1}{2a} \int_{at-x}^{x+at} g(\xi) d\xi + h\left(t - \frac{x}{a}\right) & \text{for } t > \frac{x}{a}. \end{cases}$$

In the domain  $t < x/a$  the boundary conditions have no effect on the solution and the expression of  $w(x, t)$  coincides with D'Alembert's solution for an infinite line (see Paragraph 4.1.1-2).

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

#### 4.1.1-4. Domain: $0 \leq x < \infty$ . Second boundary value problem.

1°. Problem with a homogeneous boundary condition:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= 0 & \text{at } x &= 0 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(x, t) = \begin{cases} \frac{1}{2}[f(x+at) + f(x-at)] + \frac{1}{2a}[G(x+at) - G(x-at)] & \text{for } t < \frac{x}{a}, \\ \frac{1}{2}[f(x+at) + f(at-x)] + \frac{1}{2a}[G(x+at) + G(at-x)] & \text{for } t > \frac{x}{a}, \end{cases}$$

where  $G(z) = \int_0^z g(\xi) d\xi$ .

2°. Problem with a nonhomogeneous boundary condition:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= h(t) & \text{at } x &= 0 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(x, t) = \begin{cases} \frac{1}{2}[f(x+at) + f(x-at)] + \frac{1}{2a}[G(x+at) - G(x-at)] & \text{for } t < \frac{x}{a}, \\ \frac{1}{2}[f(x+at) + f(at-x)] + \frac{1}{2a}[G(x+at) + G(at-x)] - aH\left(t - \frac{x}{a}\right) & \text{for } t > \frac{x}{a}, \end{cases}$$

where  $G(z) = \int_0^z g(\xi) d\xi$  and  $H(z) = \int_0^z h(\xi) d\xi$ . In the domain  $t < x/a$  the boundary conditions have no effect on the solution, and the expression of  $w(x, t)$  coincides with D'Alembert's solution for an infinite line (see Paragraph 4.1.1-2).

⊙ *Reference:* B. M. Budak, A. N. Tikhonov, and A. A. Samarskii (1980).

#### 4.1.1-5. Domain: $0 \leq x \leq l$ . First boundary value problem.

1°. Vibration of a string with rigidly fixed ends. The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= 0 & \text{at } x &= 0 & (\text{boundary condition}), \\ w &= 0 & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \sum_{n=1}^{\infty} [A_n \cos(\lambda_n at) + B_n \sin(\lambda_n at)] \sin(\lambda_n x), & \lambda_n &= \frac{\pi n}{l}, \\ A_n &= \frac{2}{l} \int_0^l f(x) \sin(\lambda_n x) dx, & B_n &= \frac{2}{a\pi n} \int_0^l g(x) \sin(\lambda_n x) dx. \end{aligned}$$

**Example 1.** The initial shape of the string is a triangle with base  $0 \leq x \leq l$  and height  $h$  at  $x = c$ , i.e.,

$$f(x) = \begin{cases} \frac{hx}{c} & \text{for } 0 \leq x \leq c, \\ \frac{h(l-x)}{l-c} & \text{for } c \leq x \leq l. \end{cases}$$

The initial velocities of the string points are zero,  $g(x) = 0$ .

Solution:

$$w(x, t) = \frac{2hl^2}{\pi^2 c(l-c)} \sum_{n=1}^{\infty} \frac{1}{n^2} \sin\left(\frac{n\pi c}{l}\right) \sin\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi at}{l}\right).$$

**Example 2.** Initially, the string has the shape of a parabola symmetric about the center of the string with elevation  $h$ , so that

$$f(x) = \frac{4h}{l^2} x(l-x).$$

The initial velocities of the string points are zero,  $g(x) = 0$ .

Solution:

$$w(x, t) = \frac{32h}{\pi^3} \sum_{n=0}^{\infty} \frac{1}{(2n+1)^3} \sin\left[\frac{(2n+1)\pi x}{l}\right] \cos\left[\frac{(2n+1)\pi at}{l}\right].$$

2°. For the solution of the first boundary value problem with a nonhomogeneous boundary condition, see Paragraph 4.1.2-4 with  $\Phi(x, t) \equiv 0$ .

⊙ *References:* B. M. Budak, A. N. Tikhonov, and A. A. Samarskii (1980), A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 4.1.1-6. Domain: $0 \leq x \leq l$ . Second boundary value problem.

1°. Longitudinal vibration of an elastic rod with free ends. The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= 0 & \text{at } x &= 0 & (\text{boundary condition}), \\ \partial_x w &= 0 & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= A_0 + B_0 t + \sum_{n=1}^{\infty} [A_n \cos(\lambda_n at) + B_n \sin(\lambda_n at)] \cos(\lambda_n x), \\ \lambda_n &= \frac{\pi n}{l}, \quad A_0 = \frac{1}{l} \int_0^l f(x) dx, \quad B_0 = \frac{1}{l} \int_0^l g(x) dx, \\ A_n &= \frac{2}{l} \int_0^l f(x) \cos(\lambda_n x) dx, \quad B_n = \frac{2}{a\pi n} \int_0^l g(x) \cos(\lambda_n x) dx. \end{aligned}$$

2°. For the solution of the second boundary value problem with a nonhomogeneous boundary condition, see Paragraph 4.1.2-5 with  $\Phi(x, t) \equiv 0$ .

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 4.1.1-7. Domain: $0 \leq x \leq l$ . Third boundary value problem.

1°. Longitudinal vibration of an elastic rod with clamped ends in the case of equal stiffness coefficients. The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w - kw &= 0 & \text{at } x &= 0 & (\text{boundary condition}), \\ \partial_x w + kw &= 0 & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(x, t) = \sum_{n=1}^{\infty} [A_n \cos(\lambda_n at) + B_n \sin(\lambda_n at)] \sin(\lambda_n x + \varphi_n),$$

where

$$A_n = \frac{1}{\|X_n\|^2} \int_0^l \sin(\lambda_n x + \varphi_n) f(x) dx, \quad B_n = \frac{1}{a\lambda_n \|X_n\|^2} \int_0^l \sin(\lambda_n x + \varphi_n) g(x) dx,$$

$$\varphi_n = \arctan \frac{\lambda_n}{k}, \quad \|X_n\|^2 = \int_0^l \sin^2(\lambda_n x + \varphi_n) dx = \frac{l}{2} + \frac{k}{k^2 + \lambda_n^2};$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cot(\lambda l) = \frac{1}{2} \left( \frac{\lambda}{k} - \frac{k}{\lambda} \right)$ .

2°. Longitudinal vibration of an elastic rod with clamped ends in the case of different stiffness coefficients. The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= g(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - k_1 w &= 0 & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w + k_2 w &= 0 & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$w(x, t) = \sum_{n=1}^{\infty} [A_n \cos(\lambda_n at) + B_n \sin(\lambda_n at)] \sin(\lambda_n x + \varphi_n),$$

where

$$A_n = \frac{1}{\|X_n\|^2} \int_0^l \sin(\lambda_n x + \varphi_n) f(x) dx, \quad B_n = \frac{1}{a\lambda_n \|X_n\|^2} \int_0^l \sin(\lambda_n x + \varphi_n) g(x) dx,$$

$$\varphi_n = \arctan \frac{\lambda_n}{k_1}, \quad \|X_n\|^2 = \int_0^l \sin^2(\lambda_n x + \varphi_n) dx = \frac{l}{2} + \frac{(\lambda_n^2 + k_1 k_2)(k_1 + k_2)}{2(\lambda_n^2 + k_1^2)(\lambda_n^2 + k_2^2)};$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cot(\lambda l) = \frac{\lambda^2 - k_1 k_2}{\lambda(k_1 + k_2)}$ .

3°. For the solution of the third boundary value problem with nonhomogeneous boundary conditions, see Paragraph 4.1.2-6 with  $\Phi(x, t) \equiv 0$ .

⊙ *Reference:* B. M. Budak, A. N. Tikhonov, and A. A. Samarskii (1980).

#### 4.1.1-8. Domain: $0 \leq x \leq l$ . Mixed boundary value problem.

1°. Longitudinal vibration of an elastic rod with one end rigidly fixed and the other free. The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= g(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= 0 & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w &= 0 & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$w(x, t) = \sum_{n=0}^{\infty} [A_n \cos(\lambda_n at) + B_n \sin(\lambda_n at)] \sin(\lambda_n x), \quad \lambda_n = \frac{\pi(2n+1)}{2l},$$

$$A_n = \frac{2}{l} \int_0^l f(x) \sin(\lambda_n x) dx, \quad B_n = \frac{2}{al\lambda_n} \int_0^l g(x) \sin(\lambda_n x) dx.$$

2°. For the solution of the mixed boundary value problem with nonhomogeneous boundary conditions, see Paragraph 4.1.2-7 with  $\Phi(x, t) \equiv 0$ .

⊙ *References:* M. M. Smirnov (1975), A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 4.1.1-9. Goursat problem.

The boundary conditions are prescribed to the equation characteristics:

$$w = f(x) \quad \text{for} \quad x - at = 0 \quad (0 \leq x \leq b),$$

$$w = g(x) \quad \text{for} \quad x + at = 0 \quad (0 \leq x \leq c),$$

where  $f(0) = g(0)$ .

Solution:

$$w(x, t) = f\left(\frac{x + at}{2}\right) + g\left(\frac{x - at}{2}\right) - f(0).$$

The solution propagation domain is bounded by four lines:

$$x - at = 0, \quad x + at = 0, \quad x - at = 2c, \quad x + at = 2b.$$

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 4.1.2. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} + \Phi(x, t)$

##### 4.1.2-1. Domain: $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$w = f(x) \quad \text{at} \quad t = 0,$$

$$\partial_t w = g(x) \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, t) = \frac{1}{2}[f(x - at) + f(x + at)] + \frac{1}{2a} \int_{x-at}^{x+at} g(\xi) d\xi + \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \Phi(\xi, \tau) d\xi d\tau.$$

##### 4.1.2-2. Domain: $0 \leq x < \infty$ . First boundary value problem.

The following conditions are prescribed:

$$w = f(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\partial_t w = g(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$w = h(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}).$$

Solution:

$$w(x, t) = w_1(x, t) + \frac{1}{2a} w_2(x, t),$$

where

$$w_1(x, t) = \begin{cases} \frac{1}{2}[f(x + at) + f(x - at)] + \frac{1}{2a} \int_{x-at}^{x+at} g(\xi) d\xi & \text{for } t < \frac{x}{a}, \\ \frac{1}{2}[f(x + at) - f(at - x)] + \frac{1}{2a} \int_{at-x}^{x+at} g(\xi) d\xi + h\left(t - \frac{x}{a}\right) & \text{for } t > \frac{x}{a}, \end{cases}$$

$$w_2(x, t) = \begin{cases} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \Phi(\xi, \tau) d\xi d\tau & \text{for } t < \frac{x}{a}, \\ \int_0^{t-x/a} \int_{a(t-\tau)-x}^{x+a(t-\tau)} \Phi(\xi, \tau) d\xi d\tau + \int_{t-x/a}^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \Phi(\xi, \tau) d\xi d\tau & \text{for } t > \frac{x}{a}. \end{cases}$$

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 4.1.2-3. Domain: $0 \leq x < \infty$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= h(t) & \text{at } x &= 0 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(x, t) = w_1(x, t) + \frac{1}{2a} w_2(x, t),$$

where

$$w_1(x, t) = \begin{cases} \frac{1}{2} [f(x+at) + f(x-at)] + \frac{1}{2a} \int_{x-at}^{x+at} g(\xi) d\xi & \text{for } t < \frac{x}{a}, \\ \frac{1}{2} [f(x+at) + f(at-x)] + \frac{1}{2a} \int_0^{x+at} g(\xi) d\xi \\ \quad + \frac{1}{2a} \int_0^{at-x} g(\xi) d\xi - a \int_0^{t-x/a} h(\xi) d\xi & \text{for } t > \frac{x}{a}, \end{cases}$$

$$w_2(x, t) = \begin{cases} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \Phi(\xi, \tau) d\xi d\tau & \text{for } t < \frac{x}{a}, \\ \int_0^{t-x/a} \int_0^{x+a(t-\tau)} \Phi(\xi, \tau) d\xi d\tau + \int_0^{t-x/a} \int_0^{a(t-\tau)-x} \Phi(\xi, \tau) d\xi d\tau \\ \quad + \int_{t-x/a}^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \Phi(\xi, \tau) d\xi d\tau & \text{for } t > \frac{x}{a}. \end{cases}$$

⊙ Reference: A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 4.1.2-4. Domain: $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(t) & \text{at } x &= 0 & (\text{boundary condition}), \\ w &= g_2(t) & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t-\tau) d\xi d\tau \\ &\quad + a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t-\tau) \right]_{\xi=0} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t-\tau) \right]_{\xi=l} d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{2}{a\pi} \sum_{n=1}^{\infty} \frac{1}{n} \sin\left(\frac{n\pi x}{l}\right) \sin\left(\frac{n\pi \xi}{l}\right) \sin\left(\frac{n\pi at}{l}\right).$$

#### 4.1.2-5. Domain: $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= g_1(t) & \text{at } x &= 0 & (\text{boundary condition}), \\ \partial_x w &= g_2(t) & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &\quad - a^2 \int_0^t g_1(\tau) G(x, 0, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{t}{l} + \frac{2}{a\pi} \sum_{n=1}^{\infty} \frac{1}{n} \cos\left(\frac{n\pi x}{l}\right) \cos\left(\frac{n\pi \xi}{l}\right) \sin\left(\frac{n\pi at}{l}\right).$$

#### 4.1.2-6. Domain: $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(t) & \text{at } x &= 0 & (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(t) & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, t)$  is determined by the formula in Paragraph 4.1.2-5 where

$$\begin{aligned} G(x, \xi, t) &= \frac{1}{a} \sum_{n=1}^{\infty} \frac{1}{\lambda_n \|u_n\|^2} \sin(\lambda_n x + \varphi_n) \sin(\lambda_n \xi + \varphi_n) \sin(\lambda_n at), \\ \varphi_n &= \arctan \frac{\lambda_n}{k_1}, \quad \|u_n\|^2 = \frac{l}{2} + \frac{(\lambda_n^2 + k_1 k_2)(k_1 + k_2)}{2(\lambda_n^2 + k_1^2)(\lambda_n^2 + k_2^2)}; \end{aligned}$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cot(\lambda l) = \frac{\lambda^2 - k_1 k_2}{\lambda(k_1 + k_2)}$ .

#### 4.1.2-7. Domain: $0 \leq x \leq l$ . Mixed boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(t) & \text{at } x &= 0 & (\text{boundary condition}), \\ \partial_x w &= g_2(t) & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &\quad + a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=0} d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{2}{al} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \sin(\lambda_n x) \sin(\lambda_n \xi) \sin(\lambda_n at), \quad \lambda_n = \frac{\pi(2n+1)}{2l}.$$

### 4.1.3. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} - bw + \Phi(x, t)$

This equation with  $\Phi(x, t) \equiv 0$  and  $b > 0$  is encountered in quantum field theory and a number of applications and is referred to as the *Klein–Gordon equation*.

#### 4.1.3-1. Solutions of the homogeneous equation ( $\Phi \equiv 0$ ).

1°. Particular solutions:

$$w(x, t) = \exp(\pm \mu t)(Ax + B), \quad b = -\mu^2,$$

$$w(x, t) = \exp(\pm \lambda x)(At + B), \quad b = a^2 \lambda^2,$$

$$w(x, t) = \cos(\lambda x)[A \cos(\mu t) + B \sin(\mu t)], \quad b = -a^2 \lambda^2 + \mu^2,$$

$$w(x, t) = \sin(\lambda x)[A \cos(\mu t) + B \sin(\mu t)], \quad b = -a^2 \lambda^2 + \mu^2,$$

$$w(x, t) = \exp(\pm \mu t)[A \cos(\lambda x) + B \sin(\lambda x)], \quad b = -a^2 \lambda^2 - \mu^2,$$

$$w(x, t) = \exp(\pm \lambda x)[A \cos(\mu t) + B \sin(\mu t)], \quad b = a^2 \lambda^2 + \mu^2,$$

$$w(x, t) = \exp(\pm \lambda x)[A \exp(\mu t) + B \exp(-\mu t)], \quad b = a^2 \lambda^2 - \mu^2,$$

$$w(x, t) = AJ_0(\xi) + BY_0(\xi), \quad \xi = \frac{\sqrt{b}}{a} \sqrt{a^2(t + C_1)^2 - (x + C_2)^2}, \quad b > 0,$$

$$w(x, t) = AI_0(\xi) + BK_0(\xi), \quad \xi = \frac{\sqrt{-b}}{a} \sqrt{a^2(t + C_1)^2 - (x + C_2)^2}, \quad b < 0,$$

where  $A, B, C_1$ , and  $C_2$  are arbitrary constants,  $J_0(\xi)$  and  $Y_0(\xi)$  are the Bessel functions, and  $I_0(\xi)$  and  $K_0(\xi)$  are the modified Bessel functions.

2°. Fundamental solutions:

$$\mathcal{E}(x, t) = \frac{\vartheta(at - |x|)}{2a} J_0\left(\frac{c}{a} \sqrt{a^2 t^2 - x^2}\right) \quad \text{for } b = c^2 > 0,$$

$$\mathcal{E}(x, t) = \frac{\vartheta(at - |x|)}{2a} I_0\left(\frac{c}{a} \sqrt{a^2 t^2 - x^2}\right) \quad \text{for } b = -c^2 < 0,$$

where  $\vartheta(z)$  is the Heaviside unit step function ( $\vartheta = 0$  for  $z < 0$  and  $\vartheta = 1$  for  $z \geq 0$ ),  $J_0(z)$  is the Bessel function, and  $I_0(z)$  is the modified Bessel function.

⊙ *Reference:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

#### 4.1.3-2. Some formulas and transformations of the homogeneous equation ( $\Phi \equiv 0$ ).

1°. Suppose  $w = w(x, t)$  is a solution of the Klein–Gordon equation. Then the functions

$$w_1 = Aw(x + C_1, \pm t + C_2),$$

$$w_2 = Aw(-x + C_1, \pm t + C_2),$$

$$w_3 = Aw\left(\frac{x - vt}{\sqrt{1 - (v/a)^2}}, \frac{t - va^{-2}x}{\sqrt{1 - (v/a)^2}}\right),$$

where  $A, C_1, C_2$ , and  $v$  are arbitrary constants, are also solutions of this equation.

2°. [Table 19](#) lists transformations of the independent variables that allow separation of variables in the Klein–Gordon equation.

*Notation:*  $J_\sigma(z)$  and  $Y_\sigma(z)$  are the Bessel functions,  $I_\sigma(z)$  and  $K_\sigma(z)$  are the modified Bessel functions, and  $D_\lambda(z)$  is the parabolic cylinder function.

⊙ *References:* E. Kalnins (1975), W. Miller, Jr. (1977).



TABLE 19

Orthogonal coordinates  $u = u(x, t)$ ,  $v = v(x, t)$  admitting separable solutions  $w = F(u)G(v)$  of the Klein–Gordon equation ( $a = 1$ ;  $A_1, A_2, B_1, B_2$ , and  $\lambda$  are arbitrary constants)

| No | Relation between $x, t$ and $u, v$                                                                  | Function $F = F(u)$<br>(differential equation)                                                                                                   | Function $G = G(v)$<br>(differential equation)                                                                                                      |
|----|-----------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | $x = u, \quad t = v$                                                                                | $F = A_1 e^{u\sqrt{\lambda+b}} + A_2 e^{-u\sqrt{\lambda+b}}$                                                                                     | $G = B_1 e^{v\sqrt{\lambda}} + B_2 e^{-v\sqrt{\lambda}}$                                                                                            |
| 2  | $x = u \sinh v,$<br>$t = u \cosh v$                                                                 | $F = \sqrt{u} \left[ A_1 J_\sigma(u\sqrt{b}) + A_2 Y_\sigma(u\sqrt{b}) \right],$<br>$\sigma = \frac{1}{2}\sqrt{1+\lambda^2}$                     | $G = B_1 e^{\lambda v} + B_2 e^{-\lambda v}$                                                                                                        |
| 3  | $x = uv,$<br>$t = \frac{1}{2}(u^2 + v^2)$                                                           | $F = A_1 D_\lambda(\beta u) + A_2 D_\lambda(-\beta u),$<br>$\beta = (-4b)^{1/4}$                                                                 | $G = B_1 D_\lambda(\beta v) + B_2 D_\lambda(-\beta v),$<br>$\beta = (-4b)^{1/4}$                                                                    |
| 4  | $x = \frac{1}{2}(u^2 + v^2),$<br>$t = uv$                                                           | $F = A_1 D_\lambda(\beta u) + A_2 D_\lambda(-\beta u),$<br>$\beta = (4b)^{1/4}$                                                                  | $G = B_1 D_\lambda(\beta v) + B_2 D_\lambda(-\beta v),$<br>$\beta = (4b)^{1/4}$                                                                     |
| 5  | $x = -\frac{1}{2}(u-v)^2 + u + v,$<br>$t = \frac{1}{2}(u-v)^2 + u + v$                              | $F = \sqrt{U} \left[ A_1 J_{\frac{1}{3}}(\xi) + A_2 Y_{\frac{1}{3}}(\xi) \right],$<br>$U = u + \lambda, \quad \xi = \frac{2}{3}\sqrt{b} U^{3/2}$ | $G = \sqrt{V} \left[ B_1 J_{\frac{1}{3}}(\eta) + B_2 Y_{\frac{1}{3}}(\eta) \right],$<br>$V = v + \lambda, \quad \eta = \frac{2}{3}\sqrt{b} V^{3/2}$ |
| 6  | $t + x = \cosh \left[ \frac{1}{2}(u-v) \right],$<br>$t - x = \sinh \left[ \frac{1}{2}(u+v) \right]$ | $F'' + (\lambda + b \sinh u)F = 0$                                                                                                               | $G'' + (\lambda + b \sinh v)G = 0$                                                                                                                  |
| 7  | $x = \sinh(u-v) - \frac{1}{2}e^{u+v},$<br>$t = \sinh(u-v) + \frac{1}{2}e^{u+v}$                     | $F = A_1 J_\lambda(\beta e^u) + A_2 Y_\lambda(\beta e^u),$<br>$\beta = \sqrt{b}$                                                                 | $G = B_1 I_\lambda(\beta e^v) + B_2 K_\lambda(\beta e^v),$<br>$\beta = \sqrt{b}$                                                                    |
| 8  | $x = \cosh(u-v) - \frac{1}{2}e^{u+v},$<br>$t = \cosh(u-v) + \frac{1}{2}e^{u+v}$                     | $F = A_1 J_\lambda(\beta e^u) + A_2 Y_\lambda(\beta e^u),$<br>$\beta = \sqrt{b}$                                                                 | $G = B_1 J_\lambda(\beta e^v) + B_2 Y_\lambda(\beta e^v),$<br>$\beta = \sqrt{b}$                                                                    |
| 9  | $x = \cosh u \sinh v,$<br>$t = \sinh u \cosh v$                                                     | $F'' + (\lambda + \frac{1}{2}b \cosh 2u)F = 0,$<br>modified Mathieu equation                                                                     | $G'' + (\lambda - \frac{1}{2}b \cosh 2v)G = 0,$<br>modified Mathieu equation                                                                        |
| 10 | $x = \sinh u \sinh v,$<br>$t = \cosh u \cosh v$                                                     | $F'' + (\lambda + \frac{1}{2}b \cosh 2u)F = 0,$<br>modified Mathieu equation                                                                     | $G'' + (\lambda + \frac{1}{2}b \cosh 2v)G = 0,$<br>modified Mathieu equation                                                                        |
| 11 | $x = \sin u \sin v,$<br>$t = \cos u \cos v$                                                         | $F'' + (\lambda - \frac{1}{2}b \cos 2u)F = 0,$<br>Mathieu equation                                                                               | $G'' + (\lambda - \frac{1}{2}b \cos 2v)G = 0,$<br>Mathieu equation                                                                                  |

#### 4.1.3-3. Domain: $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(x) \quad \text{at} \quad t = 0. \end{aligned}$$

Solution for  $b = -c^2 < 0$ :

$$\begin{aligned} w(x, t) &= \frac{1}{2}[f(x+at) + f(x-at)] + \frac{ct}{2a} \int_{x-at}^{x+at} \frac{I_1(c\sqrt{t^2 - (x-\xi)^2/a^2})}{\sqrt{t^2 - (x-\xi)^2/a^2}} f(\xi) d\xi \\ &\quad + \frac{1}{2a} \int_{x-at}^{x+at} I_0(c\sqrt{t^2 - (x-\xi)^2/a^2}) g(\xi) d\xi \\ &\quad + \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} I_0(c\sqrt{(t-\tau)^2 - (x-\xi)^2/a^2}) \Phi(\xi, \tau) d\xi d\tau, \end{aligned}$$

where  $I_0(z)$  and  $I_1(z)$  are the modified Bessel functions of the first kind.

Solution for  $b = c^2 > 0$ :

$$\begin{aligned} w(x, t) = & \frac{1}{2}[f(x + at) + f(x - at)] - \frac{ct}{2a} \int_{x-at}^{x+at} \frac{J_1(c\sqrt{t^2 - (x-\xi)^2/a^2})}{\sqrt{t^2 - (x-\xi)^2/a^2}} f(\xi) d\xi \\ & + \frac{1}{2a} \int_{x-at}^{x+at} J_0(c\sqrt{t^2 - (x-\xi)^2/a^2}) g(\xi) d\xi \\ & + \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} J_0(c\sqrt{(t-\tau)^2 - (x-\xi)^2/a^2}) \Phi(\xi, \tau) d\xi d\tau, \end{aligned}$$

where  $J_0(z)$  and  $J_1(z)$  are the Bessel functions of the first kind.

⊙ *Reference:* B. M. Budak, A. N. Tikhonov, and A. A. Samarskii (1980).

#### 4.1.3-4. Domain: $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) = & \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ & + a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=0} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=1}^{\infty} \sin(\lambda_n x) \sin(\lambda_n \xi) \frac{\sin(t\sqrt{a^2 \lambda_n^2 + b})}{\sqrt{a^2 \lambda_n^2 + b}}, \quad \lambda_n = \frac{\pi n}{l}.$$

*Remark.* Let  $b < 0$  and  $a^2 \lambda_n^2 + b < 0$  for  $n = 1, \dots, m$  and  $a^2 \lambda_n^2 + b > 0$  for  $n = m + 1, m + 2, \dots$ . In this case the Green's function is modified and acquires the form

$$\begin{aligned} G(x, \xi, t) = & \frac{2}{l} \sum_{n=1}^m \sin(\lambda_n x) \sin(\lambda_n \xi) \frac{\sinh(t\sqrt{|a^2 \lambda_n^2 + b|})}{\sqrt{|a^2 \lambda_n^2 + b|}} \\ & + \frac{2}{l} \sum_{n=m+1}^{\infty} \sin(\lambda_n x) \sin(\lambda_n \xi) \frac{\sin(t\sqrt{a^2 \lambda_n^2 + b})}{\sqrt{a^2 \lambda_n^2 + b}}, \quad \lambda_n = \frac{\pi n}{l}. \end{aligned}$$

Analogously, the Green's functions for the second, third, and mixed boundary value problems are modified in similar cases.

⊙ *Reference:* A. G. Butkovskiy (1979).

#### 4.1.3-5. Domain: $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(x, t) = \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ - a^2 \int_0^t g_1(\tau) G(x, 0, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau,$$

where

$$G(x, \xi, t) = \frac{1}{l\sqrt{b}} \sin(t\sqrt{b}) + \frac{2}{l} \sum_{n=1}^{\infty} \cos(\lambda_n x) \cos(\lambda_n \xi) \frac{\sin(t\sqrt{a^2 \lambda_n^2 + b})}{\sqrt{a^2 \lambda_n^2 + b}}, \quad \lambda_n = \frac{\pi n}{l}.$$

#### 4.1.3-6. Domain: $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - k_1 w &= g_1(t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w + k_2 w &= g_2(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, t)$  is determined by the formula in Paragraph 4.1.3-5 where

$$G(x, \xi, t) = \sum_{n=1}^{\infty} \frac{y_n(x) y_n(\xi) \sin(t\sqrt{a^2 \lambda_n^2 + b})}{\|y_n\|^2 \sqrt{a^2 \lambda_n^2 + b}}, \\ y_n(x) = \cos(\lambda_n x) + \frac{k_1}{\lambda_n} \sin(\lambda_n x), \quad \|y_n\|^2 = \frac{k_2}{2\lambda_n^2} \frac{\lambda_n^2 + k_1^2}{\lambda_n^2 + k_2^2} + \frac{k_1}{2\lambda_n^2} + \frac{l}{2} \left( 1 + \frac{k_1^2}{\lambda_n^2} \right).$$

Here, the  $\lambda_n$  are positive roots of the transcendental equation  $\frac{\tan(\lambda l)}{\lambda} = \frac{k_1 + k_2}{\lambda^2 - k_1 k_2}$ .

#### 4.1.3-7. Domain: $0 \leq x \leq l$ . Mixed boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w &= g_2(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$w(x, t) = \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ + a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=0} d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau,$$

where

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=0}^{\infty} \sin(\lambda_n x) \sin(\lambda_n \xi) \frac{\sin(t\sqrt{a^2 \lambda_n^2 + b})}{\sqrt{a^2 \lambda_n^2 + b}}, \quad \lambda_n = \frac{\pi(2n+1)}{2l}.$$

#### 4.1.4. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} - b \frac{\partial w}{\partial x} + \Phi(x, t)$

4.1.4-1. Reduction to the nonhomogeneous Klein–Gordon equation.

The substitution  $w(x, t) = \exp\left(\frac{1}{2}bx/a^2\right)u(x, t)$  leads the nonhomogeneous Klein–Gordon equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial x^2} - \frac{b^2}{4a^2}u + \exp\left(-\frac{bx}{2a^2}\right)\Phi(x, t),$$

which is discussed in Subsection 4.1.3.

4.1.4-2. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0, \\ \partial_t w &= g(x) & \text{at } t &= 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{1}{2}f(x+at)\exp\left(-\frac{bt}{2a}\right) + \frac{1}{2}f(x-at)\exp\left(\frac{bt}{2a}\right) \\ &\quad - \frac{\sigma t}{2a}\exp\left(\frac{bx}{2a^2}\right)\int_{x-at}^{x+at}\exp\left(-\frac{b\xi}{2a^2}\right)\frac{J_1(\sigma\sqrt{t^2-(x-\xi)^2/a^2})}{\sqrt{t^2-(x-\xi)^2/a^2}}f(\xi)d\xi \\ &\quad + \frac{1}{2a}\exp\left(\frac{bx}{2a^2}\right)\int_{x-at}^{x+at}\exp\left(-\frac{b\xi}{2a^2}\right)J_0(\sigma\sqrt{t^2-(x-\xi)^2/a^2})g(\xi)d\xi \\ &\quad + \frac{1}{2a}\int_0^t\int_{x-a(t-\tau)}^{x+a(t-\tau)}\exp\left[\frac{b(x-\xi)}{2a^2}\right]J_0(\sigma\sqrt{(t-\tau)^2-(x-\xi)^2/a^2})\Phi(\xi, \tau)d\xi d\tau, \end{aligned}$$

where  $J_0(z)$  and  $J_1(z)$  are the Bessel functions of the first kind, and  $\sigma = \frac{1}{2}|b|/a$ .

4.1.4-3. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t}\int_0^l f_0(\xi)G(x, \xi, t)d\xi + \int_0^l f_1(\xi)G(x, \xi, t)d\xi + \int_0^t\int_0^l \Phi(\xi, \tau)G(x, \xi, t-\tau)d\xi d\tau \\ &\quad + a^2\int_0^t g_1(\tau)\left[\frac{\partial}{\partial \xi}G(x, \xi, t-\tau)\right]_{\xi=0}d\tau - a^2\int_0^t g_2(\tau)\left[\frac{\partial}{\partial \xi}G(x, \xi, t-\tau)\right]_{\xi=l}d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{2}{l}\exp\left[\frac{b}{2a^2}(x-\xi)\right]\sum_{n=1}^{\infty}\sin\left(\frac{\pi nx}{l}\right)\sin\left(\frac{\pi n\xi}{l}\right)\frac{\sin(\lambda_n t)}{\lambda_n}, \quad \lambda_n = \sqrt{\frac{a^2\pi^2 n^2}{l^2} + \frac{b^2}{4a^2}}.$$

⊙ Reference: A. G. Butkovskiy (1979).

#### 4.1.4-4. Domain: $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &\quad - a^2 \int_0^t g_1(\tau) G(x, 0, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, \xi, t) &= \frac{bt}{a^2 [1 - \exp(-bl/a^2)]} \exp\left(-\frac{b\xi}{a^2}\right) + \frac{2}{l} \exp\left[\frac{b}{2a^2}(x - \xi)\right] \sum_{n=1}^{\infty} \frac{y_n(x) y_n(\xi) \sin(\lambda_n t)}{\lambda_n (1 + \mu_n^2)}, \\ y_n(x) &= \cos\left(\frac{\pi n x}{l}\right) - \frac{bl}{2a^2 \pi n} \sin\left(\frac{\pi n x}{l}\right), \quad \lambda_n = \sqrt{\frac{a^2 \pi^2 n^2}{l^2} + \frac{b^2}{4a^2}}, \quad \mu_n = \frac{bl}{2a^2 \pi n}. \end{aligned}$$

⊙ Reference: A. G. Butkovskiy (1979).

#### 4.1.4-5. Domain: $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, t)$  is determined by the formula in Paragraph 4.1.4-4 where

$$G(x, \xi, t) = \exp\left[\frac{b(x - \xi)}{2a^2}\right] \sum_{n=1}^{\infty} \frac{y_n(x) y_n(\xi) \sin(a \lambda_n t)}{a \lambda_n B_n}.$$

Here,

$$\begin{aligned} y_n(x) &= \cos(\mu_n x) + \frac{2a^2 k_1 - b}{2a^2 \mu_n} \sin(\mu_n x), \quad \lambda_n = \sqrt{\mu_n^2 + \frac{b^2}{4a^4}}, \\ B_n &= \frac{2a^2 k_2 + b}{4a^2 \mu_n^2} \frac{4a^4 \mu_n^2 + (2a^2 k_1 - b)^2}{4a^4 \mu_n^2 + (2a^2 k_2 + b)^2} + \frac{2a^2 k_1 - b}{4a^2 \mu_n^2} + \frac{l}{2} + \frac{l(2a^2 k_1 - b)^2}{8a^4 \mu_n^2}, \end{aligned}$$

where the  $\mu_n$  are positive roots of the transcendental equation

$$\frac{\tan(\mu l)}{\mu} = \frac{4a^4(k_1 + k_2)}{4a^4 \mu^2 - (2a^2 k_1 - b)(2a^2 k_2 + b)}.$$

⊙ Reference: A. G. Butkovskiy (1979).

### 4.1.5. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw + \Phi(x, t)$

4.1.5-1. Reduction to the nonhomogeneous Klein–Gordon equation.

The substitution  $w(x, t) = \exp(-\frac{1}{2}a^{-2}bx)u(x, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial x^2} + \left(c - \frac{1}{4}a^{-2}b^2\right)u + \exp\left(\frac{1}{2}a^{-2}bx\right)\Phi(x, t),$$

which is discussed in Subsection 4.1.3.

4.1.5-2. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0, \\ \partial_t w &= g(x) & \text{at } t &= 0. \end{aligned}$$

Solution for  $c - \frac{1}{4}a^{-2}b^2 = \sigma^2 > 0$ :

$$\begin{aligned} w(x, t) &= \frac{1}{2}f(x+at)\exp\left(\frac{bt}{2a}\right) + \frac{1}{2}f(x-at)\exp\left(-\frac{bt}{2a}\right) \\ &+ \frac{\sigma t}{2a}\exp\left(-\frac{bx}{2a^2}\right)\int_{x-at}^{x+at}\exp\left(\frac{b\xi}{2a^2}\right)\frac{I_1(\sigma\sqrt{t^2-(x-\xi)^2/a^2})}{\sqrt{t^2-(x-\xi)^2/a^2}}f(\xi)d\xi \\ &+ \frac{1}{2a}\exp\left(-\frac{bx}{2a^2}\right)\int_{x-at}^{x+at}\exp\left(\frac{b\xi}{2a^2}\right)I_0(\sigma\sqrt{t^2-(x-\xi)^2/a^2})g(\xi)d\xi \\ &+ \frac{1}{2a}\int_0^t\int_{x-a(t-\tau)}^{x+a(t-\tau)}\exp\left[\frac{b(\xi-x)}{2a^2}\right]I_0(\sigma\sqrt{(t-\tau)^2-(x-\xi)^2/a^2})\Phi(\xi, \tau)d\xi d\tau, \end{aligned}$$

where  $I_0(z)$  and  $I_1(z)$  are the modified Bessel functions of the first kind.

Solution for  $c - \frac{1}{4}a^{-2}b^2 = -\sigma^2 < 0$ :

$$\begin{aligned} w(x, t) &= \frac{1}{2}f(x+at)\exp\left(\frac{bt}{2a}\right) + \frac{1}{2}f(x-at)\exp\left(-\frac{bt}{2a}\right) \\ &- \frac{\sigma t}{2a}\exp\left(-\frac{bx}{2a^2}\right)\int_{x-at}^{x+at}\exp\left(\frac{b\xi}{2a^2}\right)\frac{J_1(\sigma\sqrt{t^2-(x-\xi)^2/a^2})}{\sqrt{t^2-(x-\xi)^2/a^2}}f(\xi)d\xi \\ &+ \frac{1}{2a}\exp\left(-\frac{bx}{2a^2}\right)\int_{x-at}^{x+at}\exp\left(\frac{b\xi}{2a^2}\right)J_0(\sigma\sqrt{t^2-(x-\xi)^2/a^2})g(\xi)d\xi \\ &+ \frac{1}{2a}\int_0^t\int_{x-a(t-\tau)}^{x+a(t-\tau)}\exp\left[\frac{b(\xi-x)}{2a^2}\right]J_0(\sigma\sqrt{(t-\tau)^2-(x-\xi)^2/a^2})\Phi(\xi, \tau)d\xi d\tau, \end{aligned}$$

where  $J_0(z)$  and  $J_1(z)$  are the Bessel functions of the first kind.

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

4.1.5-3. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(t) & \text{at } x &= 0 & (\text{boundary condition}), \\ w &= g_2(t) & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(x, t) = \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau + \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ + a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=0} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau.$$

Let  $a^2 \pi^2 + \frac{1}{4} a^{-2} b^2 l^2 - cl^2 > 0$ . Then

$$G(x, \xi, t) = \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} \right] \sum_{n=1}^{\infty} \sin \left( \frac{\pi n x}{l} \right) \sin \left( \frac{\pi n \xi}{l} \right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \pi^2 n^2}{l^2} + \frac{b^2}{4a^2} - c.$$

Let

$$a^2 \pi^2 n^2 + \frac{1}{4} a^{-2} b^2 l^2 - cl^2 \leq 0 \quad \text{at } n = 1, \dots, m; \\ a^2 \pi^2 n^2 + \frac{1}{4} a^{-2} b^2 l^2 - cl^2 > 0 \quad \text{at } n = m + 1, m + 2, \dots$$

Then

$$G(x, \xi, t) = \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} \right] \sum_{n=1}^m \sin \left( \frac{\pi n x}{l} \right) \sin \left( \frac{\pi n \xi}{l} \right) \frac{\sinh(t\sqrt{\beta_n})}{\sqrt{\beta_n}} \\ + \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} \right] \sum_{n=m+1}^{\infty} \sin \left( \frac{\pi n x}{l} \right) \sin \left( \frac{\pi n \xi}{l} \right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \\ \beta_n = c - \frac{a^2 \pi^2 n^2}{l^2} - \frac{b^2}{4a^2}, \quad \lambda_n = \frac{a^2 \pi^2 n^2}{l^2} + \frac{b^2}{4a^2} - c.$$

For  $\beta_n = 0$  the ratio  $\sinh(t\sqrt{\beta_n})/\sqrt{\beta_n}$  must be replaced by  $t$ .

● Reference: A. G. Butkovskiy (1979).

#### 4.1.5-4. Domain: $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$w = f_0(x) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_t w = f_1(x) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_x w = g_1(t) \quad \text{at } x = 0 \quad (\text{boundary condition}), \\ \partial_x w = g_2(t) \quad \text{at } x = l \quad (\text{boundary condition}).$$

Solution:

$$w(x, t) = \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau + \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ - a^2 \int_0^t g_1(\tau) G(x, 0, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau.$$

For  $c < 0$ ,

$$G(x, \xi, t) = \frac{b}{a^2(e^{bl/a^2} - 1)} \exp \left( \frac{b\xi}{a^2} \right) \frac{\sin(t\sqrt{|c|})}{\sqrt{|c|}} + \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} \right] \sum_{n=1}^{\infty} \frac{y_n(x)y_n(\xi)}{1 + \mu_n^2} \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \\ \lambda_n = \frac{a^2 \pi^2 n^2}{l^2} + \frac{b^2}{4a^2} - c, \quad y_n(x) = \cos \left( \frac{\pi n x}{l} \right) + \mu_n \sin \left( \frac{\pi n x}{l} \right), \quad \mu_n = \frac{bl}{2a^2 \pi n}.$$

For  $c > 0$ ,

$$G(x, \xi, t) = \frac{b}{a^2(e^{bl/a^2} - 1)} \exp \left( \frac{b\xi}{a^2} \right) \frac{\sinh(t\sqrt{c})}{\sqrt{c}} + \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} \right] \sum_{n=1}^{\infty} \frac{y_n(x)y_n(\xi)}{1 + \mu_n^2} \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}},$$

where the  $\lambda_n$ ,  $y_n(x)$ , and  $\mu_n$  were specified previously. If the inequality  $\lambda_n < 0$  holds for several first values  $n = 1, \dots, m$ , then the  $\sqrt{\lambda_n}$  in the corresponding terms of the series should be replaced by  $\sqrt{|\lambda_n|}$ , and the sines by the hyperbolic sines.

**4.1.5-5. Domain:  $0 \leq x \leq l$ . Third boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, t)$  is determined by the formula in Paragraph 4.1.5-4 where

$$G(x, \xi, t) = \exp\left[\frac{b(\xi - x)}{2a^2}\right] \sum_{n=1}^{\infty} \frac{y_n(x)y_n(\xi) \sin(t\sqrt{\lambda_n})}{B_n \sqrt{\lambda_n}}.$$

Here,

$$\begin{aligned} y_n(x) &= \cos(\mu_n x) + \frac{2a^2 k_1 + b}{2a^2 \mu_n} \sin(\mu_n x), \quad \lambda_n = a^2 \mu_n^2 + \frac{b^2}{4a^2} - c, \\ B_n &= \frac{2a^2 k_2 - b}{4a^2 \mu_n^2} \frac{4a^4 \mu_n^2 + (2a^2 k_1 + b)^2}{4a^4 \mu_n^2 + (2a^2 k_2 - b)^2} + \frac{2a^2 k_1 + b}{4a^2 \mu_n^2} + \frac{l}{2} + \frac{l(2a^2 k_1 + b)^2}{8a^4 \mu_n^2}, \end{aligned}$$

where the  $\mu_n$  are positive roots of the transcendental equation

$$\frac{\tan(\mu l)}{\mu} = \frac{4a^4(k_1 + k_2)}{4a^4 \mu^2 - (2a^2 k_1 + b)(2a^2 k_2 - b)}.$$

## 4.2. Wave Equation with Axial or Central Symmetry

### 4.2.1. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right)$

This is the one-dimensional wave equation with axial symmetry, where  $r = \sqrt{x^2 + y^2}$  is the radial coordinate. In the problems considered in Paragraphs 4.2.1-1 through 4.2.1-3, the solutions bounded at  $r = 0$  are sought (this is not specially stated below).

**4.2.1-1. Domain:  $0 \leq r \leq R$ . First boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(r, t) = \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi - a^2 \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R} d\tau,$$

where

$$G(r, \xi, t) = \frac{2\xi}{aR} \sum_{n=1}^{\infty} \frac{1}{\lambda_n J_1^2(\lambda_n)} J_0\left(\frac{\lambda_n r}{R}\right) J_0\left(\frac{\lambda_n \xi}{R}\right) \sin\left(\frac{\lambda_n a t}{R}\right).$$

Here, the  $\lambda_n$  are positive zeros of the Bessel function,  $J_0(\lambda) = 0$ . The numerical values of the first ten  $\lambda_n$  are specified in Paragraph 1.2.1-3.

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).



#### 4.2.1-2. Domain: $0 \leq r \leq R$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_r w &= g(t) & \text{at } r &= R & (\text{boundary condition}). \end{aligned}$$

Solution:

$$w(r, t) = \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi + a^2 \int_0^t g(\tau) G(r, R, t - \tau) d\tau,$$

where

$$G(r, \xi, t) = \frac{2t\xi}{R^2} + \frac{2\xi}{aR} \sum_{n=1}^{\infty} \frac{1}{\lambda_n J_0^2(\lambda_n)} J_0\left(\frac{\lambda_n r}{R}\right) J_0\left(\frac{\lambda_n \xi}{R}\right) \sin\left(\frac{\lambda_n a t}{R}\right).$$

Here, the  $\lambda_n$  are positive zeros of the first-order Bessel function,  $J_1(\lambda) = 0$ . The numerical values of the first ten roots  $\lambda_n$  are specified in Paragraph 1.2.1-4.

⊙ *References:* M. M. Smirnov (1975), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

#### 4.2.1-3. Domain: $0 \leq r \leq R$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_r w + kw &= g(t) & \text{at } r &= R & (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.1-2 where

$$G(r, \xi, t) = \frac{2\xi}{aR} \sum_{n=1}^{\infty} \frac{\lambda_n}{(k^2 R^2 + \lambda_n^2) J_0^2(\lambda_n)} J_0\left(\frac{\lambda_n r}{R}\right) J_0\left(\frac{\lambda_n \xi}{R}\right) \sin\left(\frac{\lambda_n a t}{R}\right).$$

Here, the  $\lambda_n$  are positive roots of the transcendental equation

$$\lambda J_1(\lambda) - kR J_0(\lambda) = 0.$$

The numerical values of the first six roots  $\lambda_n$  can be found in Carslaw and Jaeger (1984); see also Abramowitz and Stegun (1964).

#### 4.2.1-4. Domain: $R_1 \leq r \leq R_2$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(t) & \text{at } r &= R_1 & (\text{boundary condition}), \\ w &= g_2(t) & \text{at } r &= R_2 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\ &+ a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_1} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_2} d\tau. \end{aligned}$$

Here,

$$G(r, \xi, t) = \sum_{n=1}^{\infty} A_n \xi \Psi_n(r) \Psi_n(\xi) \sin\left(\frac{\lambda_n a t}{R_1}\right), \quad A_n = \frac{\pi^2 \lambda_n J_0^2(s \lambda_n)}{2a R_1 [J_0^2(\lambda_n) - J_0^2(s \lambda_n)]},$$

$$\Psi_n(r) = Y_0(\lambda_n) J_0\left(\frac{\lambda_n r}{R_1}\right) - J_0(\lambda_n) Y_0\left(\frac{\lambda_n r}{R_1}\right), \quad s = \frac{R_2}{R_1},$$

where  $J_0(z)$  and  $Y_0(z)$  are the Bessel functions, the  $\lambda_n$  are positive roots of the transcendental equation

$$J_0(\lambda) Y_0(s \lambda) - J_0(s \lambda) Y_0(\lambda) = 0.$$

The numerical values of the first five roots  $\lambda_n = \lambda_n(s)$  can be found in Abramowitz and Stegun (1964) and Carslaw and Jaeger (1984).

**4.2.1-5. Domain:  $R_1 \leq r \leq R_2$ . Second boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g_1(t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w &= g_2(t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\ &\quad - a^2 \int_0^t g_1(\tau) G(r, R_1, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(r, R_2, t - \tau) d\tau. \end{aligned}$$

Here,

$$G(r, \xi, t) = \frac{2t\xi}{R_2^2 - R_1^2} + \sum_{n=1}^{\infty} A_n \xi \Psi_n(r) \Psi_n(\xi) \sin\left(\frac{\lambda_n a t}{R_1}\right), \quad A_n = \frac{\pi^2 \lambda_n J_1^2(s \lambda_n)}{2a R_1 [J_1^2(\lambda_n) - J_1^2(s \lambda_n)]},$$

$$\Psi_n(r) = Y_1(\lambda_n) J_0\left(\frac{\lambda_n r}{R_1}\right) - J_1(\lambda_n) Y_0\left(\frac{\lambda_n r}{R_1}\right), \quad s = \frac{R_2}{R_1},$$

where  $J_k(z)$  and  $Y_k(z)$  are the Bessel functions ( $k = 0, 1$ ); the  $\lambda_n$  are positive roots of the transcendental equation

$$J_1(\lambda) Y_1(s \lambda) - J_1(s \lambda) Y_1(\lambda) = 0.$$

The numerical values of the first five roots  $\lambda_n = \lambda_n(s)$  can be found in Abramowitz and Stegun (1964).

**4.2.1-6. Domain:  $R_1 \leq r \leq R_2$ . Third boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.1-5 where

$$G(r, \xi, t) = \frac{\pi^2}{2a} \sum_{n=1}^{\infty} \frac{\mu_n}{B_n} [k_2 J_0(\mu_n R_2) - \mu_n J_1(\mu_n R_2)]^2 \xi H_n(r) H_n(\xi) \sin(\mu_n a t).$$

Here,

$$B_n = (\mu_n^2 + k_2^2) [k_1 J_0(\mu_n R_1) + \mu_n J_1(\mu_n R_1)]^2 - (\mu_n^2 + k_1^2) [k_2 J_0(\mu_n R_2) - \mu_n J_1(\mu_n R_2)]^2,$$

$$H_n(r) = [k_1 Y_0(\mu_n R_1) + \mu_n Y_1(\mu_n R_1)] J_0(\mu_n r) - [k_1 J_0(\mu_n R_1) + \mu_n J_1(\mu_n R_1)] Y_0(\mu_n r);$$

$J_k(z)$  and  $Y_k(z)$  are the Bessel functions ( $k=0, 1$ ); and the  $\mu_n$  are positive roots of the transcendental equation

$$[k_1 J_0(\mu R_1) + \mu J_1(\mu R_1)] [k_2 Y_0(\mu R_2) - \mu Y_1(\mu R_2)] - [k_2 J_0(\mu R_2) - \mu J_1(\mu R_2)] [k_1 Y_0(\mu R_1) + \mu Y_1(\mu R_1)] = 0.$$

## 4.2.2. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) + \Phi(r, t)$

4.2.2-1. Domain:  $0 \leq r \leq R$ . Different boundary value problems.

1°. The solution to the first boundary value problem for a circle of radius  $R$  is given by the formula from Paragraph 4.2.1-1 with the additional term

$$\int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution to the second boundary value problem for a circle of radius  $R$  is given by the formula from Paragraph 4.2.1-2 with the additional term (1).

3°. The solution to the third boundary value problem for a circle of radius  $R$  is the sum of the solution presented in Paragraph 4.2.1-3 and expression (1).

4.2.2-2. Domain:  $R_1 \leq r \leq R_2$ . Different boundary value problems.

1°. The solution to the first boundary value problem for an annular domain is given by the formula from Paragraph 4.2.1-4 with the additional term

$$\int_0^t \int_{R_1}^{R_2} \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution to the second boundary value problem for an annular domain is given by the formula from Paragraph 4.2.1-5 with the additional term (2).

3°. The solution to the third boundary value problem for an annular domain is the sum of the solution presented in Paragraph 4.2.1-6 and expression (2).

## 4.2.3. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right)$

This is the equation of one-dimensional vibration of a gas with central symmetry, where  $r = \sqrt{x^2 + y^2 + z^2}$  is the radial coordinate. In the problems considered in Paragraphs 4.2.3-1 through 4.2.3-3, the solutions bounded at  $r = 0$  are sought; this is not specially stated below.

#### 4.2.3-1. General solution:

$$w(t, r) = \frac{\varphi(r + at) + \psi(r - at)}{r},$$

where  $\varphi(r_1)$  and  $\psi(r_2)$  are arbitrary functions.

#### 4.2.3-2. Reduction to a constant coefficient equation.

The substitution  $u(r, t) = rw(r, t)$  leads to the constant coefficient equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial r^2},$$

which is discussed in Subsection 4.1.1.

#### 4.2.3-3. Domain: $0 \leq r < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(r) & \text{at } t &= 0, \\ \partial_t w &= g(r) & \text{at } t &= 0. \end{aligned}$$

Solution:

$$w(r, t) = \frac{1}{2r} [(r - at)f(|r - at|) + (r + at)f(|r + at|)] + \frac{1}{2ar} \int_{r-at}^{r+at} \xi g(|\xi|) d\xi.$$

Solution at the center  $r = 0$ :

$$w(0, t) = at f'(at) + f(at) + tg(at).$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

#### 4.2.3-4. Domain: $0 \leq r \leq R$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g(t) & \text{at } r &= R & \text{(boundary condition).} \end{aligned}$$

Solution:

$$w(r, t) = \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi - a^2 \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R} d\tau,$$

where

$$G(r, \xi, t) = \frac{2\xi}{\pi ar} \sum_{n=1}^{\infty} \frac{1}{n} \sin\left(\frac{n\pi r}{R}\right) \sin\left(\frac{n\pi \xi}{R}\right) \sin\left(\frac{an\pi t}{R}\right).$$

#### 4.2.3-5. Domain: $0 \leq r \leq R$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g(t) & \text{at } r &= R & \text{(boundary condition).} \end{aligned}$$

Solution:

$$w(r, t) = \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi + a^2 \int_0^t g(\tau) G(r, R, t - \tau) d\tau,$$

where

$$G(r, \xi, t) = \frac{3t\xi^2}{R^3} + \frac{2\xi}{ar} \sum_{n=1}^{\infty} \frac{\mu_n^2 + 1}{\mu_n^3} \sin\left(\frac{\mu_n r}{R}\right) \sin\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{\mu_n at}{R}\right).$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\tan \mu - \mu = 0$ . The numerical values of the first five roots  $\mu_n$  are specified in Paragraph 1.2.3-5.

#### 4.2.3-6. Domain: $0 \leq r \leq R$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w + kw &= g(t) & \text{at } r &= R & \text{(boundary condition).} \end{aligned}$$

The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.3-5 where

$$G(r, \xi, t) = \frac{2\xi}{ar} \sum_{n=1}^{\infty} \frac{\mu_n^2 + (kR - 1)^2}{\mu_n [\mu_n^2 + kR(kR - 1)]} \sin\left(\frac{\mu_n r}{R}\right) \sin\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{\mu_n at}{R}\right).$$

Here, the  $\mu_n$  are positive roots of the transcendental equation

$$\mu \cot \mu + kR - 1 = 0.$$

The numerical values of the first six roots  $\mu_n$  can be found in Carslaw and Jaeger (1984).

#### 4.2.3-7. Domain: $R_1 \leq r \leq R_2$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ w &= g_2(t) & \text{at } r &= R_2 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\ &+ a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_1} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_2} d\tau, \end{aligned}$$

where

$$G(r, \xi, t) = \frac{2\xi}{\pi ar} \sum_{n=1}^{\infty} \frac{1}{n} \sin\left[\frac{\pi n(r - R_1)}{R_2 - R_1}\right] \sin\left[\frac{\pi n(\xi - R_1)}{R_2 - R_1}\right] \sin\left(\frac{\pi nat}{R_2 - R_1}\right).$$

#### 4.2.3-8. Domain: $R_1 \leq r \leq R_2$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g_1(t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w &= g_2(t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\ &\quad - a^2 \int_0^t g_1(\tau) G(r, R_1, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(r, R_2, t - \tau) d\tau, \end{aligned}$$

where

$$\begin{aligned} G(r, \xi, t) &= \frac{3t\xi^2}{R_2^3 - R_1^3} + \frac{2\xi}{a(R_2 - R_1)r} \sum_{n=1}^{\infty} \frac{(1 + R_2^2 \lambda_n^2) \Psi_n(r) \Psi_n(\xi) \sin(\lambda_n a t)}{\lambda_n^3 [R_1^2 + R_2^2 + R_1 R_2 (1 + R_1 R_2 \lambda_n^2)]}, \\ \Psi_n(r) &= \sin[\lambda_n(r - R_1)] + R_1 \lambda_n \cos[\lambda_n(r - R_1)]. \end{aligned}$$

Here, the  $\lambda_n$  are positive roots of the transcendental equation

$$(\lambda^2 R_1 R_2 + 1) \tan[\lambda(R_2 - R_1)] - \lambda(R_2 - R_1) = 0.$$

#### 4.2.3-9. Domain: $R_1 \leq r \leq R_2$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.3-8 where

$$\begin{aligned} G(r, \xi, t) &= \frac{2\xi}{a(R_2 - R_1)r} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \frac{(b_2^2 + R_2^2 \lambda_n^2) \Psi_n(r) \Psi_n(\xi) \sin(\lambda_n a t)}{(b_1^2 + R_1^2 \lambda_n^2)(b_2^2 + R_2^2 \lambda_n^2) + (b_1 R_2 + b_2 R_1)(b_1 b_2 + R_1 R_2 \lambda_n^2)}, \\ \Psi_n(r) &= b_1 \sin[\lambda_n(r - R_1)] + R_1 \lambda_n \cos[\lambda_n(r - R_1)], \quad b_1 = k_1 R_1 + 1, \quad b_2 = k_2 R_2 - 1. \end{aligned}$$

Here, the  $\lambda_n$  are positive roots of the transcendental equation

$$(b_1 b_2 - R_1 R_2 \lambda^2) \sin[\lambda(R_2 - R_1)] + \lambda(R_1 b_2 + R_2 b_1) \cos[\lambda(R_2 - R_1)] = 0.$$

### 4.2.4. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right) + \Phi(r, t)$

#### 4.2.4-1. Reduction to a nonhomogeneous constant coefficient equation.

The substitution  $u(r, t) = rw(r, t)$  leads to the nonhomogeneous constant coefficient equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial r^2} + r \Phi(r, t),$$

which is discussed in Subsection 4.1.2.

#### 4.2.4-2. Domain: $0 \leq r < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(r) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(r) \quad \text{at} \quad t = 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{1}{2r} [(r - at)f(|r - at|) + (r + at)f(|r + at|)] + \frac{1}{2ar} \int_{r-at}^{r+at} \xi g(|\xi|) d\xi \\ &\quad + \frac{1}{2ar} \int_0^t d\tau \int_{r-a(t-\tau)}^{r+a(t-\tau)} \xi \Phi(|\xi|, \tau) d\xi. \end{aligned}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

#### 4.2.4-3. Domain: $0 \leq r \leq R$ . Different boundary value problems.

1°. The solution to the first boundary value problem for a sphere of radius  $R$  is given by the formula from Paragraph 4.2.3-4 with the additional term

$$\int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution to the second boundary value problem for a sphere of radius  $R$  is given by the formula from Paragraph 4.2.3-5 with the additional term (1).

3°. The solution to the third boundary value problem for a sphere of radius  $R$  is the sum of the solution presented in Paragraph 4.2.3-6 and expression (1).

#### 4.2.4-4. Domain: $R_1 \leq r \leq R_2$ . Different boundary value problems.

1°. The solution to the first boundary value problem for a spherical layer is given by the formula from Paragraph 4.2.3-7 with the additional term

$$\int_0^t \int_{R_1}^{R_2} \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution to the second boundary value problem for a spherical layer is given by the formula from Paragraph 4.2.3-8 with the additional term (2).

3°. The solution to the third boundary value problem for a spherical layer is the sum of the solution presented in Paragraph 4.2.3-9 and expression (2).

### 4.2.5. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) - bw + \Phi(r, t)$

For  $b > 0$  and  $\Phi \equiv 0$ , this is the Klein–Gordon equation describing one-dimensional wave phenomena with axial symmetry. In the problems considered in Paragraphs 4.2.5-1 through 4.2.5-3, the solutions bounded at  $r = 0$  are sought; this is not specially stated below.

#### 4.2.5-1. Domain: $0 \leq r \leq R$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi \\ &\quad - a^2 \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R} d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \xi, t) = \frac{2\xi}{R^2} \sum_{n=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{R^2} + b,$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ . The numerical values of the first ten  $\mu_n$  are specified in Paragraph 1.2.1-3.

#### 4.2.5-2. Domain: $0 \leq r \leq R$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi \\ &\quad + a^2 \int_0^t g(\tau) G(r, R, t - \tau) d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \xi, t) = \frac{2\xi \sin(t\sqrt{b})}{R^2 \sqrt{b}} + \frac{2\xi}{R^2} \sum_{n=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{R^2} + b,$$

where the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu) = 0$ . The numerical values of the first ten  $\mu_n$  are specified in Paragraph 1.2.1-4.

#### 4.2.5-3. Domain: $0 \leq r \leq R$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$



The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.5-2 where

$$G(r, \xi, t) = \frac{2}{R^2} \sum_{n=1}^{\infty} \frac{\mu_n^2 \xi}{(k^2 R^2 + \mu_n^2) J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{R^2} + b.$$

Here, the  $\mu_n$  are positive roots of the transcendental equation

$$\mu J_1(\mu) - kR J_0(\mu) = 0.$$

The numerical values of the first six roots  $\mu_n$  can be found in Abramowitz and Stegun (1964) and Carslaw and Jaeger (1984).

**4.2.5-4. Domain:  $R_1 \leq r \leq R_2$ . First boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ w &= g_2(t) & \text{at } r &= R_2 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \int_0^t \int_{R_1}^{R_2} \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\ &+ a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_1} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_2} d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \xi, t) &= \frac{\pi^2}{2R_1^2} \sum_{n=1}^{\infty} \frac{\mu_n^2 J_0^2(s\mu_n) \xi}{J_0^2(\mu_n) - J_0^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{R_1^2} + b, \\ \Psi_n(r) &= Y_0(\mu_n) J_0\left(\frac{\mu_n r}{R_1}\right) - J_0(\mu_n) Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1}, \end{aligned}$$

where  $J_0(z)$  and  $Y_0(z)$  are the Bessel functions and the  $\mu_n$  are positive roots of the transcendental equation

$$J_0(\mu) Y_0(s\mu) - J_0(s\mu) Y_0(\mu) = 0.$$

The numerical values of the first five roots  $\mu_n = \mu_n(s)$  can be found in Abramowitz and Stegun (1964) and Carslaw and Jaeger (1984).

**4.2.5-5. Domain:  $R_1 \leq r \leq R_2$ . Second boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ \partial_r w &= g_2(t) & \text{at } r &= R_2 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, t) = & \int_0^t \int_{R_1}^{R_2} \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau \\
 & + \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\
 & - a^2 \int_0^t g_1(\tau) G(r, R_1, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(r, R_2, t - \tau) d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \xi, t) = & \frac{2\xi \sin(t\sqrt{b})}{(R_2^2 - R_1^2)\sqrt{b}} + \frac{\pi^2}{2R_1^2} \sum_{n=1}^{\infty} \frac{\mu_n^2 J_1^2(s\mu_n)\xi}{J_1^2(\mu_n) - J_1^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \\
 \Psi_n(r) = & Y_1(\mu_n) J_0\left(\frac{\mu_n r}{R_1}\right) - J_1(\mu_n) Y_0\left(\frac{\mu_n r}{R_1}\right), \quad \lambda_n = \frac{a^2 \mu_n^2}{R_1^2} + b, \quad s = \frac{R_2}{R_1},
 \end{aligned}$$

where  $J_k(z)$  and  $Y_k(z)$  are the Bessel functions ( $k = 0, 1$ ); the  $\mu_n$  are positive roots of the transcendental equation

$$J_1(\mu)Y_1(s\mu) - J_1(s\mu)Y_1(\mu) = 0.$$

The numerical values of the first five roots  $\mu_n = \mu_n(s)$  can be found in Abramowitz and Stegun (1964).

4.2.5-6. Domain:  $R_1 \leq r \leq R_2$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_r w - k_1 w = g_1(t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
 \partial_r w + k_2 w = g_2(t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition).}
 \end{aligned}$$

The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.5-5 where

$$G(r, \xi, t) = \frac{\pi^2}{2} \sum_{n=1}^{\infty} \frac{\beta_n^2}{B_n \sqrt{a^2 \beta_n^2 + b}} [k_2 J_0(\beta_n R_2) - \beta_n J_1(\beta_n R_2)]^2 \xi H_n(r) H_n(\xi) \sin(t\sqrt{a^2 \beta_n^2 + b}).$$

Here,

$$\begin{aligned}
 B_n = & (\beta_n^2 + k_2^2) [k_1 J_0(\beta_n R_1) + \beta_n J_1(\beta_n R_1)]^2 - (\beta_n^2 + k_1^2) [k_2 J_0(\beta_n R_2) - \beta_n J_1(\beta_n R_2)]^2, \\
 H_n(r) = & [k_1 Y_0(\beta_n R_1) + \beta_n Y_1(\beta_n R_1)] J_0(\beta_n r) - [k_1 J_0(\beta_n R_1) + \beta_n J_1(\beta_n R_1)] Y_0(\beta_n r),
 \end{aligned}$$

where the  $\beta_n$  are positive roots of the transcendental equation

$$\begin{aligned}
 [k_1 J_0(\beta R_1) + \beta J_1(\beta R_1)] [k_2 Y_0(\beta R_2) - \beta Y_1(\beta R_2)] \\
 - [k_2 J_0(\beta R_2) - \beta J_1(\beta R_2)] [k_1 Y_0(\beta R_1) + \beta Y_1(\beta R_1)] = 0.
 \end{aligned}$$

#### 4.2.6. Equation of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right) - bw + \Phi(r, t)$

For  $b > 0$  and  $\Phi \equiv 0$ , this is the Klein–Gordon equation describing one-dimensional wave phenomena with central symmetry. In the problems considered in Paragraphs 4.2.6-1 through 4.2.6-3, the solutions bounded at  $r = 0$  are sought; this is not specially stated below.

#### 4.2.6-1. Domain: $0 \leq r \leq R$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi \\ &\quad - a^2 \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R} d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(r, \xi, t) = \frac{2\xi}{Rr} \sum_{n=1}^{\infty} \sin\left(\frac{n\pi r}{R}\right) \sin\left(\frac{n\pi \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \pi^2 n^2}{R^2} + b.$$

#### 4.2.6-2. Domain: $0 \leq r \leq R$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R f_1(\xi) G(r, \xi, t) d\xi \\ &\quad + a^2 \int_0^t g(\tau) G(r, R, t - \tau) d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(r, \xi, t) = \frac{3\xi^2 \sin(t\sqrt{b})}{R^3 \sqrt{b}} + \frac{2\xi}{Rr} \sum_{n=1}^{\infty} \frac{\mu_n^2 + 1}{\mu_n^2 \sqrt{\lambda_n}} \sin\left(\frac{\mu_n r}{R}\right) \sin\left(\frac{\mu_n \xi}{R}\right) \sin(t\sqrt{\lambda_n}), \quad \lambda_n = \frac{a^2 \mu_n^2}{R^2} + b.$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\tan \mu - \mu = 0$ ; for the numerical values of the first five roots  $\mu_n$ , see Paragraph 1.2.3-5.

#### 4.2.6-3. Domain: $0 \leq r \leq R$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.6-2 where

$$G(r, \xi, t) = \frac{2\xi}{Rr} \sum_{n=1}^{\infty} \frac{\mu_n^2 + (kR - 1)^2}{\mu_n^2 + kR(kR - 1)} \sin\left(\frac{\mu_n r}{R}\right) \sin\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{R^2} + b.$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\mu \cot \mu + kR - 1 = 0$ . The numerical values of the six five roots  $\mu_n$  can be found in Carslaw and Jaeger (1984).

**4.2.6-4. Domain:  $R_1 \leq r \leq R_2$ . First boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 && \text{(initial condition),} \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 && \text{(initial condition),} \\ w &= g_1(t) \quad \text{at} \quad r = R_1 && \text{(boundary condition),} \\ w &= g_2(t) \quad \text{at} \quad r = R_2 && \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \int_0^t \int_{R_1}^{R_2} \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\ &+ a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_1} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R_2} d\tau, \end{aligned}$$

where

$$G(r, \xi, t) = \frac{2\xi}{(R_2 - R_1)r} \sum_{n=1}^{\infty} \sin \left[ \frac{\pi n(r - R_1)}{R_2 - R_1} \right] \sin \left[ \frac{\pi n(\xi - R_1)}{R_2 - R_1} \right] \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \pi^2 n^2}{(R_2 - R_1)^2} + b.$$

**4.2.6-5. Domain:  $R_1 \leq r \leq R_2$ . Second boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 && \text{(initial condition),} \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 && \text{(initial condition),} \\ \partial_r w &= g_1(t) \quad \text{at} \quad r = R_1 && \text{(boundary condition),} \\ \partial_r w &= g_2(t) \quad \text{at} \quad r = R_2 && \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \int_0^t \int_{R_1}^{R_2} \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_{R_1}^{R_2} f_0(\xi) G(r, \xi, t) d\xi + \int_{R_1}^{R_2} f_1(\xi) G(r, \xi, t) d\xi \\ &- a^2 \int_0^t g_1(\tau) G(r, R_1, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(r, R_2, t - \tau) d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \xi, t) &= \frac{3\xi^2 \sin(t\sqrt{b})}{(R_2^3 - R_1^3)\sqrt{b}} + \frac{2\xi}{(R_2 - R_1)r} \sum_{n=1}^{\infty} \frac{(1 + R_2^2 \lambda_n^2) \Psi_n(r) \Psi_n(\xi) \sin(t\sqrt{a^2 \lambda_n^2 + b})}{\lambda_n^2 [R_1^2 + R_2^2 + R_1 R_2 (1 + R_1 R_2 \lambda_n^2)] \sqrt{a^2 \lambda_n^2 + b}}, \\ \Psi_n(r) &= \sin[\lambda_n(r - R_1)] + R_1 \lambda_n \cos[\lambda_n(r - R_1)], \end{aligned}$$

where the  $\lambda_n$  are positive roots of the transcendental equation

$$(\lambda^2 R_1 R_2 + 1) \tan[\lambda(R_2 - R_1)] - \lambda(R_2 - R_1) = 0.$$

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, t)$  is determined by the formula in Paragraph 4.2.6-5 where

$$G(r, \xi, t) = \frac{2\xi}{r} \sum_{n=1}^{\infty} \frac{(b_2^2 + R_2^2 \lambda_n^2) \Psi_n(r) \Psi_n(\xi) \sin(t\sqrt{a^2 \lambda_n^2 + b})}{[(R_2 - R_1)(b_1^2 + R_1^2 \lambda_n^2)(b_2^2 + R_2^2 \lambda_n^2) + (b_1 R_2 + b_2 R_1)(b_1 b_2 + R_1 R_2 \lambda_n^2)] \sqrt{a^2 \lambda_n^2 + b}},$$

$$\Psi_n(r) = b_1 \sin[\lambda_n(r - R_1)] + R_1 \lambda_n \cos[\lambda_n(r - R_1)], \quad b_1 = k_1 R_1 + 1, \quad b_2 = k_2 R_2 - 1.$$

Here, the  $\lambda_n$  are positive roots of the transcendental equation

$$(b_1 b_2 - R_1 R_2 \lambda^2) \sin[\lambda(R_2 - R_1)] + \lambda(R_1 b_2 + R_2 b_1) \cos[\lambda(R_2 - R_1)] = 0.$$

## 4.3. Equations Containing Power Functions and Arbitrary Parameters

### 4.3.1. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = (ax + b) \frac{\partial^2 w}{\partial x^2} + c \frac{\partial w}{\partial x} + kw + \Phi(x, t)$

$$1. \quad \frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial}{\partial x} \left( x \frac{\partial w}{\partial x} \right) + \Phi(x, t).$$

For  $\Phi(x, t) \equiv 0$ , this equation governs small-amplitude free vibration of a hanging heavy homogeneous thread ( $a^2$  is the acceleration due to gravity,  $w$  the deflection of the thread from the vertical axis, and  $x$  the vertical coordinate).

1°. The substitution  $x = \frac{1}{4}r^2$  leads to the equation

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) + \Phi\left(\frac{1}{4}r^2, t\right),$$

which is discussed in Subsections 4.2.1–4.2.2.

2°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}), \\ w &\neq \infty \quad \text{at} \quad x = 0 \quad (\text{boundedness condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ &\quad - a^2 l \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{2}{a\sqrt{l}} \sum_{n=1}^{\infty} \frac{1}{\mu_n J_1^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \sin\left(\frac{\mu_n a t}{2\sqrt{l}}\right).$$

Here, the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ . The numerical values of the first ten roots  $\mu_n$  are specified in Paragraph 1.2.1-3.

⊙ Reference: M. M. Smirnov (1975).

3°. Domain:  $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= g(t) & \text{at } x &= l & (\text{boundary condition}), \\ w &\neq \infty & \text{at } x &= 0 & (\text{boundedness condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ &+ a^2 l \int_0^t g(\tau) G(x, l, t - \tau) d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{t}{l} + \frac{2}{a\sqrt{l}} \sum_{n=1}^{\infty} \frac{1}{\mu_n J_0^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \sin\left(\frac{\mu_n a t}{2\sqrt{l}}\right).$$

Here, the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu) = 0$ . The numerical values of the first ten roots  $\mu_n$  are specified in Paragraph 1.2.1-4.

4°. Domain:  $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w + kw &= g(t) & \text{at } x &= l & (\text{boundary condition}), \\ w &\neq \infty & \text{at } x &= 0 & (\text{boundedness condition}). \end{aligned}$$

The solution  $w(x, t)$  is given by the formula in Item 3° with

$$G(x, \xi, t) = \frac{2}{a\sqrt{l}} \sum_{n=1}^{\infty} \frac{\mu_n}{(4k^2 l + \mu_n^2) J_0^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \sin\left(\frac{\mu_n a t}{2\sqrt{l}}\right).$$

Here, the  $\mu_n$  are positive roots of the transcendental equation

$$\mu J_1(\mu) - 2k\sqrt{l} J_0(\mu) = 0.$$

The numerical values of the first six roots  $\mu_n$  can be found in Carslaw and Jaeger (1984).

2. 
$$\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial}{\partial x} \left( x \frac{\partial w}{\partial x} \right) - bw + \Phi(x, t).$$

For  $b < 0$  and  $\Phi(x, t) \equiv 0$ , this equation describes small-amplitude vibration of a heavy homogeneous thread that rotates at a constant angular velocity  $\omega = \sqrt{|b|}$  about the vertical axis ( $a^2$  is the acceleration due to gravity).

1°. The substitution  $x = \frac{1}{4}r^2$  leads to the equation

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) - bw + \Phi\left(\frac{1}{4}r^2, t\right),$$

which is discussed in Subsection 4.2.5.

2°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g(t) & \text{at } x &= l & (\text{boundary condition}), \\ w &\neq \infty & \text{at } x &= 0 & (\text{boundedness condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ &\quad - a^2 l \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(x, \xi, t) = \frac{1}{l} \sum_{n=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{4l} + b,$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

⊙ *Reference:* M. M. Smirnov (1975).

3°. Domain:  $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= g(t) & \text{at } x &= l & (\text{boundary condition}), \\ w &\neq \infty & \text{at } x &= 0 & (\text{boundedness condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ &\quad + a^2 l \int_0^t g(\tau) G(x, l, t - \tau) d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \xi, t) = \frac{\sin(t\sqrt{b})}{l\sqrt{b}} + \frac{1}{l} \sum_{n=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{4l} + b,$$

where the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu) = 0$ . The numerical values of the first ten roots  $\mu_n$  are specified in Paragraph 1.2.1-4.

4°. Domain:  $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w + kw &= g(t) & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, t)$  is given by the formula in Item 3° with

$$G(r, \xi, t) = \frac{1}{l} \sum_{n=1}^{\infty} \frac{\mu_n^2}{(4k^2 l + \mu_n^2) J_0^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{4l} + b.$$

Here, the  $\mu_n$  are positive roots of the transcendental equation

$$\mu J_1(\mu) - 2k\sqrt{l} J_0(\mu) = 0.$$

The numerical values of the first six roots  $\mu_n$  can be found in Abramowitz and Stegun (1964) and Carslaw and Jaeger (1984).

$$3. \quad \frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial}{\partial x} \left[ (l-x) \frac{\partial w}{\partial x} \right].$$

This equation governs small-amplitude free vibration of a heavy homogeneous thread of length  $l$  ( $a^2$  is the acceleration due to gravity,  $w$  the deflection of the thread from the vertical axis, and  $x$  the vertical coordinate). The change of variable  $z = l - x$  leads a special case of equation 4.3.1.1 with  $b = 0$  and  $\Phi \equiv 0$ .

$$4. \quad \frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{2}{2n+1} x \frac{\partial^2 w}{\partial x^2} + \frac{\partial w}{\partial x} \right), \quad n = 1, 2, \dots$$

General solution:

$$w(x, t) = \frac{\partial^{n-1}}{\partial x^{n-1}} \left[ \frac{\Phi(\sqrt{2(2n+1)x} + at) + \Psi(\sqrt{2(2n+1)x} - at)}{\sqrt{x}} \right],$$

where  $\Phi$  and  $\Psi$  are arbitrary functions.

⊙ *Reference:* M. M. Smirnov (1975).

$$5. \quad \frac{\partial^2 w}{\partial t^2} = (ax + b) \frac{\partial^2 w}{\partial x^2} + a \frac{\partial w}{\partial x} + cw + \Phi(x, t).$$

The substitution  $z = ax + b$  leads to an equation of the form 4.3.1.2:

$$\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial}{\partial z} \left( z \frac{\partial w}{\partial z} \right) + cw + \Phi \left( \frac{z-b}{a}, t \right).$$

$$6. \quad \frac{\partial^2 w}{\partial t^2} = (ax + b) \frac{\partial^2 w}{\partial x^2} + \frac{1}{2} a \frac{\partial w}{\partial x} + cw + \Phi(x, t).$$

The substitution  $z = 2\sqrt{ax+b}$  leads to the equation

$$\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial z^2} + cw + \Phi \left( \frac{z^2 - 4b}{4a}, t \right),$$

which is considered in Subsection 4.1.3.

$$7. \quad \frac{\partial^2 w}{\partial t^2} = (a_2 x + b_2) \frac{\partial^2 w}{\partial x^2} + (a_1 x + b_1) \frac{\partial w}{\partial x} + (a_0 x + b_0) w.$$

This is a special case of equation 4.5.3.4 with  $f(x) = a_2 x + b_2$ ,  $g(x) = a_1 x + b_1$ ,  $h(x) = a_0 x + b_0$ , and  $\Phi \equiv 0$ .

Particular solutions:

$$w(x, t) = \exp(kx) F \left( \frac{x+q}{p} \right) [A \sin(t\sqrt{\mu}) + B \cos(t\sqrt{\mu})] \quad \text{for } \mu > 0,$$

$$w(x, t) = \exp(kx) F \left( \frac{x+q}{p} \right) [A \sinh(t\sqrt{-\mu}) + B \cosh(t\sqrt{-\mu})] \quad \text{for } \mu < 0.$$

Here,  $A$ ,  $B$ , and  $\mu$  are arbitrary constants; the coefficients  $k$ ,  $p$ ,  $q$  and the function  $F = F(\xi)$  are listed in [Table 20](#), where

$$\mathcal{J}(\alpha, \beta; x) = C_1 \Phi(\alpha, \beta; x) + C_2 \Psi(\alpha, \beta; x), \quad C_1, C_2 \text{ are any numbers,}$$

is an arbitrary solution of the degenerate hypergeometric equation  $xy''_{xx} + (\beta - x)y'_x - \alpha y = 0$ , and

$$Z_\nu(x) = C_1 J_\nu(x) + C_2 Y_\nu(x), \quad C_1, C_2 \text{ are any numbers,}$$

is an arbitrary solution of the Bessel equation  $x^2 y''_{xx} + xy'_x + (x^2 - \nu^2)y = 0$ .



TABLE 20

The coefficients  $k, p, q$  and the function  $F = F(\xi)$  determining the form of particular solutions to equation 4.3.1.7. Notation:  $E(k) = b_2k^2 + b_1k + b_0 + \mu$

| Conditions                                           | $k$                           | $p$                        | $q$                                       | $F = F(\xi)$                                    | Parameters                                                                    |
|------------------------------------------------------|-------------------------------|----------------------------|-------------------------------------------|-------------------------------------------------|-------------------------------------------------------------------------------|
| $a_2 \neq 0, D \neq 0$<br>$D \equiv a_1^2 - 4a_0a_2$ | $\frac{\sqrt{D} - a_1}{2a_2}$ | $-\frac{a_2}{2a_2k + a_1}$ | $\frac{b_2}{a_2}$                         | $\mathcal{J}(\alpha, \beta; \xi)$               | $\alpha = E(k)/(2a_2k + a_1),$<br>$\beta = (a_2b_1 - a_1b_2)a_2^{-2}$         |
| $a_2 = 0,$<br>$a_1 \neq 0$                           | $-\frac{a_0}{a_1}$            | 1                          | $\frac{2b_2k + b_1}{a_1}$                 | $\mathcal{J}(\alpha, \frac{1}{2}; \sigma\xi^2)$ | $\alpha = E(k)/(2a_1),$<br>$\sigma = -a_1/(2b_2)$                             |
| $a_2 \neq 0,$<br>$a_1^2 = 4a_0a_2$                   | $-\frac{a_1}{2a_2}$           | $a_2$                      | $\frac{b_2}{a_2}$                         | $\xi^\alpha Z_{2\alpha}(\sigma\sqrt{\xi})$      | $\alpha = \frac{1}{2} - \frac{2b_2k + b_1}{2a_2},$<br>$\sigma = 2\sqrt{E(k)}$ |
| $a_2 = a_1 = 0,$<br>$a_0 \neq 0$                     | $-\frac{b_1}{2b_2}$           | 1                          | $\frac{4(b_0 + \mu)b_2 - b_1^2}{4a_0b_2}$ | $\xi^{1/2} Z_{1/3}(\sigma\xi^{3/2})$            | $\sigma = \frac{2}{3} \left( \frac{a_0}{b_2} \right)^{1/2}$                   |

For the degenerate hypergeometric functions  $\Phi(a, b; x)$  and  $\Psi(a, b; x)$ , see Supplement A.9 and the books by Abramowitz and Stegun (1964) and Bateman and Erdélyi (1953, Vol. 1). For the Bessel functions  $J_\nu(x)$  and  $Y_\nu(x)$ , see Supplement A.6 and the books by Abramowitz and Stegun (1964) and Bateman and Erdélyi (1953, Vol. 2).

### 4.3.2. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = (ax^2 + b)\frac{\partial^2 w}{\partial x^2} + cx\frac{\partial w}{\partial x} + kw + \Phi(x, t)$

1.  $\frac{\partial^2 w}{\partial t^2} = x^2 \frac{\partial^2 w}{\partial x^2} + \Phi(x, t).$

This is a special case of equation 4.3.2.2 with  $a = 1$  and  $b = c = 0$ .

1°. Domain:  $1 \leq x \leq a$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(t) & \text{at } x = 1 & \quad (\text{boundary condition}), \\ w &= g_2(t) & \text{at } x = a & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \int_0^t \int_1^a \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_1^a f_0(\xi) G(x, \xi, t) d\xi + \int_1^a f_1(\xi) G(x, \xi, t) d\xi \\ &+ \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=1} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=a} d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{2\sqrt{x}}{\xi^{3/2} \ln a} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \sin(\mu_n \ln x) \sin(\mu_n \ln \xi) \sin(\lambda_n t), \quad \mu_n = \frac{\pi n}{\ln a}, \quad \lambda_n = \sqrt{\mu_n^2 + \frac{1}{4}}.$$

2°. Domain:  $1 \leq x \leq a$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_x w &= g_1(t) & \text{at } x &= 1 & (\text{boundary condition}), \\ \partial_x w &= g_2(t) & \text{at } x &= a & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \int_0^t \int_1^a \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_1^a f_0(\xi) G(x, \xi, t) d\xi + \int_1^a f_1(\xi) G(x, \xi, t) d\xi \\ &- \int_0^t g_1(\tau) G(x, 1, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(x, a, t - \tau) d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, \xi, t) &= \frac{at}{(a-1)\xi^2} + \frac{8\sqrt{x}}{\xi^{3/2} \ln a} \sum_{n=1}^{\infty} \frac{\mu_n^2}{\lambda_n(1+\mu_n^2)} \varphi_n(x) \varphi_n(\xi) \sin(\lambda_n t), \\ \varphi_n(x) &= \cos(\mu_n \ln x) - \frac{1}{2\mu_n} \sin(\mu_n \ln x), \quad \mu_n = \frac{\pi n}{\ln a}, \quad \lambda_n = \sqrt{\mu_n^2 + \frac{1}{4}}. \end{aligned}$$

⊙ *Reference:* A. G. Butkovskiy (1979).

$$2. \quad \frac{\partial^2 w}{\partial t^2} = ax^2 \frac{\partial^2 w}{\partial x^2} + bx \frac{\partial w}{\partial x} + cw + \Phi(x, t).$$

The substitution  $x = ke^z$  ( $k \neq 0$ ) leads to the constant coefficient equation  $\partial_{tt} w = a \partial_{zz} w + (b-a) \partial_z w + cw + \Phi(ke^z, t)$ , which is discussed in Subsection 4.1.5.

$$3. \quad \frac{\partial^2 w}{\partial t^2} = (ax^2 + b) \frac{\partial^2 w}{\partial x^2} + ax \frac{\partial w}{\partial x} + cw.$$

The substitution  $z = \int \frac{dx}{\sqrt{ax^2 + b}}$  leads to the constant coefficient equation  $\partial_{tt} w = \partial_{zz} w + cw$ , which is discussed in Subsection 4.1.3.

$$4. \quad \frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial}{\partial x} \left[ (l^2 - x^2) \frac{\partial w}{\partial x} \right] + \Phi(x, t).$$

Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g(t) & \text{at } x &= 0 & (\text{boundary condition}), \\ w &\neq \infty & \text{at } x &= l & (\text{boundedness condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ &+ a^2 l^2 \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=0} d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(x, \xi, t) = \frac{1}{al} \sum_{n=1}^{\infty} \frac{4n-1}{\lambda_n} P_{2n-1} \left( \frac{x}{l} \right) P_{2n-1} \left( \frac{\xi}{l} \right) \sin(\lambda_n at), \quad \lambda_n = \sqrt{2n(2n-1)},$$

where  $P_k(x) = \frac{1}{2^k k!} \frac{d^k}{dx^k} [(x^2 - 1)^k]$  are the Legendre polynomials.

⊙ Reference: M. M. Smirnov (1975).

### 4.3.3. Other Equations

$$1. \quad \frac{\partial^2 w}{\partial t^2} = a^2 \left[ \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} - \frac{n(n+1)}{r^2} w \right], \quad n = 1, 2, 3, \dots$$

General solution:

$$w(r, t) = r^n \left( \frac{1}{r} \frac{\partial}{\partial r} \right)^n \left[ \frac{\Phi(r+at) + \Psi(r-at)}{r} \right],$$

where  $\Phi(r_1)$  and  $\Psi(r_2)$  are arbitrary functions.

⊙ Reference: M. M. Smirnov (1975).

$$2. \quad \frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial x^2} + \frac{\alpha}{x} \frac{\partial w}{\partial x}.$$

The hyperbolic Euler–Poisson–Darboux equation.

1°. For  $\alpha = 1$  and  $\alpha = 2$ , see Subsections 4.2.1–4.2.4. For  $\alpha \neq 1$ , the substitution  $z = x^{1-\alpha}$  leads to an equation of the form 4.5.3.1:

$$\frac{\partial^2 w}{\partial t^2} = (1-\alpha)^2 z^{\frac{2\alpha}{\alpha-1}} \frac{\partial^2 w}{\partial z^2}.$$

2°. Suppose  $w_\alpha = w_\alpha(x, t)$  is a solution of the equation in question for a fixed value of the parameter  $\alpha$ . Then the functions  $\tilde{w}_\alpha$  defined by the relations

$$\begin{aligned} \tilde{w}_\alpha &= \frac{\partial w_\alpha}{\partial t}, \\ \tilde{w}_\alpha &= x \frac{\partial w_\alpha}{\partial x} + t \frac{\partial w_\alpha}{\partial t}, \\ \tilde{w}_\alpha &= 2xt \frac{\partial w_\alpha}{\partial x} + (x^2 + t^2) \frac{\partial w_\alpha}{\partial t} + \alpha t w_\alpha \end{aligned}$$

are also solutions of this equation.

3°. Suppose  $w_\alpha = w_\alpha(x, t)$  is a solution of the equation in question for a fixed value of the parameter  $\alpha$ . Using this  $w_\alpha$ , one can construct solutions of the equation with other values of the parameter by the formulas

$$\begin{aligned} w_{2-\alpha} &= x^{\alpha-1} w_\alpha, \\ w_{\alpha-2} &= x \frac{\partial w_\alpha}{\partial x} + (\alpha-1) w_\alpha, \\ w_{\alpha-2} &= xt \frac{\partial w_\alpha}{\partial x} + x^2 \frac{\partial w_\alpha}{\partial t} + (\alpha-1) t w_\alpha, \\ w_{\alpha-2} &= x(x^2 + t^2) \frac{\partial w_\alpha}{\partial x} + 2x^2 t \frac{\partial w_\alpha}{\partial t} + [x^2 + (\alpha-1)t^2] w_\alpha, \\ w_{\alpha+2} &= \frac{1}{x} \frac{\partial w_\alpha}{\partial x}, \\ w_{\alpha+2} &= \frac{t}{x} \frac{\partial w_\alpha}{\partial x} + \frac{\partial w_\alpha}{\partial t}, \\ w_{\alpha+2} &= \frac{x^2 + t^2}{x} \frac{\partial w_\alpha}{\partial x} + 2t \frac{\partial w_\alpha}{\partial t} + \alpha w_\alpha. \end{aligned}$$

⊙ The results of Items 2° and 3° were obtained by A. V. Aksenov (2001).

$$3. \quad \frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial x^2} + \frac{2a}{x} \frac{\partial w}{\partial x} + b^2 w, \quad 0 < 2a < 1.$$

General solution:

$$w(x, t) = \int_0^1 \frac{\Phi(t + x(2\xi - 1))}{[\xi(1 - \xi)]^{1-a}} \bar{J}_{a-1}(2bx\sqrt{\xi(1 - \xi)}) d\xi \\ + x^{1-2a} \int_0^1 \frac{\Psi(t + x(2\xi - 1))}{[\xi(1 - \xi)]^a} \bar{J}_{-a}(2bx\sqrt{\xi(1 - \xi)}) d\xi,$$

where  $\Phi(\xi_1)$  and  $\Psi(\xi_2)$  are arbitrary functions;  $\bar{J}_{-\nu}(z) = \Gamma(1 - \nu)2^{-\nu}z^{\nu}J_{-\nu}(z)$ ;  $J_{-\nu}(z)$  is the Bessel function.

⊙ *Reference:* M. M. Smirnov (1975).

$$4. \quad \frac{\partial^2 w}{\partial t^2} = ax^4 \frac{\partial^2 w}{\partial x^2} + \Phi(x, t).$$

The transformation  $z = 1/x$ ,  $u = w/x$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a \frac{\partial^2 u}{\partial z^2} + z\Phi\left(\frac{1}{z}, t\right),$$

which is discussed in Subsection 4.1.2.

$$5. \quad \frac{\partial^2 w}{\partial t^2} = (ax + b)^4 \frac{\partial^2 w}{\partial x^2}.$$

The transformation

$$u = \frac{w}{ax + b}, \quad z = at + \frac{1}{ax + b}, \quad y = -at + \frac{1}{ax + b}$$

leads to the equation  $\partial_{zy}u = 0$ . Thus, the general solution of the original equation has the form

$$w = (ax + b)[f(z) + g(y)],$$

where  $f = f(z)$  and  $g = g(y)$  are arbitrary functions.

⊙ *Reference:* N. H. Ibragimov (1994).

$$6. \quad \frac{\partial^2 w}{\partial t^2} = (a^2 - x^2)^2 \frac{\partial^2 w}{\partial x^2} + \Phi(x, t).$$

Domain:  $-l \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= 0 & \text{at } x &= l & (\text{boundary condition}), \\ w &= 0 & \text{at } x &= -l & (\text{boundary condition}). \end{aligned}$$

Solution for  $0 < l < a$ :

$$w(x, t) = \frac{\partial}{\partial t} \int_{-l}^l f(\xi)G(x, \xi, t) d\xi + \int_{-l}^l g(\xi)G(x, \xi, t) d\xi + \int_0^t \int_{-l}^l \Phi(\xi, \tau)G(x, \xi, t - \tau) d\xi d\tau,$$

where

$$G(x, \xi, t) = \frac{2a}{k(\xi^2 - a^2)^2} \sum_{n=1}^{\infty} \frac{1}{\lambda_n} \varphi_n(x) \varphi_n(\xi) \sin(\lambda_n t), \\ \varphi_n(x) = \sqrt{a^2 - x^2} \sin\left(\frac{\pi n}{2} + \frac{\pi n}{2k} \ln \frac{a+x}{a-x}\right), \quad \lambda_n = \frac{a}{k} \sqrt{\pi^2 n^2 + k^2}, \quad k = \ln \frac{a+l}{a-l}.$$

⊙ *Reference:* A. G. Butkovskiy (1979).

$$7. \quad \frac{\partial^2 w}{\partial t^2} = (x - a_1)^2(x - a_2)^2 \frac{\partial^2 w}{\partial x^2}, \quad a_1 \neq a_2.$$

The transformation

$$w(x, t) = (x - a_2)u(\xi, \tau), \quad \xi = \ln \left| \frac{x - a_1}{x - a_2} \right|, \quad \tau = |a_1 - a_2|t$$

leads to the constant coefficient equation  $\partial_{\tau\tau} u = \partial_{\xi\xi} u - \partial_{\xi} u$ , which is discussed in Subsection 4.1.4.

$$8. \quad \frac{\partial^2 w}{\partial t^2} = (ax^2 + bx + c)^2 \frac{\partial^2 w}{\partial x^2}.$$

The transformation

$$w(x, t) = u(z, t) \sqrt{|ax^2 + bx + c|}, \quad z = \int \frac{dx}{ax^2 + bx + c}$$

leads to the constant coefficient equation  $\partial_{tt} u = \partial_{zz} u + (ac - \frac{1}{4}b^2)u$ , which is discussed in Subsection 4.1.3.

$$9. \quad \frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial}{\partial x} \left( x^m \frac{\partial w}{\partial x} \right) + \Phi(x, t).$$

1°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

1.1. Case  $0 < m < 1$ :

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= 0 & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l f_1(\xi) G(x, \xi, t) d\xi \\ &\quad - a^2 l^m \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau. \end{aligned} \quad (1)$$

Here,

$$G(x, \xi, t) = \sum_{n=1}^{\infty} \frac{y_n(x) y_n(\xi) \sin(\lambda_n a t)}{a \|y_n\|^2 \lambda_n}, \quad \lambda_n = \frac{\mu_n}{2} (2 - m) l^{\frac{m-2}{2}}, \quad (2)$$

where

$$y_n(x) = x^{\frac{1-m}{2}} J_p \left( \mu_n \left( \frac{x}{l} \right)^{\frac{2-m}{2}} \right), \quad \|y_n\|^2 = \int_0^l y_n^2(x) dx, \quad p = \left| \frac{1-m}{2-m} \right|;$$

the  $\mu_n$  are positive zeros of the Bessel function,  $J_p(\mu) = 0$ .

1.2. Case  $1 \leq m < 2$ :

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &\neq \infty & \text{at } x &= 0 & \text{(boundedness condition),} \\ w &= g(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

The solution is given by the formulas presented in Item 1.1.

⊙ *Reference:* M. M. Smirnov (1975).

2°. Domain:  $0 \leq x \leq l$ . Mixed boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & (\text{initial condition}), \\ (x^m \partial_x w) &= 0 & \text{at } x &= 0 & (\text{boundary condition}), \\ w &= g(t) & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

The solution for  $0 < m < 1$  is given by relations (1) and (2) with

$$y_n(x) = x^{\frac{1-m}{2}} J_{-p} \left( \mu_n \left( \frac{x}{l} \right)^{\frac{2-m}{2}} \right), \quad \|y_n\|^2 = \int_0^l y_n^2(x) dx, \quad p = \frac{1-m}{2-m};$$

the  $\mu_n$  are positive zeros of the Bessel function,  $J_{-p}(\mu) = 0$ .

3°. For  $\Phi \equiv 0$ , the change of variable  $z = x^{1-m}$  leads to an equation of the form 4.3.3.10:

$$\frac{\partial^2 w}{\partial t^2} = a^2 (1-m)^2 z^{\frac{m}{m-1}} \frac{\partial^2 w}{\partial z^2}.$$

10. 
$$\frac{\partial^2 w}{\partial t^2} = a^2 x^m \frac{\partial^2 w}{\partial x^2}.$$

1°. Particular solutions ( $A_1, A_2, B_1, B_2$ , and  $\mu$  are arbitrary constants):

$$\begin{aligned} w(x, t) &= \sqrt{x} \left[ A_1 J_{\frac{1}{2q}}(\mu x^q) + A_2 Y_{\frac{1}{2q}}(\mu x^q) \right] [B_1 \sin(aq\mu t) + B_2 \cos(aq\mu t)], \\ w(x, t) &= \sqrt{x} \left[ A_1 I_{\frac{1}{2q}}(\mu x^q) + A_2 K_{\frac{1}{2q}}(\mu x^q) \right] [B_1 \sinh(aq\mu t) + B_2 \cosh(aq\mu t)], \end{aligned}$$

where  $q = \frac{1}{2}(2-m)$ ;  $J_\nu(z)$  and  $Y_\nu(z)$  are the Bessel functions;  $I_\nu(z)$  and  $K_\nu(z)$  are the modified Bessel functions.

2°. Below are discrete transformations that preserve the form of the original equation; what changes is the parameter  $n$ .

2.1. The point transformation

$$z = \frac{1}{x}, \quad u = \frac{w}{x} \quad (\text{transformation } \mathcal{P})$$

leads to a similar equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 z^{4-m} \frac{\partial^2 u}{\partial z^2}.$$

The transformation  $\mathcal{P}$  changes the equation parameter in accordance with the rule  $m \xrightarrow{\mathcal{P}} 4-m$ . The double application of the transformation  $\mathcal{P}$  yields the original equation.

2.2. Suppose  $w = w(x, t)$  is a solution of the original equation. Then the function  $v = v(\xi, \tau)$ , which is related to the solution  $w = w(x, t)$  by the Bäcklund transformation

$$v(\xi, \tau) = \frac{\partial}{\partial x} w(x, t), \quad x = \xi^{\frac{1}{1-m}}, \quad \tau = |1-m|t \quad (\text{transformation } \mathcal{B}),$$

is a solution of a similar equation

$$\frac{\partial^2 v}{\partial \tau^2} = a^2 \xi^{\frac{m}{m-1}} \frac{\partial^2 v}{\partial \xi^2}.$$

The transformation  $\mathcal{B}$  changes the equation parameters in accordance with the rule  $m \xrightarrow{\mathcal{B}} \frac{m}{m-1}$ . The double application of the transformation  $\mathcal{B}$  yields the original equation.

2.3. The composition of transformations  $\mathcal{F} = \mathcal{B} \circ \mathcal{P}$  changes the equation parameter as follows:

$$m \xrightarrow{\mathcal{F}} \frac{4-m}{3-m} \xrightarrow{\mathcal{F}} \frac{8-3m}{5-2m} \xrightarrow{\mathcal{F}} \frac{12-5m}{7-3m} \xrightarrow{\mathcal{F}} \frac{16-7m}{9-4m} \xrightarrow{\mathcal{F}} \dots$$

The  $n$ -fold application of the transformation  $\mathcal{F}$  yields the equation with parameter

$$m \xrightarrow{\mathcal{F}^n} \frac{4n - (2n-1)m}{2n+1 - nm}. \quad (1)$$

2.4. The composition of transformations  $\mathcal{G} = \mathcal{P} \circ \mathcal{B}$  changes the equation parameter as follows:

$$m \xrightarrow{\mathcal{G}} \frac{4-3m}{1-m} \xrightarrow{\mathcal{G}} \frac{8-5m}{3-2m} \xrightarrow{\mathcal{G}} \frac{12-7m}{5-3m} \xrightarrow{\mathcal{G}} \frac{16-9m}{7-4m} \xrightarrow{\mathcal{G}} \dots$$

The  $n$ -fold application of the transformation  $\mathcal{G}$  yields the equation with parameter

$$m \xrightarrow{\mathcal{G}^n} \frac{4n - (2n+1)m}{2n-1 - nm}. \quad (2)$$

2.5. Setting  $m = 0$  in (1) and (2), we arrive at two families of equations

$$\begin{aligned} \frac{\partial^2 w}{\partial t^2} &= a^2 x^{\frac{4n}{2n+1}} \frac{\partial^2 w}{\partial x^2} \quad \text{at } n = 1, 2, \dots; \\ \frac{\partial^2 w}{\partial t^2} &= a^2 x^{\frac{4n}{2n-1}} \frac{\partial^2 w}{\partial x^2} \quad \text{at } n = 1, 2, \dots; \end{aligned}$$

whose solutions can be obtained with the aid of the wave equation; for this constant coefficient wave equation, see Subsection 4.1.1.

3°. Below are some useful transformations that lead to other equations.

3.1. The substitution  $\xi = x^{1-m}$  leads to an equation of the form 4.3.3.9:

$$\frac{\partial^2 w}{\partial t^2} = a^2 (1-m)^2 \frac{\partial}{\partial \xi} \left( \xi^{\frac{m}{m-1}} \frac{\partial w}{\partial \xi} \right).$$

3.2. The transformation  $\tau = \frac{1}{2}a|2-m|t$ ,  $\xi = x^{\frac{2-m}{2}}$  leads to an equation of the form 4.3.3.3:

$$\frac{\partial^2 w}{\partial \tau^2} = \frac{\partial^2 w}{\partial \xi^2} + \frac{m}{m-2} \frac{1}{\xi} \frac{\partial w}{\partial \xi}.$$

11. 
$$\frac{\partial^2 w}{\partial t^2} = t^m \frac{\partial^2 w}{\partial x^2}.$$

1°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) \quad \text{at } t = 0, \\ \partial_t w &= g(x) \quad \text{at } t = 0. \end{aligned}$$

Solution for  $m > 0$ :

$$\begin{aligned} w(x, t) &= \frac{\Gamma(2\beta)}{\Gamma^2(\beta)} \int_0^1 f\left(x + \frac{2}{m+2} t^{\frac{m+2}{2}} (2\xi - 1)\right) [\xi(1-\xi)]^{\beta-1} d\xi \\ &\quad + \frac{\Gamma(2-2\beta)}{\Gamma^2(1-\beta)} t \int_0^1 g\left(x + \frac{2}{m+2} t^{\frac{m+2}{2}} (2\xi - 1)\right) [\xi(1-\xi)]^{-\beta} d\xi, \end{aligned}$$

where

$$\beta = \frac{m}{2(m+2)}, \quad \Gamma(z) = \int_0^\infty e^{-s} s^{z-1} ds.$$

⊙ Reference: M. M. Smirnov (1975).

2°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= g(x) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= 0 & \text{at } x &= 0 & (\text{boundary condition}), \\ w &= 0 & \text{at } x &= l & (\text{boundary condition}). \end{aligned}$$

Solution for  $m > -1$ :

$$\begin{aligned} w(x, t) &= \sqrt{t} \sum_{n=1}^{\infty} \left[ A_n J_{-p} \left( 2p \lambda_n t^{\frac{1}{2p}} \right) + B_n J_p \left( 2p \lambda_n t^{\frac{1}{2p}} \right) \right] \sin(\lambda_n x), \\ A_n &= \Gamma(1-p)(\lambda_n p)^p \frac{2}{l} \int_0^l f(x) \sin(\lambda_n x) dx, \quad p = \frac{1}{m+2}, \\ B_n &= \Gamma(1+p)(\lambda_n p)^{-p} \frac{2}{l} \int_0^l g(x) \sin(\lambda_n x) dx, \quad \lambda_n = \frac{\pi n}{l}, \end{aligned}$$

where  $\Gamma(p)$  is the gamma function.

⊙ Reference: M. M. Smirnov (1975).

**12.**  $\frac{\partial^2 w}{\partial t^2} = t^m \frac{\partial^2 w}{\partial x^2} + b t^{\frac{m-2}{2}} \frac{\partial w}{\partial x}, \quad m \geq 2.$

Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0, \\ \partial_t w &= g(x) & \text{at } t &= 0. \end{aligned}$$

1°. Solution for  $|b| < \frac{1}{2}m$ :

$$\begin{aligned} w(x, t) &= \frac{\Gamma(\alpha + \beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 f \left( x + \frac{2}{m+2} t^{\frac{m+2}{2}} (2\xi - 1) \right) \xi^{\beta-1} (1-\xi)^{\alpha-1} d\xi \\ &+ \frac{\Gamma(2-\alpha-\beta)}{\Gamma(1-\alpha)\Gamma(1-\beta)} t \int_0^1 g \left( x + \frac{2}{m+2} t^{\frac{m+2}{2}} (2\xi - 1) \right) \xi^{-\alpha} (1-\xi)^{-\beta} d\xi, \end{aligned}$$

where

$$\alpha = \frac{m-2b}{2(m+2)}, \quad \beta = \frac{m+2b}{2(m+2)}, \quad \Gamma(z) = \int_0^{\infty} e^{-s} s^{z-1} ds.$$

2°. Solution for  $b = \frac{1}{2}m$ :

$$w(x, t) = f \left( x + \frac{2}{m+2} t^{\frac{m+2}{2}} \right) + \frac{2t}{m+2} \int_0^1 g \left( x + \frac{2}{m+2} t^{\frac{m+2}{2}} (2\xi - 1) \right) (1-\xi)^{-\frac{m}{m+2}} d\xi.$$

3°. Solution for  $b = -\frac{1}{2}m$ :

$$w(x, t) = f \left( x - \frac{2}{m+2} t^{\frac{m+2}{2}} \right) + \frac{2t}{m+2} \int_0^1 g \left( x + \frac{2}{m+2} t^{\frac{m+2}{2}} (2\xi - 1) \right) (1-\xi)^{-\frac{m}{m+2}} d\xi.$$

⊙ Reference: M. M. Smirnov (1975).

**13.**  $(b+x)^2 \frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial}{\partial x} \left[ (b+x)^2 \frac{\partial w}{\partial x} \right].$

General solution:

$$w(x, t) = \frac{f(x+at) + g(x-at)}{b+x},$$

where  $f(y)$  and  $g(z)$  are arbitrary functions.



## 4.4. Equations Containing the First Time Derivative

### 4.4.1. Equations of the Form $\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw + \Phi(x, t)$

1. 
$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \frac{\partial^2 w}{\partial x^2} + \Phi(x, t).$$

For  $\Phi(x, t) \equiv 0$ , this equation governs free transverse vibration of a string, and also longitudinal vibration of a rod in a resisting medium with a velocity-proportional resistance coefficient.

1°. The substitution  $w(x, t) = \exp(-\frac{1}{2}kt)u(x, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial x^2} + \frac{1}{4}k^2 u + \exp(\frac{1}{2}kt)\Phi(x, t),$$

which is considered in Subsection 4.1.3.

2°. Fundamental solution:

$$\mathcal{E}(x, t) = \frac{1}{2a} \vartheta(at - |x|) \exp(-\frac{1}{2}kt) I_0\left(\frac{1}{2}k\sqrt{t^2 - x^2/a^2}\right),$$

where  $\vartheta(z)$  is the Heaviside unit step function and  $I_0(z)$  is the modified Bessel function.

⊙ *Reference:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

3°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0, \\ \partial_t w &= g(x) & \text{at } t &= 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{1}{2} \exp(-\frac{1}{2}kt) [f(x+at) + f(x-at)] \\ &+ \frac{kt}{4a} \exp(-\frac{1}{2}kt) \int_{x-at}^{x+at} \frac{I_1(\frac{1}{2}k\sqrt{t^2 - (x-\xi)^2/a^2})}{\sqrt{t^2 - (x-\xi)^2/a^2}} f(\xi) d\xi \\ &+ \frac{1}{2a} \exp(-\frac{1}{2}kt) \int_{x-at}^{x+at} I_0(\frac{1}{2}k\sqrt{t^2 - (x-\xi)^2/a^2}) [g(\xi) + \frac{1}{2}kf(\xi)] d\xi \\ &+ \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \exp[-\frac{1}{2}k(t-\tau)] I_0(\frac{1}{2}k\sqrt{(t-\tau)^2 - (x-\xi)^2/a^2}) \Phi(\xi, \tau) d\xi d\tau, \end{aligned}$$

where  $I_0(z)$  and  $I_1(z)$  are the modified Bessel functions of the first kind.

4°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t-\tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l [f_1(\xi) + kf_0(\xi)] G(x, \xi, t) d\xi \\ &+ a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t-\tau) \right]_{\xi=0} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t-\tau) \right]_{\xi=l} d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{2}{l} \exp\left(-\frac{kt}{2}\right) \sum_{n=1}^{\infty} \sin\left(\frac{\pi n x}{l}\right) \sin\left(\frac{\pi n \xi}{l}\right) \frac{\sin(\lambda_n t)}{\lambda_n}, \quad \lambda_n = \sqrt{\frac{a^2 \pi^2 n^2}{l^2} - \frac{k^2}{4}}.$$

**Example.** Consider the homogeneous equation ( $\Phi \equiv 0$ ). The initial shape of the string is a triangle with base  $0 \leq x \leq l$  and height  $h$  at  $x = c$ , that is,

$$f(x) = \begin{cases} \frac{hx}{c} & \text{for } 0 \leq x \leq c, \\ \frac{h(l-x)}{l-c} & \text{for } c \leq x \leq l. \end{cases}$$

The initial velocities of the string points are zero,  $g(x) = 0$ .

Solution:

$$w(x, t) = \frac{2hl^2}{\pi^2 c(l-c)} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \frac{1}{n^2} \sin\left(\frac{n\pi c}{l}\right) \sin\left(\frac{n\pi x}{l}\right) \Theta_n(t),$$

where

$$\Theta_n(t) = \begin{cases} \cos(\lambda_n t) + \frac{k}{2\lambda_n} \sin(\lambda_n t) & \text{for } k < \frac{2\pi n a}{l}, \\ 1 + \frac{kt}{2} & \text{for } k = \frac{2\pi n a}{l}, \\ \cosh(\lambda_n t) + \frac{k}{2\lambda_n} \sinh(\lambda_n t) & \text{for } k > \frac{2\pi n a}{l}, \end{cases} \quad \lambda_n = \sqrt{\left| \frac{a^2 n^2 \pi^2}{l^2} - \frac{k^2}{4} \right|}.$$

⊙ *References:* M. M. Smirnov (1975), B. M. Budak, A. N. Tikhonov, and A. A. Samarskii (1980).

5°. For the second and third boundary value problems on the interval  $0 \leq x \leq l$ , see equation 4.4.1.2 (Items 5° and 6° with  $b = 0$ ).

$$2. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \frac{\partial^2 w}{\partial x^2} + bw + \Phi(x, t).$$

*Telegraph equation* (with  $k > 0$ ,  $b < 0$ , and  $\Phi(x, t) \equiv 0$ ).

1°. The substitution  $w(x, t) = \exp\left(-\frac{1}{2}kt\right) u(x, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial x^2} + (b + \frac{1}{4}k^2)u + \exp\left(\frac{1}{2}kt\right) \Phi(x, t),$$

which is considered in Subsection 4.1.3.

2°. Fundamental solutions:

$$\begin{aligned} \mathcal{E}(x, t) &= \frac{1}{2a} \vartheta(at - |x|) \exp\left(-\frac{1}{2}kt\right) I_0\left(c\sqrt{t^2 - x^2/a^2}\right) \quad \text{for } b + \frac{1}{4}k^2 = c^2 > 0, \\ \mathcal{E}(x, t) &= \frac{1}{2a} \vartheta(at - |x|) \exp\left(-\frac{1}{2}kt\right) J_0\left(c\sqrt{t^2 - x^2/a^2}\right) \quad \text{for } b + \frac{1}{4}k^2 = -c^2 < 0, \end{aligned}$$

where  $\vartheta(z)$  is the Heaviside unit step function,  $J_0(z)$  and  $J_1(z)$  are the Bessel functions, and  $I_0(z)$  and  $I_1(z)$  are the modified Bessel functions.

3°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) \quad \text{at } t = 0, \\ \partial_t w &= g(x) \quad \text{at } t = 0. \end{aligned}$$

Solution for  $b + \frac{1}{4}k^2 = c^2 > 0$ :

$$\begin{aligned} w(x, t) = & \frac{1}{2} \exp\left(-\frac{1}{2}kt\right) [f(x+at) + f(x-at)] \\ & + \frac{ct}{2a} \exp\left(-\frac{1}{2}kt\right) \int_{x-at}^{x+at} \frac{I_1(c\sqrt{t^2 - (x-\xi)^2/a^2})}{\sqrt{t^2 - (x-\xi)^2/a^2}} f(\xi) d\xi \\ & + \frac{1}{2a} \exp\left(-\frac{1}{2}kt\right) \int_{x-at}^{x+at} I_0(c\sqrt{t^2 - (x-\xi)^2/a^2}) [g(\xi) + \frac{1}{2}kf(\xi)] d\xi \\ & + \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \exp\left[-\frac{1}{2}k(t-\tau)\right] I_0(c\sqrt{(t-\tau)^2 - (x-\xi)^2/a^2}) \Phi(\xi, \tau) d\xi d\tau. \end{aligned}$$

Solution for  $b + \frac{1}{4}k^2 = -c^2 < 0$ :

$$\begin{aligned} w(x, t) = & \frac{1}{2} \exp\left(-\frac{1}{2}kt\right) [f(x+at) + f(x-at)] \\ & - \frac{ct}{2a} \exp\left(-\frac{1}{2}kt\right) \int_{x-at}^{x+at} \frac{J_1(c\sqrt{t^2 - (x-\xi)^2/a^2})}{\sqrt{t^2 - (x-\xi)^2/a^2}} f(\xi) d\xi \\ & + \frac{1}{2a} \exp\left(-\frac{1}{2}kt\right) \int_{x-at}^{x+at} J_0(c\sqrt{t^2 - (x-\xi)^2/a^2}) [g(\xi) + \frac{1}{2}kf(\xi)] d\xi \\ & + \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \exp\left[-\frac{1}{2}k(t-\tau)\right] J_0(c\sqrt{(t-\tau)^2 - (x-\xi)^2/a^2}) \Phi(\xi, \tau) d\xi d\tau. \end{aligned}$$

4°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w = f_0(x) \quad & \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w = f_1(x) \quad & \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w = g_1(t) \quad & \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w = g_2(t) \quad & \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) = & \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ & + \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l [f_1(\xi) + kf_0(\xi)] G(x, \xi, t) d\xi \\ & + a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=0} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau. \end{aligned}$$

Let  $a^2\pi^2 - bl^2 - \frac{1}{4}k^2l^2 > 0$ . Then

$$G(x, \xi, t) = \frac{2}{l} \exp\left(-\frac{kt}{2}\right) \sum_{n=1}^{\infty} \sin\left(\frac{\pi nx}{l}\right) \sin\left(\frac{\pi n\xi}{l}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2\pi^2 n^2}{l^2} - b - \frac{k^2}{4}.$$

Let  $a^2\pi^2 n^2 - bl^2 - \frac{1}{4}k^2l^2 \leq 0$  for  $n = 1, \dots, m$  and  $a^2\pi^2 n^2 - bl^2 - \frac{1}{4}k^2l^2 > 0$  for  $n = m+1, m+2, \dots$ . Then

$$\begin{aligned} G(x, \xi, t) = & \frac{2}{l} \exp\left(-\frac{kt}{2}\right) \sum_{n=1}^m \sin\left(\frac{\pi nx}{l}\right) \sin\left(\frac{\pi n\xi}{l}\right) \frac{\sinh(t\sqrt{\beta_n})}{\sqrt{\beta_n}} \\ & + \frac{2}{l} \exp\left(-\frac{kt}{2}\right) \sum_{n=m+1}^{\infty} \sin\left(\frac{\pi nx}{l}\right) \sin\left(\frac{\pi n\xi}{l}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \\ & \beta_n = b + \frac{k^2}{4} - \frac{a^2\pi^2 n^2}{l^2}, \quad \lambda_n = \frac{a^2\pi^2 n^2}{l^2} - b - \frac{k^2}{4}. \end{aligned}$$

5°. Domain:  $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w &= g_1(t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w &= g_2(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l [f_1(\xi) + k f_0(\xi)] G(x, \xi, t) d\xi \\ &- a^2 \int_0^t g_1(\tau) G(x, 0, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau. \end{aligned}$$

For  $p = b + \frac{1}{4}k^2 < 0$ ,

$$G(x, \xi, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{\sin(t\sqrt{|p|})}{l\sqrt{|p|}} + \frac{2}{l} \sum_{n=1}^{\infty} \cos(\mu_n x) \cos(\mu_n \xi) \frac{\sin(t\sqrt{a^2\mu_n^2 - p})}{\sqrt{a^2\mu_n^2 - p}} \right], \quad \mu_n = \frac{\pi n}{l}.$$

For  $p = b + \frac{1}{4}k^2 > 0$ ,

$$G(x, \xi, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{\sinh(t\sqrt{p})}{l\sqrt{p}} + \frac{2}{l} \sum_{n=1}^{\infty} \cos(\mu_n x) \cos(\mu_n \xi) \frac{\sin(t\sqrt{a^2\mu_n^2 - p})}{\sqrt{a^2\mu_n^2 - p}} \right], \quad \mu_n = \frac{\pi n}{l}.$$

If the inequality  $a^2\mu_n^2 - p < 0$  holds for several first values  $n = 1, \dots, m$ , then the expressions  $\sqrt{a^2\mu_n^2 - p}$  should be replaced by  $\sqrt{|a^2\mu_n^2 - p|}$  and the sines by the hyperbolic sines in the corresponding terms of the series.

6°. Domain:  $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - s_1 w &= g_1(t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w + s_2 w &= g_2(t) & \text{at } x &= l & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, t)$  is determined by the formula in Item 5° with

$$\begin{aligned} G(x, \xi, t) &= \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \frac{y_n(x)y_n(\xi) \sin(t\sqrt{a^2\mu_n^2 - p})}{B_n \sqrt{a^2\mu_n^2 - p}}, \quad p = b + \frac{1}{4}k^2, \\ y_n(x) &= \cos(\mu_n x) + \frac{s_1}{\mu_n} \sin(\mu_n x), \quad B_n = \frac{s_2}{2\mu_n^2} \frac{\mu_n^2 + s_1^2}{\mu_n^2 + s_2^2} + \frac{s_1}{2\mu_n^2} + \frac{l}{2} \left(1 + \frac{s_1^2}{\mu_n^2}\right). \end{aligned}$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\frac{\tan(\mu l)}{\mu} = \frac{s_1 + s_2}{\mu^2 - s_1 s_2}$ .

If the inequality  $a^2\mu_n^2 - p < 0$  holds for several first values  $n = 1, \dots, m$ , then the expressions  $\sqrt{a^2\mu_n^2 - p}$  should be replaced by  $\sqrt{|a^2\mu_n^2 - p|}$  and the sines by the hyperbolic sines in the corresponding terms of the series.

$$3. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw + \Phi(x, t).$$

1°. The substitution  $w(x, t) = \exp(-\frac{1}{2}a^{-2}bx - \frac{1}{2}kt)u(x, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial^2 u}{\partial x^2} + (c + \frac{1}{4}k^2 - \frac{1}{4}a^{-2}b^2)u + \exp(\frac{1}{2}a^{-2}bx + \frac{1}{2}kt)\Phi(x, t),$$

which is discussed in Subsection 4.1.3.

2°. Fundamental solutions:

$$\mathcal{E}(x, t) = \frac{1}{2a} \vartheta(at - |x|) \exp\left(-\frac{bx}{2a^2} - \frac{kt}{2}\right) I_0\left(\sigma \sqrt{t^2 - \frac{x^2}{a^2}}\right) \quad \text{if } c + \frac{k^2}{4} - \frac{b^2}{4a^2} = \sigma^2 > 0,$$

$$\mathcal{E}(x, t) = \frac{1}{2a} \vartheta(at - |x|) \exp\left(-\frac{bx}{2a^2} - \frac{kt}{2}\right) J_0\left(\sigma \sqrt{t^2 - \frac{x^2}{a^2}}\right) \quad \text{if } c + \frac{k^2}{4} - \frac{b^2}{4a^2} = -\sigma^2 < 0,$$

where  $\vartheta(z)$  is the Heaviside unit step function,  $J_0(z)$  and  $J_1(z)$  are the Bessel functions, and  $I_0(z)$  and  $I_1(z)$  are the modified Bessel functions.

3°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) & \text{at } t &= 0, \\ \partial_t w &= g(x) & \text{at } t &= 0. \end{aligned}$$

Solution for  $c + \frac{1}{4}k^2 - \frac{1}{4}a^{-2}b^2 = \sigma^2 > 0$ :

$$\begin{aligned} w(x, t) &= \frac{1}{2} \exp\left(-\frac{kt}{2}\right) \left[ f(x+at) \exp\left(\frac{bt}{2a}\right) + f(x-at) \exp\left(-\frac{bt}{2a}\right) \right] \\ &+ \frac{\sigma t}{2a} \exp\left(-\frac{bx}{2a^2} - \frac{kt}{2}\right) \int_{x-at}^{x+at} \exp\left(\frac{b\xi}{2a^2}\right) \frac{I_1(\sigma \sqrt{t^2 - (x-\xi)^2/a^2})}{\sqrt{t^2 - (x-\xi)^2/a^2}} f(\xi) d\xi \\ &+ \frac{1}{2a} \exp\left(-\frac{bx}{2a^2} - \frac{kt}{2}\right) \int_{x-at}^{x+at} \exp\left(\frac{b\xi}{2a^2}\right) I_0(\sigma \sqrt{t^2 - (x-\xi)^2/a^2}) \left[g(\xi) + \frac{1}{2}kf(\xi)\right] d\xi \\ &+ \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \exp\left[\frac{b(\xi-x)}{2a^2} - \frac{k(t-\tau)}{2}\right] I_0(\sigma \sqrt{(t-\tau)^2 - (x-\xi)^2/a^2}) \Phi(\xi, \tau) d\xi d\tau. \end{aligned}$$

Solution for  $c + \frac{1}{4}k^2 - \frac{1}{4}a^{-2}b^2 = -\sigma^2 < 0$ :

$$\begin{aligned} w(x, t) &= \frac{1}{2} \exp\left(-\frac{kt}{2}\right) \left[ f(x+at) \exp\left(\frac{bt}{2a}\right) + f(x-at) \exp\left(-\frac{bt}{2a}\right) \right] \\ &- \frac{\sigma t}{2a} \exp\left(-\frac{bx}{2a^2} - \frac{kt}{2}\right) \int_{x-at}^{x+at} \exp\left(\frac{b\xi}{2a^2}\right) \frac{J_1(\sigma \sqrt{t^2 - (x-\xi)^2/a^2})}{\sqrt{t^2 - (x-\xi)^2/a^2}} f(\xi) d\xi \\ &+ \frac{1}{2a} \exp\left(-\frac{bx}{2a^2} - \frac{kt}{2}\right) \int_{x-at}^{x+at} \exp\left(\frac{b\xi}{2a^2}\right) J_0(\sigma \sqrt{t^2 - (x-\xi)^2/a^2}) \left[g(\xi) + \frac{1}{2}kf(\xi)\right] d\xi \\ &+ \frac{1}{2a} \int_0^t \int_{x-a(t-\tau)}^{x+a(t-\tau)} \exp\left[\frac{b(\xi-x)}{2a^2} - \frac{k(t-\tau)}{2}\right] J_0(\sigma \sqrt{(t-\tau)^2 - (x-\xi)^2/a^2}) \Phi(\xi, \tau) d\xi d\tau. \end{aligned}$$

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

4°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l [f_1(\xi) + k f_0(\xi)] G(x, \xi, t) d\xi \\ &+ a^2 \int_0^t g_1(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=0} d\tau - a^2 \int_0^t g_2(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau. \end{aligned}$$

Let  $a^2 \pi^2 + \frac{1}{4} a^{-2} b^2 l^2 - c l^2 - \frac{1}{4} k^2 l^2 > 0$ . Then

$$\begin{aligned} G(x, \xi, t) &= \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} - \frac{kt}{2} \right] \sum_{n=1}^{\infty} \sin \left( \frac{\pi n x}{l} \right) \sin \left( \frac{\pi n \xi}{l} \right) \frac{\sinh(t \sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \\ \lambda_n &= \frac{a^2 \pi^2 n^2}{l^2} + \frac{b^2}{4a^2} - c - \frac{k^2}{4}. \end{aligned}$$

Let

$$\begin{aligned} a^2 \pi^2 n^2 + \frac{1}{4} a^{-2} b^2 l^2 - c l^2 - \frac{1}{4} k^2 l^2 &\leq 0 \quad \text{for} \quad n = 1, \dots, m; \\ a^2 \pi^2 n^2 + \frac{1}{4} a^{-2} b^2 l^2 - c l^2 - \frac{1}{4} k^2 l^2 &> 0 \quad \text{for} \quad n = m + 1, m + 2, \dots \end{aligned}$$

Then

$$\begin{aligned} G(x, \xi, t) &= \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} - \frac{kt}{2} \right] \sum_{n=1}^m \sin \left( \frac{\pi n x}{l} \right) \sin \left( \frac{\pi n \xi}{l} \right) \frac{\sinh(t \sqrt{\beta_n})}{\sqrt{\beta_n}} \\ &+ \frac{2}{l} \exp \left[ \frac{b(\xi - x)}{2a^2} - \frac{kt}{2} \right] \sum_{n=m+1}^{\infty} \sin \left( \frac{\pi n x}{l} \right) \sin \left( \frac{\pi n \xi}{l} \right) \frac{\sin(t \sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \end{aligned}$$

where  $\beta_n = c + \frac{k^2}{4} - \frac{a^2 \pi^2 n^2}{l^2} - \frac{b^2}{4a^2}$  and  $\lambda_n = \frac{a^2 \pi^2 n^2}{l^2} + \frac{b^2}{4a^2} - c - \frac{k^2}{4}$ .

⊙ Reference: A. G. Butkovskiy (1979).

5°. Domain:  $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l [f_1(\xi) + k f_0(\xi)] G(x, \xi, t) d\xi \\ &- a^2 \int_0^t g_1(\tau) G(x, 0, t - \tau) d\tau + a^2 \int_0^t g_2(\tau) G(x, l, t - \tau) d\tau. \end{aligned}$$

For  $p = c + \frac{1}{4}k^2 < 0$ ,

$$G(x, \xi, t) = A \exp\left(\frac{b\xi}{a^2} - \frac{kt}{2}\right) \frac{\sin(t\sqrt{|p|})}{\sqrt{|p|}} + \frac{2}{l} \exp\left[\frac{b(\xi - x)}{2a^2} - \frac{kt}{2}\right] \sum_{n=1}^{\infty} \frac{y_n(x)y_n(\xi)}{1 + \mu_n^2} \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}},$$

where

$$A = \frac{b}{a^2(e^{bl/a^2} - 1)}, \quad \lambda_n = \frac{a^2\pi^2 n^2}{l^2} + \frac{b^2}{4a^2} - c - \frac{k^2}{4},$$

$$y_n(x) = \cos\left(\frac{\pi n x}{l}\right) + \mu_n \sin\left(\frac{\pi n x}{l}\right), \quad \mu_n = \frac{bl}{2a^2\pi n}.$$

For  $p = c + \frac{1}{4}k^2 > 0$ ,

$$G(x, \xi, t) = A \exp\left(\frac{b\xi}{a^2} - \frac{kt}{2}\right) \frac{\sinh(t\sqrt{p})}{\sqrt{p}} + \frac{2}{l} \exp\left[\frac{b(\xi - x)}{2a^2} - \frac{kt}{2}\right] \sum_{n=1}^{\infty} \frac{y_n(x)y_n(\xi)}{1 + \mu_n^2} \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}},$$

where the coefficient  $A$ ,  $\lambda_n$ ,  $\mu_n$  and the functions  $y_n(x)$  remain as before. If the inequality  $\lambda_n < 0$  holds for several first values  $n = 1, \dots, m$ , then the expressions  $\sqrt{\lambda_n}$  must be replaced by  $\sqrt{|\lambda_n|}$  and the sines by the hyperbolic sines in the corresponding terms of the series.

6°. Domain:  $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - s_1 w &= g_1(t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w + s_2 w &= g_2(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, t)$  is determined by the formula in Item 5° with

$$G(x, \xi, t) = \exp\left[\frac{b(\xi - x)}{2a^2} - \frac{kt}{2}\right] \sum_{n=1}^{\infty} \frac{y_n(x)y_n(\xi) \sin(t\sqrt{\lambda_n})}{B_n \sqrt{\lambda_n}}.$$

Here,

$$\begin{aligned} y_n(x) &= \cos(\mu_n x) + \frac{2a^2 s_1 + b}{2a^2 \mu_n} \sin(\mu_n x), \quad \lambda_n = a^2 \mu_n^2 + \frac{b^2}{4a^2} - c - \frac{k^2}{4}, \\ B_n &= \frac{2a^2 s_2 - b}{4a^2 \mu_n^2} \frac{4a^4 \mu_n^2 + (2a^2 s_1 + b)^2}{4a^4 \mu_n^2 + (2a^2 s_2 - b)^2} + \frac{2a^2 s_1 + b}{4a^2 \mu_n^2} + \frac{l}{2} + \frac{l(2a^2 s_1 + b)^2}{8a^4 \mu_n^2}, \end{aligned}$$

where the  $\mu_n$  are positive roots of the transcendental equation

$$\frac{\tan(\mu l)}{\mu} = \frac{4a^4(s_1 + s_2)}{4a^4\mu^2 - (2a^2 s_1 + b)(2a^2 s_2 - b)}.$$

#### 4.4.2. Equations of the Form

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x) \frac{\partial w}{\partial x} + h(x)w + \Phi(x, t)$$

$$1. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right).$$

This equation describes vibration of a circular membrane in a resisting medium with velocity-proportional resistance coefficient.

1°. Domain:  $0 \leq r \leq R$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= g(r) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= 0 & \text{at } x &= R & \text{(boundary condition).} \end{aligned}$$

Solution:

$$w(r, t) = \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} [A_n \cos(\lambda_n t) + B_n \sin(\lambda_n t)] J_0\left(\frac{\mu_n r}{R}\right), \quad \lambda_n = \sqrt{\frac{a^2 \mu_n^2}{R^2} - \frac{k^2}{4}}.$$

Here,

$$A_n = \frac{2}{R^2 J_1^2(\mu_n)} \int_0^R f(r) J_0\left(\frac{\mu_n r}{R}\right) r dr, \quad B_n = \frac{A_n k}{2\lambda_n} + \frac{2}{\lambda_n R^2 J_1^2(\mu_n)} \int_0^R g(r) J_0\left(\frac{\mu_n r}{R}\right) r dr,$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

2°. For the solution of the second and third boundary value problems, see equation 4.4.2.2 (Items 3° and 4° with  $b = 0$ ).

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

$$2. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} \right) - bw + \Phi(r, t).$$

1°. The substitution  $w(r, t) = \exp\left(-\frac{1}{2}kt\right)u(r, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \left( \frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} \right) - \left(b - \frac{1}{4}k^2\right)u + \exp\left(\frac{1}{2}kt\right)\Phi(r, t),$$

which is discussed in Subsection 4.2.5.

2°. Domain:  $0 \leq r \leq R$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g(t) & \text{at } r &= R & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R [f_1(\xi) + k f_0(\xi)] G(r, \xi, t) d\xi \\ &\quad - a^2 \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R} d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \xi, t) = \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \frac{2\xi}{R^2 J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \mu_n^2}{R^2} + b - \frac{k^2}{4},$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ . The numerical values of the first ten  $\mu_n$  are specified in Paragraph 1.2.1-3.



3°. Domain:  $0 \leq r \leq R$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi + \int_0^R [f_1(\xi) + k f_0(\xi)] G(r, \xi, t) d\xi \\ &\quad + a^2 \int_0^t g(\tau) G(r, R, t - \tau) d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \xi, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{2\xi \sin(t\sqrt{\lambda_0})}{R^2 \sqrt{\lambda_0}} + \frac{2}{R^2} \sum_{n=1}^{\infty} \frac{\xi}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}} \right],$$

where  $\lambda_0 = b - \frac{1}{4}k^2$ ;  $\lambda_n = a^2 \mu_n^2 R^{-2} + b - \frac{1}{4}k^2$ ; the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu) = 0$ . The numerical values of the first ten roots  $\mu_n$  are specified in Paragraph 1.2.1-4.

4°. Domain:  $0 \leq r \leq R$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + sw &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, t)$  is given by the formula in Item 3° with

$$G(r, \xi, t) = \frac{2}{R^2} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \frac{\mu_n^2 \xi}{(s^2 R^2 + \mu_n^2) J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}.$$

Here,  $\lambda_n = a^2 \mu_n^2 R^{-2} + b - \frac{1}{4}k^2$  and the  $\mu_n$  are positive roots of the transcendental equation

$$\mu J_1(\mu) - sR J_0(\mu) = 0.$$

The numerical values of the first six roots  $\mu_n$  can be found in Abramowitz and Stegun (1964) and Carslaw and Jaeger (1984).

$$3. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{2}{r} \frac{\partial w}{\partial r} \right) - bw + \Phi(r, t).$$

1°. The substitution  $w(r, t) = \exp\left(-\frac{1}{2}kt\right)u(r, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \left( \frac{\partial^2 u}{\partial r^2} + \frac{2}{r} \frac{\partial u}{\partial r} \right) - \left(b - \frac{1}{4}k^2\right)u + \exp\left(\frac{1}{2}kt\right)\Phi(r, t),$$

which is discussed in Subsection 4.2.6.

2°. Domain:  $0 \leq r \leq R$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi \int_0^R [f_1(\xi) + k f_0(\xi)] G(r, \xi, t) d\xi \\ &\quad - a^2 \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(r, \xi, t - \tau) \right]_{\xi=R} d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(r, \xi, t) = \frac{2\xi}{Rr} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \sin\left(\frac{n\pi r}{R}\right) \sin\left(\frac{n\pi \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}, \quad \lambda_n = \frac{a^2 \pi^2 n^2}{R^2} + b - \frac{k^2}{4}.$$

3°. Domain:  $0 \leq r \leq R$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \frac{\partial}{\partial t} \int_0^R f_0(\xi) G(r, \xi, t) d\xi \int_0^R [f_1(\xi) + k f_0(\xi)] G(r, \xi, t) d\xi \\ &\quad + a^2 \int_0^t g(\tau) G(r, R, t - \tau) d\tau + \int_0^t \int_0^R \Phi(\xi, \tau) G(r, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(r, \xi, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{3\xi^2 \sin(t\sqrt{\lambda_0})}{R^3 \sqrt{\lambda_0}} + \frac{2\xi}{Rr} \sum_{n=1}^{\infty} \frac{\mu_n^2 + 1}{\mu_n^2 \sqrt{\lambda_n}} \sin\left(\frac{\mu_n r}{R}\right) \sin\left(\frac{\mu_n \xi}{R}\right) \sin(t\sqrt{\lambda_n}) \right].$$

Here,  $\lambda_0 = b - \frac{1}{4}k^2$ ;  $\lambda_n = a^2 \mu_n^2 R^{-2} + b - \frac{1}{4}k^2$ ; and the  $\mu_n$  are positive roots of the transcendental equation  $\tan \mu - \mu = 0$ . The numerical values of the first five roots  $\mu_n$  are specified in Paragraph 1.2.1-5.

4°. Domain:  $0 \leq r \leq R$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + sw &= g(t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, t)$  is given by the formula in Item 3° with

$$G(r, \xi, t) = \frac{2\xi}{Rr} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \frac{\mu_n^2 + (sR - 1)^2}{\mu_n^2 + sR(sR - 1)} \sin\left(\frac{\mu_n r}{R}\right) \sin\left(\frac{\mu_n \xi}{R}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}.$$

Here,  $\lambda_n = a^2 \mu_n^2 R^{-2} + b - \frac{1}{4}k^2$  and the  $\mu_n$  are positive roots of the transcendental equation  $\mu \cot \mu + sR - 1 = 0$ . The numerical values of the first six roots  $\mu_n$  can be found in Carslaw and Jaeger (1984).

$$4. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \frac{\partial}{\partial x} \left( x \frac{\partial w}{\partial x} \right) - bw + \Phi(x, t).$$

1°. The substitution  $w(r, t) = \exp(-\frac{1}{2}kt)u(r, t)$  leads to an equation of the form 4.3.1.2:

$$\frac{\partial^2 u}{\partial t^2} = a^2 \frac{\partial}{\partial x} \left( x \frac{\partial u}{\partial x} \right) - \left(b - \frac{1}{4}k^2\right)u + \exp\left(\frac{1}{2}kt\right)\Phi(x, t).$$

2°. Domain:  $0 \leq x \leq l$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g(t) & \text{at } x &= l & \text{(boundary condition),} \\ w &\neq \infty & \text{at } x &= 0 & \text{(boundedness condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l [f_1(\xi) + k f_0(\xi)] G(x, \xi, t) d\xi \\ &\quad - a^2 l \int_0^t g(\tau) \left[ \frac{\partial}{\partial \xi} G(x, \xi, t - \tau) \right]_{\xi=l} d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(x, \xi, t) = \frac{1}{l} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}.$$

Here,  $\lambda_n = \frac{1}{4}a^2\mu_n^2 l^{-1} + b - \frac{1}{4}k^2$ ; the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

3°. Domain:  $0 \leq x \leq l$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w &= g(t) & \text{at } x &= l & \text{(boundary condition),} \\ w &\neq \infty & \text{at } x &= 0 & \text{(boundedness condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, t) &= \frac{\partial}{\partial t} \int_0^l f_0(\xi) G(x, \xi, t) d\xi + \int_0^l [f_1(\xi) + k f_0(\xi)] d\xi \\ &\quad + a^2 l \int_0^t g(\tau) G(x, l, t - \tau) d\tau + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$G(r, \xi, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{\sin(t\sqrt{\lambda_0})}{l\sqrt{\lambda_0}} + \frac{1}{l} \sum_{n=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}} \right].$$

Here,  $\lambda_0 = b - \frac{1}{4}k^2$ ;  $\lambda_n = \frac{1}{4}a^2\mu_n^2 l^{-1} + b - \frac{1}{4}k^2$ ; the  $\mu_n$  are positive zeros of the first-order Bessel function,  $J_1(\mu) = 0$ . The numerical values of the first ten roots  $\mu_n$  are specified in Paragraph 1.2.1-5.

4°. Domain:  $0 \leq x \leq l$ . Third boundary value problem.

The following conditions are prescribed:

$$w = f_0(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\partial_t w = f_1(x) \quad \text{at} \quad t = 0 \quad (\text{initial condition}),$$

$$\partial_x w + kw = g(t) \quad \text{at} \quad x = l \quad (\text{boundary condition}).$$

The solution  $w(x, t)$  is given by the formula in Item 3° with

$$G(r, \xi, t) = \frac{1}{l} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \frac{\mu_n^2}{(4k^2l + \mu_n^2)J_0^2(\mu_n)} J_0\left(\mu_n \sqrt{\frac{x}{l}}\right) J_0\left(\mu_n \sqrt{\frac{\xi}{l}}\right) \frac{\sin(t\sqrt{\lambda_n})}{\sqrt{\lambda_n}}.$$

Here,  $\lambda_n = \frac{1}{4}a^2\mu_n^2l^{-1} + b - \frac{1}{4}k^2$ , and the  $\mu_n$  are positive roots of the transcendental equation

$$\mu J_1(\mu) - 2k\sqrt{l} J_0(\mu) = 0.$$

The numerical values of the first six roots  $\mu_n$  can be found in Abramowitz and Stegun (1964) and Carslaw and Jaeger (1984).

$$5. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = (ax^m + b) \frac{\partial^2 w}{\partial x^2} + \frac{1}{2} amx^{m-1} \frac{\partial w}{\partial x} + cw.$$

The substitution  $z = \int \frac{dx}{\sqrt{ax^m + b}}$  leads to a constant coefficient equation of the form 4.4.1.2:

$$\partial_{tt}w + k\partial_t w = \partial_{zz}w + cw.$$

### 4.4.3. Other Equations

$$1. \quad \frac{\partial^2 w}{\partial t^2} + \frac{k-1}{t} \frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2}.$$

*Darboux equation.* Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$w = f(x) \quad \text{at} \quad t = 0,$$

$$\partial_t w = 0 \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, t) = \frac{\Gamma\left(\frac{k}{2}\right)}{\sqrt{\pi} \Gamma\left(\frac{k}{2} - \frac{1}{2}\right)} \int_{-1}^1 f(x + t\xi)(1 - \xi^2)^{\frac{k-3}{2}} d\xi \quad (k > 1).$$

⊙ *Reference:* R. Courant and D. Hilbert (1989).

$$2. \quad \frac{\partial^2 w}{\partial t^2} + \frac{2a}{t} \frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} - b^2 w.$$

Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$w = f(x) \quad \text{at} \quad t = 0,$$

$$t^{2a} \partial_t w = g(x) \quad \text{at} \quad t = 0.$$

Solution for  $0 < 2a < 1$ :

$$w(x, t) = \frac{\Gamma(2a)}{\Gamma^2(a)} \int_0^1 f(x + t(2\xi - 1)) \bar{J}_{a-1}(2bt\sqrt{\xi(1-\xi)}) \xi^{a-1}(1-\xi)^{a-1} d\xi \\ + \frac{\Gamma(2-2a)}{(1-2a)\Gamma^2(1-a)} t^{1-2a} \int_0^1 g(x + t(2\xi - 1)) \bar{J}_{-a}(2bt\sqrt{\xi(1-\xi)}) \xi^{-a}(1-\xi)^{-a} d\xi,$$

where

$$\bar{J}_\nu(z) = 2^\nu \Gamma(1+\nu) z^{-\nu} J_\nu(z), \quad \Gamma(\nu) = \int_0^\infty e^{-s} s^{\nu-1} ds.$$

⊙ *Reference:* M. M. Smirnov (1975).

$$3. \quad \frac{\partial^2 w}{\partial t^2} + \frac{2a}{t} \frac{\partial w}{\partial t} = t^m \frac{\partial^2 w}{\partial x^2}.$$

Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) \quad \text{at} \quad t = 0, \\ t^{2a} \partial_t w &= g(x) \quad \text{at} \quad t = 0. \end{aligned}$$

Solution for  $0 \leq 2a < 1$  and  $m > 0$ :

$$\begin{aligned} w(x, t) &= \frac{\Gamma(2\beta)}{\Gamma^2(\beta)} \int_0^1 f\left(x + \frac{2}{2+m} t^{\frac{2+m}{2}} (2\xi - 1)\right) \xi^{\beta-1} (1-\xi)^{\beta-1} d\xi \\ &+ \frac{\Gamma(2-2\beta)}{(1-2a)\Gamma^2(1-\beta)} t^{1-2a} \int_0^1 g\left(x + \frac{2}{2+m} t^{\frac{2+m}{2}} (2\xi - 1)\right) \xi^{-\beta} (1-\xi)^{-\beta} d\xi, \end{aligned}$$

where

$$\beta = \frac{m+4a}{2(m+2)}, \quad \Gamma(z) = \int_0^\infty e^{-s} s^{z-1} ds.$$

⊙ *Reference:* M. M. Smirnov (1975).

$$4. \quad t^2 \frac{\partial^2 w}{\partial t^2} + kt \frac{\partial w}{\partial t} = a^2 \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw.$$

The substitution  $t = Ae^\tau$  ( $A \neq 0$ ) leads to a constant coefficient equation of the form 4.4.1.3:

$$\frac{\partial^2 w}{\partial \tau^2} + (k-1) \frac{\partial w}{\partial \tau} = a^2 \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw.$$

$$5. \quad t^2 \frac{\partial^2 w}{\partial t^2} + kt \frac{\partial w}{\partial t} = a^2 x^2 \frac{\partial^2 w}{\partial x^2} + bx \frac{\partial w}{\partial x} + cw.$$

The transformation

$$t = Ae^\tau, \quad x = Be^\xi \quad (A \neq 0, B \neq 0)$$

leads to a constant coefficient equation of the form 4.4.1.3:

$$\frac{\partial^2 w}{\partial \tau^2} + (k-1) \frac{\partial w}{\partial \tau} = a^2 \frac{\partial^2 w}{\partial \xi^2} + (b-a^2) \frac{\partial w}{\partial \xi} + cw.$$

$$6. \quad t^m \frac{\partial^2 w}{\partial t^2} + at^{m-1} \frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2}, \quad 0 < m < 2.$$

Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x) \quad \text{at} \quad t = 0, \\ t^a \partial_t w &= g(x) \quad \text{at} \quad t = 0. \end{aligned}$$

1°. Solution for  $\frac{1}{2}m < a < 1$ :

$$\begin{aligned} w(x, t) &= \frac{\Gamma(2\beta)}{\Gamma^2(\beta)} \int_0^1 f\left(x + \frac{2}{2-m} t^{\frac{2-m}{2}} (2\xi - 1)\right) \xi^{\beta-1} (1-\xi)^{\beta-1} d\xi \\ &+ \frac{\Gamma(2-2\beta)}{(1-a)\Gamma^2(1-\beta)} t^{1-a} \int_0^1 g\left(x + \frac{2}{2-m} t^{\frac{2-m}{2}} (2\xi - 1)\right) \xi^{-\beta} (1-\xi)^{-\beta} d\xi, \end{aligned}$$

where

$$\beta = \frac{2a-m}{2(2-m)}, \quad \Gamma(z) = \int_0^\infty e^{-s} s^{z-1} ds.$$

2°. Solution for  $a = \frac{1}{2}m$ :

$$w(x, t) = \frac{f(y) + f(z)}{2} + \frac{1}{2} \int_z^y g(\xi) d\xi,$$

$$y = x - \frac{2}{2-m} t^{\frac{2-m}{2}}, \quad z = x + \frac{2}{2-m} t^{\frac{2-m}{2}}.$$

⊙ *Reference:* M. M. Smirnov (1975).

$$7. \quad (t^m + k) \frac{\partial^2 w}{\partial t^2} + \frac{1}{2} m t^{m-1} \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + c w.$$

The substitution  $\tau = \int \frac{dt}{\sqrt{t^m + k}}$  leads to the equation  $\partial_{\tau\tau} w = a \partial_{xx} w + b \partial_x w + c w$ , which is discussed in Subsection 4.1.5.

## 4.5. Equations Containing Arbitrary Functions

### 4.5.1. Equations of the Form $s(x) \frac{\partial^2 w}{\partial t^2} = \frac{\partial}{\partial x} \left[ p(x) \frac{\partial w}{\partial x} \right] - q(x)w + \Phi(x, t)$

It is assumed that the functions  $s$ ,  $p$ ,  $p'_x$ , and  $q$  are continuous and the inequalities  $s > 0$ ,  $p > 0$  hold for  $x_1 \leq x \leq x_2$ .

4.5.1-1. General relations to solve linear nonhomogeneous boundary value problems.

The solution of the equation in question under the general initial conditions

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0, \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \end{aligned} \tag{1}$$

and the arbitrary linear nonhomogeneous boundary conditions

$$\begin{aligned} a_1 \partial_x w + b_1 w &= g_1(t) \quad \text{at} \quad x = x_1, \\ a_2 \partial_x w + b_2 w &= g_2(t) \quad \text{at} \quad x = x_2 \end{aligned} \tag{2}$$

can be represented as the sum

$$\begin{aligned} w(x, t) &= \int_0^t \int_{x_1}^{x_2} \Phi(\xi, \tau) \mathcal{G}(x, \xi, t - \tau) d\xi d\tau \\ &+ \frac{\partial}{\partial t} \int_{x_1}^{x_2} s(\xi) f_0(\xi) \mathcal{G}(x, \xi, t) d\xi + \int_{x_1}^{x_2} s(\xi) f_1(\xi) \mathcal{G}(x, \xi, t) d\xi \\ &+ p(x_1) \int_0^t g_1(\tau) \Lambda_1(x, t - \tau) d\tau + p(x_2) \int_0^t g_2(\tau) \Lambda_2(x, t - \tau) d\tau. \end{aligned} \tag{3}$$

Here, the modified Green's function is determined by

$$\mathcal{G}(x, \xi, t) = \sum_{n=1}^{\infty} \frac{y_n(x) y_n(\xi) \sin(t \sqrt{\lambda_n})}{\|y_n\|^2 \sqrt{\lambda_n}}, \quad \|y_n\|^2 = \int_{x_1}^{x_2} s(x) y_n^2(x) dx, \tag{4}$$

where the  $\lambda_n$  and  $y_n(x)$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the second-order linear ordinary differential equation

$$\begin{aligned} [p(x)y'_x]'_x + [\lambda s(x) - q(x)]y &= 0, \\ a_1 y'_x + b_1 y &= 0 \quad \text{at } x = x_1, \\ a_2 y'_x + b_2 y &= 0 \quad \text{at } x = x_2. \end{aligned} \quad (5)$$

The functions  $\Lambda_1(x, t)$  and  $\Lambda_2(x, t)$  that occur in the integrands of the last two terms in solution (3) are expressed in terms of the Green's function of (4). The corresponding formulas will be specified below in studying specific boundary value problems.

General properties of the Sturm–Liouville problem (5):

1°. There are finitely many eigenvalues  $\lambda_1 < \lambda_2 < \lambda_3 < \dots$ , with  $\lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ ; hence the number of negative eigenvalues is finite.

2°. Any two eigenfunctions  $y_n(x)$  and  $y_m(x)$  for  $n \neq m$  are orthogonal to each other with weight  $s(x)$  on the interval  $x_1 \leq x \leq x_2$ ; specifically,

$$\int_{x_1}^{x_2} s(x) y_n(x) y_m(x) dx = 0 \quad \text{at } n \neq m.$$

3°. If the conditions

$$q(x) \geq 0, \quad a_1 b_1 \leq 0, \quad a_2 b_2 \geq 0 \quad (6)$$

are satisfied, then there are no negative eigenvalues. If  $q \equiv 0$  and  $b_1 = b_2 = 0$ , the least eigenvalue is  $\lambda_1 = 0$  and the corresponding eigenfunction is  $\varphi_1 = \text{const}$ . In the other cases where conditions (6) are satisfied, all eigenvalues are positive.

*Remark.* More detailed information about the properties of the Sturm–Liouville problem (5) can be found in Subsection 1.8.9. Asymptotic and approximate formulas for eigenvalues and eigenfunctions are also presented there.

#### 4.5.1-2. First boundary value problem (case $a_1 = a_2 = 0$ , $b_1 = b_2 = 1$ ).

The solution of the first boundary value problem for the equation in question with the initial conditions (1) and the boundary conditions

$$\begin{aligned} w &= g_1(t) \quad \text{at } x = x_1, \\ w &= g_2(t) \quad \text{at } x = x_2 \end{aligned}$$

is given by relations (3) and (4) in which

$$\Lambda_1(x, t) = \frac{\partial}{\partial \xi} \mathcal{G}(x, \xi, t) \Big|_{\xi=x_1}, \quad \Lambda_2(x, t) = -\frac{\partial}{\partial \xi} \mathcal{G}(x, \xi, t) \Big|_{\xi=x_2}.$$

#### 4.5.1-3. Second boundary value problem (case $a_1 = a_2 = 1$ , $b_1 = b_2 = 0$ ).

The solution of the second boundary value problem for the equation in question with the initial conditions (1) and the boundary conditions

$$\begin{aligned} \partial_x w &= g_1(t) \quad \text{at } x = x_1, \\ \partial_x w &= g_2(t) \quad \text{at } x = x_2 \end{aligned}$$

is given by relations (3) and (4) with

$$\Lambda_1(x, t) = -\mathcal{G}(x, x_1, t), \quad \Lambda_2(x, t) = \mathcal{G}(x, x_2, t).$$

#### 4.5.1-4. Third boundary value problem (case $a_1 = a_2 = 1$ , $b_1 \neq 0$ , $b_2 \neq 0$ ).

The solution of the third boundary value problem for the equation in question with the initial conditions (1) and the boundary conditions (2) with  $a_1 = a_2 = 1$  is given by relations (3) and (4) in which

$$\Lambda_1(x, t) = -\mathcal{G}(x, x_1, t), \quad \Lambda_2(x, t) = \mathcal{G}(x, x_2, t).$$

#### 4.5.1-5. Mixed boundary value problem (case $a_1 = b_2 = 0$ , $a_2 = b_1 = 1$ ).

The solution of the mixed boundary value problem for the equation in question with the initial conditions (1) and the boundary conditions

$$\begin{aligned} w &= g_1(t) \quad \text{at} \quad x = x_1, \\ \partial_x w &= g_2(t) \quad \text{at} \quad x = x_2 \end{aligned}$$

is given by relations (3) and (4) with

$$\Lambda_1(x, t) = \frac{\partial}{\partial \xi} \mathcal{G}(x, \xi, t) \Big|_{\xi=x_1}, \quad \Lambda_2(x, t) = \mathcal{G}(x, x_2, t).$$

#### 4.5.1-6. Mixed boundary value problem (case $a_1 = b_2 = 1$ , $a_2 = b_1 = 0$ ).

The solution of the mixed boundary value problem with the initial conditions (1) and the boundary conditions

$$\begin{aligned} \partial_x w &= g_1(t) \quad \text{at} \quad x = x_1, \\ w &= g_2(t) \quad \text{at} \quad x = x_2 \end{aligned}$$

is given by relations (3) and (4) with

$$\Lambda_1(x, t) = -\mathcal{G}(x, x_1, t), \quad \Lambda_2(x, t) = -\frac{\partial}{\partial \xi} \mathcal{G}(x, \xi, t) \Big|_{\xi=x_2}.$$

☉ *References for Subsection 4.5.1:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin et al. (1964), V. A. Marchenko (1986), V. S. Vladimirov (1988), A. D. Polyaniin (2000a).

## 4.5.2. Equations of the Form

$$\frac{\partial^2 w}{\partial t^2} + a(t) \frac{\partial w}{\partial t} = b(t) \left\{ \frac{\partial}{\partial x} \left[ p(x) \frac{\partial w}{\partial x} \right] - q(x) w \right\} + \Phi(x, t)$$

It is assumed that the functions  $p$ ,  $p'_x$ , and  $q$  are continuous and  $p > 0$  for  $x_1 \leq x \leq x_2$ .

#### 4.5.2-1. General relations to solve linear nonhomogeneous boundary value problems.

The solution of the equation in question under the general initial conditions

$$\begin{aligned} w &= f_0(x) \quad \text{at} \quad t = 0, \\ \partial_t w &= f_1(x) \quad \text{at} \quad t = 0 \end{aligned} \tag{1}$$

and the arbitrary linear nonhomogeneous boundary conditions

$$\begin{aligned} s_1 \partial_x w + k_1 w &= g_1(t) \quad \text{at} \quad x = x_1, \\ s_2 \partial_x w + k_2 w &= g_2(t) \quad \text{at} \quad x = x_2 \end{aligned} \tag{2}$$



can be represented as the sum

$$\begin{aligned}
 w(x, t) = & \int_0^t \int_{x_1}^{x_2} \Phi(\xi, \tau) G(x, \xi, t, \tau) d\xi d\tau \\
 & - \int_{x_1}^{x_2} f_0(\xi) \left[ \frac{\partial}{\partial \tau} G(x, \xi, t, \tau) \right]_{\tau=0} d\xi + \int_{x_1}^{x_2} [f_1(\xi) + a(0)f_0(\xi)] G(x, \xi, t, 0) d\xi \\
 & + p(x_1) \int_0^t g_1(\tau) b(\tau) \Lambda_1(x, t, \tau) d\tau + p(x_2) \int_0^t g_2(\tau) b(\tau) \Lambda_2(x, t, \tau) d\tau.
 \end{aligned} \quad (3)$$

Here, the modified Green's function is determined by

$$G(x, \xi, t, \tau) = \sum_{n=1}^{\infty} \frac{y_n(x) y_n(\xi)}{\|y_n\|^2} U_n(t, \tau), \quad \|y_n\|^2 = \int_{x_1}^{x_2} y_n^2(x) dx, \quad (4)$$

where the  $\lambda_n$  and  $y_n(x)$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the following second-order linear ordinary differential equation with homogeneous boundary conditions:

$$\begin{aligned}
 [p(x)y'_x]' + [\lambda - q(x)]y &= 0, \\
 s_1 y'_x + k_1 y &= 0 \quad \text{at } x = x_1, \\
 s_2 y'_x + k_2 y &= 0 \quad \text{at } x = x_2.
 \end{aligned} \quad (5)$$

The functions  $U_n = U_n(t, \tau)$  are determined by solving the Cauchy problem for the linear ordinary differential equation

$$\begin{aligned}
 U_n'' + a(t)U_n' + \lambda_n b(t)U_n &= 0, \\
 U_n|_{t=\tau} &= 0, \quad U_n'|_{t=\tau} = 1.
 \end{aligned} \quad (6)$$

The prime denotes the derivative with respect to  $t$ , and  $\tau$  is a free parameter occurring in the initial conditions.

The functions  $\Lambda_1(x, t)$  and  $\Lambda_2(x, t)$  that occur in the integrands of the last two terms in solution (3) are expressed in terms of the Green's function of (4). The corresponding formulas will be specified below when studying specific boundary value problems.

The properties of the Sturm–Liouville problem (5) are detailed in Subsection 1.8.9. Asymptotic and approximate formulas for eigenvalues and eigenfunctions are also presented there.

#### 4.5.2-2. First, second, third, and mixed boundary value problems.

1°. *First boundary value problem.* The solution of the equation in question with the initial conditions (1) and boundary conditions (2) for  $s_1 = s_2 = 0$  and  $k_1 = k_2 = 1$  is given by relations (3) and (4), where

$$\Lambda_1(x, t, \tau) = \frac{\partial}{\partial \xi} G(x, \xi, t, \tau) \Big|_{\xi=x_1}, \quad \Lambda_2(x, t, \tau) = -\frac{\partial}{\partial \xi} G(x, \xi, t, \tau) \Big|_{\xi=x_2}.$$

2°. *Second boundary value problem.* The solution of the equation with the initial conditions (1) and boundary conditions (2) for  $s_1 = s_2 = 1$  and  $k_1 = k_2 = 0$  is given by relations (3) and (4) with

$$\Lambda_1(x, t, \tau) = -G(x, x_1, t, \tau), \quad \Lambda_2(x, t, \tau) = G(x, x_2, t, \tau).$$

3°. *Third boundary value problem.* The solution of the equation with the initial conditions (1) and boundary conditions (2) for  $s_1 = s_2 = 1$  and  $k_1 k_2 \neq 0$  is given by relations (3) and (4) in which

$$\Lambda_1(x, t, \tau) = -G(x, x_1, t, \tau), \quad \Lambda_2(x, t, \tau) = G(x, x_2, t, \tau).$$

4°. *Mixed boundary value problem.* The solution of the equation with the initial conditions (1) and boundary conditions (2) for  $s_1 = k_2 = 0$  and  $s_2 = k_1 = 1$  is given by relations (3) and (4) with

$$\Lambda_1(x, t, \tau) = \frac{\partial}{\partial \xi} G(x, \xi, t, \tau) \Big|_{\xi=x_1}, \quad \Lambda_2(x, t, \tau) = G(x, x_2, t, \tau).$$

5°. *Mixed boundary value problem.* The solution of the equation with the initial conditions (1) and boundary conditions (2) for  $s_1 = k_2 = 1$  and  $s_2 = k_1 = 0$  is given by relations (3) and (4) with

$$\Lambda_1(x, t, \tau) = -G(x, x_1, t, \tau), \quad \Lambda_2(x, t, \tau) = -\frac{\partial}{\partial \xi} G(x, \xi, t, \tau) \Big|_{\xi=x_2}.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin et al. (1964), A. V. Bitsadze and D. F. Kalinichenko (1985), A. D. Polyaniin (2000a).

### 4.5.3. Other Equations

1. 
$$\frac{\partial^2 w}{\partial t^2} = f(x) \frac{\partial^2 w}{\partial x^2}.$$

This is a special case of the equation of Subsection 4.5.1 with  $s(x) = 1/f(x)$ ,  $p(x) = 1$ , and  $q = \Phi = 0$ .

1°. Particular solutions:

$$w = C_1 xt + C_2 t + C_3 x + C_4,$$

$$w = C_1 t^2 + C_2 xt + C_3 t + C_4 x + 2C_1 \int_a^x \frac{x-\xi}{f(\xi)} d\xi + C_5,$$

$$w = C_1 t^3 + C_2 xt + C_3 t + C_4 x + 6C_1 t \int_a^x \frac{x-\xi}{f(\xi)} d\xi + C_5,$$

$$w = (C_1 x + C_2) t^2 + C_3 xt + C_4 t + C_5 x + 2 \int_a^x (x-\xi) \frac{(C_1 \xi + C_2)}{f(\xi)} d\xi + C_6,$$

where  $C_1, C_2, C_3, C_4, C_5$ , and  $C_6$  are arbitrary constants, and  $a$  is an arbitrary real number.

2°. Separable particular solution:

$$w = (C_1 e^{\lambda t} + C_2 e^{-\lambda t}) H(x),$$

where  $C_1, C_2$ , and  $\lambda$  are arbitrary constants, and the function  $H = H(x)$  is determined by the ordinary differential equation  $f(x) H''_{xx} - \lambda^2 H = 0$ .

3°. Separable particular solution:

$$w = [C_1 \sin(\lambda t) + C_2 \cos(\lambda t)] Z(x),$$

where  $C_1, C_2$ , and  $\lambda$  are arbitrary constants, and the function  $Z = Z(x)$  is determined by the ordinary differential equation  $f(x) Z''_{xx} + \lambda^2 Z = 0$ .

4°. Particular solutions with even powers of  $t$ :

$$w = \sum_{k=0}^n \varphi_k(x) t^{2k},$$

where the functions  $\varphi_k = \varphi_k(x)$  are defined by the recurrence relations

$$\varphi_n(x) = A_n x + B_n,$$

$$\varphi_{k-1}(x) = A_k x + B_k + 2k(2k-1) \int_a^x (x-\xi) \frac{\varphi_k(\xi)}{f(\xi)} d\xi,$$

where  $A_k, B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

5°. Particular solutions with odd powers of  $t$ :

$$w = \sum_{k=0}^n \psi_k(x) t^{2k+1},$$

where the functions  $\psi_k = \psi_k(x)$  are defined by the recurrence relations

$$\psi_n(x) = A_n x + B_n,$$

$$\psi_{k-1}(x) = A_k x + B_k + 2k(2k+1) \int_a^x (x-\xi) \frac{\psi_k(\xi)}{f(\xi)} d\xi,$$

where  $A_k, B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

$$2. \quad \frac{\partial^2 w}{\partial t^2} = \frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right].$$

This is a special case of the equation of Subsection 4.5.1 with  $s(x) = 1$ ,  $p(x) = f(x)$ , and  $q = \Phi = 0$ .

1°. Particular solutions:

$$w = C_1 t^2 + C_2 t + 2 \int \frac{C_1 x + C_3}{f(x)} dx + C_4,$$

$$w = C_1 t^3 + C_2 t + 6t \int \frac{C_1 x + C_3}{f(x)} dx + C_4,$$

$$w = [C_1 \Phi(x) + C_2]t + C_3 \Phi(x) + C_4, \quad \Phi(x) = \int \frac{dx}{f(x)},$$

$$w = [C_1 \Phi(x) + C_2]t^2 + C_3 \Phi(x) + C_4 + 2 \int \left\{ \frac{1}{f(x)} \int [C_1 \Phi(x) + C_2] dx \right\} dx,$$

where  $C_1$ ,  $C_2$ ,  $C_3$ ,  $C_4$ , and  $C_5$  are arbitrary constants.

2°. Separable particular solution:

$$w = (C_1 e^{\lambda t} + C_2 e^{-\lambda t})H(x),$$

where  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants, and the function  $H = H(x)$  is determined by the ordinary differential equation  $[f(x)H'_x]'_x - \lambda^2 H = 0$ .

3°. Separable particular solution:

$$w = [C_1 \sin(\lambda t) + C_2 \cos(\lambda t)]Z(x),$$

where  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants, and the function  $Z = Z(x)$  is determined by the ordinary differential equation  $[f(x)Z'_x]'_x + \lambda^2 Z = 0$ .

4°. Particular solutions with even powers of  $t$ :

$$w = \sum_{k=0}^n \zeta_k(x) t^{2k},$$

where the functions  $\zeta_k = \zeta_k(x)$  are defined by the recurrence relations

$$\zeta_n(x) = A_n \Phi(x) + B_n, \quad \Phi(x) = \int \frac{dx}{f(x)},$$

$$\zeta_{k-1}(x) = A_k \Phi(x) + B_k + 2k(2k-1) \int \frac{1}{f(x)} \left\{ \int \zeta_k(x) dx \right\} dx,$$

where  $A_k$ ,  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

5°. Particular solutions with odd powers of  $t$ :

$$w = \sum_{k=0}^n \eta_k(x) t^{2k+1},$$

where the functions  $\eta_k = \eta_k(x)$  are defined by the recurrence relations

$$\eta_n(x) = A_n \Phi(x) + B_n, \quad \Phi(x) = \int \frac{dx}{f(x)},$$

$$\eta_{k-1}(x) = A_k \Phi(x) + B_k + 2k(2k+1) \int \frac{1}{f(x)} \left\{ \int \eta_k(x) dx \right\} dx,$$

where  $A_k$ ,  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

$$3. \quad \frac{\partial^2 w}{\partial t^2} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x) \frac{\partial w}{\partial x} + \Phi(x, t), \quad 0 < f(x) < \infty.$$

This equation can be rewritten in the form of the equation from Subsection 4.5.1 with  $q(x) \equiv 0$ :

$$s(x) \frac{\partial^2 w}{\partial t^2} = \frac{\partial}{\partial x} \left[ p(x) \frac{\partial w}{\partial x} \right] + s(x) \Phi(x, t),$$

where

$$s(x) = \frac{1}{f(x)} \exp \left[ \int \frac{g(x)}{f(x)} dx \right], \quad p(x) = \exp \left[ \int \frac{g(x)}{f(x)} dx \right].$$

$$4. \quad \frac{\partial^2 w}{\partial t^2} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x) \frac{\partial w}{\partial x} + h(x)w + \Phi(x, t).$$

This equation can be rewritten in the form of the equation from Subsection 4.5.1:

$$s(x) \frac{\partial^2 w}{\partial t^2} = \frac{\partial}{\partial x} \left[ p(x) \frac{\partial w}{\partial x} \right] - q(x)w + s(x) \Phi(x, t),$$

where

$$s(x) = \frac{1}{f(x)} \exp \left[ \int \frac{g(x)}{f(x)} dx \right], \quad p(x) = \exp \left[ \int \frac{g(x)}{f(x)} dx \right], \quad q(x) = -\frac{h(x)}{f(x)} \exp \left[ \int \frac{g(x)}{f(x)} dx \right].$$

$$5. \quad \frac{\partial^2 w}{\partial t^2} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x) \frac{\partial w}{\partial x} + [h_1(x) + h_2(t)]w.$$

1°. There are separable solutions in the product form  $w(x, t) = \varphi(x)\psi(t)$ , where the functions  $\varphi = \varphi(x)$  and  $\psi = \psi(t)$  satisfy the ordinary differential equations ( $\lambda$  is an arbitrary constant):

$$f(x)\varphi''_{xx} + g(x)\varphi'_x + [\lambda + h_1(x)]\varphi = 0, \quad \psi''_{tt} + [\lambda - h_2(t)]\psi = 0.$$

2°. For the solution of various boundary value problems for the original equation, see Subsections 0.4.1 and 0.4.2.

$$6. \quad \frac{\partial^2 w}{\partial t^2} = f(x) \frac{\partial^2 w}{\partial x^2} + \frac{1}{2}f'(x) \frac{\partial w}{\partial x} + bw.$$

The substitution  $z = \int \frac{dx}{\sqrt{f(x)}}$  leads to the constant coefficient equation  $\partial_{tt}w = \partial_{zz}w + bw$  that is discussed in Subsection 4.1.3.

$$7. \quad \frac{\partial^2 w}{\partial t^2} = f^2 \frac{\partial^2 w}{\partial x^2} + f(f'_x + 2g) \frac{\partial w}{\partial x} + (fg'_x + g^2)w, \quad f = f(x), \quad g = g(x).$$

The transformation

$$w(x, t) = u(\xi, t) \exp \left( - \int \frac{g}{f} dx \right), \quad \xi = \int \frac{dx}{f(x)}$$

leads to the wave equation  $\partial_{tt}u = \partial_{\xi\xi}u$  that is discussed in Subsection 4.1.1.

$$8. \quad \frac{\partial^2 w}{\partial t^2} + a \frac{\partial w}{\partial t} = f(x) \frac{\partial^2 w}{\partial x^2} + \frac{1}{2}f'(x) \frac{\partial w}{\partial x} + bw.$$

The substitution  $z = \int \frac{dx}{\sqrt{f(x)}}$  leads to a constant coefficient equation of the form 4.4.1.2:  $\partial_{tt}w + a\partial_t w = \partial_{zz}w + bw$ .

$$9. \quad f(t) \frac{\partial^2 w}{\partial t^2} + \frac{1}{2} f'(t) \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \frac{\partial w}{\partial x} + cw.$$

The substitution  $\tau = \int \frac{dt}{\sqrt{f(t)}}$  leads to the equation  $\partial_{\tau\tau} w = a \partial_{xx} w + b \partial_x w + cw$  that is discussed in Subsection 4.1.5.

$$10. \quad f(t) \frac{\partial^2 w}{\partial t^2} + \frac{1}{2} f'(t) \frac{\partial w}{\partial t} = g(x) \frac{\partial^2 w}{\partial x^2} + \frac{1}{2} g'(x) \frac{\partial w}{\partial x} + cw.$$

The transformation  $\tau = \int \frac{dt}{\sqrt{f(t)}}$ ,  $z = \int \frac{dx}{\sqrt{g(x)}}$  leads to the constant coefficient equation  $\partial_{\tau\tau} w = \partial_{zz} w + cw$  that is discussed in Subsection 4.1.3.

## Chapter 5

# Hyperbolic Equations with Two Space Variables

---

### 5.1. Wave Equation $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_2 w$

#### 5.1.1. Problems in Cartesian Coordinates

The wave equation with two space variables in the rectangular Cartesian system of coordinates has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right).$$

5.1.1-1. Particular solutions and some relations.

1°. Particular solutions:

$$\begin{aligned} w(x, y, t) &= A \exp \left( k_1 x + k_2 y \pm at \sqrt{k_1^2 + k_2^2} \right), \\ w(x, y, t) &= A \sin(k_1 x + C_1) \sin(k_2 y + C_2) \sin \left( at \sqrt{k_1^2 + k_2^2} \right), \\ w(x, y, t) &= A \sin(k_1 x + C_1) \sin(k_2 y + C_2) \cos \left( at \sqrt{k_1^2 + k_2^2} \right), \\ w(x, y, t) &= A \sinh(k_1 x + C_1) \sinh(k_2 y + C_2) \sinh \left( at \sqrt{k_1^2 + k_2^2} \right), \\ w(x, y, t) &= A \sinh(k_1 x + C_1) \sinh(k_2 y + C_2) \cosh \left( at \sqrt{k_1^2 + k_2^2} \right), \\ w(x, y, t) &= \varphi(x \sin \beta + y \cos \beta + at) + \psi(x \sin \beta + y \cos \beta - at), \end{aligned}$$

where  $A$ ,  $C_1$ ,  $C_2$ ,  $k_1$ ,  $k_2$ , and  $\beta$  are arbitrary constants, and  $\varphi(z)$  and  $\psi(z)$  are arbitrary functions.

2°. Particular solutions that are expressed in terms of solutions to simpler equations:

$$w(x, y, t) = [A \cos(ky) + B \sin(ky)] u(x, t), \quad \text{where} \quad \partial_{tt} u = a^2 \partial_{xx} u - a^2 k^2 u, \quad (1)$$

$$w(x, y, t) = [A \cosh(ky) + B \sinh(ky)] u(x, t), \quad \text{where} \quad \partial_{tt} u = a^2 \partial_{xx} u + a^2 k^2 u, \quad (2)$$

$$w(x, y, t) = [A \cos(kt) + B \sin(kt)] u(x, y), \quad \text{where} \quad \partial_{xx} u + \partial_{yy} u = -(k/a)^2 u, \quad (3)$$

$$w(x, y, t) = [A \cosh(kt) + B \sinh(kt)] u(x, y), \quad \text{where} \quad \partial_{xx} u + \partial_{yy} u = (k/a)^2 u, \quad (4)$$

$$w(x, y, t) = \exp \left( \frac{at \pm y}{2b} \right) u(x, \tau), \quad \tau = \frac{at \mp y}{2}, \quad \text{where} \quad \partial_\tau u = b \partial_{xx} u. \quad (5)$$

For particular solutions of equations (1) and (2) for the function  $u(x, t)$ , see the Klein–Gordon equation 4.1.3. For particular solutions of equations (3) and (4) for the function  $u(x, y)$ , see Subsection 7.3.2. For particular solutions of the heat equation (5) for the function  $u(x, \tau)$ , see Subsection 1.1.1.

3°. Fundamental solution:

$$\mathcal{E}(x, y, t) = \frac{\vartheta(at - r)}{2\pi a \sqrt{a^2 t^2 - r^2}}, \quad \vartheta(z) = \begin{cases} 1 & \text{for } z \geq 0, \\ 0 & \text{for } z < 0, \end{cases}$$

where  $r = \sqrt{x^2 + y^2}$ .

4°. Infinite series solutions that contain arbitrary functions of the space variables:

$$w(x, y, t) = f(x, y) + \sum_{n=1}^{\infty} \frac{(at)^{2n}}{(2n)!} \Delta^n f(x, y), \quad \Delta \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2},$$

$$w(x, y, t) = tg(x, y) + t \sum_{n=1}^{\infty} \frac{(at)^{2n}}{(2n+1)!} \Delta^n g(x, y),$$

where  $f(x, y)$  and  $g(x, y)$  are any infinitely differentiable functions. The first solution satisfies the initial conditions  $w(x, y, 0) = f(x, y)$ ,  $\partial_t w(x, y, 0) = 0$  and the second solution to the initial conditions  $w(x, y, 0) = 0$ ,  $\partial_t w(x, y, 0) = g(x, y)$ . The sums are finite if  $f(x, y)$  and  $g(x, y)$  are bivariate polynomials.

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

5°. A wide class of solutions to the wave equation with two space variables are described by the formulas

$$w(x, y, t) = \operatorname{Re} F(\theta) \quad \text{and} \quad w(x, y, t) = \operatorname{Im} F(\theta). \quad (6)$$

Here,  $F(\theta)$  is an arbitrary analytic function of the complex argument  $\theta$  related to the variables  $(x, y, t)$  by the implicit relation

$$at - (x - x_0)\theta + (y - y_0)\sqrt{1 - \theta^2} = G(\theta), \quad (7)$$

where  $G(\theta)$  is any analytic function and  $x_0, y_0$  are arbitrary constants. Solutions of the forms (6), (7) find wide application in the theory of diffraction. If the argument  $\theta$  obtained by solving (7) with a prescribed  $G(\theta)$  is real in some domain  $D$ , then one should set  $\operatorname{Re} F(\theta) = F(\theta)$  in relation (6) everywhere in  $D$ .

⊙ *Reference:* V. I. Smirnov (1974, Vol. 3, Pt. 2).

6°. Suppose  $w = w(x, y, t)$  is a solution of the wave equation. Then the functions

$$w_1 = Aw(\pm \lambda x + C_1, \pm \lambda y + C_2, \pm \lambda t + C_3),$$

$$w_2 = Aw\left(\frac{x - vt}{\sqrt{1 - (v/a)^2}}, y, \frac{t - va^{-2}x}{\sqrt{1 - (v/a)^2}}\right),$$

$$w_3 = \frac{A}{\sqrt{|r^2 - a^2 t^2|}} w\left(\frac{x}{r^2 - a^2 t^2}, \frac{y}{r^2 - a^2 t^2}, \frac{t}{r^2 - a^2 t^2}\right),$$

$$w_4 = \frac{A}{\sqrt{\Xi}} w\left(\frac{x + k_1(a^2 t^2 - r^2)}{\Xi}, \frac{y + k_2(a^2 t^2 - r^2)}{\Xi}, \frac{at + k_3(a^2 t^2 - r^2)}{a\Xi}\right),$$

$$r^2 = x^2 + y^2, \quad \Xi = 1 - 2(k_1 x + k_2 y - ak_3 t) + (k_1^2 + k_2^2 - k_3^2)(r^2 - a^2 t^2),$$

where  $A, C_1, C_2, C_3, k_1, k_2, k_3, v$ , and  $\lambda$  are arbitrary constants, are also solutions of the equation. The signs at  $\lambda$  in the expression of  $w_1$  can be taken independently of one another. The function  $w_2$  results from the invariance of the wave equation under the Lorentz transformation.

More detailed information about particular solutions and transformations of the wave equation with two space variables can be found in the references cited below.

⊙ *References:* E. Kalnins and W. Miller, Jr. (1975, 1976), W. Miller, Jr. (1977).

**5.1.1-2. Domain:  $-\infty < x < \infty, -\infty < y < \infty$ . Cauchy problem.**

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(x, y) \quad \text{at} \quad t = 0. \end{aligned}$$

Solution (Poisson's formula):

$$w(x, y, t) = \frac{1}{2\pi a} \frac{\partial}{\partial t} \iint_{C_{at}} \frac{f(\xi, \eta) d\xi d\eta}{\sqrt{a^2 t^2 - (\xi - x)^2 - (\eta - y)^2}} + \frac{1}{2\pi a} \iint_{C_{at}} \frac{g(\xi, \eta) d\xi d\eta}{\sqrt{a^2 t^2 - (\xi - x)^2 - (\eta - y)^2}},$$

where the integration is performed over the interior of the circle of radius  $at$  with center at  $(x, y)$ .

⊙ *References:* N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970), A. N. Tikhonov and A. A. Samarskii (1990).

**5.1.1-3. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . First boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(y, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ w &= g_3(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ w &= g_4(x, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &+ a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &- a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=l_2} d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{4}{al_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\lambda_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \sin(a \lambda_{nm} t), \\ p_n &= \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{p_n^2 + q_m^2}. \end{aligned}$$

The problem of vibration of a rectangular membrane with sides  $l_1$  and  $l_2$  rigidly fixed in its contour is characterized by homogeneous boundary conditions,  $g_s \equiv 0$  ( $s = 1, 2, 3, 4$ ).

⊙ *References:* M. M. Smirnov (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).



**5.1.1-4. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Second boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &\quad - a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\ &\quad + a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\ &\quad - a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{t}{l_1 l_2} + \frac{2}{a l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_{nm}}{\lambda_{nm}} \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) \sin(a \lambda_{nm} t), \\ p_n &= \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{p_n^2 + q_m^2}, \quad A_{nm} = \begin{cases} 0 & \text{for } n = m = 0, \\ 1 & \text{for } nm = 0 \ (n \neq m), \\ 2 & \text{for } nm \neq 0, \end{cases} \end{aligned}$$

⊙ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.1.1-5. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Third boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(y, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ \partial_y w - k_3 w &= g_3(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_y w + k_4 w &= g_4(x, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 5.1.1-4 where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{4}{a} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{E_{nm} \sqrt{\mu_n^2 + \nu_m^2}} \sin(\mu_n x + \varepsilon_n) \sin(\nu_m y + \sigma_m) \\ &\quad \times \sin(\mu_n \xi + \varepsilon_n) \sin(\nu_m \eta + \sigma_m) \sin(at \sqrt{\mu_n^2 + \nu_m^2}), \\ \varepsilon_n &= \arctan \frac{\mu_n}{l_1}, \quad \sigma_m = \arctan \frac{\nu_m}{l_2}, \quad E_{nm} = \left[ l_1 + \frac{(k_1 k_2 + \mu_n^2)(k_1 + k_2)}{(k_1^2 + \mu_n^2)(k_2^2 + \mu_n^2)} \right] \left[ l_2 + \frac{(k_3 k_4 + \nu_m^2)(k_3 + k_4)}{(k_3^2 + \nu_m^2)(k_4^2 + \nu_m^2)} \right], \end{aligned}$$

where the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\mu^2 - k_1 k_2 = (k_1 + k_2)\mu \cot(l_1 \mu), \quad \nu^2 - k_3 k_4 = (k_3 + k_4)\nu \cot(l_2 \nu).$$

⊙ *References:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.1.1-6. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Mixed boundary value problems.**

1°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(y, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ \partial_y w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ \partial_y w &= g_4(x, t) & \text{at } y &= l_2 & \text{(boundary condition).} \end{aligned}$$

**Solution:**

$$\begin{aligned} w(x, y, t) &= \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &+ a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\ &- a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\ &- a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\ &+ a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau, \end{aligned}$$

where

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{2}{al_1 l_2} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{\lambda_{nm}} \sin(p_n x) \cos(q_m y) \sin(p_n \xi) \cos(q_m \eta) \sin(a \lambda_{nm} t), \\ p_n &= \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{p_n^2 + q_m^2}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases} \end{aligned}$$

2°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w &= g_2(y, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ \partial_y w &= g_4(x, t) & \text{at } y &= l_2 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & \frac{4}{al_1l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{1}{\lambda_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \sin(a\lambda_{nm} t), \\
 p_n = & \frac{\pi(2n+1)}{2l_1}, \quad q_m = \frac{\pi(2m+1)}{2l_2}, \quad \lambda_{nm} = \sqrt{p_n^2 + q_m^2}.
 \end{aligned}$$

### 5.1.2. Problems in Polar Coordinates

The wave equation with two space variables in the polar coordinate system has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} \right), \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional solutions  $w = w(r, t)$  that are independent of the angular coordinate  $\varphi$  are considered in Subsection 4.2.1.

5.1.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A circle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 w = g(\varphi, t) \quad \text{at} \quad r = R \quad & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & - a^2 R \int_0^t \int_0^{2\pi} g(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \frac{1}{\pi a R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n}{\mu_{nm} [J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \sin(\mu_{nm} a t), \\
 A_0 = & 1, \quad A_n = 2 \quad (n = 1, 2, \dots),
 \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

The problem of vibration of a circular membrane of radius  $R$  rigidly fixed in its contour is characterized by the homogeneous boundary condition,  $g(\varphi, t) \equiv 0$ .

⊙ *References:* N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970), A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.1.2-2. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad + a^2 R \int_0^t \int_0^{2\pi} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{t}{\pi R^2} + \frac{1}{\pi a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \sin(\mu_{nm} at), \\ A_0 &= 1, \quad A_n = 2 \quad (n = 1, 2, \dots), \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

⊙ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.1.2-3. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi$ . Third boundary value problem.**

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is determined by the formula in Paragraph 5.1.2-2 where

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 + k^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \sin(\mu_{nm} at), \\ A_0 &= 1, \quad A_n = 2 \quad (n = 1, 2, \dots). \end{aligned}$$

Here, the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k J_n(\mu R) = 0.$$

**5.1.2-4. Domain:  $R_1 \leq r \leq R_2, 0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + a^2 R_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\
 & - a^2 R_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \frac{\pi}{2a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \sin(\mu_{nm} at), \\
 A_n = & \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm} J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)}, \\
 Z_n(\mu_{nm} r) = & J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),
 \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

5.1.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, \varphi) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_r w = g_1(\varphi, t) & \quad \text{at } r = R_1 \quad (\text{boundary condition}), \\
 \partial_r w = g_2(\varphi, t) & \quad \text{at } r = R_2 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & - a^2 R_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) G(r, \varphi, R_1, \eta, t - \tau) d\eta d\tau \\
 & + a^2 R_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) G(r, \varphi, R_2, \eta, t - \tau) d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \frac{t}{\pi(R_2^2 - R_1^2)} + \frac{1}{\pi a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \sin(\mu_{nm} at)}{(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\
 Z_n(\mu_{nm} r) = & J_n'(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n'(\mu_{nm} R_1) J_n(\mu_{nm} r),
 \end{aligned}$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n'(\mu R_1) Y_n'(\mu R_2) - Y_n'(\mu R_1) J_n'(\mu R_2) = 0.$$

**5.1.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is determined by the formula in Paragraph 5.1.2-5 where

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \sin(\mu_{nm} at)}{(k_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)}, \\ Z_n(\mu_{nm} r) &= [\mu_{nm} J'_n(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) \\ &\quad - [\mu_{nm} Y'_n(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r). \end{aligned}$$

Here,  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned} [\mu J'_n(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y'_n(\mu R_2) + k_2 Y_n(\mu R_2)] \\ = [\mu Y'_n(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J'_n(\mu R_2) + k_2 J_n(\mu R_2)]. \end{aligned}$$

**5.1.2-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . First boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ w &= g_2(r, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\ w &= g_3(r, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad - a^2 R \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad + a^2 \int_0^t \int_0^R g_2(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad - a^2 \int_0^t \int_0^R g_3(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{4}{a R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \sin(\mu_{nm} at)}{\mu_{nm} [J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2},$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

**5.1.2-8. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq \varphi_0$ . Second boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_2(r, t) & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_3(r, t) & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) d\xi d\eta + \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) d\xi d\eta \\ &\quad + a^2 R \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\ &\quad - a^2 \int_0^t \int_0^R g_2(\xi, \tau) G(r, \varphi, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + a^2 \int_0^t \int_0^R g_3(\xi, \tau) G(r, \varphi, \xi, \varphi_0, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{2t}{R^2 \varphi_0} + \frac{4\varphi_0}{a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_{nm} J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{(R^2 \varphi_0^2 \mu_{nm}^2 - n^2 \pi^2) [J_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \\ &\quad \times \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \sin(\mu_{nm} at), \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_{n\pi/\varphi_0}(\mu R) = 0$ .

**5.1.2-9. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq \varphi_0$ . Mixed boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) d\xi d\eta + \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) d\xi d\eta \\ &\quad + a^2 R \int_0^t \int_0^{\varphi_0} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_{nm} J_{s_n}(\mu_{nm} r) J_{s_n}(\mu_{nm} \xi) \cos(s_n \varphi) \cos(s_n \eta) \sin(\mu_{nm} at), \\ s_n &= \frac{n\pi}{\varphi_0}, \quad A_{nm} = \frac{4\mu_{nm}}{a\varphi_0(\mu_{nm}^2 R^2 + k^2 R^2 - s_n^2) [J_{s_n}(\mu_{nm} R)]^2}, \end{aligned}$$

where the  $J_{s_n}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_{s_n}(\mu R) + k J_{s_n}(\mu R) = 0.$$

### 5.1.3. Axisymmetric Problems

In the axisymmetric case the wave equation in the cylindrical system of coordinates has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} \right), \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional problems with axial symmetry that have solutions  $w = w(r, t)$  are considered in Subsection 4.2.1.

In the solution of the problems considered below, the modified Green's function  $\mathcal{G}(r, z, \xi, \eta, t) = 2\pi \xi G(r, z, \xi, \eta, t)$  is used for convenience.

5.1.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . First boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad - a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad + a^2 \int_0^t \int_0^R g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad - a^2 \int_0^t \int_0^R g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} \mathcal{G}(r, z, \xi, \eta, t) &= \frac{4\xi}{R^2 l} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nm}})}{a\sqrt{\lambda_{nm}}}, \\ \lambda_{nm} &= \frac{\mu_n^2}{R^2} + \frac{\pi^2 m^2}{l^2}, \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

5.1.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$



Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_0^R g_2(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^R g_3(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = & \frac{2t\xi}{R^2 l} + \frac{2\xi}{R^2 l} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_{nm}}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \\
 & \times \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nm}})}{a\sqrt{\lambda_{nm}}}, \\
 \lambda_{nm} = & \frac{\mu_n^2}{R^2} + \frac{\pi^2 m^2}{l^2}, \quad A_{nm} = \begin{cases} 0 & \text{for } m = 0, n = 0, \\ 1 & \text{for } m = 0, n > 0, \\ 2 & \text{for } m > 0, \end{cases}
 \end{aligned}$$

where the  $\mu_n$  are zeros of the first-order Bessel function,  $J_1(\mu) = 0$  ( $\mu_0 = 0$ ).

### 5.1.3-3. Domain: $0 \leq r \leq R$ , $0 \leq z \leq l$ . Third boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_r w + k_1 w = g_1(z, t) & \quad \text{at } r = R \quad (\text{boundary condition}), \\
 \partial_z w - k_2 w = g_2(r, t) & \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
 \partial_z w + k_3 w = g_3(r, t) & \quad \text{at } z = l \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(r, z, t)$  is determined by the formula in Paragraph 5.1.3-2 where

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = & \frac{2\xi}{R^2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_n^2}{(k_1^2 R^2 + \mu_n^2) J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\varphi_m(z) \varphi_m(\eta)}{\|\varphi_m\|^2} \frac{\sin(at\sqrt{\lambda_{nm}})}{a\sqrt{\lambda_{nm}}}, \\
 \lambda_{nm} = & \frac{\mu_n^2}{R^2} + \beta_m^2, \quad \varphi_m(z) = \cos(\beta_m z) + \frac{k_2}{\beta_m} \sin(\beta_m z), \\
 \|\varphi_m\|^2 = & \frac{k_3}{2\beta_m^2} \frac{\beta_m^2 + k_2^2}{\beta_m^2 + k_3^2} + \frac{k_2}{2\beta_m^2} + \frac{l}{2} \left(1 + \frac{k_2^2}{\beta_m^2}\right).
 \end{aligned}$$

Here, the  $\mu_n$  and  $\beta_m$  are positive roots of the transcendental equations

$$\mu J_1(\mu) - k_1 R J_0(\mu) = 0, \quad \frac{\tan(\beta l)}{\beta} = \frac{k_2 + k_3}{\beta^2 - k_2 k_3}.$$

**5.1.3-4. Domain:  $0 \leq r \leq R, 0 \leq z \leq l$ . Mixed boundary value problems.**

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(z, t) & \text{at } r &= R & \text{(boundary condition),} \\ \partial_z w &= g_2(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad - a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &\quad - a^2 \int_0^t \int_0^R g_2(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau \\ &\quad + a^2 \int_0^t \int_0^R g_3(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau. \end{aligned}$$

Here,

$$\begin{aligned} \mathcal{G}(r, z, \xi, \eta, t) &= \frac{2\xi}{R^2 l} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nm}})}{a\sqrt{\lambda_{nm}}}, \\ \lambda_{nm} &= \frac{\mu_n^2}{R^2} + \frac{\pi^2 m^2}{l^2}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 0, \end{cases} \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(z, t) & \text{at } r &= R & \text{(boundary condition),} \\ w &= g_2(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_3(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &\quad + a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R, \eta, t - \tau) d\eta d\tau \\ &\quad + a^2 \int_0^t \int_0^R g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &\quad - a^2 \int_0^t \int_0^R g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau. \end{aligned}$$

Here,

$$\mathcal{G}(r, z, \xi, \eta, t) = \frac{4\xi}{R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nm}})}{a\sqrt{\lambda_{nm}}},$$

$$\lambda_{nm} = \frac{\mu_n^2}{R^2} + \frac{\pi^2 m^2}{l^2},$$

where the  $\mu_n$  are zeros of the first-order Bessel function,  $J_1(\mu) = 0$  ( $\mu_0 = 0$ ).

**5.1.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . First boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(z, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ w &= g_2(z, t) & \text{at } r &= R_2 & \text{(boundary condition),} \\ w &= g_3(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_4(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_{R_1}^{R_2} f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + \int_0^l \int_{R_1}^{R_2} f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\ &- a^2 \int_0^t \int_0^l g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau \\ &+ a^2 \int_0^t \int_{R_1}^{R_2} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- a^2 \int_0^t \int_{R_1}^{R_2} g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau. \end{aligned}$$

Here,

$$\mathcal{G}(r, z, \xi, \eta, t) = \frac{\pi^2 \xi}{R_1^2 l} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_n^2 J_0^2(s\mu_n)}{J_0^2(\mu_n) - J_0^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nm}})}{a\sqrt{\lambda_{nm}}},$$

$$\Psi_n(r) = Y_0(\mu_n) J_0\left(\frac{\mu_n r}{R_1}\right) - J_0(\mu_n) Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1}, \quad \lambda_{nm} = \frac{\mu_n^2}{R_1^2} + \frac{\pi^2 m^2}{l^2},$$

where  $J_0(\mu)$  and  $Y_0(\mu)$  are the Bessel functions, and the  $\mu_n$  are positive roots of the transcendental equation

$$J_0(\mu) Y_0(s\mu) - J_0(s\mu) Y_0(\mu) = 0.$$

**5.1.3-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(z, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ \partial_r w &= g_2(z, t) & \text{at } r &= R_2 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_4(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_{R_1}^{R_2} f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + \int_0^l \int_{R_1}^{R_2} f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & - a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R_1, \eta, t - \tau) d\eta d\tau + a^2 \int_0^t \int_0^l g_2(\eta, \tau) \mathcal{G}(r, z, R_2, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_{R_1}^{R_2} g_3(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau + a^2 \int_0^t \int_{R_1}^{R_2} g_4(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = & \frac{2t\xi}{(R_2^2 - R_1^2)l} + \frac{4\xi}{\pi a(R_2^2 - R_1^2)} \sum_{m=1}^{\infty} \frac{1}{m} \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi\eta}{l}\right) \sin\left(\frac{m\pi at}{l}\right) \\
 & + \frac{\pi^2 \xi}{2R_1^2 l} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m \mu_n^2 J_1^2(s\mu_n)}{J_1^2(\mu_n) - J_1^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi\eta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nm}})}{a\sqrt{\lambda_{nm}}},
 \end{aligned}$$

where

$$\begin{aligned}
 \Psi_n(r) = & Y_1(\mu_n) J_0\left(\frac{\mu_n r}{R_1}\right) - J_1(\mu_n) Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1}, \\
 A_m = & \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 1, \end{cases} \quad \lambda_{nm} = \frac{\mu_n^2}{R_1^2} + \frac{\pi^2 m^2}{l^2};
 \end{aligned}$$

$J_k(\mu)$  and  $Y_k(\mu)$  are the Bessel functions ( $k=0, 1$ ); and the  $\mu_n$  are positive roots of the transcendental equation

$$J_1(\mu)Y_1(s\mu) - J_1(s\mu)Y_1(\mu) = 0.$$

## 5.2. Nonhomogeneous Wave Equation

$$\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_2 w + \Phi(x, y, t)$$

### 5.2.1. Problems in Cartesian Coordinates

5.2.1-1. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned}
 w = f(x, y) & \quad \text{at } t = 0, \\
 \partial_t w = g(x, y) & \quad \text{at } t = 0.
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{1}{2\pi a} \frac{\partial}{\partial t} \iint_{\rho \leq at} \frac{f(\xi, \eta) d\xi d\eta}{\sqrt{a^2 t^2 - \rho^2}} + \frac{1}{2\pi a} \iint_{\rho \leq at} \frac{g(\xi, \eta) d\xi d\eta}{\sqrt{a^2 t^2 - \rho^2}} \\
 & + \frac{1}{2\pi a} \int_0^t \left[ \iint_{\rho \leq a(t-\tau)} \frac{\Phi(\xi, \eta, \tau) d\xi d\eta}{\sqrt{a^2 (t-\tau)^2 - \rho^2}} \right] d\tau, \quad \rho^2 = (\xi - x)^2 + (\eta - y)^2.
 \end{aligned}$$

⊙ *Reference:* N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970).

**5.2.1-2. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . First boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is given by the formula in Paragraph 5.1.1-3 with the additional term

$$\int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,$$

which allows for the equation's nonhomogeneity; this term is the solution of the nonhomogeneous equation with homogeneous initial and boundary conditions.

☉ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.2.1-3. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Second boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_1(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_2(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is given by the formula in Paragraph 5.1.1-4 with the additional term specified in Paragraph 5.2.1-2 (the Green's function is taken from Paragraph 5.1.1-4).

☉ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.2.1-4. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Third boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_1(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_2(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_x w - k_1 w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w + k_2 w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w - k_3 w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w + k_4 w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is the sum of the solution to the homogeneous equation with nonhomogeneous initial and boundary conditions (see Paragraph 5.1.1-5) and the solution to the nonhomogeneous equation with homogeneous initial and boundary conditions. This solution is given by the formula in Paragraph 5.2.1-2 in which one should substitute the Green's function of Paragraph 5.1.1-5).

☉ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.2.1-5. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Mixed boundary value problems.**

1°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_1(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_2(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(y, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is given by the formula in Paragraph 5.1.1-6, Item 1°, with the additional term specified in Paragraph 5.2.1-2.

2°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_1(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_2(x, y) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ w &= g_3(x, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(x, y, t)$  is given by the formula in Paragraph 5.1.1-6, Item 2°, with the additional term specified in Paragraph 5.2.1-2.

## 5.2.2. Problems in Polar Coordinates

A nonhomogeneous wave equation in the polar coordinate system has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} \right) + \Phi(r, \varphi, t), \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional boundary value problems independent of the angular coordinate  $\varphi$  are considered in Subsection 4.2.2.

**5.2.2-1. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is given by the formula in Paragraph 5.1.2-1 with the additional term

$$\int_0^t \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi \, d\xi \, d\eta \, d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity; this term is the solution of the nonhomogeneous equation with homogeneous initial and boundary conditions.

⊙ *References:* N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.2.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is given by the formula in Paragraph 5.1.2-2 with the additional term (1).

⊙ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.2.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.**

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is the sum of the solution to the homogeneous equation with nonhomogeneous initial and boundary conditions (see Paragraph 5.1.2-3) and the solution to the nonhomogeneous equation with homogeneous initial and boundary conditions [this solution is given by formula (1) in which one should substitute the Green's function in Paragraph 5.1.2-3].

⊙ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.2.2-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(\varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is given by the formula in Paragraph 5.1.2-4 with the additional term

$$\int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity; this term is the solution of the nonhomogeneous equation with homogeneous initial and boundary conditions.

**5.2.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is given by the formula in Paragraph 5.1.2-5 with the additional term (2).

**5.2.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(\varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(\varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is the sum of the solution to the homogeneous equation with nonhomogeneous initial and boundary conditions (see Paragraph 5.1.2-6) and the solution to the nonhomogeneous equation with homogeneous initial and boundary conditions [this solution is given by formula (2) in which one should substitute the Green's function in Paragraph 5.1.2-6].

**5.2.2-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . First boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is given by the formula in Paragraph 5.1.2-7 with the additional term

$$\int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau, \quad (3)$$

which allows for the equation's nonhomogeneity.

**5.2.2-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Second boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_2(r, t) & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ r^{-1} \partial_\varphi w &= g_3(r, t) & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is given by the formula in Paragraph 5.1.2-8 with the additional term (3).

**5.2.2-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Mixed boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is given by the formula in Paragraph 5.1.2-9 with the additional term (3).



### 5.2.3. Axisymmetric Problems

In the axisymmetric case, a nonhomogeneous wave equation in the cylindrical system of coordinates has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} \right) + \Phi(r, z, t), \quad r = \sqrt{x^2 + y^2}.$$

**5.2.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . First boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(z, t) & \text{at } r &= R & \text{(boundary condition),} \\ w &= g_2(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_3(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

The solution  $w(r, z, t)$  is given by the formula in Paragraph 5.1.3-1 with the additional term

$$\int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity; this term is the solution of the nonhomogeneous equation with homogeneous initial and boundary conditions.

**5.2.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(z, t) & \text{at } r &= R & \text{(boundary condition),} \\ \partial_z w &= g_2(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

The solution  $w(r, z, t)$  is given by the formula in Paragraph 5.1.3-2 with the additional term (1).

**5.2.3-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . Third boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w + k_1 w &= g_1(z, t) & \text{at } r &= R & \text{(boundary condition),} \\ \partial_z w - k_2 w &= g_2(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w + k_3 w &= g_3(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

The solution  $w(r, z, t)$  is the sum of the solution to the homogeneous equation with nonhomogeneous initial and boundary conditions (see Paragraph 5.1.3-3) and the solution to the nonhomogeneous equation with homogeneous initial and boundary conditions [this solution is given by formula (1) in which one should substitute the Green's function in Paragraph 5.1.3-3].

**5.2.3-4. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.**

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, z, t)$  is given by the formula in Paragraph 5.1.3-4, Item 1°, with the additional term (1).

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, z, t)$  is given by the formula in Paragraph 5.1.3-4, Item 2°, with the additional term (1).

**5.2.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . First boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_4(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, z, t)$  is given by the formula in Paragraph 5.1.3-5 with the additional term

$$\int_0^t \int_0^l \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity; this term is the solution of the nonhomogeneous equation with homogeneous initial and boundary conditions.

**5.2.3-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_4(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, z, t)$  is given by the formula in Paragraph 5.1.3-6 with the additional term (2).

## 5.3. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_2 w - bw + \Phi(x, y, t)$

### 5.3.1. Problems in Cartesian Coordinates

The *two-dimensional nonhomogeneous Klein–Gordon equation* with two space variables in the rectangular Cartesian coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) - bw + \Phi(x, y, t).$$

#### 5.3.1-1. Fundamental solutions.

1°. Case  $b = -\sigma^2 < 0$ :

$$\mathcal{E}(x, y, t) = \frac{\vartheta(at - r)}{2\pi a^2} \frac{\cosh(\sigma \sqrt{t^2 - r^2/a^2})}{\sqrt{t^2 - r^2/a^2}}, \quad r = \sqrt{x^2 + y^2},$$

where  $\vartheta(z)$  is the Heaviside unit step function.

2°. Case  $b = \sigma^2 > 0$ :

$$\mathcal{E}(x, y, t) = \frac{\vartheta(at - r)}{2\pi a^2} \frac{\cos(\sigma \sqrt{t^2 - r^2/a^2})}{\sqrt{t^2 - r^2/a^2}}, \quad r = \sqrt{x^2 + y^2}.$$

⊙ *References:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

#### 5.3.1-2. Domain: $-\infty < x < \infty, -\infty < y < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(x, y) \quad \text{at} \quad t = 0. \end{aligned}$$

1°. Solution for  $b = -a^2 c^2 < 0$ :

$$\begin{aligned} w(x, y, t) &= \frac{1}{2\pi a} \frac{\partial}{\partial t} \iint_{\rho \leq at} f(\xi, \eta) \frac{\cosh(c\sqrt{a^2 t^2 - \rho^2})}{\sqrt{a^2 t^2 - \rho^2}} d\xi d\eta + \frac{1}{2\pi a} \iint_{\rho \leq at} g(\xi, \eta) \frac{\cosh(c\sqrt{a^2 t^2 - \rho^2})}{\sqrt{a^2 t^2 - \rho^2}} d\xi d\eta \\ &+ \frac{1}{2\pi a} \int_0^t d\tau \iint_{\rho \leq a(t-\tau)} \Phi(\xi, \eta, \tau) \frac{\cosh(c\sqrt{a^2(t-\tau)^2 - \rho^2})}{\sqrt{a^2(t-\tau)^2 - \rho^2}} d\xi d\eta, \quad \rho = \sqrt{(x-\xi)^2 + (y-\eta)^2}. \end{aligned}$$

2°. Solution for  $b = a^2 c^2 > 0$ :

$$\begin{aligned} w(x, y, t) &= \frac{1}{2\pi a} \frac{\partial}{\partial t} \iint_{\rho \leq at} f(\xi, \eta) \frac{\cos(c\sqrt{a^2 t^2 - \rho^2})}{\sqrt{a^2 t^2 - \rho^2}} d\xi d\eta + \frac{1}{2\pi a} \iint_{\rho \leq at} g(\xi, \eta) \frac{\cos(c\sqrt{a^2 t^2 - \rho^2})}{\sqrt{a^2 t^2 - \rho^2}} d\xi d\eta \\ &+ \frac{1}{2\pi a} \int_0^t d\tau \iint_{\rho \leq a(t-\tau)} \Phi(\xi, \eta, \tau) \frac{\cos(c\sqrt{a^2(t-\tau)^2 - \rho^2})}{\sqrt{a^2(t-\tau)^2 - \rho^2}} d\xi d\eta, \quad \rho = \sqrt{(x-\xi)^2 + (y-\eta)^2}. \end{aligned}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**5.3.1-3. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . First boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_t w &= f_1(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\
 w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\
 w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\
 w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) &= \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 &+ a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 &- a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 &- a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=l_2} d\xi d\tau \\
 &+ \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) &= \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\lambda_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \sin(\lambda_{nm} t), \\
 p_n &= \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b}.
 \end{aligned}$$

**5.3.1-4. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Second boundary value problem.**

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_t w &= f_1(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_x w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\
 \partial_x w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\
 \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\
 \partial_y w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & - a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & \frac{\sin(t\sqrt{b})}{l_1 l_2 \sqrt{b}} + \frac{2}{l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_{nm}}{\lambda_{nm}} \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) \sin(\lambda_{nm} t), \\
 p_n = & \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b}, \quad A_{nm} = \begin{cases} 0 & \text{for } n = m = 0, \\ 1 & \text{for } nm = 0 \text{ (} n \neq m \text{)}, \\ 2 & \text{for } nm \neq 0. \end{cases}
 \end{aligned}$$

5.3.1-5. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Third boundary value problem.

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(x, y) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(x, y) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_x w - k_1 w = g_1(y, t) & \quad \text{at } x = 0 \quad (\text{boundary condition}), \\
 \partial_x w + k_2 w = g_2(y, t) & \quad \text{at } x = l_1 \quad (\text{boundary condition}), \\
 \partial_y w - k_3 w = g_3(x, t) & \quad \text{at } y = 0 \quad (\text{boundary condition}), \\
 \partial_y w + k_4 w = g_4(x, t) & \quad \text{at } y = l_2 \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 5.3.1-3 where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & 4 \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{E_{nm} \sqrt{a^2 \mu_n^2 + a^2 \nu_m^2 + b}} \sin(\mu_n x + \varepsilon_n) \sin(\nu_m y + \sigma_m) \\
 & \times \sin(\mu_n \xi + \varepsilon_n) \sin(\nu_m \eta + \sigma_m) \sin(t \sqrt{a^2 \mu_n^2 + a^2 \nu_m^2 + b}), \\
 \varepsilon_n = & \arctan \frac{\mu_n}{l_1}, \quad \sigma_m = \arctan \frac{\nu_m}{l_2}, \quad E_{nm} = \left[ l_1 + \frac{(k_1 k_2 + \mu_n^2)(k_1 + k_2)}{(k_1^2 + \mu_n^2)(k_2^2 + \mu_n^2)} \right] \left[ l_2 + \frac{(k_3 k_4 + \nu_m^2)(k_3 + k_4)}{(k_3^2 + \nu_m^2)(k_4^2 + \nu_m^2)} \right].
 \end{aligned}$$

Here, the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\begin{aligned}
 \mu^2 - k_1 k_2 = (k_1 + k_2) \mu \cot(l_1 \mu), \\
 \nu^2 - k_3 k_4 = (k_3 + k_4) \nu \cot(l_2 \nu).
 \end{aligned}$$

⊙ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

1°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_t w &= f_1(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\
 w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\
 \partial_y w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\
 \partial_y w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) &= \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 &+ \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 &+ a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 &- a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\
 &- a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\
 &+ a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau \\
 &+ \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) &= \frac{2}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{\lambda_{nm}} \sin(p_n x) \cos(q_m y) \sin(p_n \xi) \cos(q_m \eta) \sin(\lambda_{nm} t), \\
 p_n &= \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases}
 \end{aligned}$$

2°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_t w &= f_1(x, y) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 w &= g_1(y, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\
 \partial_x w &= g_2(y, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\
 w &= g_3(x, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\
 \partial_y w &= g_4(x, t) & \text{at } y = l_2 & \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} f_1(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & \frac{4}{l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{1}{\lambda_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \sin(\lambda_{nm} t), \\
 p_n = & \frac{\pi(2n+1)}{2l_1}, \quad q_m = \frac{\pi(2m+1)}{2l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b}.
 \end{aligned}$$

### 5.3.2. Problems in Polar Coordinates

A nonhomogeneous Klein–Gordon equation with two space variables in the polar coordinate system has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} \right) - bw + \Phi(r, \varphi, t), \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional solutions  $w = w(r, t)$  independent of the angular coordinate  $\varphi$  are considered in Subsection 4.2.5.

5.3.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A circle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 w = g(\varphi, t) \quad \text{at} \quad r = R \quad & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & - a^2 R \int_0^t \int_0^{2\pi} g(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\
 & + \int_0^t \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,\*

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm}R)]^2} J_n(\mu_{nm}r) J_n(\mu_{nm}\xi) \cos[n(\varphi - \eta)] \frac{\sin(t\sqrt{a^2\mu_{nm}^2 + b})}{\sqrt{a^2\mu_{nm}^2 + b}},$$

$$A_0 = 1, \quad A_n = 2 \quad (n = 1, 2, \dots),$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

5.3.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &+ a^2 R \int_0^t \int_0^{2\pi} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\ &+ \int_0^t \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{\sin(t\sqrt{b})}{\pi R^2 \sqrt{b}} \\ &+ \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm}r) J_n(\mu_{nm}\xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm}R)]^2} \cos[n(\varphi - \eta)] \frac{\sin(t\sqrt{a^2\mu_{nm}^2 + b})}{\sqrt{a^2\mu_{nm}^2 + b}}, \end{aligned}$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(\xi)$  are the Bessel functions; and the  $\mu_m$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

5.3.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is determined by the formula in Paragraph 5.3.2-2 where

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm}r) J_n(\mu_{nm}\xi)}{(\mu_{nm}^2 R^2 + k^2 R^2 - n^2) [J_n(\mu_{nm}R)]^2} \cos[n(\varphi - \eta)] \frac{\sin(t\sqrt{a^2\mu_{nm}^2 + b})}{\sqrt{a^2\mu_{nm}^2 + b}}, \\ A_0 &= 1, \quad A_n = 2 \quad (n = 1, 2, \dots). \end{aligned}$$

Here, the  $J_n(\xi)$  are the Bessel functions and the  $\mu_m$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k J_n(\mu R) = 0.$$

---

\* In the expressions of the Green's functions specified in Subsection 5.3.2, the ratios  $\sin(t\sqrt{a^2\mu_{nm}^2 + b}) / \sqrt{a^2\mu_{nm}^2 + b}$  must be replaced by  $\sinh(t\sqrt{|a^2\mu_{nm}^2 + b|}) / \sqrt{|a^2\mu_{nm}^2 + b|}$  if  $a^2\mu_{nm}^2 + b < 0$ .



**5.3.2-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad + a^2 R_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\ &\quad - a^2 R_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau \\ &\quad + \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \frac{\pi}{2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_n B_{nm} Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \frac{\sin(t \sqrt{a^2 \mu_{nm}^2 + b})}{\sqrt{a^2 \mu_{nm}^2 + b}}, \\ A_n &= \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)}, \\ Z_n(\mu_{nm} r) &= J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r), \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

**5.3.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &\quad - a^2 R_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) G(r, \varphi, R_1, \eta, t - \tau) d\eta d\tau \\ &\quad + a^2 R_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) G(r, \varphi, R_2, \eta, t - \tau) d\eta d\tau \\ &\quad + \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{\sin(t\sqrt{b})}{\pi(R_2^2 - R_1^2)\sqrt{b}} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm} r) Z_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \sin(t\sqrt{a^2 \mu_{nm}^2 + b})}{[(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm} R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm} R_1)] \sqrt{a^2 \mu_{nm}^2 + b}},$$

where

$$Z_n(\mu_{nm} r) = J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r), \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

5.3.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, \varphi) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w - k_1 w &= g_1(\varphi, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ \partial_r w + k_2 w &= g_2(\varphi, t) & \text{at } r &= R_2 & \text{(boundary condition).} \end{aligned}$$

The solution  $w(r, \varphi, t)$  is determined by the formula in Paragraph 5.3.2-5 where

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_{nm}(r) Z_{nm}(\xi) \cos[n(\varphi - \eta)] \sin(\lambda_{nm} t)}{\lambda_{nm} [(k_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)]},$$

$$\begin{aligned} Z_{nm}(r) &= [\mu_{nm} J'_n(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) \\ &\quad - [\mu_{nm} Y'_n(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r). \end{aligned}$$

Here,  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ;  $\lambda_{nm} = \sqrt{a^2 \mu_{nm}^2 + b}$ ; the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned} &[\mu J'_n(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y'_n(\mu R_2) + k_2 Y_n(\mu R_2)] \\ &= [\mu Y'_n(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J'_n(\mu R_2) + k_2 J_n(\mu R_2)]. \end{aligned}$$

5.3.2-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . First boundary value problem.

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, \varphi) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(\varphi, t) & \text{at } r &= R & \text{(boundary condition),} \\ w &= g_2(r, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ w &= g_3(r, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & - a^2 R \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\
 & + a^2 \int_0^t \int_0^R g_2(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & - a^2 \int_0^t \int_0^R g_3(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\tau \\
 & + \int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{4}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}},$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ , and  $\lambda_{nm} = \sqrt{a^2 \mu_{nm}^2 + b}$ .

5.3.2-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Second boundary value problem.

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_r w &= g_1(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\
 r^{-1} \partial_\varphi w &= g_2(r, t) \quad \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\
 r^{-1} \partial_\varphi w &= g_3(r, t) \quad \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta + \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + a^2 R \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_0^R g_2(\xi, \tau) G(r, \varphi, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^R g_3(\xi, \tau) G(r, \varphi, \xi, \varphi_0, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \frac{2 \sin(t\sqrt{b})}{R^2 \varphi_0 \sqrt{b}} + 4\varphi_0 \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_{nm}^2 J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{(R^2 \varphi_0^2 \mu_{nm}^2 - n^2 \pi^2) [J_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \\
 & \times \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \frac{\sin(t\sqrt{a^2 \mu_{nm}^2 + b})}{\sqrt{a^2 \mu_{nm}^2 + b}},
 \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_{n\pi/\varphi_0}(\mu R) = 0$ .

**5.3.2-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Mixed boundary value problem.**

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_r w + kw &= g(\varphi, t) & \text{at } r &= R & (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi &= 0 & (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &+ \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &+ a^2 R \int_0^t \int_0^{\varphi_0} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\ &+ \int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, \xi, \eta, t) &= \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_{nm} J_{s_n}(\mu_{nm} r) J_{s_n}(\mu_{nm} \xi) \cos(s_n \varphi) \cos(s_n \eta) \sin(t \sqrt{a^2 \mu_{nm}^2 + b}), \\ s_n &= \frac{n\pi}{\varphi_0}, \quad A_{nm} = \frac{4\mu_{nm}^2}{\varphi_0(\mu_{nm}^2 R^2 + k^2 R^2 - s_n^2) [J_{s_n}(\mu_{nm} R)]^2 \sqrt{a^2 \mu_{nm}^2 + b}}, \end{aligned}$$

where the  $J_{s_n}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_{s_n}(\mu R) + k J_{s_n}(\mu R) = 0.$$

### 5.3.3. Axisymmetric Problems

In the axisymmetric case, a nonhomogeneous Klein–Gordon equation in the cylindrical system of coordinates has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} \right) - bw + \Phi(r, z, t), \quad r = \sqrt{x^2 + y^2}.$$

In the solutions of the problems considered below, the modified Green's function  $\mathcal{G}(r, z, \xi, \eta, t) = 2\pi \xi G(r, z, \xi, \eta, t)$  is used for convenience.

**5.3.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . First boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r &= R & (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } z &= 0 & (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } z &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & - a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\
 & + a^2 \int_0^t \int_0^R g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & - a^2 \int_0^t \int_0^R g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau \\
 & + \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = & \frac{4\xi}{R^2 l} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}}, \\
 \lambda_{nm} = & \frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b,
 \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

**5.3.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, z) \quad & \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, z) \quad & \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_r w = g_1(z, t) \quad & \text{at} \quad r = R \quad (\text{boundary condition}), \\
 \partial_z w = g_2(r, t) \quad & \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
 \partial_z w = g_3(r, t) \quad & \text{at} \quad z = l \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_0^R g_2(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^R g_3(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned} \mathcal{G}(r, z, \xi, \eta, t) &= \frac{2\xi \sin(t\sqrt{b})}{R^2 l \sqrt{b}} \\ &+ \frac{2\xi}{R^2 l} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_{nm}}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}}, \\ \lambda_{nm} &= \frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b, \quad A_{nm} = \begin{cases} 0 & \text{for } m=0, n=0, \\ 1 & \text{for } m=0, n>0, \\ 2 & \text{for } m>0, \end{cases} \end{aligned}$$

where the  $\mu_n$  are zeros of the first-order Bessel function,  $J_1(\mu) = 0$  ( $\mu_0 = 0$ ).

### 5.3.3-3. Domain: $0 \leq r \leq R$ , $0 \leq z \leq l$ . Third boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_r w + k_1 w &= g_1(z, t) \quad \text{at } r = R \quad (\text{boundary condition}), \\ \partial_z w - k_2 w &= g_2(r, t) \quad \text{at } z = 0 \quad (\text{boundary condition}), \\ \partial_z w + k_3 w &= g_3(r, t) \quad \text{at } z = l \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, z, t)$  is determined by the formula in Paragraph 5.3.3-2 where

$$\begin{aligned} \mathcal{G}(r, z, \xi, \eta, t) &= \frac{2\xi}{R^2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_n^2}{(k_1^2 R^2 + \mu_n^2) J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\varphi_m(z) \varphi_m(\eta)}{\|\varphi_m\|^2} \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}}, \\ \lambda_{nm} &= \frac{a^2 \mu_n^2}{R^2} + a^2 \beta_m^2 + b, \quad \varphi_m(z) = \cos(\beta_m z) + \frac{k_2}{\beta_m} \sin(\beta_m z), \\ \|\varphi_m\|^2 &= \frac{k_3}{2\beta_m^2} \frac{\beta_m^2 + k_2^2}{\beta_m^2 + k_3^2} + \frac{k_2}{2\beta_m^2} + \frac{l}{2} \left(1 + \frac{k_2^2}{\beta_m^2}\right). \end{aligned}$$

Here, the  $\mu_n$  and  $\beta_m$  are positive roots of the transcendental equations

$$\mu J_1(\mu) - k_1 R J_0(\mu) = 0, \quad \frac{\tan(\beta l)}{\beta} = \frac{k_2 + k_3}{\beta^2 - k_2 k_3}.$$

### 5.3.3-4. Domain: $0 \leq r \leq R$ , $0 \leq z \leq l$ . Mixed boundary value problems.

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ w &= g_1(z, t) \quad \text{at } r = R \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) \quad \text{at } z = 0 \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, t) \quad \text{at } z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & - a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\
 & - a^2 \int_0^t \int_0^R g_2(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^R g_3(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = & \frac{2\xi}{R^2 l} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}}, \\
 \lambda_{nm} = & \frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 0, \end{cases}
 \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_r w = g_1(z, t) & \quad \text{at } r = R \quad (\text{boundary condition}), \\
 w = g_2(r, t) & \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
 w = g_3(r, t) & \quad \text{at } z = l \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + \int_0^l \int_0^R f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R, \eta, t - \tau) d\eta d\tau \\
 & + a^2 \int_0^t \int_0^R g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & - a^2 \int_0^t \int_0^R g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau \\
 & + \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = & \frac{4\xi}{R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}}, \\
 \lambda_{nm} = & \frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b,
 \end{aligned}$$

where the  $\mu_n$  are zeros of the first-order Bessel function,  $J_1(\mu) = 0$  ( $\mu_0 = 0$ ).

**5.3.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . First boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(z, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ w &= g_2(z, t) & \text{at } r &= R_2 & \text{(boundary condition),} \\ w &= g_3(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_4(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_{R_1}^{R_2} f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ \int_0^l \int_{R_1}^{R_2} f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\ &- a^2 \int_0^t \int_0^l g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau \\ &+ a^2 \int_0^t \int_{R_1}^{R_2} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- a^2 \int_0^t \int_{R_1}^{R_2} g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau \\ &+ \int_0^t \int_0^l \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} \mathcal{G}(r, z, \xi, \eta, t) &= \frac{\pi^2 \xi}{R_1^2 l} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_n^2 J_0^2(s\mu_n)}{J_0^2(\mu_n) - J_0^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}}, \\ \Psi_n(r) &= Y_0(\mu_n) J_0\left(\frac{\mu_n r}{R_1}\right) - J_0(\mu_n) Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1}, \quad \lambda_{nm} = \frac{a^2 \mu_n^2}{R_1^2} + \frac{a^2 \pi^2 m^2}{l^2} + b, \end{aligned}$$

where  $J_0(\mu)$  and  $Y_0(\mu)$  are the Bessel functions, and the  $\mu_n$  are positive roots of the transcendental equation

$$J_0(\mu) Y_0(s\mu) - J_0(s\mu) Y_0(\mu) = 0.$$

**5.3.3-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(z, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ \partial_r w &= g_2(z, t) & \text{at } r &= R_2 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_4(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$



Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_{R_1}^{R_2} f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + \int_0^l \int_{R_1}^{R_2} f_1(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & - a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R_1, \eta, t - \tau) d\eta d\tau \\
 & + a^2 \int_0^t \int_0^l g_2(\eta, \tau) \mathcal{G}(r, z, R_2, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_{R_1}^{R_2} g_3(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_{R_1}^{R_2} g_4(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^l \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = & \frac{2\xi \sin(t\sqrt{b})}{(R_2^2 - R_1^2)l\sqrt{b}} + \frac{4\xi}{(R_2^2 - R_1^2)l} \sum_{m=1}^{\infty} \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\beta_m})}{\sqrt{\beta_m}} \\
 & + \frac{\pi^2 \xi}{2R_1^2 l} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m \mu_n^2 J_1^2(s\mu_n)}{J_1^2(\mu_n) - J_1^2(s\mu_n)} \Psi_n(r) \Psi_n(\xi) \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}},
 \end{aligned}$$

where

$$\begin{aligned}
 \Psi_n(r) = & Y_1(\mu_n) J_0\left(\frac{\mu_n r}{R_1}\right) - J_1(\mu_n) Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1}, \\
 A_m = & \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 1, \end{cases} \quad \beta_m = \frac{a^2 \pi^2 m^2}{l^2} + b, \quad \lambda_{nm} = \frac{a^2 \mu_n^2}{R_1^2} + \frac{a^2 \pi^2 m^2}{l^2} + b;
 \end{aligned}$$

$J_k(\mu)$  and  $Y_k(\mu)$  are the Bessel functions ( $k=0, 1$ ); and the  $\mu_n$  are positive roots of the transcendental equation

$$J_1(\mu)Y_1(s\mu) - J_1(s\mu)Y_1(\mu) = 0.$$

## 5.4. Telegraph Equation

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \Delta_2 w - bw + \Phi(x, y, t)$$

### 5.4.1. Problems in Cartesian Coordinates

A two-dimensional nonhomogeneous telegraph equation in the rectangular Cartesian coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) - bw + \Phi(x, y, t).$$

#### 5.4.1-1. Reduction to the two-dimensional Klein–Gordon equation.

The substitution  $w(x, y, t) = \exp(-\frac{1}{2}kt)u(x, y, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right) - \left(b - \frac{1}{4}k^2\right)u + \exp\left(\frac{1}{2}kt\right)\Phi(x, y, t),$$

which is discussed in Subsection 5.3.1.

### 5.4.1-2. Fundamental solutions.

1°. Case  $b - \frac{1}{4}k^2 = \sigma^2 > 0$ :

$$\mathcal{E}(x, y, t) = \vartheta(at - r) \exp\left(-\frac{1}{2}kt\right) \frac{\cos\left(\sigma \sqrt{t^2 - r^2/a^2}\right)}{2\pi a^2 \sqrt{t^2 - r^2/a^2}},$$

where  $r = \sqrt{x^2 + y^2}$  and  $\vartheta(z)$  is the Heaviside unit step function.

2°. Case  $b - \frac{1}{4}k^2 = -\sigma^2 < 0$ :

$$\mathcal{E}(x, y, t) = \vartheta(at - r) \exp\left(-\frac{1}{2}kt\right) \frac{\cosh\left(\sigma \sqrt{t^2 - r^2/a^2}\right)}{2\pi a^2 \sqrt{t^2 - r^2/a^2}}.$$

⊙ *Reference*: V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

### 5.4.1-3. Domain: $-\infty < x < \infty$ , $-\infty < y < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x, y) & \text{at } t &= 0, \\ \partial_t w &= g(x, y) & \text{at } t &= 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, t) &= \exp\left(-\frac{1}{2}kt\right) \frac{\partial}{\partial t} \iint_{\rho \leq at} f(\xi, \eta) H(x, y, \xi, \eta, t) d\xi d\eta \\ &+ \exp\left(-\frac{1}{2}kt\right) \iint_{\rho \leq at} [g(\xi, \eta) + \frac{1}{2}kf(\xi, \eta)] H(x, y, \xi, \eta, t) d\xi d\eta \\ &+ \int_0^t d\tau \iint_{\rho \leq a(t-\tau)} \exp\left[-\frac{1}{2}k(t-\tau)\right] \Phi(\xi, \eta, \tau) H(x, y, \xi, \eta, t-\tau) d\xi d\eta. \end{aligned}$$

Here,

$$H(x, y, \xi, \eta, t) = \begin{cases} \frac{\cos\left(\sigma \sqrt{t^2 - \rho^2/a^2}\right)}{2\pi a^2 \sqrt{t^2 - \rho^2/a^2}} & \text{for } b - \frac{1}{4}k^2 = \sigma^2 > 0, \\ \frac{\cosh\left(\sigma \sqrt{t^2 - \rho^2/a^2}\right)}{2\pi a^2 \sqrt{t^2 - \rho^2/a^2}} & \text{for } b - \frac{1}{4}k^2 = -\sigma^2 < 0, \end{cases}$$

where  $\rho = \sqrt{(x - \xi)^2 + (y - \eta)^2}$ .

### 5.4.1-4. Domain: $0 \leq x \leq l_1$ , $0 \leq y \leq l_2$ . First boundary value problem.

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(y, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ w &= g_4(x, t) & \text{at } y &= l_2 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + \int_0^{l_1} \int_0^{l_2} [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 & - a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & - a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=l_2} d\xi d\tau \\
 & + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & \frac{4}{l_1 l_2} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\lambda_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \sin(\lambda_{nm} t), \\
 p_n = & \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b - \frac{1}{4}k^2}.
 \end{aligned}$$

5.4.1-5. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Second boundary value problem.

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(x, y) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(x, y) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_x w = g_1(y, t) \quad \text{at} \quad x = 0 \quad & \text{(boundary condition),} \\
 \partial_x w = g_2(y, t) \quad \text{at} \quad x = l_1 \quad & \text{(boundary condition),} \\
 \partial_y w = g_3(x, t) \quad \text{at} \quad y = 0 \quad & \text{(boundary condition),} \\
 \partial_y w = g_4(x, t) \quad \text{at} \quad y = l_2 \quad & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + \int_0^{l_1} \int_0^{l_2} [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(x, y, \xi, \eta, t) d\eta d\xi \\
 & - a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) G(x, y, 0, \eta, t - \tau) d\eta d\tau + a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau + a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau.
 \end{aligned}$$

Here,

$$G(x, y, \xi, \eta, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{\sin(\lambda_{00}t)}{l_1 l_2 \lambda_{00}} + \frac{2}{l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_{nm}}{\lambda_{nm}} \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) \sin(\lambda_{nm} t) \right],$$

where

$$p_n = \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b - \frac{1}{4}k^2}, \quad A_{nm} = \begin{cases} 0 & \text{for } n = m = 0, \\ 1 & \text{for } nm = 0 \ (n \neq m), \\ 2 & \text{for } nm \neq 0. \end{cases}$$

5.4.1-6. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Third boundary value problem.

A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - s_1 w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w + s_2 w &= g_2(y, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ \partial_y w - s_3 w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ \partial_y w + s_4 w &= g_4(x, t) & \text{at } y &= l_2 & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, y, t)$  is determined by the formula in Paragraph 5.4.1-5 where

$$G(x, y, \xi, \eta, t) = 4 \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{E_{nm} \sqrt{a^2 \mu_n^2 + a^2 \nu_m^2 + b - \frac{1}{4}k^2}} \sin(\mu_n x + \varepsilon_n) \sin(\nu_m y + \sigma_m) \\ \times \sin(\mu_n \xi + \varepsilon_n) \sin(\nu_m \eta + \sigma_m) \sin\left(t \sqrt{a^2 \mu_n^2 + a^2 \nu_m^2 + b - \frac{1}{4}k^2}\right).$$

Here,

$$\varepsilon_n = \arctan \frac{\mu_n}{l_1}, \quad \sigma_m = \arctan \frac{\nu_m}{l_2}, \quad E_{nm} = \left[ l_1 + \frac{(s_1 s_2 + \mu_n^2)(s_1 + s_2)}{(s_1^2 + \mu_n^2)(s_2^2 + \mu_n^2)} \right] \left[ l_2 + \frac{(s_3 s_4 + \nu_m^2)(s_3 + s_4)}{(s_3^2 + \nu_m^2)(s_4^2 + \nu_m^2)} \right];$$

the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\mu^2 - s_1 s_2 = (s_1 + s_2) \mu \cot(l_1 \mu), \quad \nu^2 - s_3 s_4 = (s_3 + s_4) \nu \cot(l_2 \nu).$$

5.4.1-7. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Mixed boundary value problems.

1°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x, y) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(y, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(y, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ \partial_y w &= g_3(x, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ \partial_y w &= g_4(x, t) & \text{at } y &= l_2 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + \int_0^{l_1} \int_0^{l_2} [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 & - a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=l_1} d\eta d\tau \\
 & - a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) G(x, y, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & \frac{2}{l_1 l_2} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{\lambda_{nm}} \sin(p_n x) \cos(q_m y) \sin(p_n \xi) \cos(q_m \eta) \sin(\lambda_{nm} t), \\
 p_n = & \frac{n\pi}{l_1}, \quad q_m = \frac{m\pi}{l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b - \frac{1}{4}k^2}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases}
 \end{aligned}$$

2°. A rectangle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(x, y) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(x, y) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 w = g_1(y, t) & \quad \text{at } x = 0 \quad (\text{boundary condition}), \\
 \partial_x w = g_2(y, t) & \quad \text{at } x = l_1 \quad (\text{boundary condition}), \\
 w = g_3(x, t) & \quad \text{at } y = 0 \quad (\text{boundary condition}), \\
 \partial_y w = g_4(x, t) & \quad \text{at } y = l_2 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f_0(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + \int_0^{l_1} \int_0^{l_2} [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(x, y, \xi, \eta, t) d\eta d\xi \\
 & + a^2 \int_0^t \int_0^{l_2} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta, t - \tau) \right]_{\xi=0} d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_2} g_2(\eta, \tau) G(x, y, l_1, \eta, t - \tau) d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & + a^2 \int_0^t \int_0^{l_1} g_4(\xi, \tau) G(x, y, \xi, l_2, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau,
 \end{aligned}$$

where

$$G(x, y, \xi, \eta, t) = \frac{4}{l_1 l_2} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{1}{\lambda_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \sin(\lambda_{nm} t),$$

$$p_n = \frac{\pi(2n+1)}{2l_1}, \quad q_m = \frac{\pi(2m+1)}{2l_2}, \quad \lambda_{nm} = \sqrt{a^2 p_n^2 + a^2 q_m^2 + b - \frac{1}{4}k^2}.$$

## 5.4.2. Problems in Polar Coordinates

A two-dimensional nonhomogeneous telegraph equation in the polar coordinate system has the form

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} \right) - bw + \Phi(r, \varphi, t), \quad r = \sqrt{x^2 + y^2}.$$

For one-dimensional solutions  $w = w(r, t)$ , see equation 4.4.2.2.

5.4.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &+ \int_0^{2\pi} \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &- a^2 R \int_0^t \int_0^{2\pi} g(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &+ \int_0^t \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{\pi R^2} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{[J'_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}},$$

$$\lambda_{nm} = a^2 \mu_{nm}^2 + b - \frac{1}{4}k^2, \quad A_0 = 1, \quad A_n = 2 \quad (n = 1, 2, \dots),$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

5.4.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A circle is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + \int_0^{2\pi} \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + a^2 R \int_0^t \int_0^{2\pi} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\
 & + \int_0^t \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \exp\left(-\frac{1}{2}kt\right) \left[ \frac{\sin\left(t\sqrt{b - k^2/4}\right)}{\pi R^2 \sqrt{b - k^2/4}} \right. \\
 & + \left. \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}} \right], \\
 \lambda_{nm} = & a^2 \mu_{nm}^2 + b - \frac{1}{4}k^2, \quad A_0 = 1, \quad A_n = 2 \quad (n = 1, 2, \dots),
 \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_m$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

5.4.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A circle is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_r w + sw = g(\varphi, t) \quad \text{at} \quad r = R \quad & \text{(boundary condition).}
 \end{aligned}$$

The solution  $w(r, \varphi, t)$  is determined by the formula in Paragraph 5.4.2-2 where

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \frac{1}{\pi} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \sin(t\sqrt{\lambda_{nm}})}{(\mu_{nm}^2 R^2 + s^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nm}}}, \\
 \lambda_{nm} = & a^2 \mu_{nm}^2 + b - \frac{1}{4}k^2, \quad A_0 = 1, \quad A_n = 2 \quad (n = 1, 2, \dots).
 \end{aligned}$$

Here, the  $J_n(\xi)$  are the Bessel functions and the  $\mu_m$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + s J_n(\mu R) = 0.$$

5.4.2-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 w = g_1(\varphi, t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
 w = g_2(\varphi, t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + \int_0^{2\pi} \int_{R_1}^{R_2} [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + a^2 R_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\
 & - a^2 R_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau \\
 & + \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \frac{\pi}{2} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_n B_{nm} Z_n(\mu_{nm}r) Z_n(\mu_{nm}\xi) \cos[n(\varphi - \eta)] \frac{\sin\left(t\sqrt{\lambda_{nm}}\right)}{\sqrt{\lambda_{nm}}}, \\
 A_n = & \begin{cases} 1/2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0, \end{cases} \quad B_{nm} = \frac{\mu_{nm}^2 J_n^2(\mu_{nm}R_2)}{J_n^2(\mu_{nm}R_1) - J_n^2(\mu_{nm}R_2)}, \\
 Z_n(\mu_{nm}r) = & J_n(\mu_{nm}R_1)Y_n(\mu_{nm}r) - Y_n(\mu_{nm}R_1)J_n(\mu_{nm}r), \quad \lambda_{nm} = a^2\mu_{nm}^2 + b - \frac{1}{4}k^2,
 \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1)Y_n(\mu R_2) - Y_n(\mu R_1)J_n(\mu R_2) = 0.$$

5.4.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, \varphi) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_r w = g_1(\varphi, t) & \quad \text{at } r = R_1 \quad (\text{boundary condition}), \\
 \partial_r w = g_2(\varphi, t) & \quad \text{at } r = R_2 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + \int_0^{2\pi} \int_{R_1}^{R_2} [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & - a^2 R_1 \int_0^t \int_0^{2\pi} g_1(\eta, \tau) G(r, \varphi, R_1, \eta, t - \tau) d\eta d\tau \\
 & + a^2 R_2 \int_0^t \int_0^{2\pi} g_2(\eta, \tau) G(r, \varphi, R_2, \eta, t - \tau) d\eta d\tau \\
 & + \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$



Here,

$$G(r, \varphi, \xi, \eta, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{\sin\left(t\sqrt{b-k^2/4}\right)}{\pi(R_2^2 - R_1^2)\sqrt{b-k^2/4}} + \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_n(\mu_{nm}r) Z_n(\mu_{nm}\xi) \cos[n(\varphi - \eta)] \sin\left(t\sqrt{a^2 \mu_{nm}^2 + b - k^2/4}\right)}{[(\mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm}R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm}R_1)] \sqrt{a^2 \mu_{nm}^2 + b - k^2/4}} \right],$$

where

$$Z_n(\mu_{nm}r) = J'_n(\mu_{nm}R_1)Y_n(\mu_{nm}r) - Y'_n(\mu_{nm}R_1)J_n(\mu_{nm}r), \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1)Y'_n(\mu R_2) - Y'_n(\mu R_1)J'_n(\mu R_2) = 0.$$

5.4.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

An annular domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w - s_1 w &= g_1(\varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w + s_2 w &= g_2(\varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, t)$  is determined by the formula in Paragraph 5.4.2-5 where

$$G(r, \varphi, \xi, \eta, t) = \frac{1}{\pi} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2}{B_{nm} \lambda_{nm}} Z_n(\mu_{nm}r) Z_n(\mu_{nm}\xi) \cos[n(\varphi - \eta)] \sin(\lambda_{nm}t).$$

Here,

$$\begin{aligned} A_n &= \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \quad \lambda_{nm} = \sqrt{a^2 \mu_{nm}^2 + b - \frac{1}{4}k^2}, \\ B_{nm} &= (s_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_n^2(\mu_{nm}R_2) - (s_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_n^2(\mu_{nm}R_1), \\ Z_n(\mu_{nm}r) &= [\mu_{nm} J'_n(\mu_{nm}R_1) - s_1 J_n(\mu_{nm}R_1)] Y_n(\mu_{nm}r) \\ &\quad - [\mu_{nm} Y'_n(\mu_{nm}R_1) - s_1 Y_n(\mu_{nm}R_1)] J_n(\mu_{nm}r), \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned} &[\mu J'_n(\mu R_1) - s_1 J_n(\mu R_1)] [\mu Y'_n(\mu R_2) + s_2 Y_n(\mu R_2)] \\ &= [\mu Y'_n(\mu R_1) - s_1 Y_n(\mu R_1)] [\mu J'_n(\mu R_2) + s_2 J_n(\mu R_2)]. \end{aligned}$$

5.4.2-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . First boundary value problem.

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + \int_0^{\varphi_0} \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & - a^2 R \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\
 & + a^2 \int_0^t \int_0^R g_2(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\
 & - a^2 \int_0^t \int_0^R g_3(\xi, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\tau \\
 & + \int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, \xi, \eta, t) = & \frac{4}{R^2 \varphi_0} \exp(-\tfrac{1}{2} k t) \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \\
 & \times \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \frac{\sin\left(t\sqrt{a^2\mu_{nm}^2 + b - k^2/4}\right)}{\sqrt{a^2\mu_{nm}^2 + b - k^2/4}},
 \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

5.4.2-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ . Second boundary value problem.

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi) \quad & \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, \varphi) \quad & \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_r w = g_1(\varphi, t) \quad & \text{at} \quad r = R \quad (\text{boundary condition}), \\
 r^{-1} \partial_\varphi w = g_2(r, t) \quad & \text{at} \quad \varphi = 0 \quad (\text{boundary condition}), \\
 r^{-1} \partial_\varphi w = g_3(r, t) \quad & \text{at} \quad \varphi = \varphi_0 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + \int_0^{\varphi_0} \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\
 & + a^2 R \int_0^t \int_0^{\varphi_0} g_1(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_0^R g_2(\xi, \tau) G(r, \varphi, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^R g_3(\xi, \tau) G(r, \varphi, \xi, \varphi_0, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \exp\left(-\frac{1}{2}kt\right) \left[ \frac{2 \sin(t\sqrt{b-k^2/4})}{R^2 \varphi_0 \sqrt{b-k^2/4}} + 4\varphi_0 \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_{nm}^2 J_{n\pi/\varphi_0}(\mu_{nm}r) J_{n\pi/\varphi_0}(\mu_{nm}\xi)}{(R^2 \varphi_0^2 \mu_{nm}^2 - n^2 \pi^2) J_{n\pi/\varphi_0}^2(\mu_{nm}R)} \right. \\ \left. \times \cos\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{n\pi\eta}{\varphi_0}\right) \frac{\sin(t\sqrt{a^2 \mu_{nm}^2 + b - k^2/4})}{\sqrt{a^2 \mu_{nm}^2 + b - k^2/4}} \right],$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_{n\pi/\varphi_0}(\mu R) = 0$ .

#### 5.4.2-9. Domain: $0 \leq r \leq R$ , $0 \leq \varphi \leq \varphi_0$ . Mixed boundary value problem.

A circular sector is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_r w + \beta w &= g(\varphi, t) & \text{at } r &= R & (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi &= 0 & (\text{boundary condition}), \\ \partial_\varphi w &= 0 & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta) G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &+ \int_0^{\varphi_0} \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] G(r, \varphi, \xi, \eta, t) \xi d\xi d\eta \\ &+ a^2 R \int_0^t \int_0^{\varphi_0} g(\eta, \tau) G(r, \varphi, R, \eta, t - \tau) d\eta d\tau \\ &+ \int_0^t \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \tau) G(r, \varphi, \xi, \eta, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, \xi, \eta, t) = \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} A_{nm} J_{s_n}(\mu_{nm}r) J_{s_n}(\mu_{nm}\xi) \cos(s_n\varphi) \cos(s_n\eta) \sin(\lambda_{nm}t),$$

$$s_n = \frac{n\pi}{\varphi_0}, \quad A_{nm} = \frac{4\mu_{nm}^2}{\varphi_0(\mu_{nm}^2 R^2 + \beta^2 R^2 - s_n^2) [J_{s_n}(\mu_{nm}R)]^2 \lambda_{nm}}, \quad \lambda_{nm} = \sqrt{a^2 \mu_{nm}^2 + b - \frac{1}{4}k^2},$$

where the  $J_{s_n}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_{s_n}(\mu R) + \beta J_{s_n}(\mu R) = 0.$$

### 5.4.3. Axisymmetric Problems

In the axisymmetric case, a nonhomogeneous telegraph equation in the cylindrical coordinate system has the form

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} \right) - bw + \Phi(r, z, t), \quad r = \sqrt{x^2 + y^2}.$$

In the solutions of the problems considered below, the modified Green's function  $\mathcal{G}(r, z, \xi, \eta, t) = 2\pi\xi G(r, z, \xi, \eta, t)$  is used for convenience.

**5.4.3-1. Domain:  $0 \leq r \leq R, 0 \leq z \leq l$ . First boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ \int_0^l \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &- a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &+ a^2 \int_0^t \int_0^R g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- a^2 \int_0^t \int_0^R g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau \\ &+ \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} \mathcal{G}(r, z, \xi, \eta, t) &= \frac{4\xi e^{-kt/2}}{R^2 l} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}}, \\ \lambda_{nm} &= \sqrt{\frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b - \frac{k^2}{4}}, \end{aligned}$$

where the  $\mu_n$  are positive zeros of the Bessel function,  $J_0(\mu) = 0$ .

**5.4.3-2. Domain:  $0 \leq r \leq R, 0 \leq z \leq l$ . Second boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + \int_0^l \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\
 & + a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R, \eta, t - \tau) d\eta d\tau \\
 & - a^2 \int_0^t \int_0^R g_2(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau \\
 & + a^2 \int_0^t \int_0^R g_3(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau \\
 & + \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 \mathcal{G}(r, z, \xi, \eta, t) = 2\xi \exp(-\tfrac{1}{2}kt) & \left[ \frac{\sin(t\sqrt{c})}{R^2 l \sqrt{c}} + \frac{1}{R^2 l} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_{nm}}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \right. \\
 & \left. \times \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}} \right],
 \end{aligned}$$

where

$$c = b - \frac{k^2}{4}, \quad \lambda_{nm} = \frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b - \frac{k^2}{4}, \quad A_{nm} = \begin{cases} 0 & \text{for } m = 0, n = 0, \\ 1 & \text{for } m = 0, n > 0, \\ 2 & \text{for } m > 0, \end{cases}$$

and the  $\mu_n$  are zeros of the first-order Bessel function,  $J_1(\mu) = 0$  ( $\mu_0 = 0$ ).

5.4.3-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq z \leq l$ . Third boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(r, z) \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w &= f_1(r, z) \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_r w + s_1 w &= g_1(z, t) \quad \text{at } r = R \quad (\text{boundary condition}), \\
 \partial_z w - s_2 w &= g_2(r, t) \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
 \partial_z w + s_3 w &= g_3(r, t) \quad \text{at } z = l \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(r, z, t)$  is determined by the formula in Paragraph 5.4.3-2 where

$$\mathcal{G}(r, z, \xi, \eta, t) = \frac{2\xi}{R^2} \exp(-\tfrac{1}{2}kt) \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} A_n J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \frac{\varphi_m(z) \varphi_m(\eta)}{\|\varphi_m\|^2} \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}}.$$

Here,

$$\begin{aligned}
 A_n &= \frac{\mu_n^2}{(s_1^2 R^2 + \mu_n^2) J_0^2(\mu_n)}, \quad \lambda_{nm} = \frac{a^2 \mu_n^2}{R^2} + a^2 \beta_m^2 + b - \frac{k^2}{4}, \\
 \varphi_m(z) &= \cos(\beta_m z) + \frac{s_2}{\beta_m} \sin(\beta_m z), \quad \|\varphi_m\|^2 = \frac{s_3}{2\beta_m^2} \frac{\beta_m^2 + s_2^2}{\beta_m^2 + s_3^2} + \frac{s_2}{2\beta_m^2} + \frac{l}{2} \left(1 + \frac{s_2^2}{\beta_m^2}\right);
 \end{aligned}$$

the  $\mu_n$  and  $\beta_m$  are positive roots of the transcendental equations

$$\mu J_1(\mu) - s_1 R J_0(\mu) = 0, \quad \frac{\tan(\beta l)}{\beta} = \frac{s_2 + s_3}{\beta^2 - s_2 s_3}.$$

**5.4.3-4. Domain:  $0 \leq r \leq R, 0 \leq z \leq l$ . Mixed boundary value problems.**

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ \int_0^l \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &- a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R} d\eta d\tau \\ &- a^2 \int_0^t \int_0^R g_2(\xi, \tau) \mathcal{G}(r, z, \xi, 0, t - \tau) d\xi d\tau + a^2 \int_0^t \int_0^R g_3(\xi, \tau) \mathcal{G}(r, z, \xi, l, t - \tau) d\xi d\tau \\ &+ \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} \mathcal{G}(r, z, \xi, \eta, t) &= \frac{2\xi e^{-kt/2}}{R^2 l} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{J_1^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \cos\left(\frac{m\pi z}{l}\right) \cos\left(\frac{m\pi \eta}{l}\right) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}}, \\ \lambda_{nm} &= \sqrt{\frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b - \frac{k^2}{4}}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 0, \end{cases} \end{aligned}$$

where the  $\mu_n$  are zeros of the Bessel function,  $J_0(\mu) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^R f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ \int_0^l \int_0^R [f_1(\xi, \eta) + k f_0(\xi, \eta)] \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta + a^2 \int_0^t \int_0^l g_1(\eta, \tau) \mathcal{G}(r, z, R, \eta, t - \tau) d\eta d\tau \\ &+ a^2 \int_0^t \int_0^R g_2(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- a^2 \int_0^t \int_0^R g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau \\ &+ \int_0^t \int_0^l \int_0^R \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\mathcal{G}(r, z, \xi, \eta, t) = \frac{4\xi e^{-kt/2}}{R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{1}{J_0^2(\mu_n)} J_0\left(\frac{\mu_n r}{R}\right) J_0\left(\frac{\mu_n \xi}{R}\right) \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}},$$

$$\lambda_{nm} = \sqrt{\frac{a^2 \mu_n^2}{R^2} + \frac{a^2 \pi^2 m^2}{l^2} + b - \frac{k^2}{4}},$$

where the  $\mu_n$  are zeros of the first-order Bessel function,  $J_1(\mu) = 0$  ( $\mu_0 = 0$ ).

5.4.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq z \leq l$ . First boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(z, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ w &= g_2(z, t) & \text{at } r &= R_2 & \text{(boundary condition),} \\ w &= g_3(r, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_4(r, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_{R_1}^{R_2} f_0(\xi, \eta) \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ \int_0^l \int_{R_1}^{R_2} [f_1(\xi, \eta) + k f_0(\xi, \eta)] \mathcal{G}(r, z, \xi, \eta, t) d\xi d\eta \\ &+ a^2 \int_0^t \int_0^l g_1(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_1} d\eta d\tau \\ &- a^2 \int_0^t \int_0^l g_2(\eta, \tau) \left[ \frac{\partial}{\partial \xi} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\xi=R_2} d\eta d\tau \\ &+ a^2 \int_0^t \int_{R_1}^{R_2} g_3(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=0} d\xi d\tau \\ &- a^2 \int_0^t \int_{R_1}^{R_2} g_4(\xi, \tau) \left[ \frac{\partial}{\partial \eta} \mathcal{G}(r, z, \xi, \eta, t - \tau) \right]_{\eta=l} d\xi d\tau \\ &+ \int_0^t \int_0^l \int_{R_1}^{R_2} \Phi(\xi, \eta, \tau) \mathcal{G}(r, z, \xi, \eta, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\mathcal{G}(r, z, \xi, \eta, t) = \frac{\pi^2 \xi}{R_1^2 l} e^{-kt/2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\mu_n^2 J_0^2(s\mu_n) \Psi_n(r) \Psi_n(\xi)}{J_0^2(\mu_n) - J_0^2(s\mu_n)} \sin\left(\frac{m\pi z}{l}\right) \sin\left(\frac{m\pi \eta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nm}})}{\sqrt{\lambda_{nm}}},$$

$$\Psi_n(r) = Y_0(\mu_n) J_0\left(\frac{\mu_n r}{R_1}\right) - J_0(\mu_n) Y_0\left(\frac{\mu_n r}{R_1}\right), \quad s = \frac{R_2}{R_1}, \quad \lambda_{nm} = \frac{a^2 \mu_n^2}{R_1^2} + \frac{a^2 \pi^2 m^2}{l^2} + b - \frac{k^2}{4},$$

where  $J_0(\mu)$  and  $Y_0(\mu)$  are the Bessel functions, and the  $\mu_n$  are positive roots of the transcendental equation

$$J_0(\mu) Y_0(s\mu) - J_0(s\mu) Y_0(\mu) = 0.$$

## 5.5. Other Equations with Two Space Variables

$$1. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) + b_1 \frac{\partial w}{\partial x} + b_2 \frac{\partial w}{\partial y} + cw.$$

The transformation

$$w(x, y, t) = u(x, y, \tau) \exp \left( -\frac{1}{2}kt - \frac{b_1x + b_2y}{2a^2} \right), \quad \tau = at$$

leads to the equation from Subsection 5.1.3:

$$\frac{\partial^2 u}{\partial \tau^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \beta u, \quad \beta = \frac{c}{a^2} + \frac{k^2}{4a^2} - \frac{1}{4a^4}(b_1^2 + b_2^2).$$

$$2. \quad t^m \frac{\partial^2 w}{\partial t^2} + \frac{m}{2} t^{m-1} \frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2}.$$

Domain:  $-\infty < x < \infty, -\infty < y < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x, y) \quad \text{at} \quad t = 0, \\ t^{m/2} \partial_t w &= g(x, y) \quad \text{at} \quad t = 0. \end{aligned}$$

Solution for  $1 \leq m < 2$ :

$$\begin{aligned} w(x, y, t) &= \frac{1}{2\pi} t^{m/2} \frac{\partial}{\partial t} \iint_{C_t} \frac{f(\xi, \eta) d\xi d\eta}{\sqrt{k_m^2 t^{2-m} - \rho^2}} + \frac{1}{2\pi} \iint_{C_t} \frac{g(\xi, \eta) d\xi d\eta}{\sqrt{k_m^2 t^{2-m} - \rho^2}}, \\ k_m &= \frac{2}{2-m}, \quad \rho = \sqrt{(x-\xi)^2 + (y-\eta)^2}, \end{aligned}$$

where  $C_t = \{\rho^2 \leq k_m^2 t^{2-m}\}$  is the circle with center at  $(x, y)$  and radius  $k_m t^{1/k_m}$ .

⊙ Reference: M. M. Smirnov (1975).



## Chapter 6

# Hyperbolic Equations with Three or More Space Variables

---

### 6.1. Wave Equation $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_3 w$

#### 6.1.1. Problems in Cartesian Coordinates

The wave equation with three space variables in the rectangular Cartesian coordinate system has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right).$$

This equation is of fundamental importance in sound propagation theory, the propagation of electromagnetic fields theory, and a number of other areas of physics and mechanics.

##### 6.1.1-1. Particular solutions and their properties.

1°. Particular solutions:

$$w(x, y, z, t) = A \exp \left( k_1 x + k_2 y + k_3 z \pm at \sqrt{k_1^2 + k_2^2 + k_3^2} \right),$$

$$w(x, y, z, t) = A \sin(k_1 x + C_1) \sin(k_2 y + C_2) \sin(k_3 z + C_3) \sin \left( at \sqrt{k_1^2 + k_2^2 + k_3^2} \right),$$

$$w(x, y, z, t) = A \sin(k_1 x + C_1) \sin(k_2 y + C_2) \sin(k_3 z + C_3) \cos \left( at \sqrt{k_1^2 + k_2^2 + k_3^2} \right),$$

$$w(x, y, z, t) = A \sinh(k_1 x + C_1) \sinh(k_2 y + C_2) \sinh(k_3 z + C_3) \sinh \left( at \sqrt{k_1^2 + k_2^2 + k_3^2} \right),$$

$$w(x, y, z, t) = A \sinh(k_1 x + C_1) \sinh(k_2 y + C_2) \sinh(k_3 z + C_3) \cosh \left( at \sqrt{k_1^2 + k_2^2 + k_3^2} \right),$$

where  $A$ ,  $C_1$ ,  $C_2$ ,  $C_3$ ,  $k_1$ ,  $k_2$ , and  $k_3$  are arbitrary constants.

2°. Fundamental solution:

$$\mathcal{E}(x, y, z, t) = \frac{1}{2\pi a} \delta(a^2 t^2 - r^2), \quad r = \sqrt{x^2 + y^2 + z^2},$$

where  $\delta(\xi)$  is the Dirac delta function.

⊙ *Reference:* V. S. Vladimirov (1988).

3°. Infinite series solutions containing arbitrary functions of space variables:

$$w(x, y, z, t) = f(x, y, z) + \sum_{n=1}^{\infty} \frac{(at)^{2n}}{(2n)!} \Delta^n f(x, y, z), \quad \Delta \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2},$$

$$w(x, y, z, t) = tg(x, y, z) + t \sum_{n=1}^{\infty} \frac{(at)^{2n}}{(2n+1)!} \Delta^n g(x, y, z),$$

where  $f(x, y, z)$  and  $g(x, y, z)$  are any infinitely differentiable functions. The first solution satisfies the initial conditions  $w(x, y, z, 0) = f(x, y, z)$ ,  $\partial_t w(x, y, z, 0) = 0$ , and the second solution initial conditions  $w(x, y, z, 0) = 0$ ,  $\partial_t w(x, y, z, 0) = g(x, y, z)$ . The sums are finite if  $f(x, y, z)$  and  $g(x, y, z)$  are polynomials in  $x, y, z$ .

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

4°. Suppose  $w = w(x, y, z, t)$  is a solution of the wave equation. Then the functions

$$\begin{aligned} w_1 &= Aw(\pm\lambda x + C_1, \pm\lambda y + C_2, \pm\lambda z + C_3, \pm\lambda t + C_4), \\ w_2 &= Aw\left(\frac{x - vt}{\sqrt{1 - (v/a)^2}}, y, z, \frac{t - va^{-2}x}{\sqrt{1 - (v/a)^2}}\right), \\ w_3 &= \frac{A}{r^2 - a^2t^2}w\left(\frac{x}{r^2 - a^2t^2}, \frac{y}{r^2 - a^2t^2}, \frac{z}{r^2 - a^2t^2}, \frac{t}{r^2 - a^2t^2}\right), \end{aligned}$$

where  $A, C_n, v$ , and  $\lambda$  are arbitrary constants, are also solutions of the equation. The signs at  $\lambda$  in the expression of  $w_1$  can be taken independently of one another. The function  $w_2$  is a consequence of the invariance of the wave equation under the Lorentz transformation.

⊙ *References:* G. N. Polozhii (1964), W. Miller, Jr. (1977), A. V. Bitsadze and D. F. Kalinichenko (1985).

6.1.1-2. Domain:  $-\infty < x < \infty, -\infty < y < \infty, -\infty < z < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(x, y, z) \quad \text{at} \quad t = 0. \end{aligned}$$

Solution (Kirchhoff's formula):

$$\begin{aligned} w(x, y, z, t) &= \frac{1}{4\pi a} \frac{\partial}{\partial t} \iint_{S_{at}} \frac{f(\xi, \eta, \zeta)}{r} dS + \frac{1}{4\pi a} \iint_{S_{at}} \frac{g(\xi, \eta, \zeta)}{r} dS, \\ r &= \sqrt{(\xi - x)^2 + (\eta - y)^2 + (\zeta - z)^2}, \end{aligned}$$

where the integration is performed over the surface of the sphere of radius  $at$  with center at  $(x, y, z)$ .

⊙ *References:* N. S. Koshlyakov, E. B. Glizer, and M. M. Smirnov (1970), A. N. Tikhonov and A. A. Samarskii (1990).

6.1.1-3. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2, 0 \leq z \leq l_3$ . First boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\ w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_6(x, y, t) \quad \text{at} \quad z = l_3 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
w(x, y, z, t) = & \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
& + \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
& + a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\
& - a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l_3} d\xi d\eta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(x, y, z, \xi, \eta, \zeta, t) = & \frac{8}{al_1l_2l_3} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{\lambda_{nmk}} \sin(\alpha_n x) \sin(\beta_m y) \sin(\gamma_k z) \\
& \times \sin(\alpha_n \xi) \sin(\beta_m \eta) \sin(\gamma_k \zeta) \sin(a\lambda_{nmk} t),
\end{aligned}$$

where

$$\begin{aligned}
\alpha_n &= \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_k = \frac{k\pi}{l_3}, \\
\lambda_{nmk} &= \sqrt{\alpha_n^2 + \beta_m^2 + \gamma_k^2}.
\end{aligned}$$

6.1.1-4. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Second boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned}
w &= f_0(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
\partial_t w &= f_1(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
\partial_x w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\
\partial_x w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\
\partial_y w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\
\partial_y w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\
\partial_z w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
\partial_z w &= g_6(x, y, t) \quad \text{at} \quad z = l_3 \quad (\text{boundary condition}).
\end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) = & \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & - a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & - a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & + a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\
 & - a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta, t) = & \frac{t}{l_1 l_2 l_3} + \frac{1}{a l_1 l_2 l_3} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_m A_k}{\lambda_{nmk}} \cos(\alpha_n x) \cos(\beta_m y) \cos(\gamma_k z) \\
 & \times \cos(\alpha_n \xi) \cos(\beta_m \eta) \cos(\gamma_k \zeta) \sin(a \lambda_{nmk} t), \\
 \alpha_n = & \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_k = \frac{k\pi}{l_3}, \quad \lambda_{nmk} = \sqrt{\alpha_n^2 + \beta_m^2 + \gamma_k^2}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0. \end{cases}
 \end{aligned}$$

The summation here is performed over the indices satisfying the condition  $n + m + k > 0$ ; the term corresponding to  $n = m = k = 0$  is singled out.

6.1.1-5. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Third boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(x, y, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(x, y, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_x w - s_1 w = g_1(y, z, t) & \quad \text{at } x = 0 \quad (\text{boundary condition}), \\
 \partial_x w + s_2 w = g_2(y, z, t) & \quad \text{at } x = l_1 \quad (\text{boundary condition}), \\
 \partial_y w - s_3 w = g_3(x, z, t) & \quad \text{at } y = 0 \quad (\text{boundary condition}), \\
 \partial_y w + s_4 w = g_4(x, z, t) & \quad \text{at } y = l_2 \quad (\text{boundary condition}), \\
 \partial_z w - s_5 w = g_5(x, y, t) & \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
 \partial_z w + s_6 w = g_6(x, y, t) & \quad \text{at } z = l_3 \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 6.1.1-4 where

$$\begin{aligned}
 G(x, y, \xi, \eta, t) = & \frac{8}{a} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{E_{nmk} \sqrt{\alpha_n^2 + \beta_m^2 + \gamma_k^2}} \sin(\alpha_n x + \varepsilon_n) \sin(\beta_m y + \sigma_m) \sin(\gamma_k z + \nu_k) \\
 & \times \sin(\alpha_n \xi + \varepsilon_n) \sin(\beta_m \eta + \sigma_m) \sin(\gamma_k \zeta + \nu_k) \sin\left(at \sqrt{\alpha_n^2 + \beta_m^2 + \gamma_k^2}\right)
 \end{aligned}$$

with

$$\varepsilon_n = \arctan \frac{\alpha_n}{l_1}, \quad \sigma_m = \arctan \frac{\beta_m}{l_2}, \quad \nu_k = \arctan \frac{\gamma_k}{l_3},$$

$$E_{nmk} = \left[ l_1 + \frac{(s_1 s_2 + \alpha_n^2)(s_1 + s_2)}{(s_1^2 + \alpha_n^2)(s_2^2 + \alpha_n^2)} \right] \left[ l_2 + \frac{(s_3 s_4 + \beta_m^2)(s_3 + s_4)}{(s_3^2 + \beta_m^2)(s_4^2 + \beta_m^2)} \right] \left[ l_3 + \frac{(s_5 s_6 + \gamma_k^2)(s_5 + s_6)}{(s_5^2 + \gamma_k^2)(s_6^2 + \gamma_k^2)} \right].$$

Here, the  $\alpha_n$ ,  $\beta_m$ , and  $\gamma_k$  are positive roots of the transcendental equations

$$\alpha^2 - s_1 s_2 = (s_1 + s_2) \alpha \cot(l_1 \alpha), \quad \beta^2 - s_3 s_4 = (s_3 + s_4) \beta \cot(l_2 \beta), \quad \gamma^2 - s_5 s_6 = (s_5 + s_6) \gamma \cot(l_3 \gamma).$$

6.1.1-6. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Mixed boundary value problems.

1°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{2}{al_1 l_2 l_3} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{A_m A_k}{\lambda_{nmk}} \sin(\alpha_n x) \cos(\beta_m y) \cos(\gamma_k z) \\ &\quad \times \sin(\alpha_n \xi) \cos(\beta_m \eta) \cos(\gamma_k \zeta) \sin(a \lambda_{nmk} t), \end{aligned}$$

where

$$\alpha_n = \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_k = \frac{k\pi}{l_3},$$

$$\lambda_{nmk} = \sqrt{\alpha_n^2 + \beta_m^2 + \gamma_k^2}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 0. \end{cases}$$

2°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{8}{al_1 l_2 l_3} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{\lambda_{nmk}} \sin(\alpha_n x) \sin(\beta_m y) \sin(\gamma_k z) \\ &\quad \times \sin(\alpha_n \xi) \sin(\beta_m \eta) \sin(\gamma_k \zeta) \sin(a \lambda_{nmk} t), \end{aligned}$$

where

$$\alpha_n = \frac{\pi(2n+1)}{2l_1}, \quad \beta_m = \frac{\pi(2m+1)}{2l_2}, \quad \gamma_k = \frac{\pi(2k+1)}{2l_3}, \quad \lambda_{nmk} = \sqrt{\alpha_n^2 + \beta_m^2 + \gamma_k^2}.$$

## 6.1.2. Problems in Cylindrical Coordinates

The three-dimensional wave equation in the cylindrical coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left[ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} \right], \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional problems with axial symmetry that have solutions  $w = w(r, t)$  are considered in Subsection 4.2.1. Two-dimensional problems whose solutions have the form  $w = w(r, \varphi, t)$  or  $w = w(r, z, t)$  are discussed in Subsections 5.1.2 and 5.1.3.

6.1.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{2}{\pi a R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nmk}}} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\ &\quad \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \sin(at \sqrt{\lambda_{nmk}}), \\ \lambda_{nmk} &= \mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

6.1.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + a^2 R \int_0^t \int_0^{2\pi} \int_0^R g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & - a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\
 & + a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{t}{\pi R^2 l} + \frac{2}{\pi^2 a R^2} \sum_{k=1}^{\infty} \frac{1}{k} \cos\left(\frac{k\pi x}{l}\right) \cos\left(\frac{k\pi \xi}{l}\right) \sin\left(\frac{ak\pi t}{l}\right) \\
 & + \frac{1}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi x}{l}\right) \cos\left(\frac{k\pi \xi}{l}\right) \frac{\sin(\lambda_{nmk} t)}{\lambda_{nmk}}, \\
 & \lambda_{nmk} = a \sqrt{\mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}
 \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

6.1.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, \varphi, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_r w + k_1 w = g(\varphi, z, t) & \quad \text{at } r = R \quad (\text{boundary condition}), \\
 \partial_z w - k_2 w = g_2(r, \varphi, t) & \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
 \partial_z w + k_3 w = g_3(r, \varphi, t) & \quad \text{at } z = l \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 6.1.2-2 where

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] h_s(z) h_s(\zeta) \sin(\lambda_{nms} t)}{(\mu_{nm}^2 R^2 + k_1^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2 \|h_s\|^2 \lambda_{nms}}, \\
 \lambda_{nms} = & a \sqrt{\mu_{nm}^2 + \beta_s^2}, \quad h_s(z) = \cos(\beta_s z) + \frac{k_2}{\beta_s} \sin(\beta_s z), \quad \|h_s\|^2 = \frac{k_3}{2\beta_s^2} \frac{\beta_s^2 + k_2^2}{\beta_s^2 + k_3^2} + \frac{k_2}{2\beta_s^2} + \frac{l}{2} \left(1 + \frac{k_2^2}{\beta_s^2}\right).
 \end{aligned}$$

Here,  $A_0 = 1$  and  $A_n = 2$  for  $n = 1, 2, \dots$ ; the  $J_n(\xi)$  are the Bessel functions; and the  $\mu_{nm}$  and  $\beta_s$  are positive roots of the transcendental equations

$$\mu J'_n(\mu R) + k_1 J_n(\mu R) = 0, \quad \frac{\tan(\beta l)}{\beta} = \frac{k_2 + k_3}{\beta^2 - k_2 k_3}.$$



1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{\pi a R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k}{[J'_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nmk}}} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\ &\quad \times \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi \zeta}{l}\right) \sin(at \sqrt{\lambda_{nmk}}), \\ \lambda_{nmk} &= \mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{2}{\pi^2 a R^2} \sum_{k=1}^{\infty} \frac{1}{k} \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \sin\left(\frac{k\pi at}{l}\right) \\
 & + \frac{2}{\pi al} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n \mu_{nm}^2}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nmk}}} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\
 & \quad \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \sin(at \sqrt{\lambda_{nmk}}), \\
 \lambda_{nmk} = & \mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}
 \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

6.1.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi, z) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, \varphi, z) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
 w = g_1(\varphi, z, t) \quad & \text{at } r = R_1 \quad (\text{boundary condition}), \\
 w = g_2(\varphi, z, t) \quad & \text{at } r = R_2 \quad (\text{boundary condition}), \\
 w = g_3(r, \varphi, t) \quad & \text{at } z = 0 \quad (\text{boundary condition}), \\
 w = g_4(r, \varphi, t) \quad & \text{at } z = l \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & + \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 & + a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\
 & - a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\
 & + a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\
 & - a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{\pi}{2l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)} Z_{nm}(r) Z_{nm}(\xi) \\
 & \quad \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \frac{\sin(at \sqrt{\lambda_{nmk}})}{a \sqrt{\lambda_{nmk}}}, \\
 A_n = & \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nmk} = \mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}, \\
 Z_{nm}(r) = & J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),
 \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1)Y_n(\mu R_2) - Y_n(\mu R_1)J_n(\mu R_2) = 0.$$

6.1.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r &= R_1 & \text{(boundary condition),} \\ \partial_r w &= g_2(\varphi, z, t) & \text{at } r &= R_2 & \text{(boundary condition),} \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{t}{\pi(R_2^2 - R_1^2)l} + \frac{2}{\pi^2 a(R_2^2 - R_1^2)} \sum_{k=1}^{\infty} \frac{1}{k} \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi\zeta}{l}\right) \sin\left(\frac{k\pi at}{l}\right) \\ &+ \frac{1}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k \mu_{nm}^2 Z_{nm}(r) Z_{nm}(\xi)}{(\mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)} \\ &\quad \times \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi\zeta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nmk}})}{a\sqrt{\lambda_{nmk}}}, \end{aligned}$$

where

$$\begin{aligned} A_n &= \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nmk} = \mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}, \\ Z_{nm}(r) &= J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r); \end{aligned}$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

**6.1.2-7. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.**

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(\varphi, z, t) & \text{at } r &= R_1 & (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(\varphi, z, t) & \text{at } r &= R_2 & (\text{boundary condition}), \\ \partial_z w - k_3 w &= g_3(r, \varphi, t) & \text{at } z &= 0 & (\text{boundary condition}), \\ \partial_z w + k_4 w &= g_4(r, \varphi, t) & \text{at } z &= l & (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 6.1.2-6 where

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{\pi a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2}{\|h_s\|^2 \sqrt{\mu_{nm}^2 + \lambda_s^2}} \\ &\times \frac{Z_{nm}(r) Z_{nm}(\xi) \cos[n(\varphi - \eta)] h_s(z) h_s(\zeta) \sin(at \sqrt{\mu_{nm}^2 + \lambda_s^2})}{(k_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)}. \end{aligned}$$

Here,

$$\begin{aligned} A_n &= \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad Z_{nm}(r) = [\mu_{nm} J'_n(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) \\ &\quad - [\mu_{nm} Y'_n(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r), \\ h_s(z) &= \cos(\lambda_s z) + \frac{k_3}{\lambda_s} \sin(\lambda_s z), \quad \|h_s\|^2 = \frac{k_4}{2\lambda_s^2} \frac{\lambda_s^2 + k_3^2}{\lambda_s^2 + k_4^2} + \frac{k_3}{2\lambda_s^2} + \frac{l}{2} \left( 1 + \frac{k_3^2}{\lambda_s^2} \right), \end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions; the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\begin{aligned} [\mu J'_n(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y'_n(\mu R_2) + k_2 Y_n(\mu R_2)] \\ = [\mu Y'_n(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J'_n(\mu R_2) + k_2 J_n(\mu R_2)]; \end{aligned}$$

and the  $\lambda_s$  are positive roots of the transcendental equation  $\frac{\tan(\lambda l)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}$ .

**6.1.2-8. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.**

1°. A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r &= R_1 & (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) & \text{at } r &= R_2 & (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z &= 0 & (\text{boundary condition}), \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\
& - a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
& + a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{\pi}{4l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k \mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)} Z_{nm}(r) Z_{nm}(\xi) \\
& \times \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi \zeta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nmk}})}{a\sqrt{\lambda_{nmk}}}, \\
A_n = & \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nmk} = \mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}, \\
Z_{nm}(r) = & J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),
\end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

2°. A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(r, \varphi, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_t w = f_1(r, \varphi, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_r w = g_1(\varphi, z, t) & \quad \text{at } r = R_1 \quad (\text{boundary condition}), \\
\partial_r w = g_2(\varphi, z, t) & \quad \text{at } r = R_2 \quad (\text{boundary condition}), \\
w = g_3(r, \varphi, t) & \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
w = g_4(r, \varphi, t) & \quad \text{at } z = l \quad (\text{boundary condition}).
\end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& - a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& + a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\
& - a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau.
\end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{2}{\pi^2 a(R_2^2 - R_1^2)} \sum_{k=1}^{\infty} \frac{1}{k} \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \sin\left(\frac{k\pi at}{l}\right) \\ + \frac{2}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_{nm}(r) Z_{nm}(\xi)}{(\mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)} \\ \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \frac{\sin(at\sqrt{\lambda_{nmk}})}{a\sqrt{\lambda_{nmk}}},$$

where

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nmk} = \mu_{nm}^2 + \frac{k^2 \pi^2}{l^2}, \\ Z_{nm}(r) = J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r);$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

6.1.2-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . First boundary value problem.

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(\varphi, z, t) & \text{at } r &= R & \text{(boundary condition),} \\ w &= g_2(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\ w &= g_4(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_5(r, \varphi, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{8}{R^2 l \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \\ \times \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi\zeta}{l}\right) \frac{\sin(at\sqrt{\mu_{nm}^2 + k^2\pi^2/l^2})}{a\sqrt{\mu_{nm}^2 + k^2\pi^2/l^2}},$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

6.1.2-10. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . Mixed boundary value problem.

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(\varphi, z, t) & \text{at } r &= R & \text{(boundary condition),} \\ w &= g_2(r, z, t) & \text{at } \varphi &= 0 & \text{(boundary condition),} \\ w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & \text{(boundary condition),} \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w &= g_5(r, \varphi, t) & \text{at } z &= l & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_5(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{4}{R^2 l \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_k J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \\ \times \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi\zeta}{l}\right) \frac{\sin(at\sqrt{\mu_{nm}^2 + k^2\pi^2/l^2})}{a\sqrt{\mu_{nm}^2 + k^2\pi^2/l^2}},$$

where  $A_0 = 1$  and  $A_k = 2$  for  $k \geq 1$ ; the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

### 6.1.3. Problems in Spherical Coordinates

The three-dimensional wave equation in the spherical coordinate system is represented as

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} \right], \quad r = \sqrt{x^2 + y^2 + z^2}.$$

One-dimensional problems with central symmetry that have solutions  $w = w(r, t)$  are considered in Subsection 4.2.3.

6.1.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_0^R f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &\quad + \int_0^{2\pi} \int_0^\pi \int_0^R f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &\quad - a^2 R^2 \int_0^t \int_0^{2\pi} \int_0^\pi g(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} \sin \eta \, d\eta \, d\zeta \, d\tau, \end{aligned}$$

where

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= \frac{1}{2\pi a R^2 \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\ &\quad \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \sin(\lambda_{nm} a t), \\ A_k &= \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{(2n+1)(n-k)!}{(n+k)! [J'_{n+1/2}(\lambda_{nm} R)]^2 \lambda_{nm}}. \end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as

$$P_n^k(\mu) = (1 - \mu^2)^{k/2} \frac{d^k}{d\mu^k} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n,$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $J_{n+1/2}(\lambda R) = 0$ .

6.1.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$



Solution:

$$\begin{aligned}
 w(r, \theta, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_0^R f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
 & + \int_0^{2\pi} \int_0^\pi \int_0^R f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
 & + a^2 R^2 \int_0^t \int_0^{2\pi} \int_0^\pi g(\eta, \zeta, \tau) G(r, \theta, \varphi, R, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(r, \theta, \varphi, \xi, \eta, \zeta, t) = & \frac{3t}{4\pi R^3} + \frac{1}{2\pi a \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\
 & \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \sin(\lambda_{nm} at), \\
 A_k = & \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{\lambda_{nm} (2n+1)(n-k)!}{(n+k)! [R^2 \lambda_{nm}^2 - n(n+1)] [J_{n+1/2}(\lambda_{nm} R)]^2}.
 \end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions (see Paragraph 6.1.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$2\lambda R J'_{n+1/2}(\lambda R) - J_{n+1/2}(\lambda R) = 0.$$

⊙ *Reference:* M. M. Smirnov (1975).

6.1.3-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_r w + kw = g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad & \text{(boundary condition).}
 \end{aligned}$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 6.1.3-2 where

$$\begin{aligned}
 G(r, \theta, \varphi, \xi, \eta, \zeta, t) = & \frac{1}{2\pi a \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n A_s B_{nms} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\
 & \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \sin(\lambda_{nm} at), \\
 A_s = & \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{\lambda_{nm} (2n+1)(n-s)!}{(n+s)! [R^2 \lambda_{nm}^2 + (kR+n)(kR-n-1)] [J_{n+1/2}(\lambda_{nm} R)]^2}.
 \end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 6.1.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda R J'_{n+1/2}(\lambda R) + \left(kR - \frac{1}{2}\right) J_{n+1/2}(\lambda R) = 0.$$

6.1.3-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 w = g_1(\theta, \varphi, t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
 w = g_2(\theta, \varphi, t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \theta, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + a^2 R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} \sin \eta \, d\eta \, d\zeta \, d\tau \\
& - a^2 R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} \sin \eta \, d\eta \, d\zeta \, d\tau,
\end{aligned}$$

where

$$\begin{aligned}
G(r, \theta, \varphi, \xi, \eta, \zeta, t) = & \frac{\pi}{8a\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} Z_{n+1/2}(\lambda_{nm} r) Z_{n+1/2}(\lambda_{nm} \xi) \\
& \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \sin(\lambda_{nm} at).
\end{aligned}$$

Here,

$$\begin{aligned}
Z_{n+1/2}(\lambda_{nm} r) &= J_{n+1/2}(\lambda_{nm} R_1) Y_{n+1/2}(\lambda_{nm} r) - Y_{n+1/2}(\lambda_{nm} R_1) J_{n+1/2}(\lambda_{nm} r), \\
A_k &= \begin{cases} 1 & \text{for } k=0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{\lambda_{nm} (2n+1)(n-k)! J_{n+1/2}^2(\lambda_{nm} R_2)}{(n+k)! [J_{n+1/2}^2(\lambda_{nm} R_1) - J_{n+1/2}^2(\lambda_{nm} R_2)]},
\end{aligned}$$

where the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as

$$P_n^k(\mu) = (1 - \mu^2)^{k/2} \frac{d^k}{d\mu^k} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n,$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $Z_{n+1/2}(\lambda R_2) = 0$ .

6.1.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned}
w &= f_0(r, \theta, \varphi) \quad \text{at } t=0 \quad (\text{initial condition}), \\
\partial_t w &= f_1(r, \theta, \varphi) \quad \text{at } t=0 \quad (\text{initial condition}), \\
\partial_r w &= g_1(\theta, \varphi, t) \quad \text{at } r=R_1 \quad (\text{boundary condition}), \\
\partial_r w &= g_2(\theta, \varphi, t) \quad \text{at } r=R_2 \quad (\text{boundary condition}).
\end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \theta, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& - a^2 R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) G(r, \theta, \varphi, R_1, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\
& + a^2 R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) G(r, \theta, \varphi, R_2, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau,
\end{aligned}$$

where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{3t}{4\pi(R_2^3 - R_1^3)} + \frac{1}{4\pi a \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n \frac{A_k}{B_{nmk}} Z_{n+1/2}(\lambda_{nm} r) Z_{n+1/2}(\lambda_{nm} \xi) \\ \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \sin(\lambda_{nm} at),$$

$$A_k = \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{\lambda_{nm}(n+k)!}{(2n+1)(n-k)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm} r) dr,$$

$$Z_{n+1/2}(\lambda_{nm} r) = \left[ \lambda_{nm} J'_{n+1/2}(\lambda_{nm} R_1) - \frac{1}{2R_1} J_{n+1/2}(\lambda_{nm} R_1) \right] Y_{n+1/2}(\lambda_{nm} r) \\ - \left[ \lambda_{nm} Y'_{n+1/2}(\lambda_{nm} R_1) - \frac{1}{2R_1} Y_{n+1/2}(\lambda_{nm} R_1) \right] J_{n+1/2}(\lambda_{nm} r).$$

Here, the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions (see Paragraph 6.1.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) - \frac{1}{2R_2} Z_{n+1/2}(\lambda R_2) = 0.$$

6.1.3-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(\theta, \varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(\theta, \varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 6.1.3-5 where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{1}{4\pi a \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n \frac{A_s}{B_{nms}} Z_{n+1/2}(\lambda_{nm} r) Z_{n+1/2}(\lambda_{nm} \xi) \\ \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \sin(\lambda_{nm} at).$$

Here,

$$A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{\lambda_{nm}(n+s)!}{(2n+1)(n-s)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm} r) dr,$$

$$Z_{n+1/2}(\lambda r) = \left[ \lambda J'_{n+1/2}(\lambda R_1) - \left( k_1 + \frac{1}{2R_1} \right) J_{n+1/2}(\lambda R_1) \right] Y_{n+1/2}(\lambda r) \\ - \left[ \lambda Y'_{n+1/2}(\lambda R_1) - \left( k_1 + \frac{1}{2R_1} \right) Y_{n+1/2}(\lambda R_1) \right] J_{n+1/2}(\lambda r),$$

where the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 6.1.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) + \left( k_2 - \frac{1}{2R_2} \right) Z_{n+1/2}(\lambda R_2) = 0.$$

## 6.2. Nonhomogeneous Wave Equation

$$\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_3 w + \Phi(x, y, z, t)$$

### 6.2.1. Problems in Cartesian Coordinates

6.2.1-1. Domain:  $-\infty < x < \infty, -\infty < y < \infty, -\infty < z < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(x, y, z) \quad \text{at} \quad t = 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \frac{1}{4\pi a} \frac{\partial}{\partial t} \iint_{r=at} \frac{f(\xi, \eta, \zeta)}{r} dS + \frac{1}{4\pi a} \iint_{r=at} \frac{g(\xi, \eta, \zeta)}{r} dS \\ &\quad + \frac{1}{4\pi a^2} \iiint_{r \leq at} \frac{1}{r} \Phi\left(\xi, \eta, \zeta, t - \frac{r}{a}\right) d\xi d\eta d\zeta, \quad r = \sqrt{(\xi - x)^2 + (\eta - y)^2 + (\zeta - z)^2}, \end{aligned}$$

where the integration is performed over the surface of the sphere ( $r = at$ ) and the volume of the sphere ( $r \leq at$ ) with center at  $(x, y, z)$ .

⊙ *Reference:* N. S. Koshlyakov, E. B. Glizer, and M. M. Smirnov (1970).

6.2.1-2. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2, 0 \leq z \leq l_3$ . Different boundary value problems.

1°. The solution of the first boundary value problem for a parallelepiped is given by the formula from Paragraph 6.1.1-3 with the additional term

$$\int_0^t \int_0^{l_1} \int_0^{l_2} \int_0^{l_3} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\zeta d\eta d\xi d\tau,$$

which allows for the equation's nonhomogeneity; this term is the solution of the nonhomogeneous equation with homogeneous initial and boundary conditions.

2°. The solution of the second boundary value problem for a parallelepiped is given by the formula from Paragraph 6.1.1-4 with the additional term specified in Paragraph 6.2.1-2, Item 1°; the Green's function is taken from Paragraph 6.1.1-4.

3°. The solution of the third boundary value problem for a parallelepiped is the sum of the solution of the homogeneous equation with nonhomogeneous initial and boundary conditions (see Paragraph 6.1.1-5) and the solution of the nonhomogeneous equation with homogeneous initial and boundary conditions. The latter solution is given by the formula from Paragraph 6.2.1-2, Item 1°, in which one should substitute the Green's function from Paragraph 6.1.1-5.

4°. The solutions of mixed boundary value problems for a parallelepiped are given by the formulas from Paragraph 6.1.1-6 to which one should add the term specified in Paragraph 6.2.1-2, Item 1°.

### 6.2.2. Problems in Cylindrical Coordinates

A three-dimensional nonhomogeneous wave equation in the cylindrical coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left[ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} \right] + \Phi(r, \varphi, z, t).$$

**6.2.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a circular cylinder of finite length is given by the formula from Paragraph 6.1.2-1 with the additional term

$$\int_0^t \int_0^l \int_0^{2\pi} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a circular cylinder of finite length is given by the formula from Paragraph 6.1.2-2 with the additional term (1).

3°. The solution of the third boundary value problem for a circular cylinder of finite length is the sum of the solution specified in Paragraph 6.1.2-3 and expression (1).

4°. The solutions of mixed boundary value problems for a circular cylinder of finite length are given by the formulas from Paragraph 6.1.2-4 with additional terms of the form (1).

**6.2.2-2. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a hollow cylinder of finite dimensions is given by the formula from Paragraph 6.1.2-5 with the additional term

$$\int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a hollow cylinder of finite dimensions is given by the formula from Paragraph 6.1.2-6 with the additional term (2).

3°. The solution of the third boundary value problem for a hollow cylinder of finite dimensions is the sum of the solution specified in Paragraph 6.1.2-7 and expression (2).

4°. The solutions of mixed boundary value problems for a hollow cylinder of finite dimensions are given by the formulas from Paragraph 6.1.2-8 with additional terms of the form (2).

**6.2.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . Different boundary value problems.**

1°. The solution of the first boundary value problem for a cylindrical sector of finite thickness is given by the formula from Paragraph 6.1.2-9 with the additional term

$$\int_0^t \int_0^l \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau, \quad (3)$$

which allows for the equation's nonhomogeneity.

2°. The solution of a mixed boundary value problem for a cylindrical sector of finite thickness is given by the formula from Paragraph 6.1.2-10 with the additional term (2).

## 6.2.3. Problems in Spherical Coordinates

A three-dimensional nonhomogeneous wave equation in the spherical coordinate system is represented as

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} \right] + \Phi(r, \theta, \varphi, t).$$

**6.2.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Boundary value problem.**

1°. The solution of the first boundary value problem for a sphere is given by the formula from Paragraph 6.1.3-1 with the additional term

$$\int_0^t \int_0^{2\pi} \int_0^\pi \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau, \quad (1)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a sphere is given by the formula from Paragraph 6.1.3-2 with the additional term (1).

3°. The solution of the third boundary value problem for a sphere is the sum of the solution specified in Paragraph 6.1.3-3 and expression (1).

**6.2.3-2. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Boundary value problems.**

1°. The solution of the first boundary value problem for a spherical layer is given by the formula from Paragraph 6.1.3-4 with the additional term

$$\int_0^t \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau, \quad (2)$$

which allows for the equation's nonhomogeneity.

2°. The solution of the second boundary value problem for a spherical layer is given by the formula from Paragraph 6.1.3-5 with the additional term (2).

3°. The solution of the third boundary value problem for a spherical layer is the sum of the solution specified in Paragraph 6.1.3-6 and expression (2).

## 6.3. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_3 w - bw + \Phi(x, y, z, t)$

### 6.3.1. Problems in Cartesian Coordinates

A three-dimensional nonhomogeneous Klein–Gordon equation in the rectangular Cartesian system of coordinates has the form

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) - bw + \Phi(x, y, z, t).$$

**6.3.1-1. Fundamental solutions.**

1°. For  $b = -c^2 < 0$ ,

$$\mathcal{E}(x, y, z, t) = \frac{1}{4\pi a^2} \left[ \frac{\delta(t - r/a)}{r} - \frac{c}{a} \frac{I_1(c\sqrt{t^2 - r^2/a^2})}{\sqrt{t^2 - r^2/a^2}} \vartheta(t - r/a) \right],$$

where  $r = \sqrt{x^2 + y^2 + z^2}$ ,  $\delta(\xi)$  is the Dirac delta function,  $\vartheta(\xi)$  is the Heaviside unit step function, and  $I_1(z)$  is the modified Bessel function.

2°. For  $b = c^2 > 0$ ,

$$\mathcal{E}(x, y, z, t) = \frac{1}{4\pi a^2} \left[ \frac{\delta(t - r/a)}{r} - \frac{c}{a} \frac{J_1(c\sqrt{t^2 - r^2/a^2})}{\sqrt{t^2 - r^2/a^2}} \vartheta(t - r/a) \right],$$

where  $J_1(z)$  is the Bessel function.

☉ *Reference:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

6.3.1-2. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(x, y, z) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(x, y, z) \quad \text{at} \quad t = 0. \end{aligned}$$

Let  $a = 1$  and  $\Phi(x, y, z, t) \equiv 0$ .

1°. Solution for  $b = -c^2 < 0$ :

$$\begin{aligned} w(x, y, z, t) &= \frac{\partial}{\partial t} \left[ \frac{1}{t} \frac{\partial}{\partial t} \int_0^t r^2 I_0(c\sqrt{t^2 - r^2}) T_r[f(x, y, z)] dr \right] \\ &\quad + \frac{1}{t} \frac{\partial}{\partial t} \int_0^t r^2 I_0(c\sqrt{t^2 - r^2}) T_r[g(x, y, z)] dr. \end{aligned}$$

Here,  $I_0(z)$  is the modified Bessel function and  $T_r[h(x, y, z)]$  is the average of  $h(x, y, z)$  over the spherical surface with center at  $(x, y, z)$  and radius  $r$ :

$$T_r[h(x, y, z)] = \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi h(x + r \sin \theta \cos \varphi, y + r \sin \theta \sin \varphi, z + r \cos \theta) \sin \theta d\theta d\varphi.$$

2°. Solution for  $b = c^2 > 0$ :

$$\begin{aligned} w(x, y, z, t) &= \frac{\partial}{\partial t} \left[ \frac{1}{t} \frac{\partial}{\partial t} \int_0^t r^2 J_0(c\sqrt{t^2 - r^2}) T_r[f(x, y, z)] dr \right] \\ &\quad + \frac{1}{t} \frac{\partial}{\partial t} \int_0^t r^2 J_0(c\sqrt{t^2 - r^2}) T_r[g(x, y, z)] dr, \end{aligned}$$

where  $J_0(z)$  is the Bessel function.

☉ *Reference:* V. I. Smirnov (1974, Vol. 2).

6.3.1-3. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . First boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\ w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\ w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_6(x, y, t) \quad \text{at} \quad z = l_3 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
w(x, y, z, t) = & \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
& + \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
& + a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\
& - a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_6(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l_3} d\xi d\eta d\tau \\
& + \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(x, y, z, \xi, \eta, \zeta, t) = & \frac{8}{l_1 l_2 l_3} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{\sqrt{\lambda_{nmk}}} \sin(\alpha_n x) \sin(\beta_m y) \sin(\gamma_k z) \\
& \times \sin(\alpha_n \xi) \sin(\beta_m \eta) \sin(\gamma_k \zeta) \sin(t \sqrt{\lambda_{nmk}}),
\end{aligned}$$

where

$$\alpha_n = \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_k = \frac{k\pi}{l_3}, \quad \lambda_{nmk} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_k^2) + b.$$

6.3.1-4. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Second boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned}
w &= f_0(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
\partial_t w &= f_1(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
\partial_x w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\
\partial_x w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\
\partial_y w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\
\partial_y w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\
\partial_z w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
\partial_z w &= g_6(x, y, t) \quad \text{at} \quad z = l_3 \quad (\text{boundary condition}).
\end{aligned}$$



Solution:

$$\begin{aligned}
w(x, y, z, t) = & \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
& + \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
& - a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\
& + a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau \\
& + \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau,
\end{aligned}$$

where

$$\begin{aligned}
G(x, y, z, \xi, \eta, \zeta, t) = & \frac{\sin(t\sqrt{b})}{l_1 l_2 l_3 \sqrt{b}} + \frac{1}{l_1 l_2 l_3} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_m A_k}{\sqrt{\lambda_{nmk}}} \cos(\alpha_n x) \cos(\beta_m y) \cos(\gamma_k z) \\
& \times \cos(\alpha_n \xi) \cos(\beta_m \eta) \cos(\gamma_k \zeta) \sin(t\sqrt{\lambda_{nmk}}), \\
\alpha_n = & \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_k = \frac{k\pi}{l_3}, \quad \lambda_{nmk} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_k^2) + b, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0. \end{cases}
\end{aligned}$$

The summation is performed over the indices satisfying the condition  $n + m + k > 0$ ; the term corresponding to  $n = m = k = 0$  is singled out.

6.3.1-5. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2, 0 \leq z \leq l_3$ . Third boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(x, y, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_t w = f_1(x, y, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_x w - s_1 w = g_1(y, z, t) & \quad \text{at } x = 0 \quad (\text{boundary condition}), \\
\partial_x w + s_2 w = g_2(y, z, t) & \quad \text{at } x = l_1 \quad (\text{boundary condition}), \\
\partial_y w - s_3 w = g_3(x, z, t) & \quad \text{at } y = 0 \quad (\text{boundary condition}), \\
\partial_y w + s_4 w = g_4(x, z, t) & \quad \text{at } y = l_2 \quad (\text{boundary condition}), \\
\partial_z w - s_5 w = g_5(x, y, t) & \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
\partial_z w + s_6 w = g_6(x, y, t) & \quad \text{at } z = l_3 \quad (\text{boundary condition}).
\end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 6.3.1-4 where

$$\begin{aligned}
G(x, y, \xi, \eta, t) = & 8 \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{E_{nmk} \sqrt{\lambda_{nmk}}} \sin(\alpha_n x + \varepsilon_n) \sin(\beta_m y + \sigma_m) \sin(\gamma_k z + \nu_k) \\
& \times \sin(\alpha_n \xi + \varepsilon_n) \sin(\beta_m \eta + \sigma_m) \sin(\gamma_k \zeta + \nu_k) \sin(t\sqrt{\lambda_{nmk}}),
\end{aligned}$$

$$\varepsilon_n = \arctan \frac{\alpha_n}{l_1}, \quad \sigma_m = \arctan \frac{\beta_m}{l_2}, \quad \nu_k = \arctan \frac{\gamma_k}{l_3}, \quad \lambda_{nmk} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_k^2) + b,$$

$$E_{nmk} = \left[ l_1 + \frac{(s_1 s_2 + \alpha_n^2)(s_1 + s_2)}{(s_1^2 + \alpha_n^2)(s_2^2 + \alpha_n^2)} \right] \left[ l_2 + \frac{(s_3 s_4 + \beta_m^2)(s_3 + s_4)}{(s_3^2 + \beta_m^2)(s_4^2 + \beta_m^2)} \right] \left[ l_3 + \frac{(s_5 s_6 + \gamma_k^2)(s_5 + s_6)}{(s_5^2 + \gamma_k^2)(s_6^2 + \gamma_k^2)} \right].$$

Here, the  $\alpha_n$ ,  $\beta_m$ , and  $\gamma_k$  are positive roots of the transcendental equations

$$\alpha^2 - s_1 s_2 = (s_1 + s_2) \alpha \cot(l_1 \alpha), \quad \beta^2 - s_3 s_4 = (s_3 + s_4) \beta \cot(l_2 \beta), \quad \gamma^2 - s_5 s_6 = (s_5 + s_6) \gamma \cot(l_3 \gamma).$$

6.3.1-6. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Mixed boundary value problems.

1°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ \partial_y w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau \\ &+ \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{2}{l_1 l_2 l_3} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{A_m A_k}{\sqrt{\lambda_{nmk}}} \sin(\alpha_n x) \cos(\beta_m y) \cos(\gamma_k z) \\ &\quad \times \sin(\alpha_n \xi) \cos(\beta_m \eta) \cos(\gamma_k \zeta) \sin(t \sqrt{\lambda_{nmk}}), \end{aligned}$$

where

$$A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 0, \end{cases} \quad A_k = \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k > 0, \end{cases}$$

$$\alpha_n = \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_k = \frac{k\pi}{l_3}, \quad \lambda_{nmk} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_k^2) + b.$$

2°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\ \partial_x w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\ w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\ \partial_y w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\ w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_1(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau \\ &+ \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) &= \frac{8}{l_1 l_2 l_3} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{\sqrt{\lambda_{nmk}}} \sin(\alpha_n x) \sin(\beta_m y) \sin(\gamma_k z) \\ &\quad \times \sin(\alpha_n \xi) \sin(\beta_m \eta) \sin(\gamma_k \zeta) \sin(t\sqrt{\lambda_{nmk}}), \end{aligned}$$

where

$$\alpha_n = \frac{\pi(2n+1)}{2l_1}, \quad \beta_m = \frac{\pi(2m+1)}{2l_2}, \quad \gamma_k = \frac{\pi(2k+1)}{2l_3},$$

$$\lambda_{nmk} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_k^2) + b.$$

### 6.3.2. Problems in Cylindrical Coordinates

A nonhomogeneous Klein–Gordon equation in the cylindrical coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left[ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} \right] - bw + \Phi(r, \varphi, z, t), \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional problems with axial symmetry that have solutions  $w = w(r, t)$  are treated in Subsection 4.2.5. Two-dimensional problems whose solutions have the form  $w = w(r, \varphi, t)$  or  $w = w(r, z, t)$  are considered in Subsections 5.3.2 and 5.3.3.

6.3.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) \quad \text{at} \quad z = l \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{2}{\pi R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nmk}}} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\ &\quad \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \sin(t\sqrt{\lambda_{nmk}}), \end{aligned}$$

where

$$\lambda_{nmk} = a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}$$

the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument), and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

**6.3.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{\sin(t\sqrt{b})}{\pi R^2 l \sqrt{b}} + \frac{2}{\pi R^2 l} \sum_{k=1}^{\infty} \frac{1}{\sqrt{\beta_k}} \cos\left(\frac{k\pi x}{l}\right) \cos\left(\frac{k\pi \xi}{l}\right) \sin(t\sqrt{\beta_k}) \\ &+ \frac{1}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi x}{l}\right) \cos\left(\frac{k\pi \xi}{l}\right) \frac{\sin(\lambda_{nmk} t)}{\lambda_{nmk}}, \\ \beta_k &= \frac{a^2 k^2 \pi^2}{l^2} + b, \quad \lambda_{nmk} = \sqrt{a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

**6.3.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.**

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w + k_1 w &= g(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w - k_2 w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w + k_3 w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 6.3.2-2 where

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] h_s(z) h_s(\zeta) \sin(\lambda_{nms} t)}{(\mu_{nm}^2 R^2 + k_1^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2 \|h_s\|^2 \lambda_{nms}}.$$

Here,

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \quad \lambda_{nms} = \sqrt{a^2 \mu_{nm}^2 + a^2 \beta_s^2 + b},$$

$$h_s(z) = \cos(\beta_s z) + \frac{k_2}{\beta_s} \sin(\beta_s z), \quad \|h_s\|^2 = \frac{k_3}{2\beta_s^2} \frac{\beta_s^2 + k_2^2}{\beta_s^2 + k_3^2} + \frac{k_2}{2\beta_s^2} + \frac{l}{2} \left(1 + \frac{k_2^2}{\beta_s^2}\right),$$

the  $J_n(\xi)$  are the Bessel functions, and the  $\mu_{nm}$  and  $\beta_s$  are positive roots of the transcendental equations

$$\mu J'_n(\mu R) + k_1 J_n(\mu R) = 0, \quad \frac{\tan(\beta l)}{\beta} = \frac{k_2 + k_3}{\beta^2 - k_2 k_3}.$$

6.3.2-4. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{\pi R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k}{[J'_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nmk}}} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\ &\quad \times \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi \zeta}{l}\right) \sin(t\sqrt{\lambda_{nmk}}), \\ \lambda_{nmk} &= a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\
 w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\
 w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 &+ \int_0^l \int_0^{2\pi} \int_0^R \xi f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 &+ a^2 R \int_0^t \int_0^{2\pi} \int_0^R g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 &+ a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\
 &- a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau \\
 &+ \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{2}{\pi R^2 l} \sum_{k=1}^{\infty} \frac{1}{\sqrt{\beta_k}} \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \sin(t\sqrt{\beta_k}) \\
 &+ \frac{2}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n \mu_{nm}^2}{(\mu_{nm}^2 R^2 - n^2)[J_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nmk}}} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\
 &\quad \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \sin(t\sqrt{\lambda_{nmk}}), \\
 \beta_k &= \frac{a^2 k^2 \pi^2}{l^2} + b, \quad \lambda_{nmk} = a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}
 \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

6.3.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\
 w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\
 w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\
 w &= g_4(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\
& - a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\
& + a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\
& - a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau \\
& + \int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{\pi}{2l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)} Z_{nm}(r) Z_{nm}(\xi) \\
& \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nmk}})}{\sqrt{\lambda_{nmk}}},
\end{aligned}$$

where

$$\begin{aligned}
A_n = & \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nmk} = a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b, \\
Z_{nm}(r) = & J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r);
\end{aligned}$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

6.3.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(r, \varphi, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_t w = f_1(r, \varphi, z) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_r w = g_1(\varphi, z, t) & \quad \text{at } r = R_1 \quad (\text{boundary condition}), \\
\partial_r w = g_2(\varphi, z, t) & \quad \text{at } r = R_2 \quad (\text{boundary condition}), \\
\partial_z w = g_3(r, \varphi, t) & \quad \text{at } z = 0 \quad (\text{boundary condition}), \\
\partial_z w = g_4(r, \varphi, t) & \quad \text{at } z = l \quad (\text{boundary condition}).
\end{aligned}$$



Solution:

$$\begin{aligned}
w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& - a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& + a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& - a^2 \int_0^t \int_0^l \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
& + a^2 \int_0^t \int_0^l \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau \\
& + \int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{\sin(t\sqrt{b})}{\pi(R_2^2 - R_1^2)l\sqrt{b}} + \frac{2}{\pi(R_2^2 - R_1^2)l} \sum_{k=1}^{\infty} \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi\zeta}{l}\right) \frac{\sin(t\sqrt{\beta_k})}{\sqrt{\beta_k}} \\
& + \frac{1}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k \mu_{nm}^2 Z_{nm}(r) Z_{nm}(\xi)}{(\mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)} \\
& \times \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi\zeta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nmk}})}{\sqrt{\lambda_{nmk}}},
\end{aligned}$$

where

$$\begin{aligned}
A_n = & \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \beta_k = \frac{a^2 k^2 \pi^2}{l^2} + b, \quad \lambda_{nmk} = a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b, \\
Z_{nm}(r) = & J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r);
\end{aligned}$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

6.3.2-7. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(r, \varphi, z) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_t w = f_1(r, \varphi, z) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_r w - k_1 w = g_1(\varphi, z, t) \quad & \text{at } r = R_1 \quad (\text{boundary condition}), \\
\partial_r w + k_2 w = g_2(\varphi, z, t) \quad & \text{at } r = R_2 \quad (\text{boundary condition}), \\
\partial_z w - k_3 w = g_3(r, \varphi, t) \quad & \text{at } z = 0 \quad (\text{boundary condition}), \\
\partial_z w + k_4 w = g_4(r, \varphi, t) \quad & \text{at } z = l \quad (\text{boundary condition}).
\end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 6.3.2-6 where

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2}{\|h_s\|^2 \sqrt{a^2 \mu_{nm}^2 + a^2 \lambda_s^2 + b}} \times \frac{Z_{nm}(r) Z_{nm}(\xi) \cos[n(\varphi - \eta)] h_s(z) h_s(\zeta) \sin(t \sqrt{a^2 \mu_{nm}^2 + a^2 \lambda_s^2 + b})}{(k_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (k_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)}.$$

Here,

$$Z_{nm}(r) = [\mu_{nm} J'_n(\mu_{nm} R_1) - k_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) - [\mu_{nm} Y'_n(\mu_{nm} R_1) - k_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r),$$

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad h_s(z) = \cos(\lambda_s z) + \frac{k_3}{\lambda_s} \sin(\lambda_s z), \quad \|h_s\|^2 = \frac{k_4}{2\lambda_s^2} \frac{\lambda_s^2 + k_3^2}{\lambda_s^2 + k_4^2} + \frac{k_3}{2\lambda_s^2} + \frac{l}{2} \left(1 + \frac{k_3^2}{\lambda_s^2}\right),$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$[\mu J'_n(\mu R_1) - k_1 J_n(\mu R_1)] [\mu Y'_n(\mu R_2) + k_2 Y_n(\mu R_2)] = [\mu Y'_n(\mu R_1) - k_1 Y_n(\mu R_1)] [\mu J'_n(\mu R_2) + k_2 J_n(\mu R_2)],$$

and the  $\lambda_s$  are positive roots of the transcendental equation  $\frac{\tan(\lambda l)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}$ .

6.3.2-8. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.

1°. A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\ &- a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{\pi}{4l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_n A_k \mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)} Z_{nm}(r) Z_{nm}(\xi) \\ \times \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nmk}})}{\sqrt{\lambda_{nmk}}},$$

where

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nmk} = a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b, \\ Z_{nm}(r) = J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r);$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

2°. A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_4(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R_1 \int_0^t \int_0^{2\pi} \int_0^l g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 R_2 \int_0^t \int_0^{2\pi} \int_0^l g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{2}{\pi(R_2^2 - R_1^2)l} \sum_{k=1}^{\infty} \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\beta_k})}{\sqrt{\beta_k}} \\ &+ \frac{2}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_{nm}(r) Z_{nm}(\xi)}{(\mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)} \\ &\times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nmk}})}{\sqrt{\lambda_{nmk}}}, \end{aligned}$$

where

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \beta_k = \frac{a^2 k^2 \pi^2}{l^2} + b, \quad \lambda_{nmk} = a^2 \mu_{nm}^2 + \frac{a^2 k^2 \pi^2}{l^2} + b,$$

$$Z_{nm}(r) = J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r);$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

6.3.2-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . First boundary value problem.

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, z, t) & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, z, t) & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}), \\ w &= g_4(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_5(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{8}{R^2 l \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \\ &\quad \times \sin\left(\frac{k\pi z}{l}\right) \sin\left(\frac{k\pi\zeta}{l}\right) \frac{\sin\left(t\sqrt{a^2\mu_{nm}^2 + a^2k^2\pi^2l^{-2} + b}\right)}{\sqrt{a^2\mu_{nm}^2 + a^2k^2\pi^2l^{-2} + b}}, \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\
 \partial_t w &= f_1(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\
 w &= g_1(\varphi, z, t) & \text{at } r &= R & (\text{boundary condition}), \\
 w &= g_2(r, z, t) & \text{at } \varphi &= 0 & (\text{boundary condition}), \\
 w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}), \\
 \partial_z w &= g_4(r, \varphi, t) & \text{at } z &= 0 & (\text{boundary condition}), \\
 \partial_z w &= g_5(r, \varphi, t) & \text{at } z &= l & (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 &+ \int_0^l \int_0^{\varphi_0} \int_0^R f_1(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
 &- a^2 R \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\
 &+ a^2 \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 &- a^2 \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\
 &- a^2 \int_0^t \int_0^l \int_0^{\varphi_0} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
 &+ a^2 \int_0^t \int_0^l \int_0^{\varphi_0} g_5(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau \\
 &+ \int_0^t \int_0^l \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{4}{R^2 l \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_k J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \\
 &\times \cos\left(\frac{k\pi z}{l}\right) \cos\left(\frac{k\pi\zeta}{l}\right) \frac{\sin\left(t\sqrt{a^2\mu_{nm}^2 + a^2k^2\pi^2l^{-2} + b}\right)}{\sqrt{a^2\mu_{nm}^2 + a^2k^2\pi^2l^{-2} + b}},
 \end{aligned}$$

where  $A_0 = 1$  and  $A_k = 2$  for  $k \geq 1$ ; the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

### 6.3.3. Problems in Spherical Coordinates

A nonhomogeneous Klein–Gordon equation in the spherical coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} = a^2 \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} \right] - bw + \Phi(r, \theta, \varphi, t).$$

One-dimensional problems with central symmetry that have solutions of the form  $w = w(r, t)$  are treated in Subsection 4.2.6.

**6.3.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_0^R f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &+ \int_0^{2\pi} \int_0^\pi \int_0^R f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &- a^2 R^2 \int_0^t \int_0^{2\pi} \int_0^\pi g(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} \sin \eta \, d\eta \, d\zeta \, d\tau \\ &+ \int_0^t \int_0^{2\pi} \int_0^\pi \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= \frac{1}{2\pi R^2 \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\ &\quad \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \sin(t \sqrt{a^2 \lambda_{nm}^2 + b}), \end{aligned}$$

where

$$A_k = \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{(2n+1)(n-k)!}{(n+k)! [J'_{n+1/2}(\lambda_{nm} R)]^2 \sqrt{a^2 \lambda_{nm}^2 + b}};$$

the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as

$$P_n^k(\mu) = (1 - \mu^2)^{k/2} \frac{d^k}{d\mu^k} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n,$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $J_{n+1/2}(\lambda R) = 0$ .

**6.3.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_0^R f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &+ \int_0^{2\pi} \int_0^\pi \int_0^R f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &+ a^2 R^2 \int_0^t \int_0^{2\pi} \int_0^\pi g(\eta, \zeta, \tau) G(r, \theta, \varphi, R, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\ &+ \int_0^t \int_0^{2\pi} \int_0^\pi \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau. \end{aligned}$$

Here,

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{3 \sin(t\sqrt{b})}{4\pi R^3 \sqrt{b}} + \frac{1}{2\pi \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n \frac{A_k B_{nmk}}{\sqrt{a^2 \lambda_{nm}^2 + b}} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\ \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \sin(t\sqrt{a^2 \lambda_{nm}^2 + b}),$$

where

$$A_k = \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{\lambda_{nm}^2 (2n+1)(n-k)!}{(n+k)! [R^2 \lambda_{nm}^2 - n(n+1)] [J_{n+1/2}(\lambda_{nm} R)]^2};$$

the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions (see Paragraph 6.3.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$2\lambda R J'_{n+1/2}(\lambda R) - J_{n+1/2}(\lambda R) = 0.$$

6.3.3-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_r w + kw &= g(\theta, \varphi, t) \quad \text{at } r = R \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 6.3.3-2 where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{1}{2\pi \sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n \frac{A_s B_{nms}}{\sqrt{a^2 \lambda_{nm}^2 + b}} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \\ \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \sin(t\sqrt{a^2 \lambda_{nm}^2 + b}).$$

Here,

$$A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{\lambda_{nm}^2 (2n+1)(n-s)!}{(n+s)! [R^2 \lambda_{nm}^2 + (kR+n)(kR-n-1)] [J_{n+1/2}(\lambda_{nm} R)]^2};$$

the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 6.3.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda R J'_{n+1/2}(\lambda R) + (kR - \frac{1}{2}) J_{n+1/2}(\lambda R) = 0.$$

6.3.3-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at } t = 0 \quad (\text{initial condition}), \\ w &= g_1(\theta, \varphi, t) \quad \text{at } r = R_1 \quad (\text{boundary condition}), \\ w &= g_2(\theta, \varphi, t) \quad \text{at } r = R_2 \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \theta, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + a^2 R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} \sin \eta \, d\eta \, d\zeta \, d\tau \\
& - a^2 R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} \sin \eta \, d\eta \, d\zeta \, d\tau \\
& + \int_0^t \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau,
\end{aligned}$$

where

$$\begin{aligned}
G(r, \theta, \varphi, \xi, \eta, \zeta, t) = & \frac{\pi}{8\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n \frac{A_k B_{nmk}}{\sqrt{a^2 \lambda_{nm}^2 + b}} Z_{n+1/2}(\lambda_{nm} r) Z_{n+1/2}(\lambda_{nm} \xi) \\
& \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \sin(t \sqrt{a^2 \lambda_{nm}^2 + b}).
\end{aligned}$$

Here,

$$\begin{aligned}
Z_{n+1/2}(\lambda_{nm} r) = & J_{n+1/2}(\lambda_{nm} R_1) Y_{n+1/2}(\lambda_{nm} r) - Y_{n+1/2}(\lambda_{nm} R_1) J_{n+1/2}(\lambda_{nm} r), \\
A_k = & \begin{cases} 1 & \text{for } k=0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{\lambda_{nm} (2n+1)(n-k)! J_{n+1/2}^2(\lambda_{nm} R_2)}{(n+k)! [J_{n+1/2}^2(\lambda_{nm} R_1) - J_{n+1/2}^2(\lambda_{nm} R_2)]},
\end{aligned}$$

where the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as

$$P_n^k(\mu) = (1 - \mu^2)^{k/2} \frac{d^k}{d\mu^k} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n,$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $Z_{n+1/2}(\lambda R_2) = 0$ .

6.3.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(r, \theta, \varphi) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_t w = f_1(r, \theta, \varphi) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
\partial_r w = g_1(\theta, \varphi, t) \quad & \text{at } r = R_1 \quad (\text{boundary condition}), \\
\partial_r w = g_2(\theta, \varphi, t) \quad & \text{at } r = R_2 \quad (\text{boundary condition}).
\end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \theta, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_1(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& - a^2 R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) G(r, \theta, \varphi, R_1, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\
& + a^2 R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) G(r, \theta, \varphi, R_2, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\
& + \int_0^t \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau,
\end{aligned}$$



where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{3 \sin(t\sqrt{b})}{4\pi(R_2^3 - R_1^3)\sqrt{b}} + \frac{1}{4\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n \frac{A_k}{B_{nmk}} Z_{n+1/2}(\lambda_{nm}r) Z_{n+1/2}(\lambda_{nm}\xi) \\ \times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2\lambda_{nm}^2 + b})}{\sqrt{a^2\lambda_{nm}^2 + b}}.$$

Here,

$$A_k = \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{(n+k)!}{(2n+1)(n-k)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm}r) dr, \\ Z_{n+1/2}(\lambda_{nm}r) = \left[ \lambda_{nm} J'_{n+1/2}(\lambda_{nm}R_1) - \frac{1}{2R_1} J_{n+1/2}(\lambda_{nm}R_1) \right] Y_{n+1/2}(\lambda_{nm}r) \\ - \left[ \lambda_{nm} Y'_{n+1/2}(\lambda_{nm}R_1) - \frac{1}{2R_1} Y_{n+1/2}(\lambda_{nm}R_1) \right] J_{n+1/2}(\lambda_{nm}r),$$

where the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions (see Paragraph 6.3.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) - \frac{1}{2R_2} Z_{n+1/2}(\lambda R_2) = 0.$$

6.3.3-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w - k_1 w &= g_1(\theta, \varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w + k_2 w &= g_2(\theta, \varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 6.3.3-5 where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{1}{4\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n \frac{A_s}{B_{nms}} Z_{n+1/2}(\lambda_{nm}r) Z_{n+1/2}(\lambda_{nm}\xi) \\ \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2\lambda_{nm}^2 + b})}{\sqrt{a^2\lambda_{nm}^2 + b}}.$$

Here,

$$A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{(n+s)!}{(2n+1)(n-s)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm}r) dr, \\ Z_{n+1/2}(\lambda r) = \left[ \lambda J'_{n+1/2}(\lambda R_1) - \left( k_1 + \frac{1}{2R_1} \right) J_{n+1/2}(\lambda R_1) \right] Y_{n+1/2}(\lambda r) \\ - \left[ \lambda Y'_{n+1/2}(\lambda R_1) - \left( k_1 + \frac{1}{2R_1} \right) Y_{n+1/2}(\lambda R_1) \right] J_{n+1/2}(\lambda r),$$

where the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 6.3.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) + \left( k_2 - \frac{1}{2R_2} \right) Z_{n+1/2}(\lambda R_2) = 0.$$

## 6.4. Telegraph Equation

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \Delta_3 w - bw + \Phi(x, y, z, t)$$

### 6.4.1. Problems in Cartesian Coordinates

A three-dimensional nonhomogeneous telegraph equation in the rectangular Cartesian system of coordinates has the form

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) - bw + \Phi(x, y, z, t).$$

#### 6.4.1-1. Reduction to the three-dimensional Klein–Gordon equation.

The substitution  $w(x, y, z, t) = \exp(-\frac{1}{2}kt)u(x, y, z, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right) - \left(b - \frac{1}{4}k^2\right)u + \exp\left(\frac{1}{2}kt\right)\Phi(x, y, z, t),$$

which is discussed in Subsection 6.3.1.

#### 6.4.1-2. Domain: $0 \leq x \leq l_1$ , $0 \leq y \leq l_2$ , $0 \leq z \leq l_3$ . First boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x, y, z) & \text{at } t &= 0 & \text{(initial condition),} \\ w &= g_1(y, z, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ w &= g_2(y, z, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ w &= g_3(x, z, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ w &= g_4(x, z, t) & \text{at } y &= l_2 & \text{(boundary condition),} \\ w &= g_5(x, y, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ w &= g_6(x, y, t) & \text{at } z &= l_3 & \text{(boundary condition).} \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=l_2} d\xi d\zeta d\tau \end{aligned}$$

$$\begin{aligned}
& + a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\
& - a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l_3} d\xi d\eta d\tau \\
& + \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(x, y, z, \xi, \eta, \zeta, t) = \frac{8}{l_1 l_2 l_3} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{1}{\sqrt{\lambda_{nms}}} \sin(\alpha_n x) \sin(\beta_m y) \sin(\gamma_s z) \\
\times \sin(\alpha_n \xi) \sin(\beta_m \eta) \sin(\gamma_s \zeta) \sin\left(t\sqrt{\lambda_{nms}}\right),
\end{aligned}$$

where

$$\alpha_n = \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_s = \frac{s\pi}{l_3}, \quad \lambda_{nms} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_s^2) + b - \frac{1}{4}k^2.$$

6.4.1-3. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Second boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned}
w &= f_0(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
\partial_t w &= f_1(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
\partial_x w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\
\partial_x w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\
\partial_y w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\
\partial_y w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\
\partial_z w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
\partial_z w &= g_6(x, y, t) \quad \text{at} \quad z = l_3 \quad (\text{boundary condition}).
\end{aligned}$$

Solution:

$$\begin{aligned}
w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
&+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
&- a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) G(x, y, z, 0, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
&+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
&- a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
&+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\
&- a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\
&+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau \\
&+ \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau,
\end{aligned}$$

where

$$G(x, y, z, \xi, \eta, \zeta, t) = \frac{e^{-kt/2}}{l_1 l_2 l_3} \left[ \frac{\sin(t\sqrt{c})}{\sqrt{c}} + \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{s=0}^{\infty} \frac{A_n A_m A_s}{\sqrt{\lambda_{nms}}} \cos(\alpha_n x) \cos(\beta_m y) \cos(\gamma_s z) \right. \\ \left. \times \cos(\alpha_n \xi) \cos(\beta_m \eta) \cos(\gamma_s \zeta) \sin(t\sqrt{\lambda_{nms}}) \right],$$

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \quad \alpha_n = \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_s = \frac{s\pi}{l_3},$$

$$c = b - \frac{1}{4}k^2, \quad \lambda_{nms} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_s^2) + b - \frac{1}{4}k^2.$$

The summation is performed over the indices satisfying the condition  $n + m + s > 0$ ; the term corresponding to  $n = m = s = 0$  is singled out.

6.4.1-4. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ ,  $0 \leq z \leq l_3$ . Third boundary value problem.

A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(x, y, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(x, y, z) & \text{at } t &= 0 & \text{(initial condition),} \\ \partial_x w - s_1 w &= g_1(y, z, t) & \text{at } x &= 0 & \text{(boundary condition),} \\ \partial_x w + s_2 w &= g_2(y, z, t) & \text{at } x &= l_1 & \text{(boundary condition),} \\ \partial_y w - s_3 w &= g_3(x, z, t) & \text{at } y &= 0 & \text{(boundary condition),} \\ \partial_y w + s_4 w &= g_4(x, z, t) & \text{at } y &= l_2 & \text{(boundary condition),} \\ \partial_z w - s_5 w &= g_5(x, y, t) & \text{at } z &= 0 & \text{(boundary condition),} \\ \partial_z w + s_6 w &= g_6(x, y, t) & \text{at } z &= l_3 & \text{(boundary condition).} \end{aligned}$$

The solution  $w(x, y, z, t)$  is determined by the formula in Paragraph 6.4.1-3 where

$$G(x, y, \xi, \eta, t) = 8 \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \sum_{p=1}^{\infty} \sum_{q=1}^{\infty} \frac{1}{E_{npq} \sqrt{\lambda_{npq}}} \sin(\alpha_n x + \varepsilon_n) \sin(\beta_p y + \sigma_p) \sin(\gamma_q z + \nu_q) \\ \times \sin(\alpha_n \xi + \varepsilon_n) \sin(\beta_p \eta + \sigma_p) \sin(\gamma_q \zeta + \nu_q) \sin(t\sqrt{\lambda_{npq}}).$$

Here,

$$\varepsilon_n = \arctan \frac{\alpha_n}{l_1}, \quad \sigma_p = \arctan \frac{\beta_p}{l_2}, \quad \nu_q = \arctan \frac{\gamma_q}{l_3}, \quad \lambda_{npq} = a^2(\alpha_n^2 + \beta_p^2 + \gamma_q^2) + b - \frac{1}{4}k^2,$$

$$E_{npq} = \left[ l_1 + \frac{(s_1 s_2 + \alpha_n^2)(s_1 + s_2)}{(s_1^2 + \alpha_n^2)(s_2^2 + \alpha_n^2)} \right] \left[ l_2 + \frac{(s_3 s_4 + \beta_p^2)(s_3 + s_4)}{(s_3^2 + \beta_p^2)(s_4^2 + \beta_p^2)} \right] \left[ l_3 + \frac{(s_5 s_6 + \gamma_q^2)(s_5 + s_6)}{(s_5^2 + \gamma_q^2)(s_6^2 + \gamma_q^2)} \right],$$

where the  $\alpha_n$ ,  $\beta_p$ , and  $\gamma_q$  are positive roots of the transcendental equations

$$\begin{aligned} \alpha^2 - s_1 s_2 &= (s_1 + s_2) \alpha \cot(l_1 \alpha), \\ \beta^2 - s_3 s_4 &= (s_3 + s_4) \beta \cot(l_2 \beta), \\ \gamma^2 - s_5 s_6 &= (s_5 + s_6) \gamma \cot(l_3 \gamma). \end{aligned}$$

1°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 \partial_t w &= f_1(x, y, z) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\
 w &= g_1(y, z, t) \quad \text{at} \quad x = 0 \quad (\text{boundary condition}), \\
 w &= g_2(y, z, t) \quad \text{at} \quad x = l_1 \quad (\text{boundary condition}), \\
 \partial_y w &= g_3(x, z, t) \quad \text{at} \quad y = 0 \quad (\text{boundary condition}), \\
 \partial_y w &= g_4(x, z, t) \quad \text{at} \quad y = l_2 \quad (\text{boundary condition}), \\
 \partial_z w &= g_5(x, y, t) \quad \text{at} \quad z = 0 \quad (\text{boundary condition}), \\
 \partial_z w &= g_6(x, y, t) \quad \text{at} \quad z = l_3 \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 &+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\
 &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=l_1} d\eta d\zeta d\tau \\
 &- a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) G(x, y, z, \xi, 0, \zeta, t - \tau) d\xi d\zeta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\
 &- a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_5(\xi, \eta, \tau) G(x, y, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_2} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau \\
 &+ \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta, t) &= \frac{2}{l_1 l_2 l_3} \exp\left(-\frac{1}{2}kt\right) \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \sum_{s=0}^{\infty} \frac{A_m A_s}{\sqrt{\lambda_{nms}}} \sin(\alpha_n x) \cos(\beta_m y) \cos(\gamma_s z) \\
 &\quad \times \sin(\alpha_n \xi) \cos(\beta_m \eta) \cos(\gamma_s \zeta) \sin\left(t\sqrt{\lambda_{nms}}\right), \\
 A_m &= \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m > 0, \end{cases} \quad \alpha_n = \frac{n\pi}{l_1}, \quad \beta_m = \frac{m\pi}{l_2}, \quad \gamma_s = \frac{s\pi}{l_3}, \\
 \lambda_{nms} &= a^2(\alpha_n^2 + \beta_m^2 + \gamma_s^2) + b - \frac{1}{4}k^2.
 \end{aligned}$$

2°. A rectangular parallelepiped is considered. The following conditions are prescribed:

$$\begin{aligned}
 w &= f_0(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 \partial_t w &= f_1(x, y, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\
 w &= g_1(y, z, t) & \text{at } x = 0 & \quad (\text{boundary condition}), \\
 \partial_x w &= g_2(y, z, t) & \text{at } x = l_1 & \quad (\text{boundary condition}), \\
 w &= g_3(x, z, t) & \text{at } y = 0 & \quad (\text{boundary condition}), \\
 \partial_y w &= g_4(x, z, t) & \text{at } y = l_2 & \quad (\text{boundary condition}), \\
 w &= g_5(x, y, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\
 \partial_z w &= g_6(x, y, t) & \text{at } z = l_3 & \quad (\text{boundary condition}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z, t) &= \frac{\partial}{\partial t} \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} f_0(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 &+ \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(x, y, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=0} d\eta d\zeta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_2} g_2(\eta, \zeta, \tau) G(x, y, z, l_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_3(\xi, \zeta, \tau) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_4(\xi, \zeta, \tau) G(x, y, z, \xi, l_2, \zeta, t - \tau) d\xi d\zeta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\
 &+ a^2 \int_0^t \int_0^{l_3} \int_0^{l_1} g_6(\xi, \eta, \tau) G(x, y, z, \xi, \eta, l_3, t - \tau) d\xi d\eta d\tau \\
 &+ \int_0^t \int_0^{l_3} \int_0^{l_2} \int_0^{l_1} \Phi(\xi, \eta, \zeta, \tau) G(x, y, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta, t) &= \frac{8}{l_1 l_2 l_3} \exp(-\tfrac{1}{2}kt) \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{1}{\sqrt{\lambda_{nms}}} \sin(\alpha_n x) \sin(\beta_m y) \sin(\gamma_s z) \\
 &\quad \times \sin(\alpha_n \xi) \sin(\beta_m \eta) \sin(\gamma_s \zeta) \sin(t\sqrt{\lambda_{nms}}), \\
 \alpha_n &= \frac{\pi(2n+1)}{2l_1}, \quad \beta_m = \frac{\pi(2m+1)}{2l_2}, \quad \gamma_s = \frac{\pi(2s+1)}{2l_3}, \quad \lambda_{nms} = a^2(\alpha_n^2 + \beta_m^2 + \gamma_s^2) + b - \tfrac{1}{4}k^2.
 \end{aligned}$$

## 6.4.2. Problems in Cylindrical Coordinates

A three-dimensional nonhomogeneous telegraph equation in the cylindrical coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left[ \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} \right] - bw + \Phi(r, \varphi, z, t), \quad r = \sqrt{x^2 + y^2}.$$

One-dimensional problems with axial symmetry that have solutions  $w = w(r, t)$  are treated in Subsection 4.4.2. Two-dimensional problems whose solutions have the form  $w = w(r, \varphi, t)$  or  $w = w(r, z, t)$  are considered in Subsections 5.4.2 and 5.4.3.

6.4.2-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{2e^{-kt/2}}{\pi R^2 l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] \\ &\quad \times \sin\left(\frac{s\pi z}{l}\right) \sin\left(\frac{s\pi \zeta}{l}\right) \frac{\sin\left(t\sqrt{\lambda_{nms}}\right)}{\sqrt{\lambda_{nms}}}, \end{aligned}$$

where

$$\lambda_{nms} = a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4} k^2, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}$$

the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument), and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

6.4.2-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + \int_0^l \int_0^{2\pi} \int_0^R \xi [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & - a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\
 & + a^2 \int_0^t \int_0^l \int_0^{2\pi} \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau \\
 & + \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \exp\left(-\frac{1}{2}kt\right) \left[ \frac{\sin(t\sqrt{c})}{\pi R^2 l \sqrt{c}} + \frac{2}{\pi R^2 l} \sum_{s=1}^{\infty} \frac{1}{\sqrt{\beta_s}} \cos\left(\frac{s\pi x}{l}\right) \cos\left(\frac{s\pi \xi}{l}\right) \sin(t\sqrt{\beta_s}) \right. \\
 & \left. + \frac{1}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^{\infty} \frac{A_n A_s \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)] \cos\left(\frac{s\pi x}{l}\right) \cos\left(\frac{s\pi \xi}{l}\right) \frac{\sin(\lambda_{nms} t)}{\lambda_{nms}} \right], \\
 c = b - \frac{k^2}{4}, \quad \beta_s = \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{k^2}{4}, \quad \lambda_{nms} = \sqrt{a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{k^2}{4}}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases}
 \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

6.4.2-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.

A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi, z) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(r, \varphi, z) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_r w + s_1 w = g(\varphi, z, t) \quad & \text{at } r = R \quad (\text{boundary condition}), \\
 \partial_z w - s_2 w = g_2(r, \varphi, t) \quad & \text{at } z = 0 \quad (\text{boundary condition}), \\
 \partial_z w + s_3 w = g_3(r, \varphi, t) \quad & \text{at } z = l \quad (\text{boundary condition}).
 \end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 6.4.2-2 where

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{1}{\pi} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{p=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 + s_1^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \\
 & \times \cos[n(\varphi - \eta)] \frac{h_p(z) h_p(\zeta)}{\|h_p\|^2} \frac{\sin(t\sqrt{\lambda_{nmp}})}{\sqrt{\lambda_{nmp}}}.
 \end{aligned}$$

Here, the  $J_n(\xi)$  are the Bessel functions,

$$\begin{aligned}
 A_n = & \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \quad \lambda_{nmp} = a^2 \mu_{nm}^2 + a^2 \beta_p^2 + b - \frac{1}{4}k^2, \\
 h_p(z) = & \cos(\beta_p z) + \frac{s_2}{\beta_p} \sin(\beta_p z), \quad \|h_p\|^2 = \frac{s_3}{2\beta_p^2} \frac{\beta_p^2 + s_2^2}{\beta_p^2 + s_3^2} + \frac{s_2}{2\beta_p^2} + \frac{l}{2} \left(1 + \frac{s_2^2}{\beta_p^2}\right);
 \end{aligned}$$



the  $\mu_{nm}$  and  $\beta_p$  are positive roots of the transcendental equations

$$\mu J'_n(\mu R) + s_1 J_n(\mu R) = 0, \quad \frac{\tan(\beta l)}{\beta} = \frac{s_2 + s_3}{\beta^2 - s_2 s_3}.$$

6.4.2-4. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.

1°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ \partial_z w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_0^R \xi [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{1}{\pi R^2 l} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^{\infty} \frac{A_n A_s}{[J'_n(\mu_{nm} R)]^2 \sqrt{\lambda_{nms}}} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\ &\quad \times \cos[n(\varphi - \eta)] \cos\left(\frac{s\pi z}{l}\right) \cos\left(\frac{s\pi \zeta}{l}\right) \sin\left(t\sqrt{\lambda_{nms}}\right), \\ \lambda_{nms} &= a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

2°. A circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_0^R \xi f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + \int_0^l \int_0^{2\pi} \int_0^R \xi [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) d\xi d\eta d\zeta \\
 & + a^2 R \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
 & + a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_2(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} d\xi d\eta d\tau \\
 & - a^2 \int_0^t \int_0^{2\pi} \int_0^R \xi g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} d\xi d\eta d\tau \\
 & + \int_0^t \int_0^l \int_0^{2\pi} \int_0^R \xi \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) d\xi d\eta d\zeta d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{2}{\pi R^2 l} \exp(-\tfrac{1}{2}kt) \sum_{s=1}^{\infty} \frac{1}{\sqrt{\beta_s}} \sin\left(\frac{s\pi z}{l}\right) \sin\left(\frac{s\pi \zeta}{l}\right) \sin(t\sqrt{\beta_s}) \\
 & + \frac{2}{\pi l} \exp(-\tfrac{1}{2}kt) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2}{(\mu_{nm}^2 R^2 - n^2)[J_n(\mu_{nm} R)]^2} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\
 & \quad \times \cos[n(\varphi - \eta)] \sin\left(\frac{s\pi z}{l}\right) \sin\left(\frac{s\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nms}})}{\sqrt{\lambda_{nms}}}, \\
 \beta_s = & \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n > 0, \end{cases} \\
 \lambda_{nms} = & a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2,
 \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

6.4.2-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . First boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(r, \varphi, z) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 \partial_t w = f_1(r, \varphi, z) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
 w = g_1(\varphi, z, t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
 w = g_2(\varphi, z, t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition),} \\
 w = g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad & \text{(boundary condition),} \\
 w = g_4(r, \varphi, t) \quad \text{at} \quad z = l \quad & \text{(boundary condition).}
 \end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + a^2 R_1 \int_0^t \int_0^l \int_0^{2\pi} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\
& - a^2 R_2 \int_0^t \int_0^l \int_0^{2\pi} g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\
& + a^2 \int_0^t \int_0^l \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\
& - a^2 \int_0^t \int_0^l \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau \\
& + \int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{\pi}{2l} \exp(-\tfrac{1}{2}kt) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)} Z_{nm}(r) Z_{nm}(\xi) \\
& \times \cos[n(\varphi - \eta)] \sin\left(\frac{s\pi z}{l}\right) \sin\left(\frac{s\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nms}})}{\sqrt{\lambda_{nms}}}, \\
A_n = & \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nms} = a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4} k^2, \\
Z_{nm}(r) = & J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),
\end{aligned}$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

6.4.2-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Second boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(r, \varphi, z) & \quad \text{at } t = 0 & \text{(initial condition),} \\
\partial_t w = f_1(r, \varphi, z) & \quad \text{at } t = 0 & \text{(initial condition),} \\
\partial_r w = g_1(\varphi, z, t) & \quad \text{at } r = R_1 & \text{(boundary condition),} \\
\partial_r w = g_2(\varphi, z, t) & \quad \text{at } r = R_2 & \text{(boundary condition),} \\
\partial_z w = g_3(r, \varphi, t) & \quad \text{at } z = 0 & \text{(boundary condition),} \\
\partial_z w = g_4(r, \varphi, t) & \quad \text{at } z = l & \text{(boundary condition).}
\end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \varphi, z, t) = & \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& + \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\
& - a^2 R_1 \int_0^t \int_0^{2\pi} \int_0^{R_2} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& + a^2 R_2 \int_0^t \int_0^{2\pi} \int_0^{R_2} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\
& - a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\
& + a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau \\
& + \int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau.
\end{aligned}$$

Here,

$$\begin{aligned}
G(r, \varphi, z, \xi, \eta, \zeta, t) = & \frac{e^{-kt/2}}{\pi(R_2^2 - R_1^2)l} \left[ \frac{\sin(t\sqrt{c})}{\sqrt{c}} + 2 \sum_{s=1}^{\infty} \cos\left(\frac{s\pi z}{l}\right) \cos\left(\frac{s\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\beta_s})}{\sqrt{\beta_s}} \right] \\
& + \frac{e^{-kt/2}}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^{\infty} \frac{A_n A_s \mu_{nm}^2 Z_{nm}(r) Z_{nm}(\xi)}{(\mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)} \\
& \times \cos[n(\varphi - \eta)] \cos\left(\frac{s\pi z}{l}\right) \cos\left(\frac{s\pi \zeta}{l}\right) \frac{\sin(t\sqrt{\lambda_{nms}})}{\sqrt{\lambda_{nms}}},
\end{aligned}$$

where

$$\begin{aligned}
A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad c = b - \frac{1}{4}k^2, \quad \beta_s = \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2, \quad \lambda_{nms} = a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2, \\
Z_{nm}(r) = J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r);
\end{aligned}$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

6.4.2-7. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Third boundary value problem.

A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(r, \varphi, z) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
\partial_t w = f_1(r, \varphi, z) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
\partial_r w - s_1 w = g_1(\varphi, z, t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
\partial_r w + s_2 w = g_2(\varphi, z, t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition),} \\
\partial_z w - s_3 w = g_3(r, \varphi, t) \quad \text{at} \quad z = 0 \quad & \text{(boundary condition),} \\
\partial_z w + s_4 w = g_4(r, \varphi, t) \quad \text{at} \quad z = l \quad & \text{(boundary condition).}
\end{aligned}$$

The solution  $w(r, \varphi, z, t)$  is determined by the formula in Paragraph 6.4.2-6 where

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{1}{\pi} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{p=1}^{\infty} \frac{A_n \mu_{nm}^2}{\|h_p\|^2 \sqrt{a^2 \mu_{nm}^2 + a^2 \lambda_p^2 + b - k^2/4}} \\ \times \frac{Z_{nm}(r) Z_{nm}(\xi) \cos[n(\varphi - \eta)] h_p(z) h_p(\zeta) \sin\left(t \sqrt{a^2 \mu_{nm}^2 + a^2 \lambda_p^2 + b - k^2/4}\right)}{(s_2^2 R_2^2 + \mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (s_1^2 R_1^2 + \mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)}.$$

Here,

$$Z_{nm}(r) = [\mu_{nm} J'_n(\mu_{nm} R_1) - s_1 J_n(\mu_{nm} R_1)] Y_n(\mu_{nm} r) - [\mu_{nm} Y'_n(\mu_{nm} R_1) - s_1 Y_n(\mu_{nm} R_1)] J_n(\mu_{nm} r), \\ A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad h_p(z) = \cos(\lambda_p z) + \frac{s_3}{\lambda_p} \sin(\lambda_p z), \quad \|h_p\|^2 = \frac{s_4}{2\lambda_p^2} \frac{\lambda_p^2 + s_3^2}{\lambda_p^2 + s_4^2} + \frac{s_3}{2\lambda_p^2} + \frac{l}{2} \left(1 + \frac{s_3^2}{\lambda_p^2}\right),$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, the  $\mu_{nm}$  are positive roots of the transcendental equation

$$[\mu J'_n(\mu R_1) - s_1 J_n(\mu R_1)] [\mu Y'_n(\mu R_2) + s_2 Y_n(\mu R_2)] = [\mu Y'_n(\mu R_1) - s_1 Y_n(\mu R_1)] [\mu J'_n(\mu R_2) + s_2 J_n(\mu R_2)],$$

and the  $\lambda_p$  are positive roots of the transcendental equation  $\frac{\tan(\lambda l)}{\lambda} = \frac{s_3 + s_4}{\lambda^2 - s_3 s_4}$ .

6.4.2-8. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq l$ . Mixed boundary value problems.

1°. A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(\varphi, z, t) & \text{at } r = R_2 & \quad (\text{boundary condition}), \\ \partial_z w &= g_3(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ a^2 R_1 \int_0^t \int_0^{2\pi} \int_0^l g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} d\eta d\zeta d\tau \\ &- a^2 R_2 \int_0^t \int_0^{2\pi} \int_0^l g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} d\eta d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau \\ &+ \int_0^t \int_0^{2\pi} \int_0^l \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta, t) = \frac{\pi}{4l} \exp\left(-\frac{1}{2}kt\right) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^{\infty} \frac{A_n A_s \mu_{nm}^2 J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)} Z_{nm}(r) Z_{nm}(\xi) \\ \times \cos[n(\varphi - \eta)] \cos\left(\frac{s\pi z}{l}\right) \cos\left(\frac{s\pi \zeta}{l}\right) \frac{\sin\left(t\sqrt{\lambda_{nms}}\right)}{\sqrt{\lambda_{nms}}}, \\ A_n = \begin{cases} 1 & \text{for } n=0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \lambda_{nms} = a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2, \\ Z_{nm}(r) = J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y_n(\mu_{nm} R_1) J_n(\mu_{nm} r),$$

where the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - Y_n(\mu R_1) J_n(\mu R_2) = 0.$$

2°. A hollow circular cylinder of finite length is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t=0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t=0 & \quad (\text{initial condition}), \\ \partial_r w &= g_1(\varphi, z, t) & \text{at } r=R_1 & \quad (\text{boundary condition}), \\ \partial_r w &= g_2(\varphi, z, t) & \text{at } r=R_2 & \quad (\text{boundary condition}), \\ w &= g_3(r, \varphi, t) & \text{at } z=0 & \quad (\text{boundary condition}), \\ w &= g_4(r, \varphi, t) & \text{at } z=l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R_1 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_1(\eta, \zeta, \tau) G(r, \varphi, z, R_1, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 R_2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_2(\eta, \zeta, \tau) G(r, \varphi, z, R_2, \eta, \zeta, t - \tau) d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_3(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^{2\pi} \int_{R_1}^{R_2} g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{2e^{-kt/2}}{\pi(R_2^2 - R_1^2)l} \sum_{s=1}^{\infty} \sin\left(\frac{s\pi z}{l}\right) \sin\left(\frac{s\pi \zeta}{l}\right) \frac{\sin\left(t\sqrt{\beta_s}\right)}{\sqrt{\beta_s}} \\ &+ \frac{2e^{-kt/2}}{\pi l} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2 Z_{nm}(r) Z_{nm}(\xi)}{(\mu_{nm}^2 R_2^2 - n^2) Z_{nm}^2(R_2) - (\mu_{nm}^2 R_1^2 - n^2) Z_{nm}^2(R_1)} \\ &\times \cos[n(\varphi - \eta)] \sin\left(\frac{s\pi z}{l}\right) \sin\left(\frac{s\pi \zeta}{l}\right) \frac{\sin\left(t\sqrt{\lambda_{nms}}\right)}{\sqrt{\lambda_{nms}}}, \end{aligned}$$

where

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad \beta_s = \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2, \quad \lambda_{nms} = a^2 \mu_{nm}^2 + \frac{a^2 s^2 \pi^2}{l^2} + b - \frac{1}{4}k^2,$$

$$Z_{nm}(r) = J'_n(\mu_{nm} R_1) Y_n(\mu_{nm} r) - Y'_n(\mu_{nm} R_1) J_n(\mu_{nm} r);$$

the  $J_n(r)$  and  $Y_n(r)$  are the Bessel functions, and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J'_n(\mu R_1) Y'_n(\mu R_2) - Y'_n(\mu R_1) J'_n(\mu R_2) = 0.$$

6.4.2-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq l$ . First boundary value problem.

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r &= R & (\text{boundary condition}), \\ w &= g_2(r, z, t) & \text{at } \varphi &= 0 & (\text{boundary condition}), \\ w &= g_3(r, z, t) & \text{at } \varphi &= \varphi_0 & (\text{boundary condition}), \\ w &= g_4(r, \varphi, t) & \text{at } z &= 0 & (\text{boundary condition}), \\ w &= g_5(r, \varphi, t) & \text{at } z &= l & (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{\varphi_0} \int_0^R [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=0} \xi d\xi d\eta d\tau \\ &- a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_5(\xi, \eta, \tau) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\zeta=l} \xi d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{8e^{-kt/2}}{R^2 l \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \\ &\times \sin\left(\frac{s\pi z}{l}\right) \sin\left(\frac{s\pi\zeta}{l}\right) \frac{\sin\left(t\sqrt{a^2\mu_{nm}^2 + a^2s^2\pi^2l^{-2} + b - k^2/4}\right)}{\sqrt{a^2\mu_{nm}^2 + a^2s^2\pi^2l^{-2} + b - k^2/4}}, \end{aligned}$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

A cylindrical sector of finite thickness is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \varphi, z) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\varphi, z, t) & \text{at } r = R & \quad (\text{boundary condition}), \\ w &= g_2(r, z, t) & \text{at } \varphi = 0 & \quad (\text{boundary condition}), \\ w &= g_3(r, z, t) & \text{at } \varphi = \varphi_0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_4(r, \varphi, t) & \text{at } z = 0 & \quad (\text{boundary condition}), \\ \partial_z w &= g_5(r, \varphi, t) & \text{at } z = l & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \varphi, z, t) &= \frac{\partial}{\partial t} \int_0^l \int_0^{\varphi_0} \int_0^R f_0(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &+ \int_0^l \int_0^{\varphi_0} \int_0^R [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \varphi, z, \xi, \eta, \zeta, t) \xi d\xi d\eta d\zeta \\ &- a^2 R \int_0^t \int_0^l \int_0^{\varphi_0} g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} d\eta d\zeta d\tau \\ &+ a^2 \int_0^t \int_0^l \int_0^R g_2(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^l \int_0^R g_3(\xi, \zeta, \tau) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \right]_{\eta=\varphi_0} d\xi d\zeta d\tau \\ &- a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_4(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, 0, t - \tau) \xi d\xi d\eta d\tau \\ &+ a^2 \int_0^t \int_0^{\varphi_0} \int_0^R g_5(\xi, \eta, \tau) G(r, \varphi, z, \xi, \eta, l, t - \tau) \xi d\xi d\eta d\tau \\ &+ \int_0^t \int_0^l \int_0^{\varphi_0} \int_0^R \Phi(\xi, \eta, \zeta, \tau) G(r, \varphi, z, \xi, \eta, \zeta, t - \tau) \xi d\xi d\eta d\zeta d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta, t) &= \frac{4e^{-kt/2}}{R^2 l \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^{\infty} \frac{A_s J_{n\pi/\varphi_0}(\mu_{nm} r) J_{n\pi/\varphi_0}(\mu_{nm} \xi) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right)}{[J'_{n\pi/\varphi_0}(\mu_{nm} R)]^2} \\ &\times \cos\left(\frac{s\pi z}{l}\right) \cos\left(\frac{s\pi\zeta}{l}\right) \frac{\sin\left(t\sqrt{a^2\mu_{nm}^2 + a^2s^2\pi^2l^{-2} + b - k^2/4}\right)}{\sqrt{a^2\mu_{nm}^2 + a^2s^2\pi^2l^{-2} + b - k^2/4}}, \end{aligned}$$

where  $A_0 = 1$  and  $A_s = 2$  for  $s \geq 1$ ; the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

### 6.4.3. Problems in Spherical Coordinates

A three-dimensional nonhomogeneous telegraph equation in the spherical coordinate system is written as

$$\frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} \right] - bw + \Phi(r, \theta, \varphi, t).$$



**6.4.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_0^R f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &+ \int_0^{2\pi} \int_0^\pi \int_0^R [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &- a^2 R^2 \int_0^t \int_0^{2\pi} \int_0^\pi g(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R} \sin \eta \, d\eta \, d\zeta \, d\tau \\ &+ \int_0^t \int_0^{2\pi} \int_0^\pi \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau, \end{aligned}$$

where

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta, t) &= \frac{1}{2\pi R^2 \sqrt{r\xi}} \exp(-\tfrac{1}{2}kt) \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n A_s B_{nms} J_{n+1/2}(\lambda_{nm}r) J_{n+1/2}(\lambda_{nm}\xi) \\ &\quad \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2\lambda_{nm}^2 + b - k^2/4})}{\sqrt{a^2\lambda_{nm}^2 + b - k^2/4}}, \\ A_s &= \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{(2n+1)(n-s)!}{(n+s)! [J'_{n+1/2}(\lambda_{nm}R)]^2}. \end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as

$$P_n^s(\mu) = (1 - \mu^2)^{s/2} \frac{d^s}{d\mu^s} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n,$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $J_{n+1/2}(\lambda R) = 0$ .

**6.4.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_r w &= g(\theta, \varphi, t) \quad \text{at} \quad r = R \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned} w(r, \theta, \varphi, t) &= \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_0^R f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &+ \int_0^{2\pi} \int_0^\pi \int_0^R [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\ &+ a^2 R^2 \int_0^t \int_0^{2\pi} \int_0^\pi g(\eta, \zeta, \tau) G(r, \theta, \varphi, R, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\ &+ \int_0^t \int_0^{2\pi} \int_0^\pi \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau, \end{aligned}$$

where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{3e^{-kt/2}}{4\pi R^3} \frac{\sin(t\sqrt{c})}{\sqrt{c}} + \frac{e^{-kt/2}}{2\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n A_s B_{nms} J_{n+1/2}(\lambda_{nm}r) \\ \times J_{n+1/2}(\lambda_{nm}\xi) P_n^s(\cos\theta) P_n^s(\cos\eta) \cos[s(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2\lambda_{nm}^2 + c})}{\sqrt{a^2\lambda_{nm}^2 + c}}, \\ A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{\lambda_{nm}^2 (2n+1)(n-s)!}{(n+s)! [R^2\lambda_{nm}^2 - n(n+1)] [J_{n+1/2}(\lambda_{nm}R)]^2}, \quad c = b - \frac{1}{4}k^2.$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 6.4.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$2\lambda R J'_{n+1/2}(\lambda R) - J_{n+1/2}(\lambda R) = 0.$$

6.4.3-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical domain is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w + sw &= g(\theta, \varphi, t) & \text{at } r = R & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 6.4.3-2 where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{e^{-kt/2}}{2\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{l=0}^n A_l B_{nml} J_{n+1/2}(\lambda_{nm}r) J_{n+1/2}(\lambda_{nm}\xi) \\ \times P_n^l(\cos\theta) P_n^l(\cos\eta) \cos[l(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2\lambda_{nm}^2 + b - k^2/4})}{\sqrt{a^2\lambda_{nm}^2 + b - k^2/4}}, \\ A_l = \begin{cases} 1 & \text{for } l = 0, \\ 2 & \text{for } l \neq 0, \end{cases} \quad B_{nml} = \frac{\lambda_{nm}^2 (2n+1)(n-l)!}{(n+l)! [R^2\lambda_{nm}^2 + (sR+n)(sR-n-1)] [J_{n+1/2}(\lambda_{nm}R)]^2}.$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^l(\mu)$  are the associated Legendre functions (see Paragraph 6.4.3-1), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda R J'_{n+1/2}(\lambda R) + \left(sR - \frac{1}{2}\right) J_{n+1/2}(\lambda R) = 0.$$

6.4.3-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ w &= g_1(\theta, \varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ w &= g_2(\theta, \varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \theta, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + a^2 R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_1} \sin \eta \, d\eta \, d\zeta \, d\tau \\
& - a^2 R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) \left[ \frac{\partial}{\partial \xi} G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \right]_{\xi=R_2} \sin \eta \, d\eta \, d\zeta \, d\tau \\
& + \int_0^t \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau,
\end{aligned}$$

where

$$\begin{aligned}
G(r, \theta, \varphi, \xi, \eta, \zeta, t) = & \frac{\pi e^{-kt/2}}{8\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n A_s B_{nms} Z_{n+1/2}(\lambda_{nm} r) Z_{n+1/2}(\lambda_{nm} \xi) \\
& \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2 \lambda_{nm}^2 + b - k^2/4})}{\sqrt{a^2 \lambda_{nm}^2 + b - k^2/4}}, \\
Z_{n+1/2}(\lambda_{nm} r) = & J_{n+1/2}(\lambda_{nm} R_1) Y_{n+1/2}(\lambda_{nm} r) - Y_{n+1/2}(\lambda_{nm} R_1) J_{n+1/2}(\lambda_{nm} r), \\
A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = & \frac{\lambda_{nm}(2n+1)(n-s)! J_{n+1/2}^2(\lambda_{nm} R_2)}{(n+s)! [J_{n+1/2}^2(\lambda_{nm} R_1) - J_{n+1/2}^2(\lambda_{nm} R_2)]}.
\end{aligned}$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions expressed in terms of the Legendre polynomials  $P_n(\mu)$  as

$$P_n^s(\mu) = (1 - \mu^2)^{s/2} \frac{d^s}{d\mu^s} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n,$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $Z_{n+1/2}(\lambda R_2) = 0$ .

6.4.3-5. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned}
w = f_0(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
\partial_t w = f_1(r, \theta, \varphi) \quad \text{at} \quad t = 0 \quad & \text{(initial condition),} \\
\partial_r w = g_1(\theta, \varphi, t) \quad \text{at} \quad r = R_1 \quad & \text{(boundary condition),} \\
\partial_r w = g_2(\theta, \varphi, t) \quad \text{at} \quad r = R_2 \quad & \text{(boundary condition).}
\end{aligned}$$

Solution:

$$\begin{aligned}
w(r, \theta, \varphi, t) = & \frac{\partial}{\partial t} \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} f_0(\xi, \eta, \zeta) G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& + \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} [f_1(\xi, \eta, \zeta) + k f_0(\xi, \eta, \zeta)] G(r, \theta, \varphi, \xi, \eta, \zeta, t) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \\
& - a^2 R_1^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_1(\eta, \zeta, \tau) G(r, \theta, \varphi, R_1, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\
& + a^2 R_2^2 \int_0^t \int_0^{2\pi} \int_0^\pi g_2(\eta, \zeta, \tau) G(r, \theta, \varphi, R_2, \eta, \zeta, t - \tau) \sin \eta \, d\eta \, d\zeta \, d\tau \\
& + \int_0^t \int_0^{2\pi} \int_0^\pi \int_{R_1}^{R_2} \Phi(\xi, \eta, \zeta, \tau) G(r, \theta, \varphi, \xi, \eta, \zeta, t - \tau) \xi^2 \sin \eta \, d\xi \, d\eta \, d\zeta \, d\tau,
\end{aligned}$$

where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{3e^{-kt/2} \sin(t\sqrt{c})}{4\pi(R_2^3 - R_1^3)\sqrt{c}} + \frac{e^{-kt/2}}{4\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n \frac{A_s}{B_{nms}} Z_{n+1/2}(\lambda_{nm}r) Z_{n+1/2}(\lambda_{nm}\xi) \\ \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2\lambda_{nm}^2 + c})}{\sqrt{a^2\lambda_{nm}^2 + c}}.$$

Here,

$$A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{(n+s)!}{(2n+1)(n-s)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm}r) dr, \quad c = b - \frac{1}{4}k^2, \\ Z_{n+1/2}(\lambda_{nm}r) = \left[ \lambda_{nm} J'_{n+1/2}(\lambda_{nm}R_1) - \frac{1}{2R_1} J_{n+1/2}(\lambda_{nm}R_1) \right] Y_{n+1/2}(\lambda_{nm}r) \\ - \left[ \lambda_{nm} Y'_{n+1/2}(\lambda_{nm}R_1) - \frac{1}{2R_1} Y_{n+1/2}(\lambda_{nm}R_1) \right] J_{n+1/2}(\lambda_{nm}r),$$

where the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions (see Paragraph 6.4.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) - \frac{1}{2R_2} Z_{n+1/2}(\lambda R_2) = 0.$$

6.4.3-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical layer is considered. The following conditions are prescribed:

$$\begin{aligned} w &= f_0(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_t w &= f_1(r, \theta, \varphi) & \text{at } t = 0 & \quad (\text{initial condition}), \\ \partial_r w - s_1 w &= g_1(\theta, \varphi, t) & \text{at } r = R_1 & \quad (\text{boundary condition}), \\ \partial_r w + s_2 w &= g_2(\theta, \varphi, t) & \text{at } r = R_2 & \quad (\text{boundary condition}). \end{aligned}$$

The solution  $w(r, \theta, \varphi, t)$  is determined by the formula in Paragraph 6.4.3-5 where

$$G(r, \theta, \varphi, \xi, \eta, \zeta, t) = \frac{e^{-kt/2}}{4\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{l=0}^n \frac{A_l}{B_{nml}} Z_{n+1/2}(\lambda_{nm}r) Z_{n+1/2}(\lambda_{nm}\xi) \\ \times P_n^l(\cos \theta) P_n^l(\cos \eta) \cos[l(\varphi - \zeta)] \frac{\sin(t\sqrt{a^2\lambda_{nm}^2 + c})}{\sqrt{a^2\lambda_{nm}^2 + c}}.$$

Here,

$$A_l = \begin{cases} 1 & \text{for } l = 0, \\ 2 & \text{for } l \neq 0, \end{cases} \quad B_{nml} = \frac{(n+l)!}{(2n+1)(n-l)!} \int_{R_1}^{R_2} r Z_{n+1/2}^2(\lambda_{nm}r) dr, \quad c = b - \frac{1}{4}k^2, \\ Z_{n+1/2}(\lambda r) = \left[ \lambda J'_{n+1/2}(\lambda R_1) - \left( s_1 + \frac{1}{2R_1} \right) J_{n+1/2}(\lambda R_1) \right] Y_{n+1/2}(\lambda r) \\ - \left[ \lambda Y'_{n+1/2}(\lambda R_1) - \left( s_1 + \frac{1}{2R_1} \right) Y_{n+1/2}(\lambda R_1) \right] J_{n+1/2}(\lambda r),$$

where the  $J_{n+1/2}(r)$  and  $Y_{n+1/2}(r)$  are the Bessel functions, the  $P_n^l(\mu)$  are the associated Legendre functions (see Paragraph 6.4.3-4), and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda Z'_{n+1/2}(\lambda R_2) + \left( s_2 - \frac{1}{2R_2} \right) Z_{n+1/2}(\lambda R_2) = 0.$$

## 6.5. Other Equations with Three Space Variables

### 6.5.1. Equations Containing Arbitrary Parameters

$$1. \quad \frac{\partial^2 w}{\partial t^2} = \frac{\partial}{\partial x} \left( a x^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( b y^m \frac{\partial w}{\partial y} \right) + \frac{\partial}{\partial z} \left( c z^k \frac{\partial w}{\partial z} \right).$$

This equation admits separable solutions. In addition, for  $n \neq 2$ ,  $m \neq 2$ , and  $k \neq 2$ , there are particular solutions of the form

$$w = w(\xi, t), \quad \xi^2 = 4 \left[ \frac{x^{2-n}}{a(2-n)^2} + \frac{y^{2-m}}{b(2-m)^2} + \frac{z^{2-k}}{c(2-k)^2} \right],$$

where  $w(\xi, t)$  is determined by the one-dimensional nonstationary equation

$$\frac{\partial^2 w}{\partial t^2} = \frac{\partial^2 w}{\partial \xi^2} + \frac{A}{\xi} \frac{\partial w}{\partial \xi}, \quad A = 2 \left( \frac{1}{2-n} + \frac{1}{2-m} + \frac{1}{2-k} \right) - 1.$$

$$2. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} = a^2 \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} \right) + b_1 \frac{\partial w}{\partial x} + b_2 \frac{\partial w}{\partial y} + b_3 \frac{\partial w}{\partial z} + cw.$$

The transformation

$$w(x, y, z, t) = u(x, y, z, \tau) \exp \left( -\frac{1}{2} k t - \frac{b_1 x + b_2 y + b_3 z}{2a^2} \right), \quad \tau = at$$

leads to the equation in Subsection 6.3.1:

$$\frac{\partial^2 u}{\partial \tau^2} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} + \beta u, \quad \beta = \frac{c}{a^2} + \frac{k^2}{4a^2} - \frac{1}{4a^4} (b_1^2 + b_2^2 + b_3^2).$$

### 6.5.2. Equation of the Form

$$\rho(x, y, z) \frac{\partial^2 w}{\partial t^2} = \operatorname{div}[a(x, y, z) \nabla w] - q(x, y, z)w + \Phi(x, y, z, t)$$

Such equations are encountered when studying vibration of finite volumes. The equation is written using the notation

$$\operatorname{div}[a(\mathbf{r}) \nabla w] = \frac{\partial}{\partial x} \left[ a(\mathbf{r}) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ a(\mathbf{r}) \frac{\partial w}{\partial y} \right] + \frac{\partial}{\partial z} \left[ a(\mathbf{r}) \frac{\partial w}{\partial z} \right], \quad \mathbf{r} = \{x, y, z\}.$$

The problems for the equation in question are considered below for the interior of a bounded domain  $V$  with smooth surface  $S$ . In what follows, it is assumed that  $\rho(\mathbf{r}) > 0$ ,  $a(\mathbf{r}) > 0$ , and  $q(\mathbf{r}) \geq 0$ .

#### 6.5.2-1. First boundary value problem.

The solution of the equation in question with the initial conditions

$$\begin{aligned} w &= f_0(\mathbf{r}) \quad \text{at} \quad t = 0, \\ \partial_t w &= f_1(\mathbf{r}) \quad \text{at} \quad t = 0 \end{aligned} \tag{1}$$

and the nonhomogeneous boundary conditions of the first kind

$$w = g(\mathbf{r}, t) \quad \text{for} \quad \mathbf{r} \in S \tag{2}$$

can be written as the sum

$$w(\mathbf{r}, t) = \frac{\partial}{\partial t} \int_V f_0(\xi) \rho(\xi) \mathcal{G}(\mathbf{r}, \xi, t) dV_\xi + \int_V f_1(\xi) \rho(\xi) \mathcal{G}(\mathbf{r}, \xi, t) dV_\xi \\ - \int_0^t \int_S g(\xi, \tau) a(\xi) \left[ \frac{\partial}{\partial N_\xi} \mathcal{G}(\mathbf{r}, \xi, t - \tau) \right] dS_\xi d\tau + \int_0^t \int_V \Phi(\xi, \tau) \mathcal{G}(\mathbf{r}, \xi, t - \tau) dV_\xi d\tau. \quad (3)$$

Here, the modified Green's function is expressed as

$$\mathcal{G}(\mathbf{r}, \xi, t) = \sum_{n=1}^{\infty} \frac{1}{\sqrt{\lambda_n} \|u_n\|^2} u_n(\mathbf{r}) u_n(\xi) \sin(\sqrt{\lambda_n} t), \quad (4) \\ \|u_n\|^2 = \int_V \rho(\mathbf{r}) u_n^2(\mathbf{r}) dV, \quad \xi = \{\xi_1, \xi_2, \xi_3\},$$

where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the following second-order elliptic equation with homogeneous boundary conditions of the first kind:

$$\operatorname{div} [a(\mathbf{r}) \nabla u] - q(\mathbf{r}) u + \lambda \rho(\mathbf{r}) u = 0, \quad (5)$$

$$u = 0 \quad \text{for } \mathbf{r} \in S. \quad (6)$$

The integration in solution (3) is performed with respect to  $\xi_1, \xi_2$ , and  $\xi_3$ ;  $\frac{\partial}{\partial N_\xi}$  is the derivative along the outward normal to the surface  $S$  with respect to  $\xi_1, \xi_2$ , and  $\xi_3$ .

General properties of the Sturm–Liouville problem (5)–(6):

1°. There are finitely many eigenvalues. All eigenvalues are real and can be ordered so that  $\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$  and  $\lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ ; therefore the number of negative eigenvalues is finite.

2°. If  $\rho(\mathbf{r}) > 0$ ,  $a(\mathbf{r}) > 0$ , and  $q(\mathbf{r}) \geq 0$ , then all eigenvalues are positive,  $\lambda_n > 0$ .

3°. An eigenfunction is determined up to a constant multiplier. Two eigenfunctions  $u_n(\mathbf{r})$  and  $u_m(\mathbf{r})$  corresponding to different eigenvalues  $\lambda_n$  and  $\lambda_m$  are orthogonal with weight  $\rho(\mathbf{r})$  in the domain  $V$ , that is,

$$\int_V \rho(\mathbf{r}) u_n(\mathbf{r}) u_m(\mathbf{r}) dV = 0 \quad \text{for } n \neq m.$$

4°. An arbitrary function  $F(\mathbf{r})$  twice continuously differentiable and satisfying the boundary condition of the Sturm–Liouville problem ( $F = 0$  for  $\mathbf{r} \in S$ ) can be expanded into an absolutely and uniformly convergent series in the eigenfunctions:

$$F(\mathbf{r}) = \sum_{n=1}^{\infty} F_n u_n(\mathbf{r}), \quad F_n = \frac{1}{\|u_n\|^2} \int_V F(\mathbf{r}) \rho(\mathbf{r}) u_n(\mathbf{r}) dV,$$

where  $\|u_n\|^2$  is defined in (4).

*Remark.* In a three-dimensional problem, finitely many linearly independent eigenfunctions  $u_n^{(1)}, \dots, u_n^{(m)}$  generally correspond to each eigenvalue  $\lambda_n$ . These functions can always be replaced by their linear combinations

$$\tilde{u}_n^{(k)} = A_{k,1} u_n^{(1)} + \dots + A_{k,k-1} u_n^{(k-1)} + u_n^{(k)}, \quad k = 1, \dots, m,$$

so that  $\tilde{u}_n^{(1)}, \dots, \tilde{u}_n^{(m)}$  are now orthogonal pairwise. For this reason, without loss of generality, all eigenfunctions can be assumed orthogonal.

### 6.5.2-2. Second boundary value problem.

The solution of the equation with the initial conditions (1) and nonhomogeneous boundary conditions of the second kind,

$$\frac{\partial w}{\partial N} = g(\mathbf{r}, t) \quad \text{for } \mathbf{r} \in S,$$

can be represented as the sum

$$\begin{aligned} w(\mathbf{r}, t) = & \frac{\partial}{\partial t} \int_V f_0(\boldsymbol{\xi}) \rho(\boldsymbol{\xi}) \mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t) dV_{\boldsymbol{\xi}} + \int_V f_1(\boldsymbol{\xi}) \rho(\boldsymbol{\xi}) \mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t) dV_{\boldsymbol{\xi}} \\ & + \int_0^t \int_S g(\boldsymbol{\xi}, \tau) a(\boldsymbol{\xi}) \mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t - \tau) dS_{\boldsymbol{\xi}} d\tau + \int_0^t \int_V \Phi(\boldsymbol{\xi}, \tau) \mathcal{G}(\mathbf{r}, \boldsymbol{\xi}, t - \tau) dV_{\boldsymbol{\xi}} d\tau. \end{aligned} \quad (7)$$

Here, the modified Green's function  $\mathcal{G}$  is given by relation (4), the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and corresponding eigenfunctions of the Sturm–Liouville problem for the second-order elliptic equation (5) with homogeneous boundary conditions of the second kind,

$$\frac{\partial u}{\partial N} = 0 \quad \text{for } \mathbf{r} \in S. \quad (8)$$

For  $q(\mathbf{r}) > 0$ , the general properties of the eigenvalue problem (5), (8) are the same as those of the first boundary value problem (all  $\lambda_n$  are positive).

### 6.5.2-3. Third boundary value problem.

The solution of the equation with the initial conditions (1) and nonhomogeneous boundary conditions of the third kind,

$$\frac{\partial w}{\partial N} + k(\mathbf{r})w = g(\mathbf{r}, t) \quad \text{for } \mathbf{r} \in S,$$

is determined by relations (7) and (4), where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and eigenfunctions of the Sturm–Liouville problem for the second-order elliptic equation (5) with homogeneous boundary conditions of the third kind,

$$\frac{\partial u}{\partial N} + k(\mathbf{r})u = 0 \quad \text{for } \mathbf{r} \in S. \quad (9)$$

If  $q(\mathbf{r}) \geq 0$  and  $k(\mathbf{r}) > 0$ , the general properties of the eigenvalue problem (5), (9) are the same as those of the first boundary value problem (see Paragraph 6.5.2-1).

Suppose  $k(\mathbf{r}) = k = \text{const}$ . Denote the Green's functions of the second and third boundary value problems by  $G_2(\mathbf{r}, \boldsymbol{\xi}, t)$  and  $G_3(\mathbf{r}, \boldsymbol{\xi}, t, k)$ , respectively. If  $q(\mathbf{r}) > 0$ , the limit relation  $G_2(\mathbf{r}, \boldsymbol{\xi}, t) = \lim_{k \rightarrow 0} G_3(\mathbf{r}, \boldsymbol{\xi}, t, k)$  holds.

⊙ *References for Subsection 6.5.2:* V. S. Vladimirov (1988), A. D. Polyanin (2000a).

## 6.6. Equations with $n$ Space Variables

Throughout this section the following notation is used:

$$\Delta_n w = \sum_{k=1}^n \frac{\partial^2 w}{\partial x_k^2}, \quad \mathbf{x} = \{x_1, \dots, x_n\}, \quad \mathbf{y} = \{y_1, \dots, y_n\}, \quad |\mathbf{x}| = \sqrt{x_1^2 + \dots + x_n^2}.$$

## 6.6.1. Wave Equation $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_n w$

### 6.6.1-1. Fundamental solution:

$$\mathcal{E}(\mathbf{x}, t) = \begin{cases} \frac{(-1)^{\frac{n-2}{2}}}{2a\pi^{\frac{n+1}{2}}} \Gamma\left(\frac{n-1}{2}\right) \frac{\vartheta(at - |\mathbf{x}|)}{(a^2 t^2 - |\mathbf{x}|^2)^{\frac{n-1}{2}}} & \text{if } n \geq 2 \text{ is even;} \\ \frac{1}{2\pi a} \left(\frac{1}{2\pi a^2 t} \frac{\partial}{\partial t}\right)^{\frac{n-3}{2}} \delta(a^2 t^2 - |\mathbf{x}|^2) & \text{if } n \geq 3 \text{ is odd;} \end{cases}$$

where  $\vartheta(z)$  is the Heaviside unit step function and  $\delta(z)$  is the Dirac delta function.

⊙ *Reference:* V. S. Vladimirov (1988).

### 6.6.1-2. Properties of solutions.

Suppose  $w(x_1, \dots, x_n, t)$  is a solution of the wave equation. Then the functions

$$\begin{aligned} w_1 &= Aw(\pm\lambda x_1 + C_1, \dots, \pm\lambda x_n + C_n, \pm\lambda t + C_{n+1}), \\ w_2 &= Aw\left(\frac{x_1 - vt}{\sqrt{1 - (v/a)^2}}, x_2, \dots, x_n, \frac{t - va^{-2}x_1}{\sqrt{1 - (v/a)^2}}\right), \\ w_3 &= A|r^2 - a^2 t^2|^{-\frac{n-1}{2}} w\left(\frac{x_1}{r^2 - a^2 t^2}, \dots, \frac{x_n}{r^2 - a^2 t^2}, \frac{t}{r^2 - a^2 t^2}\right), \quad r = |\mathbf{x}|, \end{aligned}$$

are also solutions of this equation everywhere they are defined;  $A, C_1, \dots, C_{n+1}, v$ , and  $\lambda$  are arbitrary constants. The signs at  $\lambda$  in the expression of  $w_1$  can be taken independently of one another.

### 6.6.1-3. Domain: $-\infty < x_k < \infty$ ; $k = 1, \dots, n$ . Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) \quad \text{at } t = 0, \\ \partial_t w &= g(\mathbf{x}) \quad \text{at } t = 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \frac{1}{a^{n-1}(n-2)!} \frac{\partial^{n-1}}{\partial t^{n-1}} \int_0^{at} (a^2 t^2 - r^2)^{\frac{n-3}{2}} r T_r[f(\mathbf{x})] dr \\ &\quad + \frac{1}{a^{n-1}(n-2)!} \frac{\partial^{n-2}}{\partial t^{n-2}} \int_0^{at} (a^2 t^2 - r^2)^{\frac{n-3}{2}} r T_r[g(\mathbf{x})] dr. \end{aligned}$$

Here,  $T_r[f(\mathbf{x})]$  is the average of  $f$  over the surface of the sphere of radius  $r$  with center at  $\mathbf{x}$ :

$$T_r[f(\mathbf{x})] \equiv \frac{1}{\sigma_n r^{n-1}} \int_{|\mathbf{x}-\mathbf{y}|=r} f(\mathbf{y}) dS_y, \quad \sigma_n = \frac{2\pi^{n/2}}{\Gamma(n/2)},$$

where  $\sigma_n r^{n-1}$  is the area of the surface of an  $n$ -dimensional sphere of radius  $r$ ,  $dS_y$  is the area element of this surface, and  $|\mathbf{x} - \mathbf{y}|^2 = (x_1 - y_1)^2 + \dots + (x_n - y_n)^2$ .

For odd  $n$ , the solution can be alternatively represented as

$$\begin{aligned} w(\mathbf{x}, t) &= \frac{1}{1 \times 3 \dots (n-2)} \frac{\partial}{\partial t} \left( \frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-3}{2}} (t^{n-2} T_{at}[f(\mathbf{x})]) \\ &\quad + \frac{1}{1 \times 3 \dots (n-2)} \left( \frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-3}{2}} (t^{n-2} T_{at}[g(\mathbf{x})]). \end{aligned}$$



For even  $n$ , the solution can be alternatively represented as

$$w(\mathbf{x}, t) = \frac{1}{2 \times 4 \dots (n-2)a^{n-1}} \frac{\partial}{\partial t} \left( \frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \int_0^{at} T_r[f(\mathbf{x})] \frac{r^{n-1} dr}{\sqrt{a^2 t^2 - r^2}} \\ + \frac{1}{2 \times 4 \dots (n-2)a^{n-1}} \left( \frac{1}{t} \frac{\partial}{\partial t} \right)^{\frac{n-2}{2}} \int_0^{at} T_r[g(\mathbf{x})] \frac{r^{n-1} dr}{\sqrt{a^2 t^2 - r^2}}.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), R. Courant and D. Hilbert (1989), D. Zwilling (1998).

6.6.1-4. Domain:  $0 \leq x_k \leq l_k$ ;  $k = 1, \dots, n$ . Boundary value problems.

For solutions of the first, second, third, and mixed boundary value problems with nonhomogeneous conditions of general form, see Paragraphs 6.6.2-2, 6.6.2-3, 6.6.2-4, and 6.6.2-5 for  $\Phi \equiv 0$ , respectively.

## 6.6.2. Nonhomogeneous Wave Equation

$$\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_n w + \Phi(x_1, \dots, x_n, t)$$

6.6.2-1. Domain:  $-\infty < x_k < \infty$ ;  $k = 1, \dots, n$ . Cauchy problem.

Initial conditions are prescribed:

$$w = f(\mathbf{x}) \quad \text{at} \quad t = 0, \\ \partial_t w = g(\mathbf{x}) \quad \text{at} \quad t = 0.$$

Solution:

$$w(\mathbf{x}, t) = \frac{1}{a^{n-1}(n-2)!} \frac{\partial^{n-1}}{\partial t^{n-1}} \int_0^{at} (a^2 t^2 - r^2)^{\frac{n-3}{2}} r T_r[f(\mathbf{x})] dr \\ + \frac{1}{a^{n-1}(n-2)!} \frac{\partial^{n-2}}{\partial t^{n-2}} \int_0^{at} (a^2 t^2 - r^2)^{\frac{n-3}{2}} r T_r[g(\mathbf{x})] dr \\ + \frac{1}{a^{n-1}(n-2)!} \frac{\partial^{n-2}}{\partial t^{n-2}} \int_0^{at} d\tau \int_0^{a\tau} (a^2 \tau^2 - r^2)^{\frac{n-3}{2}} r T_r[\Phi(\mathbf{x}, t - \tau)] dr.$$

Here,  $T_r[f(\mathbf{x})]$  is the average of  $f$  over the spherical surface of radius  $r$  with center at  $\mathbf{x}$ :

$$T_r[f(\mathbf{x})] \equiv \frac{1}{\sigma_n r^{n-1}} \int_{|\mathbf{x}-\mathbf{y}|=r} f(\mathbf{y}) dS_y, \quad \sigma_n = \frac{2\pi^{n/2}}{\Gamma(n/2)},$$

where  $\sigma_n r^{n-1}$  is the area of the surface of an  $n$ -dimensional sphere of radius  $r$  and  $dS_y$  is the area element of this surface.

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), R. Courant and D. Hilbert (1989).

6.6.2-2. Domain:  $V = \{0 \leq x_k \leq l_k$ ;  $k = 1, \dots, n\}$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at} \quad t &= 0 & \text{(initial condition),} \\ \partial_t w &= f_1(\mathbf{x}) & \text{at} \quad t &= 0 & \text{(initial condition),} \\ w &= g_k(\mathbf{x}, t) & \text{at} \quad x_k &= 0 & \text{(boundary conditions),} \\ w &= h_k(\mathbf{x}, t) & \text{at} \quad x_k &= l_k & \text{(boundary conditions).} \end{aligned}$$

Solution:

$$\begin{aligned}
 w(\mathbf{x}, t) = & \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\
 & + \frac{\partial}{\partial t} \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V f_1(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\
 & + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\
 & - a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ h_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau,
 \end{aligned}$$

where

$$\begin{aligned}
 \mathbf{x} = \{x_1, \dots, x_n\}, \quad \mathbf{y} = \{y_1, \dots, y_n\}, \quad d\mathbf{y} = dy_1 dy_2 \dots dy_n, \quad dS_y^{(k)} = dy_1 \dots dy_{k-1} dy_{k+1} \dots dy_n, \\
 S^{(k)} = \{0 \leq y_m \leq l_m \text{ for } m = 1, \dots, k-1, k+1, \dots, n\}.
 \end{aligned}$$

Green's function:

$$\begin{aligned}
 G(\mathbf{x}, \mathbf{y}, t) = & \frac{2^n}{l_1 l_2 \dots l_n} \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \sin(\lambda_{s_1} x_1) \sin(\lambda_{s_2} x_2) \dots \sin(\lambda_{s_n} x_n) \\
 & \times \sin(\lambda_{s_1} y_1) \sin(\lambda_{s_2} y_2) \dots \sin(\lambda_{s_n} y_n) \frac{\sin(at \sqrt{\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2})}{a \sqrt{\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2}},
 \end{aligned}$$

where

$$\lambda_{s_1} = \frac{s_1 \pi}{l_1}, \quad \lambda_{s_2} = \frac{s_2 \pi}{l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{s_n \pi}{l_n}.$$

6.6.2-3. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(\mathbf{x}) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(\mathbf{x}) \quad & \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_{x_k} w = g_k(\mathbf{x}, t) \quad & \text{at } x_k = 0 \quad (\text{boundary conditions}), \\
 \partial_{x_k} w = h_k(\mathbf{x}, t) \quad & \text{at } x_k = l_k \quad (\text{boundary conditions}).
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(\mathbf{x}, t) = & \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\
 & + \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V f_1(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\
 & - a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [g_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=0} dS_y^{(k)} d\tau \\
 & + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=l_k} dS_y^{(k)} d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(\mathbf{x}, \mathbf{y}, t) = & \frac{t}{l_1 l_2 \dots l_n} + \frac{1}{l_1 l_2 \dots l_n} \sum_{s_1=0}^{\infty} \sum_{s_2=0}^{\infty} \dots \sum_{s_n=0}^{\infty} \frac{A_{s_1} A_{s_2} \dots A_{s_n}}{a \sqrt{\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2}} \sin(at \sqrt{\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2}) \\
 & \times \cos(\lambda_{s_1} x_1) \cos(\lambda_{s_2} x_2) \dots \cos(\lambda_{s_n} x_n) \cos(\lambda_{s_1} y_1) \cos(\lambda_{s_2} y_2) \dots \cos(\lambda_{s_n} y_n),
 \end{aligned}$$

where

$$\lambda_{s_1} = \frac{s_1\pi}{l_1}, \quad \lambda_{s_2} = \frac{s_2\pi}{l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{s_n\pi}{l_n}; \quad A_{s_m} = \begin{cases} 1 & \text{for } s_m = 0, \\ 2 & \text{for } s_m \neq 0, \end{cases} \quad m = 1, 2, \dots, n.$$

The summation is performed over the indices satisfying the condition  $s_1 + \dots + s_n > 0$ ; the term corresponding to  $s_1 = \dots = s_n = 0$  is singled out.

6.6.2-4. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_{x_k} w - b_k w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w + c_k w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

The solution  $w(\mathbf{x}, t)$  is determined by the formula in Paragraph 6.6.2-3 where

$$\begin{aligned} G(\mathbf{x}, \mathbf{y}, t) &= 2^n \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \frac{\sin(at \sqrt{\lambda_{s_1}^2 + \lambda_{s_2}^2 + \dots + \lambda_{s_n}^2})}{a E_{s_1} E_{s_2} \dots E_{s_n} \sqrt{\lambda_{s_1}^2 + \lambda_{s_2}^2 + \dots + \lambda_{s_n}^2}} \\ &\quad \times \sin(\lambda_{s_1} x_1 + \varphi_{s_1}) \sin(\lambda_{s_2} x_2 + \varphi_{s_2}) \dots \sin(\lambda_{s_n} x_n + \varphi_{s_n}) \\ &\quad \times \sin(\lambda_{s_1} y_1 + \varphi_{s_1}) \sin(\lambda_{s_2} y_2 + \varphi_{s_2}) \dots \sin(\lambda_{s_n} y_n + \varphi_{s_n}). \end{aligned}$$

Here,

$$\varphi_{s_m} = \arctan \frac{\lambda_{s_m}}{l_m}, \quad E_{s_m} = l_m + \frac{(b_m c_m + \lambda_{s_m}^2)(b_m + c_m)}{(b_m^2 + \lambda_{s_m}^2)(c_m^2 + \lambda_{s_m}^2)}, \quad m = 1, 2, \dots, n;$$

the  $\lambda_{s_m}$  are positive roots of the transcendental equations

$$\frac{1}{b_m + c_m} \left( \lambda - \frac{b_m c_m}{\lambda} \right) = \cot(l_m \lambda), \quad m = 1, 2, \dots, n.$$

6.6.2-5. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Mixed boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\ &\quad + \frac{\partial}{\partial t} \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V f_1(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ &\quad + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\ &\quad + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau, \end{aligned}$$

where

$$G(\mathbf{x}, \mathbf{y}, t) = \frac{2^n}{l_1 l_2 \dots l_n} \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \sin(\lambda_{s_1} x_1) \sin(\lambda_{s_2} x_2) \dots \sin(\lambda_{s_n} x_n) \\ \times \sin(\lambda_{s_1} y_1) \sin(\lambda_{s_2} y_2) \dots \sin(\lambda_{s_n} y_n) \frac{\sin(at \sqrt{\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2})}{a \sqrt{\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2}}, \\ \lambda_{s_1} = \frac{\pi(2s_1 + 1)}{2l_1}, \quad \lambda_{s_2} = \frac{\pi(2s_2 + 1)}{2l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{\pi(2s_n + 1)}{2l_n}.$$

### 6.6.3. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = a^2 \Delta_n w - bw + \Phi(x_1, \dots, x_n, t)$

6.6.3-1. Domain:  $-\infty < x_k < \infty$ ;  $k = 1, \dots, n$ . Cauchy problem.

Initial conditions are prescribed:

$$w = f(\mathbf{x}) \quad \text{at} \quad t = 0, \\ \partial_t w = g(\mathbf{x}) \quad \text{at} \quad t = 0,$$

where  $\mathbf{x} = \{x_1, \dots, x_n\}$ .

1°. Let  $b = -c^2 < 0$  and  $\Phi \equiv 0$ . The solution is sought by the descent method in the form

$$w(\mathbf{x}, t) = \frac{1}{\exp(cx_{n+1})} u(\mathbf{x}, x_{n+1}, t), \quad (1)$$

where  $u$  is the solution of the Cauchy problem for the auxiliary  $(n + 1)$ -dimensional wave equation

$$\frac{\partial^2 u}{\partial t^2} = \Delta_{n+1} u \quad (2)$$

with the initial conditions

$$u = \exp(cx_{n+1}) f(\mathbf{x}) \quad \text{at} \quad t = 0, \\ \partial_t u = \exp(cx_{n+1}) g(\mathbf{x}) \quad \text{at} \quad t = 0. \quad (3)$$

For the solution of problem (2), (3), see Paragraph 6.6.1-3.

2°. Let  $b = c^2 > 0$  and  $\Phi \equiv 0$ . In this case the function  $\exp(cx_{n+1})$  in (1) and (3) must be replaced by  $\cos(cx_{n+1})$ .

⊙ *Reference:* R. Courant and D. Hilbert (1989).

6.6.3-2. Domain:  $V = \{0 \leq x_k \leq l_k$ ;  $k = 1, \dots, n\}$ . First boundary value problem.

The following conditions are prescribed:

$$w = f_0(\mathbf{x}) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ \partial_t w = f_1(\mathbf{x}) \quad \text{at} \quad t = 0 \quad (\text{initial condition}), \\ w = g_k(\mathbf{x}, t) \quad \text{at} \quad x_k = 0 \quad (\text{boundary conditions}), \\ w = h_k(\mathbf{x}, t) \quad \text{at} \quad x_k = l_k \quad (\text{boundary conditions}).$$

Solution:

$$w(\mathbf{x}, t) = \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\ + \frac{\partial}{\partial t} \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V f_1(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\ - a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ h_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau,$$

where

$$\mathbf{x} = \{x_1, \dots, x_n\}, \quad \mathbf{y} = \{y_1, \dots, y_n\}, \quad d\mathbf{y} = dy_1 dy_2 \dots dy_n, \quad dS_y^{(k)} = dy_1 \dots dy_{k-1} dy_{k+1} \dots dy_n, \\ S^{(k)} = \{0 \leq y_m \leq l_m \text{ for } m = 1, \dots, k-1, k+1, \dots, n\}.$$

Green's function:

$$G(\mathbf{x}, \mathbf{y}, t) = \frac{2^n}{l_1 l_2 \dots l_n} \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \sin(\lambda_{s_1} x_1) \sin(\lambda_{s_2} x_2) \dots \sin(\lambda_{s_n} x_n) \\ \times \sin(\lambda_{s_1} y_1) \sin(\lambda_{s_2} y_2) \dots \sin(\lambda_{s_n} y_n) \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b})}{\sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b}},$$

where

$$\lambda_{s_1} = \frac{s_1 \pi}{l_1}, \quad \lambda_{s_2} = \frac{s_2 \pi}{l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{s_n \pi}{l_n}.$$

6.6.3-3. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_{x_k} w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\ &+ \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V f_1(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ &- a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [g_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=0} dS_y^{(k)} d\tau \\ &+ a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=l_k} dS_y^{(k)} d\tau. \end{aligned}$$

Here,

$$G(\mathbf{x}, \mathbf{y}, t) = \frac{1}{l_1 l_2 \dots l_n} \sum_{s_1=0}^{\infty} \sum_{s_2=0}^{\infty} \dots \sum_{s_n=0}^{\infty} A_{s_1} A_{s_2} \dots A_{s_n} \cos(\lambda_{s_1} x_1) \cos(\lambda_{s_2} x_2) \dots \cos(\lambda_{s_n} x_n) \\ \times \cos(\lambda_{s_1} y_1) \cos(\lambda_{s_2} y_2) \dots \cos(\lambda_{s_n} y_n) \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b})}{\sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b}},$$

where

$$\lambda_{s_1} = \frac{s_1 \pi}{l_1}, \quad \lambda_{s_2} = \frac{s_2 \pi}{l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{s_n \pi}{l_n}; \quad A_{s_m} = \begin{cases} 1 & \text{for } s_m = 0, \\ 2 & \text{for } s_m \neq 0, \end{cases} \quad m = 1, 2, \dots, n.$$

**6.6.3-4. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Third boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_{x_k} w - b_k w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w + c_k w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

The solution  $w(\mathbf{x}, t)$  is determined by the formula in Paragraph 6.6.3-3 where

$$\begin{aligned} G(\mathbf{x}, \mathbf{y}, t) &= 2^n \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \lambda_{s_2}^2 + \dots + \lambda_{s_n}^2) + b})}{E_{s_1} E_{s_2} \dots E_{s_n} \sqrt{a^2(\lambda_{s_1}^2 + \lambda_{s_2}^2 + \dots + \lambda_{s_n}^2) + b}} \\ &\quad \times \sin(\lambda_{s_1} x_1 + \varphi_{s_1}) \sin(\lambda_{s_2} x_2 + \varphi_{s_2}) \dots \sin(\lambda_{s_n} x_n + \varphi_{s_n}) \\ &\quad \times \sin(\lambda_{s_1} y_1 + \varphi_{s_1}) \sin(\lambda_{s_2} y_2 + \varphi_{s_2}) \dots \sin(\lambda_{s_n} y_n + \varphi_{s_n}). \end{aligned}$$

Here,

$$\varphi_{s_m} = \arctan \frac{\lambda_{s_m}}{l_m}, \quad E_{s_m} = l_m + \frac{(b_m c_m + \lambda_{s_m}^2)(b_m + c_m)}{(b_m^2 + \lambda_{s_m}^2)(c_m^2 + \lambda_{s_m}^2)}, \quad m = 1, 2, \dots, n;$$

the  $\lambda_{s_m}$  are positive roots of the transcendental equations

$$\frac{1}{b_m + c_m} \left( \lambda - \frac{b_m c_m}{\lambda} \right) = \cot(l_m \lambda), \quad m = 1, 2, \dots, n.$$

**6.6.3-5. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Mixed boundary value problem.**

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\ &\quad + \frac{\partial}{\partial t} \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V f_1(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ &\quad + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\ &\quad + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau. \end{aligned}$$

Here,

$$\begin{aligned} G(\mathbf{x}, \mathbf{y}, t) &= \frac{2^n}{l_1 l_2 \dots l_n} \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \sin(\lambda_{s_1} x_1) \sin(\lambda_{s_2} x_2) \dots \sin(\lambda_{s_n} x_n) \\ &\quad \times \sin(\lambda_{s_1} y_1) \sin(\lambda_{s_2} y_2) \dots \sin(\lambda_{s_n} y_n) \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b})}{\sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b}}, \end{aligned}$$

where

$$\lambda_{s_1} = \frac{\pi(2s_1 + 1)}{2l_1}, \quad \lambda_{s_2} = \frac{\pi(2s_2 + 1)}{2l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{\pi(2s_n + 1)}{2l_n}.$$

## 6.6.4. Equations Containing the First Time Derivative

1. 
$$\frac{\partial^2 w}{\partial t^2} + \beta \frac{\partial w}{\partial t} = a^2 \Delta_n w - bw + \Phi(x_1, \dots, x_n, t).$$

*Nonhomogeneous telegraph equation with  $n$  space variables.*

1°. The substitution  $w = \exp(-\frac{1}{2}\beta t)u$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} = a^2 \Delta_n u - (b - \frac{1}{4}\beta^2)u + \exp(\frac{1}{2}\beta t)\Phi(x_1, \dots, x_n, t),$$

which is considered in Subsection 6.6.3.

2°. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . First boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\ &+ \frac{\partial}{\partial t} \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V [f_1(\mathbf{y}) + \beta f_0(\mathbf{y})] G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ &+ a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\ &- a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ h_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau, \end{aligned}$$

where

$$\begin{aligned} \mathbf{x} &= \{x_1, \dots, x_n\}, \quad \mathbf{y} = \{y_1, \dots, y_n\}, \quad d\mathbf{y} = dy_1 dy_2 \dots dy_n, \quad dS_y^{(k)} = dy_1 \dots dy_{k-1} dy_{k+1} \dots dy_n, \\ S^{(k)} &= \{0 \leq y_m \leq l_m \text{ for } m = 1, \dots, k-1, k+1, \dots, n\}. \end{aligned}$$

Green's function:

$$\begin{aligned} G(\mathbf{x}, \mathbf{y}, t) &= \frac{2^n e^{-\beta t/2}}{l_1 l_2 \dots l_n} \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \sin(\lambda_{s_1} x_1) \sin(\lambda_{s_2} x_2) \dots \sin(\lambda_{s_n} x_n) \\ &\times \sin(\lambda_{s_1} y_1) \sin(\lambda_{s_2} y_2) \dots \sin(\lambda_{s_n} y_n) \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4})}{\sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4}}, \end{aligned}$$

where

$$\lambda_{s_1} = \frac{s_1 \pi}{l_1}, \quad \lambda_{s_2} = \frac{s_2 \pi}{l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{s_n \pi}{l_n}.$$

3°. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Second boundary value problem.

The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_{x_k} w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned}
 w(\mathbf{x}, t) = & \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\
 & + \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V [f_1(\mathbf{y}) + \beta f_0(\mathbf{y})] G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\
 & - a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [g_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=0} dS_y^{(k)} d\tau \\
 & + a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} [h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau)]_{y_k=l_k} dS_y^{(k)} d\tau.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(\mathbf{x}, \mathbf{y}, t) = & \frac{e^{-\beta t/2}}{l_1 l_2 \dots l_n} \sum_{s_1=0}^{\infty} \sum_{s_2=0}^{\infty} \dots \sum_{s_n=0}^{\infty} A_{s_1} A_{s_2} \dots A_{s_n} \cos(\lambda_{s_1} x_1) \cos(\lambda_{s_2} x_2) \dots \cos(\lambda_{s_n} x_n) \\
 & \times \cos(\lambda_{s_1} y_1) \cos(\lambda_{s_2} y_2) \dots \cos(\lambda_{s_n} y_n) \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4})}{\sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4}},
 \end{aligned}$$

where

$$\lambda_{s_1} = \frac{s_1 \pi}{l_1}, \quad \lambda_{s_2} = \frac{s_2 \pi}{l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{s_n \pi}{l_n}; \quad A_{s_m} = \begin{cases} 1 & \text{for } s_m = 0, \\ 2 & \text{for } s_m \neq 0, \end{cases} \quad m = 1, 2, \dots, n.$$

4°. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Third boundary value problem.

The following conditions are prescribed:

$$\begin{aligned}
 w = f_0(\mathbf{x}) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_t w = f_1(\mathbf{x}) & \quad \text{at } t = 0 \quad (\text{initial condition}), \\
 \partial_{x_k} w - b_k w = g_k(\mathbf{x}, t) & \quad \text{at } x_k = 0 \quad (\text{boundary conditions}), \\
 \partial_{x_k} w + c_k w = h_k(\mathbf{x}, t) & \quad \text{at } x_k = l_k \quad (\text{boundary conditions}).
 \end{aligned}$$

The solution  $w(\mathbf{x}, t)$  is given by the formula in Item 3° with

$$\begin{aligned}
 G(\mathbf{x}, \mathbf{y}, t) = & 2^n e^{-\beta t/2} \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \lambda_{s_2}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4})}{E_{s_1} E_{s_2} \dots E_{s_n} \sqrt{a^2(\lambda_{s_1}^2 + \lambda_{s_2}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4}} \\
 & \times \sin(\lambda_{s_1} x_1 + \varphi_{s_1}) \sin(\lambda_{s_2} x_2 + \varphi_{s_2}) \dots \sin(\lambda_{s_n} x_n + \varphi_{s_n}) \\
 & \times \sin(\lambda_{s_1} y_1 + \varphi_{s_1}) \sin(\lambda_{s_2} y_2 + \varphi_{s_2}) \dots \sin(\lambda_{s_n} y_n + \varphi_{s_n}).
 \end{aligned}$$

Here,

$$\varphi_{s_m} = \arctan \frac{\lambda_{s_m}}{l_m}, \quad E_{s_m} = l_m + \frac{(b_m c_m + \lambda_{s_m}^2)(b_m + c_m)}{(b_m^2 + \lambda_{s_m}^2)(c_m^2 + \lambda_{s_m}^2)}, \quad m = 1, 2, \dots, n;$$

the  $\lambda_{s_m}$  are positive roots of the transcendental equation

$$\frac{1}{b_m + c_m} \left( \lambda - \frac{b_m c_m}{\lambda} \right) = \cot(l_m \lambda), \quad m = 1, 2, \dots, n.$$



5°. Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, \dots, n\}$ . Mixed boundary value problem.  
The following conditions are prescribed:

$$\begin{aligned} w &= f_0(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ \partial_t w &= f_1(\mathbf{x}) & \text{at } t &= 0 & (\text{initial condition}), \\ w &= g_k(\mathbf{x}, t) & \text{at } x_k &= 0 & (\text{boundary conditions}), \\ \partial_{x_k} w &= h_k(\mathbf{x}, t) & \text{at } x_k &= l_k & (\text{boundary conditions}). \end{aligned}$$

Solution:

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_V \Phi(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) d\mathbf{y} d\tau \\ &+ \frac{\partial}{\partial t} \int_V f_0(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V [f_1(\mathbf{y}) + \beta f_0(\mathbf{y})] G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} \\ &+ a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ g_k(\mathbf{y}, \tau) \frac{\partial}{\partial y_k} G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=0} dS_y^{(k)} d\tau \\ &+ a^2 \sum_{k=1}^n \int_0^t \int_{S^{(k)}} \left[ h_k(\mathbf{y}, \tau) G(\mathbf{x}, \mathbf{y}, t - \tau) \right]_{y_k=l_k} dS_y^{(k)} d\tau, \end{aligned}$$

where

$$\begin{aligned} G(\mathbf{x}, \mathbf{y}, t) &= \frac{2^n e^{-\beta t/2}}{l_1 l_2 \dots l_n} \sum_{s_1=1}^{\infty} \sum_{s_2=1}^{\infty} \dots \sum_{s_n=1}^{\infty} \sin(\lambda_{s_1} x_1) \sin(\lambda_{s_2} x_2) \dots \sin(\lambda_{s_n} x_n) \\ &\times \sin(\lambda_{s_1} y_1) \sin(\lambda_{s_2} y_2) \dots \sin(\lambda_{s_n} y_n) \frac{\sin(t \sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4})}{\sqrt{a^2(\lambda_{s_1}^2 + \dots + \lambda_{s_n}^2) + b - \beta^2/4}}, \\ \lambda_{s_1} &= \frac{\pi(2s_1 + 1)}{2l_1}, \quad \lambda_{s_2} = \frac{\pi(2s_2 + 1)}{2l_2}, \quad \dots, \quad \lambda_{s_n} = \frac{\pi(2s_n + 1)}{2l_n}. \end{aligned}$$

$$2. \quad \frac{\partial^2 w}{\partial t^2} + \beta \frac{\partial w}{\partial t} = a^2 \Delta_n w + \sum_{k=1}^n b_k \frac{\partial w}{\partial x_k} + c w.$$

The transformation

$$w(x_1, \dots, x_n, t) = u(x_1, \dots, x_n, \tau) \exp\left(-\frac{1}{2}\beta t - \frac{1}{2a^2} \sum_{k=1}^n b_k x_k\right), \quad \tau = at$$

leads to the equation

$$\frac{\partial^2 u}{\partial \tau^2} = \Delta_n u + \lambda u, \quad \lambda = \frac{c}{a^2} + \frac{\beta^2}{4a^2} - \frac{1}{4a^4} \sum_{k=1}^n b_k^2,$$

which is considered in Subsection 6.6.3.

⊙ *Reference:* R. Courant and D. Hilbert (1989).

$$3. \quad \frac{\partial^2 w}{\partial t^2} + \frac{n-1}{t} \frac{\partial w}{\partial t} = \Delta_n w.$$

*Darboux equation.* Cauchy problem.

Initial conditions are prescribed:

$$\begin{aligned} w &= f(\mathbf{x}) & \text{at } t &= 0, \\ \partial_t w &= 0 & \text{at } t &= 0. \end{aligned}$$

Solution:

$$w(\mathbf{x}, t) = \frac{1}{\sigma_n t^{n-1}} \int_{|\mathbf{x}-\mathbf{y}|=t} f(\mathbf{y}) dS_y, \quad \sigma_n = \frac{2\pi^{n/2}}{\Gamma(n/2)},$$

where  $\sigma_n t^{n-1}$  is the area of the surface of an  $n$ -dimensional sphere of radius  $t$ , and  $dS_y$  is the area element of this surface (i.e., the solution  $w$  is the average of the function  $f$  over the sphere for radius  $t$  with center at  $\mathbf{x}$ ).

⊙ *Reference:* R. Courant and D. Hilbert (1989).

## Chapter 7

# Elliptic Equations with Two Space Variables

---

### 7.1. Laplace Equation $\Delta_2 w = 0$

The Laplace equation is often encountered in heat and mass transfer theory, fluid mechanics, elasticity, electrostatics, and other areas of mechanics and physics. For example, in heat and mass transfer theory, this equation describes steady-state temperature distribution in the absence of heat sources and sinks in the domain under study.

A regular solution of the Laplace equation is called a harmonic function. The first boundary value problem for the Laplace equation is often referred to as the Dirichlet problem, and the second boundary value problem as the Neumann problem.

*Extremum principle:* Given a domain  $D$ , a harmonic function  $w$  in  $D$  that is not identically constant in  $D$  cannot attain its maximum or minimum value at any interior point of  $D$ .

#### 7.1.1. Problems in Cartesian Coordinate System

The Laplace equation with two space variables in the rectangular Cartesian system of coordinates is written as

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = 0.$$

|                                                                    |
|--------------------------------------------------------------------|
| 7.1.1-1. Particular solutions and a method for their construction. |
|--------------------------------------------------------------------|

1°. Particular solutions:

$$w(x, y) = Ax + By + C,$$

$$w(x, y) = A(x^2 - y^2) + Bxy,$$

$$w(x, y) = A(x^3 - 3xy^2) + B(3x^2y - y^3),$$

$$w(x, y) = \frac{Ax + By}{x^2 + y^2} + C,$$

$$w(x, y) = \exp(\pm \mu x)(A \cos \mu y + B \sin \mu y),$$

$$w(x, y) = (A \cos \mu x + B \sin \mu x) \exp(\pm \mu y),$$

$$w(x, y) = (A \sinh \mu x + B \cosh \mu x)(C \cos \mu y + D \sin \mu y),$$

$$w(x, y) = (A \cos \mu x + B \sin \mu x)(C \sinh \mu y + D \cosh \mu y),$$

$$w(x, y) = A \ln[(x - x_0)^2 + (y - y_0)^2] + B,$$

where  $A, B, C, D, x_0, y_0$ , and  $\mu$  are arbitrary constants.

2°. Fundamental solution:

$$\mathcal{E}(x, y) = \frac{1}{2\pi} \ln \frac{1}{r}, \quad r = \sqrt{x^2 + y^2}.$$

3°. If  $w(x, y)$  is a solution of the Laplace equation, then the functions

$$\begin{aligned}w_1 &= Aw(\pm\lambda x + C_1, \pm\lambda y + C_2), \\w_2 &= Aw(x \cos \beta + y \sin \beta, -x \sin \beta + y \cos \beta), \\w_3 &= Aw\left(\frac{x}{x^2 + y^2}, \frac{y}{x^2 + y^2}\right),\end{aligned}$$

are also solutions everywhere they are defined;  $A$ ,  $C_1$ ,  $C_2$ ,  $\beta$ , and  $\lambda$  are arbitrary constants. The signs at  $\lambda$  in  $w_1$  are taken independently of each other.

4°. A fairly general method for constructing particular solutions involves the following. Let  $f(z) = u(x, y) + iv(x, y)$  be any analytic function of the complex variable  $z = x + iy$  ( $u$  and  $v$  are real functions of the real variables  $x$  and  $y$ ;  $i^2 = -1$ ). Then the real and imaginary parts of  $f$  both satisfy the two-dimensional Laplace equation,

$$\Delta_2 u = 0, \quad \Delta_2 v = 0.$$

Recall that the Cauchy–Riemann conditions

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

are necessary and sufficient conditions for the function  $f$  to be analytic. Thus, by specifying analytic functions  $f(z)$  and taking their real and imaginary parts, one obtains various solutions of the two-dimensional Laplace equation.

☉ *References:* M. A. Lavrent'ev and B. V. Shabat (1973), A. G. Sveshnikov and A. N. Tikhonov (1974), A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 7.1.1-2. Specific features of stating boundary value problems for the Laplace equation.

1°. For outer boundary value problems on the plane, it is (usually) required to set the additional condition that the solution of the Laplace equation must be bounded at infinity.

2°. The solution of the second boundary value problem is determined up to an arbitrary additive term.

3°. Let the second boundary value problem in a closed bounded domain  $D$  with piecewise smooth boundary  $\Sigma$  be characterized by the boundary condition\*

$$\frac{\partial w}{\partial N} = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in \Sigma,$$

where  $\frac{\partial w}{\partial N}$  is the derivative along the (outward) normal to  $\Sigma$ . The necessary and sufficient condition of solvability of the problem has the form

$$\int_{\Sigma} f(\mathbf{r}) d\Sigma = 0.$$

*Remark.* The same solvability condition occurs for the outer second boundary value problem if the domain is infinite but has a finite boundary.

☉ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

---

\* More rigorously,  $\Sigma$  must satisfy the Lyapunov condition [see Babich, Kapilevich, Mikhlin, et al. (1964) and Tikhonov and Samarskii (1990)].

**7.1.1-3. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.**

A half-plane is considered. A boundary condition is prescribed:

$$w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y f(\xi) d\xi}{(x - \xi)^2 + y^2} = \frac{1}{\pi} \int_{-\pi/2}^{\pi/2} f(x + y \tan \theta) d\theta.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), H. S. Carslaw and J. C. Jaeger (1984).

**7.1.1-4. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ . Second boundary value problem.**

A half-plane is considered. A boundary condition is prescribed:

$$\partial_y w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(\xi) \ln \sqrt{(x - \xi)^2 + y^2} d\xi + C,$$

where  $C$  is an arbitrary constant.

⊙ *Reference:* V. S. Vladimirov (1988).

**7.1.1-5. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.**

A quadrant of the plane is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at} \quad x = 0, \quad w = f_2(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \frac{4}{\pi} xy \int_0^{\infty} \frac{f_1(\eta) \eta d\eta}{[x^2 + (y - \eta)^2][x^2 + (y + \eta)^2]} + \frac{4}{\pi} xy \int_0^{\infty} \frac{f_2(\xi) \xi d\xi}{[(x - \xi)^2 + y^2][(x + \xi)^2 + y^2]}.$$

⊙ *Reference:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

**7.1.1-6. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . First boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(x) \quad \text{at} \quad y = 0, \quad w = f_2(x) \quad \text{at} \quad y = a.$$

Solution:

$$w(x, y) = \frac{1}{2a} \sin\left(\frac{\pi y}{a}\right) \int_{-\infty}^{\infty} \frac{f_1(\xi) d\xi}{\cosh[\pi(x - \xi)/a] - \cos(\pi y/a)} + \frac{1}{2a} \sin\left(\frac{\pi y}{a}\right) \int_{-\infty}^{\infty} \frac{f_2(\xi) d\xi}{\cosh[\pi(x - \xi)/a] + \cos(\pi y/a)}.$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**7.1.1-7. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . Second boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$\partial_y w = f_1(x) \quad \text{at} \quad y = 0, \quad \partial_y w = f_2(x) \quad \text{at} \quad y = a.$$

Solution:

$$w(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} f_1(\xi) \ln\{\cosh[\pi(x - \xi)/a] - \cos(\pi y/a)\} d\xi - \frac{1}{2\pi} \int_{-\infty}^{\infty} f_2(\xi) \ln\{\cosh[\pi(x - \xi)/a] + \cos(\pi y/a)\} d\xi + C,$$

where  $C$  is an arbitrary constant.

**7.1.1-8. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . First boundary value problem.**

A semiinfinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at} \quad x = 0, \quad w = f_2(x) \quad \text{at} \quad y = 0, \quad w = f_3(x) \quad \text{at} \quad y = a.$$

**Solution:**

$$\begin{aligned} w(x, y) = & \frac{2}{a} \sum_{n=1}^{\infty} \exp\left(-\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi y}{a}\right) \int_0^a f_1(\eta) \sin\left(\frac{n\pi \eta}{a}\right) d\eta \\ & + \frac{1}{2a} \sin\left(\frac{\pi y}{a}\right) \int_0^{\infty} \left\{ \frac{1}{\cosh[\pi(x-\xi)/a] - \cos(\pi y/a)} - \frac{1}{\cosh[\pi(x+\xi)/a] - \cos(\pi y/a)} \right\} f_2(\xi) d\xi \\ & + \frac{1}{2a} \sin\left(\frac{\pi y}{a}\right) \int_0^{\infty} \left\{ \frac{1}{\cosh[\pi(x-\xi)/a] + \cos(\pi y/a)} - \frac{1}{\cosh[\pi(x+\xi)/a] + \cos(\pi y/a)} \right\} f_3(\xi) d\xi. \end{aligned}$$

**Example.** Consider the first boundary value problem for the Laplace equation in a semiinfinite strip with  $f_1(y) = 1$  and  $f_2(x) = f_3(x) = 0$ .

Using the general formula and carrying out transformations, we obtain the solution

$$w(x, y) = \frac{2}{\pi} \arctan \left[ \frac{\sin(\pi y/a)}{\sinh(\pi x/a)} \right].$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**7.1.1-9. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . First boundary value problem.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y) \quad \text{at} \quad x = 0, \quad w = f_2(y) \quad \text{at} \quad x = a, \\ w = f_3(x) \quad \text{at} \quad y = 0, \quad w = f_4(x) \quad \text{at} \quad y = b. \end{aligned}$$

**Solution:**

$$\begin{aligned} w(x, y) = & \sum_{n=1}^{\infty} A_n \sinh\left[\frac{n\pi}{b}(a-x)\right] \sin\left(\frac{n\pi}{b}y\right) + \sum_{n=1}^{\infty} B_n \sinh\left(\frac{n\pi}{b}x\right) \sin\left(\frac{n\pi}{b}y\right) \\ & + \sum_{n=1}^{\infty} C_n \sin\left(\frac{n\pi}{a}x\right) \sinh\left[\frac{n\pi}{a}(b-y)\right] + \sum_{n=1}^{\infty} D_n \sin\left(\frac{n\pi}{a}x\right) \sinh\left(\frac{n\pi}{a}y\right), \end{aligned}$$

where the coefficients  $A_n$ ,  $B_n$ ,  $C_n$ , and  $D_n$  are expressed as

$$\begin{aligned} A_n &= \frac{2}{\lambda_n} \int_0^b f_1(\xi) \sin\left(\frac{n\pi \xi}{b}\right) d\xi, \quad B_n = \frac{2}{\lambda_n} \int_0^b f_2(\xi) \sin\left(\frac{n\pi \xi}{b}\right) d\xi, \\ C_n &= \frac{2}{\mu_n} \int_0^a f_3(\xi) \sin\left(\frac{n\pi \xi}{a}\right) d\xi, \quad D_n = \frac{2}{\mu_n} \int_0^a f_4(\xi) \sin\left(\frac{n\pi \xi}{a}\right) d\xi, \\ \lambda_n &= b \sinh\left(\frac{n\pi a}{b}\right), \quad \mu_n = a \sinh\left(\frac{n\pi b}{a}\right). \end{aligned}$$

⊙ *References:* M. M. Smirnov (1975), H. S. Carslaw and J. C. Jaeger (1984).

**7.1.1-10. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . Second boundary value problem.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w &= f_1(y) \quad \text{at} \quad x = 0, & \partial_x w &= f_2(y) \quad \text{at} \quad x = a, \\ \partial_y w &= f_3(x) \quad \text{at} \quad y = 0, & \partial_y w &= f_4(x) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y) = & -\frac{A_0}{4a}(x-a)^2 + \frac{B_0}{4a}x^2 - \frac{C_0}{4b}(x-b)^2 + \frac{D_0}{4b}y^2 + K \\ & -b \sum_{n=1}^{\infty} \frac{A_n}{\lambda_n} \cosh\left[\frac{n\pi}{b}(a-x)\right] \cos\left(\frac{n\pi}{b}y\right) + b \sum_{n=1}^{\infty} \frac{B_n}{\lambda_n} \cosh\left(\frac{n\pi}{b}x\right) \cos\left(\frac{n\pi}{b}y\right) \\ & -a \sum_{n=1}^{\infty} \frac{C_n}{\mu_n} \cos\left(\frac{n\pi}{a}x\right) \cosh\left[\frac{n\pi}{a}(b-y)\right] + a \sum_{n=1}^{\infty} \frac{D_n}{\mu_n} \cos\left(\frac{n\pi}{a}x\right) \cosh\left(\frac{n\pi}{a}y\right), \end{aligned}$$

where  $K$  is an arbitrary constant, and the coefficients  $A_n$ ,  $B_n$ ,  $C_n$ ,  $D_n$ ,  $\lambda_n$ , and  $\mu_n$  are expressed as

$$\begin{aligned} A_n &= \frac{2}{b} \int_0^b f_1(\xi) \cos\left(\frac{n\pi\xi}{b}\right) d\xi, & B_n &= \frac{2}{b} \int_0^b f_2(\xi) \cos\left(\frac{n\pi\xi}{b}\right) d\xi, \\ C_n &= \frac{2}{a} \int_0^a f_3(\xi) \cos\left(\frac{n\pi\xi}{a}\right) d\xi, & D_n &= \frac{2}{a} \int_0^a f_4(\xi) \cos\left(\frac{n\pi\xi}{a}\right) d\xi, \\ \lambda_n &= n\pi \sinh\left(\frac{n\pi a}{b}\right), & \mu_n &= n\pi \sinh\left(\frac{n\pi b}{a}\right). \end{aligned}$$

The solvability condition for the problem in question has the form (see Paragraph 7.1.1-2, Item 3°)

$$\int_0^b f_1(y) dy + \int_0^b f_2(y) dy - \int_0^a f_3(x) dx - \int_0^a f_4(x) dx = 0.$$

**7.1.1-11. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . Third boundary value problem.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w - k_1 w &= f_1(y) \quad \text{at} \quad x = 0, & \partial_x w + k_2 w &= f_2(y) \quad \text{at} \quad x = a, \\ \partial_y w - k_3 w &= f_3(x) \quad \text{at} \quad y = 0, & \partial_y w + k_4 w &= f_4(x) \quad \text{at} \quad y = b. \end{aligned}$$

For the solution, see Paragraph 7.2.2-14 with  $\Phi \equiv 0$ .

**7.1.1-12. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . Mixed boundary value problems.**

1°. A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w &= f(y) \quad \text{at} \quad x = 0, & \partial_x w &= g(y) \quad \text{at} \quad x = a, \\ w &= h(x) \quad \text{at} \quad y = 0, & w &= s(x) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y) = & -\frac{b}{\pi} \sum_{n=1}^{\infty} \frac{f_n}{n\lambda_n} \cosh\left[\frac{\pi n}{b}(a-x)\right] \sin\left(\frac{\pi n y}{b}\right) + \frac{b}{\pi} \sum_{n=1}^{\infty} \frac{g_n}{n\lambda_n} \cosh\left(\frac{\pi n x}{b}\right) \sin\left(\frac{\pi n y}{b}\right) \\ & + \sum_{n=1}^{\infty} \frac{h_n}{\mu_n} \cos\left(\frac{\pi n x}{a}\right) \sinh\left[\frac{\pi n}{a}(b-y)\right] + \sum_{n=1}^{\infty} \frac{s_n}{\mu_n} \cos\left(\frac{\pi n x}{a}\right) \sinh\left(\frac{\pi n y}{a}\right) \\ & + \frac{b-y}{ab} \int_0^a h(x) dx + \frac{y}{ab} \int_0^a s(x) dx, \end{aligned}$$

where

$$\begin{aligned} f_n &= \frac{2}{b} \int_0^b f(\xi) \sin\left(\frac{\pi n \xi}{b}\right) d\xi, & g_n &= \frac{2}{b} \int_0^b g(\xi) \sin\left(\frac{\pi n \xi}{b}\right) d\xi, \\ h_n &= \frac{2}{a} \int_0^a h(\xi) \cos\left(\frac{\pi n \xi}{a}\right) d\xi, & s_n &= \frac{2}{a} \int_0^a s(\xi) \cos\left(\frac{\pi n \xi}{a}\right) d\xi, \\ \lambda_n &= \sinh\left(\frac{\pi n a}{b}\right), & \mu_n &= \sinh\left(\frac{\pi n b}{a}\right), \end{aligned}$$

⊙ *Reference:* M. M. Smirnov (1975).

2°. A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f(y) \quad \text{at} \quad x = 0, & \partial_x w &= g(y) \quad \text{at} \quad x = a, \\ w &= h(x) \quad \text{at} \quad y = 0, & \partial_y w &= s(x) \quad \text{at} \quad y = b, \end{aligned}$$

where  $f(0) = h(0)$ .

Solution:

$$\begin{aligned} w(x, y) &= \sum_{n=0}^{\infty} \frac{f_n}{\cosh \lambda_n} \cosh\left(\lambda_n \frac{a-x}{a}\right) \sin\left(\lambda_n \frac{y}{a}\right) + a \sum_{n=0}^{\infty} \frac{g_n}{\lambda_n \cosh \lambda_n} \sinh\left(\lambda_n \frac{x}{a}\right) \sin\left(\lambda_n \frac{y}{a}\right) \\ &+ \sum_{n=0}^{\infty} \frac{h_n}{\cosh \mu_n} \sin\left(\mu_n \frac{x}{b}\right) \cosh\left(\mu_n \frac{b-y}{b}\right) + b \sum_{n=0}^{\infty} \frac{s_n}{\mu_n \cosh \mu_n} \sin\left(\mu_n \frac{x}{b}\right) \sinh\left(\mu_n \frac{y}{b}\right), \end{aligned}$$

where

$$\begin{aligned} f_n &= \frac{2}{b} \int_0^b f(\xi) \sin\left[\frac{\pi(2n+1)}{b}\xi\right] d\xi, & g_n &= \frac{2}{b} \int_0^b g(\xi) \sin\left[\frac{\pi(2n+1)}{b}\xi\right] d\xi, \\ h_n &= \frac{2}{a} \int_0^a h(\xi) \sin\left[\frac{\pi(2n+1)}{a}\xi\right] d\xi, & s_n &= \frac{2}{a} \int_0^a s(\xi) \sin\left[\frac{\pi(2n+1)}{a}\xi\right] d\xi, \\ \lambda_n &= \frac{\pi(2n+1)a}{2b}, & \mu_n &= \frac{\pi(2n+1)b}{2a}. \end{aligned}$$

⊙ *Reference:* M. M. Smirnov (1975).

## 7.1.2. Problems in Polar Coordinate System

The two-dimensional Laplace equation in the polar coordinate system is written as

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} = 0, \quad r = \sqrt{x^2 + y^2}.$$

**7.1.2-1. Particular solutions:**

$$\begin{aligned} w(r) &= A \ln r + B, \\ w(r, \varphi) &= \left( Ar^m + \frac{B}{r^m} \right) (C \cos m\varphi + D \sin m\varphi), \end{aligned}$$

where  $m = 1, 2, \dots$ ;  $A, B, C$ , and  $D$  are arbitrary constants.



**7.1.2-2. Domain:  $0 \leq r \leq R$  or  $R \leq r < \infty$ . First boundary value problem.**

The condition

$$w = f(\varphi) \quad \text{at} \quad r = R$$

is set at the boundary of the circle;  $f(\varphi)$  is a given function.

1°. Solution of the inner problem ( $r \leq R$ ):

$$w(r, \varphi) = \frac{1}{2\pi} \int_0^{2\pi} f(\psi) \frac{R^2 - r^2}{r^2 - 2Rr \cos(\varphi - \psi) + R^2} d\psi.$$

This formula is conventionally referred to as the Poisson integral.

Solution of the outer problem in series form:

$$w(r, \varphi) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( \frac{r}{R} \right)^n (a_n \cos n\varphi + b_n \sin n\varphi),$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(\psi) \cos(n\psi) d\psi, \quad n = 0, 1, 2, \dots,$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(\psi) \sin(n\psi) d\psi, \quad n = 1, 2, 3, \dots$$

2°. Bounded solution of the outer problem ( $r \geq R$ ):

$$w(r, \varphi) = \frac{1}{2\pi} \int_0^{2\pi} f(\psi) \frac{r^2 - R^2}{r^2 - 2Rr \cos(\varphi - \psi) + R^2} d\psi.$$

Bounded solution of the outer problem in series form:

$$w(r, \varphi) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( \frac{R}{r} \right)^n (a_n \cos n\varphi + b_n \sin n\varphi),$$

where the coefficients  $a_0$ ,  $a_n$ , and  $b_n$  are defined by the same relations as in the inner problem.

In hydrodynamics and other applications, outer problems are sometimes encountered in which one has to consider unbounded solutions for  $r \rightarrow \infty$ .

**Example.** The potential flow of an ideal (inviscid) incompressible fluid about a circular cylinder of radius  $R$  with a constant incident velocity  $U$  at infinity is characterized by the following boundary conditions for the stream function:

$$w = 0 \quad \text{at} \quad r = R, \quad w \rightarrow Ur \sin \varphi \quad \text{as} \quad r \rightarrow \infty.$$

Solution:

$$w(r, \varphi) = U \left( r - \frac{R^2}{r} \right) \sin \varphi.$$

☉ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), A. N. Tikhonov and A. A. Samarskii (1990).

**7.1.2-3. Domain:  $0 \leq r \leq R$  or  $R \leq r < \infty$ . Second boundary value problem.**

The condition

$$\partial_r w = f(\varphi) \quad \text{at} \quad r = R$$

is set at the boundary of the circle. The function  $f(\varphi)$  must satisfy the solvability condition

$$\int_0^{2\pi} f(\varphi) d\varphi = 0.$$

1°. Solution of the inner problem ( $r \leq R$ ):

$$w(r, \varphi) = \frac{R}{2\pi} \int_0^{2\pi} f(\psi) \ln \frac{r^2 - 2Rr \cos(\varphi - \psi) + R^2}{R^2} d\psi + C,$$

where  $C$  is an arbitrary constant; this formula is known as the Dini integral.

Series solution of the inner problem:

$$w(r, \varphi) = \sum_{n=1}^{\infty} \frac{R}{n} \left( \frac{r}{R} \right)^n (a_n \cos n\varphi + b_n \sin n\varphi) + C,$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(\psi) \cos(n\psi) d\psi, \quad b_n = \frac{1}{\pi} \int_0^{2\pi} f(\psi) \sin(n\psi) d\psi,$$

where  $C$  is an arbitrary constant.

2°. Solution of the outer problem ( $r \geq R$ ):

$$w(r, \varphi) = -\frac{R}{2\pi} \int_0^{2\pi} f(\psi) \ln \frac{r^2 - 2Rr \cos(\varphi - \psi) + R^2}{r^2} d\psi + C,$$

where  $C$  is an arbitrary constant.

Series solution of the outer problem:

$$w(r, \varphi) = -\sum_{n=1}^{\infty} \frac{R}{n} \left( \frac{R}{r} \right)^n (a_n \cos n\varphi + b_n \sin n\varphi) + C,$$

where the coefficient  $a_n$  and  $b_n$  are defined by the same relations as in the inner problem, and  $C$  is an arbitrary constant.

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

7.1.2-4. Domain:  $0 \leq r \leq R$  or  $R \leq r < \infty$ . Third boundary value problem.

The condition

$$\partial_r w + kw = f(\varphi) \quad \text{at} \quad r = R.$$

is set at the circle boundary;  $f(\varphi)$  is a given function.

1°. Solution of the inner problem ( $r \leq R$ ):

$$w(r, \varphi) = \frac{a_0}{2k} + \sum_{n=1}^{\infty} \frac{R}{kR + n} \left( \frac{r}{R} \right)^n (a_n \cos n\varphi + b_n \sin n\varphi),$$

$$a_n = \frac{1}{\pi} \int_0^{2\pi} f(\psi) \cos(n\psi) d\psi, \quad n = 0, 1, 2, \dots,$$

$$b_n = \frac{1}{\pi} \int_0^{2\pi} f(\psi) \sin(n\psi) d\psi, \quad n = 1, 2, 3, \dots$$

2°. Solution of the outer problem ( $r \geq R$ ):

$$w(r, \varphi) = \frac{a_0}{2k} + \sum_{n=1}^{\infty} \frac{R}{kR - n} \left( \frac{R}{r} \right)^n (a_n \cos n\varphi + b_n \sin n\varphi),$$

where the coefficient  $a_0$ ,  $a_n$ , and  $b_n$  are defined by the same relations as in the inner problem.

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

**7.1.2-5. Domain:  $R_1 \leq r \leq R_2$ . First boundary value problem.**

An annular domain is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi) \quad \text{at} \quad r = R_1, \quad w = f_2(\varphi) \quad \text{at} \quad r = R_2.$$

Solution:

$$w(r, \varphi) = A_0 + B_0 \ln r + \sum_{n=1}^{\infty} r^n (A_n \cos n\varphi + B_n \sin n\varphi) + \sum_{n=1}^{\infty} \frac{1}{r^n} (C_n \cos n\varphi + D_n \sin n\varphi),$$

where the coefficient  $A_0$ ,  $B_0$ ,  $A_n$ ,  $B_n$ ,  $C_n$ , and  $D_n$  are expressed as

$$\begin{aligned} A_0 &= \frac{1}{2} \frac{a_0^{(1)} \ln R_2 - a_0^{(2)} \ln R_1}{\ln R_2 - \ln R_1}, & B_0 &= \frac{1}{2} \frac{a_0^{(2)} - a_0^{(1)}}{\ln R_2 - \ln R_1}, \\ A_n &= \frac{R_2^n a_n^{(2)} - R_1^n a_n^{(1)}}{R_2^{2n} - R_1^{2n}}, & B_n &= \frac{R_2^n b_n^{(2)} - R_1^n b_n^{(1)}}{R_2^{2n} - R_1^{2n}}, \\ C_n &= (R_1 R_2)^n \frac{R_2^n a_n^{(1)} - R_1^n a_n^{(2)}}{R_2^{2n} - R_1^{2n}}, & D_n &= (R_1 R_2)^n \frac{R_2^n b_n^{(1)} - R_1^n b_n^{(2)}}{R_2^{2n} - R_1^{2n}}. \end{aligned}$$

Here, the  $a_n^{(i)}$  and  $b_n^{(i)}$  ( $i = 1, 2$ ) are the coefficients of the Fourier series expansions of the functions  $f_1(\varphi)$  and  $f_2(\varphi)$ :

$$\begin{aligned} a_n^{(i)} &= \frac{1}{\pi} \int_0^{2\pi} f_i(\psi) \cos(n\psi) d\psi, & n &= 0, 1, 2, \dots, \\ b_n^{(i)} &= \frac{1}{\pi} \int_0^{2\pi} f_i(\psi) \sin(n\psi) d\psi, & n &= 1, 2, 3, \dots \end{aligned}$$

☉ *Reference:* M. M. Smirnov (1975).

**7.1.2-6. Domain:  $R_1 \leq r \leq R_2$ . Second boundary value problem.**

An annular domain is considered. Boundary conditions are prescribed:

$$\partial_r w = f_1(\varphi) \quad \text{at} \quad r = R_1, \quad \partial_r w = f_2(\varphi) \quad \text{at} \quad r = R_2.$$

Solution:

$$w(r, \varphi) = B \ln r + \sum_{n=1}^{\infty} r^n (A_n \cos n\varphi + B_n \sin n\varphi) + \sum_{n=1}^{\infty} \frac{1}{r^n} (C_n \cos n\varphi + D_n \sin n\varphi) + K.$$

Here, the coefficients  $B$ ,  $A_n$ ,  $B_n$ ,  $C_n$ , and  $D_n$  are expressed as

$$\begin{aligned} B &= \frac{1}{2} R_1 a_0^{(1)}, & A_n &= \frac{R_2^{n+1} a_n^{(2)} - R_1^{n+1} a_n^{(1)}}{n(R_2^{2n} - R_1^{2n})}, & B_n &= \frac{R_2^{n+1} b_n^{(2)} - R_1^{n+1} b_n^{(1)}}{n(R_2^{2n} - R_1^{2n})}, \\ C_n &= (R_1 R_2)^{n+1} \frac{R_1^{n-1} a_n^{(2)} - R_2^{n-1} a_n^{(1)}}{n(R_2^{2n} - R_1^{2n})}, & D_n &= (R_1 R_2)^{n+1} \frac{R_1^{n-1} b_n^{(2)} - R_2^{n-1} b_n^{(1)}}{n(R_2^{2n} - R_1^{2n})}, \end{aligned}$$

where the constants  $a_n^{(i)}$  and  $b_n^{(i)}$  ( $i = 1, 2$ ) are defined by the same relations as in the first boundary value problem;  $K$  is an arbitrary constant.

*Remark.* Note that the condition  $a_0^{(1)} R_1 = a_0^{(2)} R_2$  must hold; this relation is a consequence of the solvability condition for the problem,

$$\int_{r=R_1} f_1 dS - \int_{r=R_2} f_2 dS = 0.$$

TABLE 21

Two-dimensional Laplace operator in some curvilinear orthogonal systems of coordinates

| Coordinates                           | Transformation ( $c > 0$ )                                                                                                                                 | Laplace operator, $\Delta_2 w$                                                                                                           |
|---------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------|
| Parabolic coordinates<br>$u, v$       | $x = cuv, y = \frac{1}{2}c(v^2 - u^2)$<br>$-\infty < u < \infty, 0 \leq v < \infty$                                                                        | $\frac{1}{c^2(u^2 + v^2)} \left( \frac{\partial^2 w}{\partial u^2} + \frac{\partial^2 w}{\partial v^2} \right)$                          |
| Elliptic coordinates<br>$\xi, \eta$   | $x = c \cosh \xi \cos \eta, y = c \sinh \xi \sin \eta$<br>$0 \leq \xi < \infty, 0 \leq \eta < 2\pi$                                                        | $\frac{1}{c^2(\sinh^2 \xi + \sin^2 \eta)} \left( \frac{\partial^2 w}{\partial \xi^2} + \frac{\partial^2 w}{\partial \eta^2} \right)$     |
| Bipolar coordinates<br>$\sigma, \tau$ | $x = \frac{c \sinh \tau}{\cosh \tau - \cos \sigma}, y = \frac{c \sin \sigma}{\cosh \tau - \cos \sigma}$<br>$0 \leq \sigma < 2\pi, -\infty < \tau < \infty$ | $\frac{1}{c^2(\cosh \tau - \cos \sigma)^2} \left( \frac{\partial^2 w}{\partial \sigma^2} + \frac{\partial^2 w}{\partial \tau^2} \right)$ |

7.1.2-7. Domain:  $R_1 \leq r \leq R_2$ . Mixed boundary value problem.

An annular domain is considered. Boundary conditions are prescribed:

$$\partial_r w = f_1(\varphi) \quad \text{at} \quad r = R_1, \quad w = f_2(\varphi) \quad \text{at} \quad r = R_2.$$

Solution:

$$w(r, \varphi) = \frac{1}{2}a_0^{(2)} + \frac{1}{2}a_0^{(1)}R_1 \ln \frac{r}{R_2} + \sum_{n=1}^{\infty} r^n (A_n \cos n\varphi + B_n \sin n\varphi) + \sum_{n=1}^{\infty} \frac{1}{r^n} (C_n \cos n\varphi + D_n \sin n\varphi).$$

Here, the coefficients  $A_n, B_n, C_n$ , and  $D_n$  are expressed as

$$A_n = \frac{nR_2^n a_n^{(2)} + R_1^{n+1} a_n^{(1)}}{n(R_2^{2n} + R_1^{2n})}, \quad B_n = \frac{nR_2^n b_n^{(2)} + R_1^{n+1} b_n^{(1)}}{n(R_2^{2n} + R_1^{2n})},$$

$$C_n = R_1^{n+1} R_2^n \frac{nR_1^{n-1} a_n^{(2)} - R_2^n a_n^{(1)}}{n(R_2^{2n} + R_1^{2n})}, \quad D_n = R_1^{n+1} R_2^n \frac{nR_1^{n-1} b_n^{(2)} - R_2^n b_n^{(1)}}{n(R_2^{2n} + R_1^{2n})},$$

where the constants  $a_n^{(i)}$  and  $b_n^{(i)}$  ( $i = 1, 2$ ) are defined by the same formulas as in the first boundary value problem.

⊙ Reference: M. M. Smirnov (1975).

### 7.1.3. Other Coordinate Systems. Conformal Mappings Method

7.1.3-1. Parabolic, elliptic, and bipolar coordinate systems.

In a number of applications, it is convenient to solve the Laplace equation in other orthogonal systems of coordinates. Some of those commonly encountered are displayed in Table 21. In all the coordinate systems presented, the Laplace equation  $\Delta_2 w = 0$  is reduced to the equation considered in Paragraph 7.1.1-1 in detail (particular solutions and solutions to boundary value problems are given there).

The orthogonal transformations presented in Table 21 can be written in the language of complex variables as follows:

$$x + iy = -\frac{1}{2}ic(u + iv)^2 \quad (\text{parabolic coordinates}),$$

$$x + iy = c \cosh(\xi + i\eta) \quad (\text{elliptic coordinates}),$$

$$x + iy = ic \cot\left[\frac{1}{2}(\sigma + i\tau)\right] \quad (\text{bipolar coordinates}).$$

The real parts, as well as the imaginary parts, in both sides of these relations must be equated to each other ( $i^2 = -1$ ).

**Example.** Plane hydrodynamic problems of potential flows of ideal (inviscid) incompressible fluid are reduced to the Laplace equation for the stream function. In particular, the motion of an elliptic cylinder with semiaxes  $a$  and  $b$  at a velocity  $U$  in the direction parallel to the major semiaxis ( $a > b$ ) in ideal fluid is described by the stream function

$$w(\xi, \eta) = -Ub \left( \frac{a+b}{a-b} \right)^{1/2} e^{\xi} \sin \eta, \quad c^2 = a^2 - b^2,$$

where  $\xi$  and  $\eta$  are the elliptic coordinates.

⊙ *References:* G. Lamb (1945), J. Happel and H. Brenner (1965), G. Korn and T. Korn (1968).

### 7.1.3-2. Domain of arbitrary shape. Method of conformal mappings.

1°. Let  $\zeta = \zeta(z)$  be an analytic function that defines a conformal mapping from the complex plane  $z = x + iy$  into a complex plane  $\zeta = u + iv$ , where  $u = u(x, y)$  and  $v = v(x, y)$  are new independent variables. With reference to the fact that the real and imaginary parts of an analytic function satisfy the Cauchy–Riemann conditions, we have  $\partial_x u = \partial_y v$  and  $\partial_y u = -\partial_x v$ , and hence

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = |\zeta'(z)|^2 \left( \frac{\partial^2 w}{\partial u^2} + \frac{\partial^2 w}{\partial v^2} \right).$$

Therefore, the Laplace equation in the  $xy$ -plane transforms under a conformal mapping into the Laplace equation in the  $uv$ -plane.

2°. Any simply connected domain  $D$  in the  $xy$ -plane with a piecewise smooth boundary can be mapped, with appropriate conformal mappings, onto the upper half-plane or into a unit circle in the  $uv$ -plane. Consequently, a first and a second boundary value problem for the Laplace equation in  $D$  can be reduced, respectively, to a first and a second boundary value problem for the upper half-space or a circle; such problems are considered in Subsections 7.1.1 and 7.1.2.

Subsection 7.2.4 presents conformal mappings of some domains onto the upper half-plane or a unit circle. Moreover, examples of solving specific boundary value problems for the Poisson equation by the conformal mappings method are given there; the Green's functions for a semicircle and a quadrant of a circle are obtained.

A large number of conformal mappings of various domains can be found, for example, in the references cited below.

⊙ *References:* V. I. Lavrik and V. N. Savenkov (1970), M. A. Lavrent'ev and B. V. Shabat (1973), V. I. Ivanov and M. K. Trubetskov (1994).

### 7.1.3-3. Reduction of the two-dimensional Neumann problem to the Dirichlet problem.

Let the position of any point  $(x_*, y_*)$  located on the boundary  $\Sigma$  of a domain  $D$  be specified by a parameter  $s$ , so that  $x_* = x_*(s)$  and  $y_* = y_*(s)$ . Then a function of two variables,  $f(x, y)$ , is determined on  $\Sigma$  by the parameter  $s$  as well,  $f(x, y)|_{\Sigma} = f(x_*(s), y_*(s)) = f_*(s)$ .

The solution of the two-dimensional Neumann problem for the Laplace equation  $\Delta_2 w = 0$  in  $D$  with the boundary condition of the second kind

$$\frac{\partial w}{\partial N} = f_*(s) \quad \text{for } \mathbf{r} \in \Sigma$$

can be expressed in terms of the solution of the two-dimensional Dirichlet problem for the Laplace equation  $\Delta_2 u = 0$  in  $D$  with the boundary condition of the first kind

$$u = F_*(s) \quad \text{for } \mathbf{r} \in \Sigma,$$

where  $F_*(s) = \int f_*(s) ds$ , as follows:

$$w(x, y) = \int_{x_0}^x \frac{\partial u}{\partial y}(t, y_0) dt - \int_{y_0}^y \frac{\partial u}{\partial x}(x, t) dt + C.$$

Here,  $(x_0, y_0)$  are the coordinates of any point in  $D$ , and  $C$  is an arbitrary constant.

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

## 7.2. Poisson Equation $\Delta_2 w = -\Phi(\mathbf{x})$

### 7.2.1. Preliminary Remarks. Solution Structure

Just as the Laplace equation, the Poisson equation is often encountered in heat and mass transfer theory, fluid mechanics, elasticity, electrostatics, and other areas of mechanics and physics. For example, it describes steady-state temperature distribution in the presence of heat sources or sinks in the domain under study.

The Laplace equation is a special case of the Poisson equation with  $\Phi \equiv 0$ .

In what follows, we consider a finite domain  $S$  with a sufficiently smooth boundary  $L$ . Let  $\mathbf{r} \in S$  and  $\boldsymbol{\rho} \in S$ , where  $\mathbf{r} = \{x, y\}$ ,  $\boldsymbol{\rho} = \{\xi, \eta\}$ ,  $|\mathbf{r} - \boldsymbol{\rho}|^2 = (x - \xi)^2 + (y - \eta)^2$ .

#### 7.2.1-1. First boundary value problem.

The solution of the first boundary value problem for the Poisson equation

$$\Delta_2 w = -\Phi(\mathbf{r}) \quad (1)$$

in the domain  $S$  with the nonhomogeneous boundary condition

$$w = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in L$$

can be represented as

$$w(\mathbf{r}) = \int_S \Phi(\boldsymbol{\rho}) G(\mathbf{r}, \boldsymbol{\rho}) dS_\rho - \int_L f(\boldsymbol{\rho}) \frac{\partial G}{\partial N_\rho} dL_\rho. \quad (2)$$

Here,  $G(\mathbf{r}, \boldsymbol{\rho})$  is the Green's function of the first boundary value problem,  $\frac{\partial G}{\partial N_\rho}$  is the derivative of the Green's function with respect to  $\xi, \eta$  along the outward normal  $\mathbf{N}$  to the boundary  $L$ . The integration is performed with respect to  $\xi, \eta$ , with  $dS_\rho = d\xi d\eta$ .

The Green's function  $G = G(\mathbf{r}, \boldsymbol{\rho})$  of the first boundary value problem is determined by the following conditions.

1°. The function  $G$  satisfies the Laplace equation in  $x, y$  in the domain  $S$  everywhere except for the point  $(\xi, \eta)$ , at which  $G$  has a singularity of the form  $\frac{1}{2\pi} \ln \frac{1}{|\mathbf{r} - \boldsymbol{\rho}|}$ .

2°. With respect to  $x, y$ , the function  $G$  satisfies the homogeneous boundary condition of the first kind at the domain boundary, i.e., the condition  $G|_L = 0$ .

The Green's function can be represented in the form

$$G(\mathbf{r}, \boldsymbol{\rho}) = \frac{1}{2\pi} \ln \frac{1}{|\mathbf{r} - \boldsymbol{\rho}|} + u, \quad (3)$$

where the auxiliary function  $u = u(\mathbf{r}, \boldsymbol{\rho})$  is determined by solving the first boundary value problem for the Laplace equation  $\Delta_2 u = 0$  with the boundary condition  $u|_L = -\frac{1}{2\pi} \ln \frac{1}{|\mathbf{r} - \boldsymbol{\rho}|}$ ; in this problem,  $\boldsymbol{\rho}$  is treated as a two-dimensional free parameter.

The Green's function is symmetric with respect to its arguments:  $G(\mathbf{r}, \boldsymbol{\rho}) = G(\boldsymbol{\rho}, \mathbf{r})$ .

*Remark 1.* When using the polar coordinate system, one should set

$$\mathbf{r} = \{r, \varphi\}, \quad \boldsymbol{\rho} = \{\xi, \eta\}, \quad |\mathbf{r} - \boldsymbol{\rho}|^2 = r^2 + \xi^2 - 2r\xi \cos(\varphi - \eta), \quad dS_\rho = \xi d\xi d\eta$$

in relations (2) and (3).

### 7.2.1-2. Second boundary value problem.

The second boundary value problem for the Poisson equation (1) is characterized by the boundary condition

$$\frac{\partial w}{\partial N} = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in L.$$

The necessary solvability condition for this problem is

$$\int_S \Phi(\mathbf{r}) dS + \int_L f(\mathbf{r}) dL = 0. \quad (4)$$

The solution of the second boundary value problem, provided that condition (4) is satisfied, can be represented as

$$w(\mathbf{r}) = \int_S \Phi(\boldsymbol{\rho}) G(\mathbf{r}, \boldsymbol{\rho}) dS_\rho + \int_L f(\boldsymbol{\rho}) G(\mathbf{r}, \boldsymbol{\rho}) dL_\rho + C, \quad (5)$$

where  $C$  is an arbitrary constant.

The Green's function  $G = G(\mathbf{r}, \boldsymbol{\rho})$  of the second boundary value problem is determined by the following conditions:

- 1°. The function  $G$  satisfies the Laplace equation in  $x, y$  in the domain  $S$  everywhere except for the point  $(\xi, \eta)$ , at which  $G$  has a singularity of the form  $\frac{1}{2\pi} \ln \frac{1}{|\mathbf{r}-\boldsymbol{\rho}|}$ .
- 2°. With respect to  $x, y$ , the function  $G$  satisfies the homogeneous boundary condition of the second kind at the domain boundary:

$$\left. \frac{\partial G}{\partial N} \right|_L = \frac{1}{L_0},$$

where  $L_0$  is the length of the boundary of  $S$ .

The Green's function is unique up to an additive constant.

*Remark 2.* The Green's function cannot be determined by condition 1° and the homogeneous boundary condition  $\left. \frac{\partial G}{\partial N} \right|_L = 0$ . The point is that the problem is unsolvable for  $G$  in this case, because, on representing  $G$  in the form (3), for  $u$  we obtain a problem with a nonhomogeneous boundary condition of the second kind for which the solvability condition (4) now is not satisfied.

### 7.2.1-3. Third boundary value problem.

The solution of the third boundary value problem for the Poisson equation (1) in the domain  $S$  with the nonhomogeneous boundary condition

$$\frac{\partial w}{\partial N} + kw = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in L$$

is given by formula (5) with  $C = 0$ , where  $G = G(\mathbf{r}, \boldsymbol{\rho})$  is the Green's function of the third boundary value problem and is determined by the following conditions:

- 1°. The function  $G$  satisfies the Laplace equation in  $x, y$  in the domain  $S$  everywhere except for the point  $(\xi, \eta)$ , at which  $G$  has a singularity of the form  $\frac{1}{2\pi} \ln \frac{1}{|\mathbf{r}-\boldsymbol{\rho}|}$ .
- 2°. With respect to  $x, y$ , the function  $G$  satisfies the homogeneous boundary condition of the third kind at the domain boundary, i.e., the condition  $\left[ \frac{\partial G}{\partial N} + kG \right]_L = 0$ .

The Green's function can be represented in the form (3); the auxiliary function  $u$  is identified by solving the corresponding third boundary value problem for the Laplace equation  $\Delta_2 u = 0$ .

The Green's function is symmetric with respect to its arguments:  $G(\mathbf{r}, \boldsymbol{\rho}) = G(\boldsymbol{\rho}, \mathbf{r})$ .

⊙ *References for Subsection 7.2.1:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970).

## 7.2.2. Problems in Cartesian Coordinate System

The two-dimensional Poisson equation in the rectangular Cartesian coordinate system has the form

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \Phi(x, y) = 0.$$

7.2.2-1. Particular solutions of the Poisson equation with a special right-hand side.

1°. If  $\Phi(x, y) = \sum_{i=1}^n \sum_{j=1}^n a_{ij} \exp(b_i x + c_j y)$ , the equation has solutions of the form

$$w(x, y) = - \sum_{i=1}^n \sum_{j=1}^n \frac{a_{ij}}{b_i^2 + c_j^2} \exp(b_i x + c_j y).$$

2°. If  $\Phi(x, y) = \sum_{i=1}^n \sum_{j=1}^n a_{ij} \sin(b_i x + p_i) \sin(c_j y + q_j)$ , the equation admits solutions of the form

$$w(x, y) = \sum_{i=1}^n \sum_{j=1}^n \frac{a_{ij}}{b_i^2 + c_j^2} \sin(b_i x + p_i) \sin(c_j y + q_j).$$

7.2.2-2. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ .

Solution:

$$w(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) \ln \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2}} d\xi d\eta.$$

7.2.2-3. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.

A half-plane is considered. A boundary condition is prescribed:

$$w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{y f(\xi) d\xi}{(x-\xi)^2 + y^2} + \frac{1}{2\pi} \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) \ln \frac{\sqrt{(x-\xi)^2 + (y+\eta)^2}}{\sqrt{(x-\xi)^2 + (y-\eta)^2}} d\xi d\eta.$$

⊙ *Reference:* A. G. Butkovskiy (1979).

7.2.2-4. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ . Second boundary value problem.

A half-plane is considered. A boundary condition is prescribed:

$$\partial_y w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(\xi) \ln \sqrt{(x-\xi)^2 + y^2} d\xi + \frac{1}{2\pi} \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) \left[ \ln \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2}} + \ln \frac{1}{\sqrt{(x-\xi)^2 + (y+\eta)^2}} \right] d\xi d\eta + C,$$

where  $C$  is an arbitrary constant.

⊙ *Reference:* V. S. Vladimirov (1988).



**7.2.2-5. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . First boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(x) \quad \text{at} \quad y = 0, \quad w = f_2(x) \quad \text{at} \quad y = a.$$

Solution:

$$\begin{aligned} w(x, y) = & \frac{1}{2a} \sin\left(\frac{\pi y}{a}\right) \int_{-\infty}^{\infty} \frac{f_1(\xi) d\xi}{\cosh[\pi(x-\xi)/a] - \cos(\pi y/a)} \\ & + \frac{1}{2a} \sin\left(\frac{\pi y}{a}\right) \int_{-\infty}^{\infty} \frac{f_2(\xi) d\xi}{\cosh[\pi(x-\xi)/a] + \cos(\pi y/a)} \\ & + \frac{1}{4\pi} \int_0^a \int_{-\infty}^{\infty} \Phi(\xi, \eta) \ln \frac{\cosh[\pi(x-\xi)/a] - \cos[\pi(y+\eta)/a]}{\cosh[\pi(x-\xi)/a] - \cos[\pi(y-\eta)/a]} d\xi d\eta. \end{aligned}$$

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

**7.2.2-6. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . Second boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$\partial_y w = f_1(x) \quad \text{at} \quad y = 0, \quad \partial_y w = f_2(x) \quad \text{at} \quad y = a.$$

Solution:

$$\begin{aligned} w(x, y) = & - \int_{-\infty}^{\infty} f_1(\xi) G(x, y, \xi, 0) d\xi + \int_{-\infty}^{\infty} f_2(\xi) G(x, y, \xi, a) d\xi \\ & + \int_0^a \int_{-\infty}^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta + C. \end{aligned}$$

Here,

$$G(x, y, \xi, \eta) = \frac{1}{4\pi} \ln \frac{1}{\cosh[\pi(x-\xi)/a] - \cos[\pi(y-\eta)/a]} + \frac{1}{4\pi} \ln \frac{1}{\cosh[\pi(x-\xi)/a] - \cos[\pi(y+\eta)/a]},$$

where  $C$  is an arbitrary constant.

**7.2.2-7. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . Third boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$\partial_y w - k_1 w = f_1(x) \quad \text{at} \quad y = 0, \quad \partial_y w + k_2 w = f_2(x) \quad \text{at} \quad y = a.$$

The solution  $w(x, y)$  is determined by the formula in Paragraph 7.2.2-6 where

$$\begin{aligned} G(x, y, \xi, \eta) = & \frac{1}{2} \sum_{n=1}^{\infty} \frac{\varphi_n(y) \varphi_n(\eta)}{\|\varphi_n\|^2 \mu_n} \exp(-\mu_n |x - \xi|), \\ \varphi_n(y) = & \mu_n \cos(\mu_n y) + k_1 \sin(\mu_n y), \quad \|\varphi_n\|^2 = \frac{1}{2} (\mu_n^2 + k_1^2) \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right]. \end{aligned}$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}$ .

**7.2.2-8. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . Mixed boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(x) \quad \text{at} \quad y = 0, \quad \partial_y w = f_2(x) \quad \text{at} \quad y = a.$$

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} f_1(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi + \int_{-\infty}^{\infty} f_2(\xi) G(x, y, \xi, a) d\xi \\ + \int_0^a \int_{-\infty}^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta,$$

where

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=0}^{\infty} \frac{1}{\mu_n} \exp(-\mu_n |x - \xi|) \sin(\mu_n y) \sin(\mu_n \eta), \quad \mu_n = \frac{\pi(2n+1)}{2a}.$$

**7.2.2-9. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . First boundary value problem.**

A semiinfinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at} \quad x = 0, \quad w = f_2(x) \quad \text{at} \quad y = 0, \quad w = f_3(x) \quad \text{at} \quad y = a.$$

Solution:

$$w(x, y) = \int_0^a f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta + \int_0^{\infty} f_2(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi \\ - \int_0^{\infty} f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=a} d\xi + \int_0^a \int_0^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta,$$

where

$$G(x, y, \xi, \eta) = \frac{1}{4\pi} \ln \frac{\cosh[\pi(x-\xi)/a] - \cos[\pi(y+\eta)/a]}{\cosh[\pi(x-\xi)/a] - \cos[\pi(y-\eta)/a]} - \frac{1}{4\pi} \ln \frac{\cosh[\pi(x+\xi)/a] - \cos[\pi(y+\eta)/a]}{\cosh[\pi(x+\xi)/a] - \cos[\pi(y-\eta)/a]}.$$

Alternatively, the Green's function can be represented in the series form

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=1}^{\infty} \frac{1}{q_n} [\exp(-q_n |x - \xi|) - \exp(-q_n |x + \xi|)] \sin(q_n y) \sin(q_n \eta), \quad q_n = \frac{\pi n}{a}.$$

☉ *References:* N. N. Lebedev, I. P. Skal'skaya, and Ya. S. Uflyand (1955), A. G. Butkovskiy (1979).

**7.2.2-10. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . Third boundary value problem.**

A semiinfinite strip is considered. Boundary conditions are prescribed:

$$\partial_x w - k_1 w = f_1(y) \quad \text{at} \quad x = 0, \quad \partial_y w - k_2 w = f_2(x) \quad \text{at} \quad y = 0, \quad \partial_y w + k_3 w = f_3(x) \quad \text{at} \quad y = a.$$

Solution:

$$w(x, y) = \int_0^a \int_0^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta - \int_0^a f_1(\eta) G(x, y, 0, \eta) d\eta \\ - \int_0^{\infty} f_2(\xi) G(x, y, \xi, 0) d\xi + \int_0^{\infty} f_3(\xi) G(x, y, \xi, a) d\xi,$$

where

$$G(x, y, \xi, \eta) = \sum_{n=1}^{\infty} \frac{\varphi_n(y)\varphi_n(\eta)}{\|\varphi_n\|^2 \mu_n(\mu_n + k_1)} H_n(x, \xi),$$

$$\varphi_n(y) = \mu_n \cos(\mu_n y) + k_2 \sin(\mu_n y), \quad \|\varphi_n\|^2 = \frac{1}{2}(\mu_n^2 + k_2^2) \left[ a + \frac{(k_2 + k_3)(\mu_n^2 + k_2 k_3)}{(\mu_n^2 + k_2^2)(\mu_n^2 + k_3^2)} \right],$$

$$H_n(x, \xi) = \begin{cases} \exp(-\mu_n x) [\mu_n \cosh(\mu_n \xi) + k_1 \sinh(\mu_n \xi)] & \text{for } x > \xi, \\ \exp(-\mu_n \xi) [\mu_n \cosh(\mu_n x) + k_1 \sinh(\mu_n x)] & \text{for } \xi > x. \end{cases}$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\tan(\mu a) = \frac{(k_2 + k_3)\mu}{\mu^2 - k_2 k_3}$ .

7.2.2-11. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . Mixed boundary value problems.

1°. A semiinfinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at } x = 0, \quad \partial_y w = f_2(x) \quad \text{at } y = 0, \quad \partial_y w = f_3(x) \quad \text{at } y = a.$$

Solution:

$$\begin{aligned} w(x, y) = & \int_0^a f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta - \int_0^\infty f_2(\xi) G(x, y, \xi, 0) d\xi \\ & + \int_0^\infty f_3(\xi) G(x, y, \xi, a) d\xi + \int_0^a \int_0^\infty \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta, \end{aligned}$$

where

$$G(x, y, \xi, \eta) = \frac{1}{2a} \sum_{n=0}^{\infty} \frac{\varepsilon_n}{q_n} [\exp(-q_n|x - \xi|) - \exp(-q_n|x + \xi|)] \cos(q_n y) \cos(q_n \eta),$$

$$q_n = \frac{\pi n}{a}, \quad \varepsilon = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}$$

2°. A semiinfinite strip is considered. Boundary conditions are prescribed:

$$\partial_x w = f_1(y) \quad \text{at } x = 0, \quad w = f_2(x) \quad \text{at } y = 0, \quad w = f_3(x) \quad \text{at } y = a.$$

Solution:

$$\begin{aligned} w(x, y) = & - \int_0^a f_1(\eta) G(x, y, 0, \eta) d\eta + \int_0^\infty f_2(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi \\ & - \int_0^\infty f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=a} d\xi + \int_0^a \int_0^\infty \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta, \end{aligned}$$

where

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=1}^{\infty} \frac{1}{q_n} [\exp(-q_n|x - \xi|) + \exp(-q_n|x + \xi|)] \sin(q_n y) \sin(q_n \eta), \quad q_n = \frac{\pi n}{a}.$$

7.2.2-12. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.

A quadrant of the plane is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at } x = 0, \quad w = f_2(x) \quad \text{at } y = 0.$$

Solution:

$$\begin{aligned} w(x, y) = & \frac{4}{\pi} xy \int_0^\infty \frac{f_1(\eta) \eta d\eta}{[x^2 + (y - \eta)^2][x^2 + (y + \eta)^2]} + \frac{4}{\pi} xy \int_0^\infty \frac{f_2(\xi) \xi d\xi}{[(x - \xi)^2 + y^2][(x + \xi)^2 + y^2]} \\ & + \frac{1}{2\pi} \int_0^\infty \int_0^\infty \Phi(\xi, \eta) \ln \frac{\sqrt{(x - \xi)^2 + (y + \eta)^2} \sqrt{(x + \xi)^2 + (y - \eta)^2}}{\sqrt{(x - \xi)^2 + (y - \eta)^2} \sqrt{(x + \xi)^2 + (y + \eta)^2}} d\xi d\eta. \end{aligned}$$

⊙ References: V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), A. G. Butkovskiy (1979).

**7.2.2-13. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . First boundary value problem.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y) \quad \text{at} \quad x = 0, & w &= f_2(y) \quad \text{at} \quad x = a, \\ w &= f_3(x) \quad \text{at} \quad y = 0, & w &= f_4(x) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y) &= \int_0^a \int_0^b \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi \\ &+ \int_0^b f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta - \int_0^b f_2(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=a} d\eta \\ &+ \int_0^a f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi - \int_0^a f_4(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=b} d\xi. \end{aligned}$$

Two forms of representation of the Green's function:

$$G(x, y, \xi, \eta) = \frac{2}{a} \sum_{n=1}^{\infty} \frac{\sin(p_n x) \sin(p_n \xi)}{p_n \sinh(p_n b)} H_n(y, \eta) = \frac{2}{b} \sum_{m=1}^{\infty} \frac{\sin(q_m y) \sin(q_m \eta)}{q_m \sinh(q_m a)} Q_m(x, \xi),$$

where

$$\begin{aligned} p_n &= \frac{\pi n}{a}, \quad H_n(y, \eta) = \begin{cases} \sinh(p_n \eta) \sinh[p_n(b-y)] & \text{for } b \geq y > \eta \geq 0, \\ \sinh(p_n y) \sinh[p_n(b-\eta)] & \text{for } b \geq \eta > y \geq 0, \end{cases} \\ q_m &= \frac{\pi m}{b}, \quad Q_m(x, \xi) = \begin{cases} \sinh(q_m \xi) \sinh[q_m(a-x)] & \text{for } a \geq x > \xi \geq 0, \\ \sinh(q_m x) \sinh[q_m(a-\xi)] & \text{for } a \geq \xi > x \geq 0. \end{cases} \end{aligned}$$

The Green's function can be written in form of a double series:

$$G(x, y, \xi, \eta) = \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta)}{p_n^2 + q_m^2}, \quad p_n = \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}.$$

☉ Reference: A. G. Butkovskiy (1979).

**7.2.2-14. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . Third boundary value problem.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w - k_1 w &= f_1(y) \quad \text{at} \quad x = 0, & \partial_x w + k_2 w &= f_2(y) \quad \text{at} \quad x = a, \\ \partial_y w - k_3 w &= f_3(x) \quad \text{at} \quad y = 0, & \partial_y w + k_4 w &= f_4(x) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y) &= \int_0^a \int_0^b \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi \\ &- \int_0^b f_1(\eta) G(x, y, 0, \eta) d\eta + \int_0^b f_2(\eta) G(x, y, a, \eta) d\eta \\ &- \int_0^a f_3(\xi) G(x, y, \xi, 0) d\xi + \int_0^a f_4(\xi) G(x, y, \xi, b) d\xi. \end{aligned}$$

Here,

$$\begin{aligned} G(x, y, \xi, \eta) &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi) \psi_m(y) \psi_m(\eta)}{\|\varphi_n\|^2 \|\psi_m\|^2 (\mu_n^2 + \lambda_m^2)}, \\ \varphi_n(x) &= \cos(\mu_n x) + \frac{k_1}{\mu_n} \sin(\mu_n x), \quad \|\varphi_n\|^2 = \frac{k_2}{2\mu_n^2} \frac{\mu_n^2 + k_1^2}{\mu_n^2 + k_2^2} + \frac{k_1}{2\mu_n^2} + \frac{a}{2} \left( 1 + \frac{k_1^2}{\mu_n^2} \right), \\ \psi_m(y) &= \cos(\lambda_m y) + \frac{k_3}{\lambda_m} \sin(\lambda_m y), \quad \|\psi_m\|^2 = \frac{k_4}{2\lambda_m^2} \frac{\lambda_m^2 + k_3^2}{\lambda_m^2 + k_4^2} + \frac{k_3}{2\lambda_m^2} + \frac{b}{2} \left( 1 + \frac{k_3^2}{\lambda_m^2} \right), \end{aligned}$$

where the  $\mu_n$  and  $\lambda_m$  are positive roots of the transcendental equations

$$\frac{\tan(\mu a)}{\mu} = \frac{k_1 + k_2}{\mu^2 - k_1 k_2}, \quad \frac{\tan(\lambda b)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}.$$

7.2.2-15. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . Mixed boundary value problem.

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y) & \text{at } x &= 0, & \partial_x w &= f_2(y) & \text{at } x &= a, \\ w &= f_3(x) & \text{at } y &= 0, & \partial_y w &= f_4(x) & \text{at } y &= b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y) &= \int_0^a \int_0^b \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi \\ &+ \int_0^b f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta + \int_0^b f_2(\eta) G(x, y, a, \eta) d\eta \\ &+ \int_0^a f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi + \int_0^a f_4(\xi) G(x, y, \xi, b) d\xi. \end{aligned}$$

Two forms of representation of the Green's function:

$$G(x, y, \xi, \eta) = \frac{2}{a} \sum_{n=0}^{\infty} \frac{\sin(p_n x) \sin(p_n \xi)}{p_n \cosh(p_n b)} H_n(y, \eta) = \frac{2}{b} \sum_{m=0}^{\infty} \frac{\sin(q_m y) \sin(q_m \eta)}{q_m \cosh(q_m a)} Q_m(x, \xi),$$

where

$$\begin{aligned} p_n &= \frac{\pi(2n+1)}{a}, & H_n(y, \eta) &= \begin{cases} \sinh(p_n \eta) \cosh[p_n(b-y)] & \text{for } b \geq y > \eta \geq 0, \\ \sinh(p_n y) \cosh[p_n(b-\eta)] & \text{for } b \geq \eta > y \geq 0, \end{cases} \\ q_m &= \frac{\pi(2m+1)}{b}, & Q_m(x, \xi) &= \begin{cases} \sinh(q_m \xi) \cosh[q_m(a-x)] & \text{for } a \geq x > \xi \geq 0, \\ \sinh(q_m x) \cosh[q_m(a-\xi)] & \text{for } a \geq \xi > x \geq 0. \end{cases} \end{aligned}$$

The Green's function can be written in form of a double series:

$$\begin{aligned} G(x, y, \xi, \eta) &= \frac{4}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{\sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta)}{p_n^2 + q_m^2}, \\ p_n &= \frac{\pi(2n+1)}{2a}, & q_m &= \frac{\pi(2m+1)}{2b}. \end{aligned}$$

### 7.2.3. Problems in Polar Coordinate System

The two-dimensional Poisson equation in the polar coordinate system is written as

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \Phi(r, \varphi) = 0, \quad r = \sqrt{x^2 + y^2}.$$

7.2.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

A circle is considered. A boundary condition is prescribed:

$$w = f(\varphi) \quad \text{at } r = R.$$

Solution:

$$w(r, \varphi) = \frac{1}{2\pi} \int_0^{2\pi} f(\eta) \frac{R^2 - r^2}{r^2 - 2Rr \cos(\varphi - \eta) + R^2} d\eta + \int_0^{2\pi} \int_0^R \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta,$$

where

$$\begin{aligned} G(r, \varphi, \xi, \eta) &= \frac{1}{2\pi} \ln \frac{1}{|\mathbf{r} - \mathbf{r}_0|} - \frac{1}{2\pi} \ln \frac{R}{r_0 |(R/r_0)^2 \mathbf{r}_0 - \mathbf{r}|}, \\ \mathbf{r} &= \{x, y\}, \quad x = r \cos \varphi, \quad y = r \sin \varphi, \\ \mathbf{r}_0 &= \{x_0, y_0\}, \quad x_0 = \xi \cos \eta, \quad y_0 = \xi \sin \eta. \end{aligned}$$

The magnitude of a vector difference is calculated as  $|\mathbf{a}\mathbf{r} - \mathbf{b}\mathbf{r}_0|^2 = a^2 r^2 - 2abr\xi \cos(\varphi - \eta) + b^2 \xi^2$  ( $a$  and  $b$  are any scalars). Thus, we obtain

$$G(r, \varphi, \xi, \eta) = \frac{1}{4\pi} \ln \frac{r^2 \xi^2 - 2R^2 r \xi \cos(\varphi - \eta) + R^4}{R^2 [r^2 - 2r\xi \cos(\varphi - \eta) + \xi^2]}.$$

☉ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), A. G. Butkovskiy (1979).

7.2.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A circle is considered. A boundary condition is prescribed:

$$\partial_r w + kw = f(\varphi) \quad \text{at} \quad r = R.$$

Solution:

$$w(r, \varphi) = R \int_0^{2\pi} f(\eta) G(r, \varphi, R, \eta) d\eta + \int_0^{2\pi} \int_0^R \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta,$$

where

$$\begin{aligned} G(r, \varphi, \xi, \eta) &= \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{(\mu_{nm}^2 R^2 + k^2 R^2 - n^2) [J_n(\mu_{nm} R)]^2} \cos[n(\varphi - \eta)], \\ A_0 &= 1, \quad A_n = 2 \quad (n = 1, 2, \dots). \end{aligned}$$

Here, the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k J_n(\mu R) = 0.$$

7.2.3-3. Domain:  $R \leq r < \infty$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.

The exterior of a circle is considered. A boundary condition is prescribed:

$$w = f(\varphi) \quad \text{at} \quad r = R.$$

Solution:

$$w(r, \varphi) = \frac{1}{2\pi} \int_0^{2\pi} f(\eta) \frac{r^2 - R^2}{r^2 - 2Rr \cos(\varphi - \eta) + R^2} d\eta + \int_0^{2\pi} \int_R^{\infty} \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta,$$

where the Green's function  $G(r, \varphi, \xi, \eta)$  is defined by the formula presented in Paragraph 7.2.3-1.

☉ *Reference:* A. G. Butkovskiy (1979).

**7.2.3-4. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

An annular domain is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi) \quad \text{at} \quad r = R_1, \quad w = f_2(\varphi) \quad \text{at} \quad r = R_2.$$

Solution:

$$w(r, \varphi) = R_1 \int_0^{2\pi} f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta) \right]_{\xi=R_1} d\eta - R_2 \int_0^{2\pi} f_2(\eta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta) \right]_{\xi=R_2} d\eta \\ + \int_0^{2\pi} \int_{R_1}^{R_2} \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta.$$

Here,

$$G(r, \varphi, \xi, \eta) = \frac{1}{2\pi} \sum_{n=0}^{\infty} \left( \ln \frac{1}{r_n} - \ln \frac{R_1}{\xi r_n^*} \right),$$

where

$$r_n^2 = r^2 + \rho_n^2 - 2r\rho_n \cos(\varphi - \eta), \quad (r_n^*)^2 = r^2 + (\rho_n^*)^2 - 2r\rho_n^* \cos(\varphi - \eta), \\ \rho_n = \begin{cases} (R_1/R_2)^{2k} \xi & \text{for } n = 2k, \\ (R_2/R_1)^{2k+2} \xi & \text{for } n = 2k+1, \end{cases} \quad \rho_n^* = \frac{R_1^2}{\rho_n}.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.2.3-5. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \pi$ . First boundary value problem.**

A semicircle is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi) \quad \text{at} \quad r = R, \quad w = f_2(r) \quad \text{at} \quad \varphi = 0, \quad w = f_3(r) \quad \text{at} \quad \varphi = \pi.$$

Solution:

$$w(r, \varphi) = -R \int_0^{\pi} f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta) \right]_{\xi=R} d\eta + \int_0^R f_2(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=0} d\xi \\ - \int_0^R f_3(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=\pi} d\xi + \int_0^{\pi} \int_0^R \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta,$$

where

$$G(r, \varphi, \xi, \eta) = \frac{1}{4\pi} \ln \frac{r^2 \xi^2 - 2R^2 r \xi \cos(\varphi - \eta) + R^4}{R^2 [r^2 - 2r \xi \cos(\varphi - \eta) + \xi^2]} - \frac{1}{4\pi} \ln \frac{r^2 \xi^2 - 2R^2 r \xi \cos(\varphi + \eta) + R^4}{R^2 [r^2 - 2r \xi \cos(\varphi + \eta) + \xi^2]}.$$

See also Example 2 in Paragraph 7.2.4-2.

⊙ *References:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.2.3-6. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \pi/2$ . First boundary value problem.**

A quadrant of a circle is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi) \quad \text{at} \quad r = R, \quad w = f_2(r) \quad \text{at} \quad \varphi = 0, \quad w = f_3(r) \quad \text{at} \quad \varphi = \pi/2.$$

Solution:

$$w(r, \varphi) = -R \int_0^{\pi/2} f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta) \right]_{\xi=R} d\eta + \int_0^R f_2(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=0} d\xi \\ - \int_0^R f_3(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=\pi/2} d\xi + \int_0^{\pi/2} \int_0^R \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta,$$

where

$$G(r, \varphi, \xi, \eta) = G_1(r, \varphi, \xi, \eta) - G_1(r, \varphi, \xi, 2\pi - \eta) - G_1(r, \varphi, \xi, \pi - \eta) + G_1(r, \varphi, \xi, \pi + \eta),$$

$$G_1(r, \varphi, \xi, \eta) = \frac{1}{4\pi} \ln \frac{r^2 \xi^2 - 2R^2 r \xi \cos(\varphi - \eta) + R^4}{R^2[r^2 - 2r\xi \cos(\varphi - \eta) + \xi^2]}.$$

See also Example 3 in Paragraph 7.2.4-2.

⊙ *References:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 7.2.3-7. Domain: $0 \leq r \leq R$ , $0 \leq \varphi \leq \beta$ . First boundary value problem.

A circular sector is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi) \quad \text{at} \quad r = R, \quad w = f_2(r) \quad \text{at} \quad \varphi = 0, \quad w = f_3(r) \quad \text{at} \quad \varphi = \beta.$$

**Solution:**

$$\begin{aligned} w(r, \varphi) = & -R \int_0^\beta f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, \xi, \eta) \right]_{\xi=R} d\eta + \int_0^R f_2(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=0} d\xi \\ & - \int_0^R f_3(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=\beta} d\xi + \int_0^\beta \int_0^R \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta. \end{aligned}$$

1°. For  $\beta = \pi/n$ , where  $n$  is a positive integer, the Green's function is expressed as

$$G(r, \varphi, \xi, \eta) = \sum_{k=0}^{n-1} [G_1(r, \varphi, \xi, 2k\beta + \eta) - G_1(r, \varphi, \xi, 2k\beta - \eta)],$$

$$G_1(r, \varphi, \xi, \eta) = \frac{1}{4\pi} \ln \frac{r^2 \xi^2 - 2R^2 r \xi \cos(\varphi - \eta) + R^4}{R^2[r^2 - 2r\xi \cos(\varphi - \eta) + \xi^2]}.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

2°. For arbitrary  $\beta$ , the Green's function is given by

$$G(r, \varphi, \xi, \eta) = \frac{1}{2\pi} \ln \frac{|z^{\pi/\beta} - \bar{\zeta}^{\pi/\beta}| |R^{2\pi/\beta} - (\bar{\zeta}z)^{\pi/\beta}|}{|z^{\pi/\beta} - \zeta^{\pi/\beta}| |R^{2\pi/\beta} - (\zeta z)^{\pi/\beta}|},$$

where  $z = re^{i\varphi}$ ,  $\zeta = \xi e^{i\eta}$ ,  $\bar{\zeta} = \xi e^{-i\eta}$ , and  $i^2 = -1$ .

### 7.2.3-8. Domain: $0 \leq r < \infty$ , $0 \leq \varphi \leq \beta$ . First boundary value problem.

A wedge domain is considered. Boundary conditions are prescribed:

$$w = f_1(r) \quad \text{at} \quad \varphi = 0, \quad w = f_2(r) \quad \text{at} \quad \varphi = \beta.$$

**Solution:**

$$\begin{aligned} w(r, \varphi) = & \int_0^\infty f_1(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=0} d\xi - \int_0^\infty f_2(\xi) \frac{1}{\xi} \left[ \frac{\partial}{\partial \eta} G(r, \varphi, \xi, \eta) \right]_{\eta=\beta} d\xi \\ & + \int_0^\beta \int_0^\infty \Phi(\xi, \eta) G(r, \varphi, \xi, \eta) \xi d\xi d\eta, \end{aligned}$$

where

$$G(r, \varphi, \xi, \eta) = \frac{1}{4\pi} \ln \frac{r^{2\pi/\beta} - 2(r\xi)^{\pi/\beta} \cos[\pi(\varphi + \eta)/\beta] + \xi^{2\pi/\beta}}{r^{2\pi/\beta} - 2(r\xi)^{\pi/\beta} \cos[\pi(\varphi - \eta)/\beta] + \xi^{2\pi/\beta}}.$$

Alternatively, the Green's function can be represented in the complex form

$$G(r, \varphi, \xi, \eta) = \frac{1}{2\pi} \ln \frac{|z^{\pi/\beta} - \bar{\zeta}^{\pi/\beta}|}{|z^{\pi/\beta} - \zeta^{\pi/\beta}|}, \quad z = re^{i\varphi}, \quad \zeta = \xi e^{i\eta}, \quad \bar{\zeta} = \xi e^{-i\eta}, \quad i^2 = -1.$$



## 7.2.4. Arbitrary Shape Domain. Conformal Mappings Method

### 7.2.4-1. Description of the method. Tables of conformal mappings.

Any simply connected domain  $D$  in the  $xy$ -plane with a piecewise smooth boundary can be mapped in a mutually unique way, with an appropriate conformal mapping, onto the upper half-plane or into a unit circle in a  $uv$ -plane. Under a conformal mapping, a Poisson equation in the  $xy$ -plane transforms into a Poisson equation in the  $uv$ -plane; what is changed is the function  $\Phi$ , as well as the function  $f$  in the boundary condition. Consequently, a first and a second boundary value problem for the plane domain  $D$  can be reduced, respectively, to a first and a second boundary value problem for the upper half-plane or a unit circle. The latter problems are considered above (see Subsections 7.2.2 and 7.2.3).

A large number of conformal mappings (mappings defined by analytic functions) of various domains onto the upper half-plane or a unit circle can be found, for example, in Lavrik and Savenkov (1970), Lavrent'ev and Shabat (1973), and Ivanov and Trubetskov (1994).

**Table 22** presents conformal mappings of some domains  $D$  in the complex plane  $z$  onto the upper half-plane  $\text{Im } \omega \geq 0$  in the complex plane  $\omega$ . In the relations involving square roots, it is assumed that  $\sqrt{\zeta} = \sqrt{|\zeta|} [\cos(\frac{1}{2}\varphi) + i \sin(\frac{1}{2}\varphi)]$ , where  $\varphi = \arg \zeta$  (i.e., the first branch of  $\sqrt{\zeta}$  is taken).

**Table 23** presents conformal mappings of some domains  $D$  in the complex plane  $z$  onto the unit circle  $|\omega| \leq 1$  in the complex plane  $\omega$ .

### 7.2.4-2. General formula for the Green's function. Example boundary value problems.

Let a function  $\omega = \omega(z)$  define a conformal mapping of a domain  $D$  in the complex plane  $z$  onto the upper half-plane in the complex plane  $\omega$ . Then the Green's function of the first boundary value problem in  $D$  for the Poisson (Laplace) equation is expressed as

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} \ln \left| \frac{\omega(z) - \bar{\omega}(\zeta)}{\omega(z) - \omega(\zeta)} \right|, \quad z = x + iy, \quad \zeta = \xi + i\eta, \quad (1)$$

where  $\omega(z) = u(x, y) + iv(x, y)$  and  $\bar{\omega}(z) = u(x, y) - iv(x, y)$ .

The solution of the first boundary value problem for the Poisson equation is determined by the above Green's function in accordance with formula (2) specified in Paragraph 7.2.1-1.

**Example 1.** Consider the first boundary value problem for the Poisson equation in the strip  $-\infty < x < \infty, 0 \leq y \leq a$ . The function that maps this strip onto the upper half-plane has the form  $\omega(z) = \exp(\pi x/a)$  (see the second row of **Table 22**). Substituting this expression into relation (1) and performing elementary transformations, we obtain the Green's function

$$G(x, y, \xi, \eta) = \frac{1}{4\pi} \ln \frac{\cosh[\pi(x - \xi)/a] - \cos[\pi(y + \eta)/a]}{\cosh[\pi(x - \xi)/a] - \cos[\pi(y - \eta)/a]}.$$

**Example 2.** Consider the first boundary value problem for the Poisson equation in a semicircle of radius  $a$  such that  $D = \{x^2 + y^2 \leq a^2, y \geq 0\}$ . The domain  $D$  is conformally mapped onto the upper half-plane by the function  $\omega(z) = -(z/a + a/z)$  (see the sixth row in **Table 22**). Substituting this expression into (1), we arrive at the Green's function

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} \ln \frac{|z - \bar{\zeta}| |a^2 - z\bar{z}|}{|z - \zeta| |a^2 - z\bar{z}|}, \quad z = x + iy, \quad \zeta = \xi + i\eta.$$

**Example 3.** Consider the first boundary value problem for the Poisson equation in a quadrant of a circle of radius  $a$ , so that  $D = \{x^2 + y^2 \leq a^2, x \geq 0, y \geq 0\}$ . The conformal mapping of the domain  $D$  onto the upper half-plane is performed with the function  $\omega(z) = -(a/z)^2 - (a/\bar{z})^2$  (see the seventh row of **Table 22**). Substituting this expression into (1) yields

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} \ln \frac{|z^2 - \bar{\zeta}^2| |a^4 - z^2 \bar{\zeta}^2|}{|z^2 - \zeta^2| |a^4 - z^2 \bar{\zeta}^2|}, \quad z = x + iy, \quad \zeta = \xi + i\eta.$$

3°. Let a function  $\omega = \omega(z)$  define a conformal mapping of a domain  $D$  in the complex plane  $z$  onto the unit circle  $|\omega| \leq 1$  in the complex plane  $\omega$ . Then the Green's function of the first boundary value problem in  $D$  for the Laplace equation is given by

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} \ln \left| \frac{1 - \bar{\omega}(\zeta)\omega(z)}{\omega(z) - \omega(\zeta)} \right|, \quad z = x + iy, \quad \zeta = \xi + i\eta. \quad (2)$$

© *References for Subsection 7.2.4:* N. N. Lebedev, I. P. Skalk'skaya, and Ya. S. Uflyand (1955), A. G. Sveshnikov and A. N. Tikhonov (1974).

TABLE 22

Conformal mapping of some domains  $D$  in the  $z$ -plane onto the upper half-plane  $\text{Im } \omega \geq 0$  in the  $\omega$ -plane. Notation:  $z = x + iy$  and  $\omega = u + iv$

| No | Domain $D$ in the $z$ -plane                                                                                                     | Transformation                                                                          |
|----|----------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------|
| 1  | First quadrant:<br>$0 \leq x < \infty, 0 \leq y < \infty$                                                                        | $\omega = a^2 z^2 + b$ ,<br>$a, b$ are real numbers                                     |
| 2  | Infinite strip of width $a$ :<br>$-\infty < x < \infty, 0 \leq y \leq a$                                                         | $\omega = \exp(\pi z/a)$                                                                |
| 3  | Semiinfinite strip of width $a$ :<br>$0 \leq x < \infty, 0 \leq y \leq a$                                                        | $\omega = \cosh(\pi z/a)$                                                               |
| 4  | Plane with the cut<br>in the real axis                                                                                           | $\omega = \sqrt{z}$                                                                     |
| 5  | Interior of an infinite sector with angle $\beta$ :<br>$0 \leq \arg z \leq \beta, 0 \leq  z  < \infty$ ( $0 < \beta \leq 2\pi$ ) | $\omega = z^{\pi/\beta}$                                                                |
| 6  | Upper half of a circle of radius $a$ :<br>$x^2 + y^2 \leq a^2, y \geq 0$                                                         | $\omega = -\frac{z}{a} - \frac{a}{z}$                                                   |
| 7  | Quadrant of a circle of radius $a$ :<br>$x^2 + y^2 \leq a^2, x \geq 0, y \geq 0$                                                 | $\omega = -\frac{z^2}{a^2} - \frac{a^2}{z^2}$                                           |
| 8  | Sector of a circle of radius $a$ with angle $\beta$ :<br>$x^2 + y^2 \leq a^2, 0 \leq \arg z \leq \beta$                          | $\omega = -\left(\frac{z}{a}\right)^{\pi/\beta} - \left(\frac{a}{z}\right)^{\pi/\beta}$ |
| 9  | Upper half-plane with a circular domain<br>or radius $a$ removed: $y \geq 0, x^2 + y^2 \geq a^2$                                 | $\omega = \frac{z}{a} + \frac{a}{z}$                                                    |
| 10 | Exterior of a parabola:<br>$y^2 - 2px \geq 0$                                                                                    | $\omega = \sqrt{z - \frac{1}{2}p} - i\sqrt{\frac{1}{2}p}$                               |
| 11 | Interior of a parabola:<br>$y^2 - 2px \leq 0$                                                                                    | $\omega = i \cosh\left(\pi \sqrt{\frac{1}{2}z/p - \frac{1}{4}}\right)$                  |

## 7.3. Helmholtz Equation $\Delta_2 w + \lambda w = -\Phi(\mathbf{x})$

Many problems related to steady-state oscillations (mechanical, acoustical, thermal, electromagnetic, etc.) lead to the two-dimensional Helmholtz equation. For  $\lambda < 0$ , this equation describes mass transfer processes with volume chemical reactions of the first order. Moreover, any elliptic equation with constant coefficients can be reduced to the Helmholtz equation.

### 7.3.1. General Remarks, Results, and Formulas

#### 7.3.1-1. Some definitions.

The Helmholtz equation is called homogeneous if  $\Phi = 0$  and nonhomogeneous if  $\Phi \neq 0$ . A homogeneous boundary value problem is a boundary value problem for the homogeneous Helmholtz equation with homogeneous boundary conditions; a particular solution of a homogeneous boundary value problem is  $w = 0$ .

The values  $\lambda_n$  of the parameter  $\lambda$  for which there are nontrivial solutions (solutions other

TABLE 23

Conformal mapping of some domains  $D$  in the  $z$ -plane onto the unit circle  $|\omega| \leq 1$ . Notation:  $z = x + iy$ ,  $\omega = u + iv$ ,  $z_0 = x_0 + iy_0$ , and  $\bar{z}_0 = x_0 - iy_0$

| No | Domain $D$ in $z$ -plane                                                               | Transformation                                                                                                   |
|----|----------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------|
| 1  | Upper half-plane:<br>$-\infty < x < \infty, 0 \leq y < \infty$                         | $\omega = e^{i\lambda} \frac{z - z_0}{z - \bar{z}_0}$ ,<br>$\lambda$ is a real number                            |
| 2  | A circle of unit radius:<br>$x^2 + y^2 \leq 1$                                         | $\omega = e^{i\lambda} \frac{z - z_0}{1 - \bar{z}_0 z}$ ,<br>$\lambda$ is a real number                          |
| 3  | Exterior of a circle of radius $a$ :<br>$x^2 + y^2 \geq a^2$                           | $\omega = \frac{a}{z}$                                                                                           |
| 4  | Infinite strip of width $a$ :<br>$-\infty < x < \infty, 0 \leq y \leq a$               | $\omega = \frac{\exp(\pi z/a) - \exp(\pi z_0/a)}{\exp(\pi z/a) - \exp(\pi \bar{z}_0/a)}$                         |
| 5  | Semicircle of radius $a$ :<br>$x^2 + y^2 \leq a^2, x \geq 0$                           | $\omega = i \frac{z^2 + 2az - a^2}{z^2 - 2az - a^2}$                                                             |
| 6  | Sector of a unit circle with angle $\beta$ :<br>$ z  \leq 1, 0 \leq \arg z \leq \beta$ | $\omega = \frac{(1 + z^{\pi/\beta})^2 - i(1 - z^{\pi/\beta})^2}{(1 + z^{\pi/\beta})^2 + i(1 - z^{\pi/\beta})^2}$ |
| 7  | Exterior of an ellipse with semiaxes $a$ and $b$ :<br>$(x/a)^2 + (y/b)^2 \geq 1$       | $z = \frac{1}{2} \left[ (a-b)\omega + \frac{a+b}{\omega} \right]$                                                |

than identical zero) of the homogeneous boundary value problem are called eigenvalues and the corresponding solutions,  $w = w_n$ , are called eigenfunctions of the boundary value problem.

In what follows, the first, second, and third boundary value problems for the two-dimensional Helmholtz equation in a finite two-dimensional domain  $S$  with boundary  $L$  are considered. For the third boundary value problem with the boundary condition

$$\frac{\partial w}{\partial N} + kw = 0 \quad \text{for } \mathbf{r} \in L,$$

it is assumed that  $k > 0$ . Here,  $\frac{\partial w}{\partial N}$  is the derivative along the outward normal to the contour  $L$ , and  $\mathbf{r} = \{x, y\}$ .

#### 7.3.1-2. Properties of eigenvalues and eigenfunctions.

1°. There are infinitely many eigenvalues  $\{\lambda_n\}$ ; the set of eigenvalues forms a discrete spectrum for the given boundary value problem.

2°. All eigenvalues are positive, except for the eigenvalue  $\lambda_0 = 0$  existing in the second boundary value problem (the corresponding eigenfunction is  $w_0 = \text{const}$ ). We number the eigenvalues in order of increasing magnitudes,  $\lambda_1 < \lambda_2 < \lambda_3 < \dots$ .

3°. The eigenvalues tend to infinity as the number  $n$  increases. The following asymptotic estimate holds:

$$\lim_{n \rightarrow \infty} \frac{n}{\lambda_n} = \frac{S_2}{4\pi},$$

where  $S_2$  is the area of the two-dimensional domain under study.

4°. The eigenfunctions  $w_n = w_n(x, y)$  are defined up to a constant multiplier. Any two eigenfunctions corresponding to different eigenvalues,  $\lambda_n \neq \lambda_m$ , are orthogonal:

$$\int_S w_n w_m dS = 0.$$

5°. Any twice continuously differentiable function  $f = f(\mathbf{r})$  that satisfies the boundary conditions of a boundary value problem can be expanded into a uniformly convergent series in the eigenfunctions of the boundary value problem:

$$f = \sum_{n=1}^{\infty} f_n w_n, \quad \text{where} \quad f_n = \frac{1}{\|w_n\|^2} \int_S f w_n dS, \quad \|w_n\|^2 = \int_S w_n^2 dS.$$

If  $f$  is square summable, then the series converges in mean.

6°. The eigenvalues of the first boundary value problem do not increase if the domain is extended.

*Remark 1.* In a two-dimensional problem, generally correspond to each eigenvalue  $\lambda_n$  finitely many linearly independent eigenfunctions  $w_n^{(1)}, w_n^{(2)}, \dots, w_n^{(p)}$ . These functions can always be replaced by their linear combinations

$$\bar{w}_n^{(j)} = c_{j,1} w_n^{(1)} + \dots + c_{j,j-1} w_n^{(j-1)} + w_n^{(j)}, \quad j = 1, 2, \dots, p,$$

so that the new eigenfunctions  $\bar{w}_n^{(1)}, \bar{w}_n^{(2)}, \dots, \bar{w}_n^{(p)}$  now are pairwise orthogonal. Therefore, without loss of generality, we assume that all the eigenfunctions are orthogonal.

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

### 7.3.1-3. Nonhomogeneous Helmholtz equation with homogeneous boundary conditions.

Three cases are possible.

1°. If the equation parameter  $\lambda$  is not equal to any one of the eigenvalues, then there exists the series solution

$$w = \sum_{n=1}^{\infty} \frac{A_n}{\lambda_n - \lambda} w_n, \quad \text{where} \quad A_n = \frac{1}{\|w_n\|^2} \int_S \Phi w_n dS, \quad \|w_n\|^2 = \int_S w_n^2 dS.$$

2°. If  $\lambda$  is equal to some eigenvalue,  $\lambda = \lambda_m$ , then the solution of the nonhomogeneous problem exists only if the function  $\Phi$  is orthogonal to  $w_m$ , i.e.,

$$\int_S \Phi w_m dS = 0.$$

In this case the system is expressed as

$$w = \sum_{n=1}^{m-1} \frac{A_n}{\lambda_n - \lambda_m} w_n + \sum_{n=m+1}^{\infty} \frac{A_n}{\lambda_n - \lambda_m} w_n + C w_m, \quad A_n = \frac{1}{\|w_n\|^2} \int_S \Phi w_n dS,$$

where  $\|w_n\|^2 = \int_S w_n^2 dS$ , and  $C$  is an arbitrary constant.

3°. If  $\lambda = \lambda_m$  and  $\int_S \Phi w_m dS \neq 0$ , then the boundary value problem for the nonhomogeneous equation does not have solutions.

*Remark 2.* If  $p_n$  mutually orthogonal eigenfunctions  $w_n^{(j)}$  ( $j = 1, 2, \dots, p_n$ ) correspond to each eigenvalue  $\lambda_n$ , then, for  $\lambda \neq \lambda_n$ , the solution is written as

$$w = \sum_{n=1}^{\infty} \sum_{j=1}^{p_n} \frac{A_n^{(j)}}{\lambda_n - \lambda} w_n^{(j)}, \quad \text{where} \quad A_n^{(j)} = \frac{1}{\|w_n^{(j)}\|^2} \int_S \Phi w_n^{(j)} dS, \quad \|w_n^{(j)}\|^2 = \int_S [w_n^{(j)}]^2 dS.$$

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

### 7.3.1-4. Solution of nonhomogeneous boundary value problem of general form.

1°. The solution of the first boundary value problem for the Helmholtz equation with the boundary condition

$$w = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in L$$

can be represented in the form

$$w(\mathbf{r}) = \int_S \Phi(\rho) G(\mathbf{r}, \rho) dS_\rho - \int_L f(\rho) \frac{\partial}{\partial N_\rho} G(\mathbf{r}, \rho) dL_\rho. \quad (1)$$

Here,  $\mathbf{r} = \{x, y\}$  and  $\rho = \{\xi, \eta\}$  ( $\mathbf{r} \in S$ ,  $\rho \in S$ );  $\frac{\partial}{\partial N_\rho}$  denotes the derivative along the outward normal to the contour  $L$  with respect to the variables  $\xi$  and  $\eta$ . The Green's function is given by the series

$$G(\mathbf{r}, \rho) = \sum_{n=1}^{\infty} \frac{w_n(\mathbf{r}) w_n(\rho)}{\|w_n\|^2 (\lambda_n - \lambda)}, \quad \lambda \neq \lambda_n, \quad (2)$$

where the  $w_n$  and  $\lambda_n$  are the eigenfunctions and eigenvalues of the homogeneous first boundary value problem.

2°. The solution of the second boundary value problem with the boundary condition

$$\frac{\partial w}{\partial N} = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in L$$

can be written as

$$w(\mathbf{r}) = \int_S \Phi(\rho) G(\mathbf{r}, \rho) dS_\rho + \int_L f(\rho) G(\mathbf{r}, \rho) dL_\rho. \quad (3)$$

Here, the Green's function is given by the series

$$G(\mathbf{r}, \rho) = -\frac{1}{S_2 \lambda} + \sum_{n=1}^{\infty} \frac{w_n(\mathbf{r}) w_n(\rho)}{\|w_n\|^2 (\lambda_n - \lambda)}, \quad \lambda \neq \lambda_n, \quad (4)$$

where  $S_2$  is the area of the two-dimensional domain under consideration, and the  $\lambda_n$  and  $w_n$  are the positive eigenvalues and the corresponding eigenfunctions of the homogeneous second boundary value problem. For clarity, the term corresponding to the zero eigenvalue  $\lambda_0 = 0$  ( $w_0 = \text{const}$ ) is singled out in (4).

3°. The solution of the third boundary value problem for the Helmholtz equation with the boundary condition

$$\frac{\partial w}{\partial N} + kw = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in L$$

is given by formula (3), where the Green's function is defined by series (2), which involves the eigenfunctions  $w_n$  and eigenvalues  $\lambda_n$  of the homogeneous third boundary value problem.

### 7.3.1-5. Boundary conditions at infinity in the case of an infinite domain.

In what follows, the function  $\Phi$  is assumed to be finite or sufficiently rapidly decaying as  $r \rightarrow \infty$ .

1°. For  $\lambda < 0$ , in the case of an infinite domain, the vanishing condition of the solution at infinity is set,

$$w \rightarrow 0 \quad \text{as } r \rightarrow \infty.$$

2°. For  $\lambda > 0$ , if the domain is unbounded, the radiation conditions (Sommerfeld conditions) at infinity are used. In two-dimensional problems, these conditions are written as

$$\lim_{r \rightarrow \infty} \sqrt{r} w = \text{const}, \quad \lim_{r \rightarrow \infty} \sqrt{r} \left( \frac{\partial w}{\partial r} + i \sqrt{\lambda} w \right) = 0,$$

where  $i^2 = -1$ .

To identify a single solution, the principle of limit absorption and the principle of limit amplitude are also used.

◎ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

### 7.3.2. Problems in Cartesian Coordinate System

A two-dimensional nonhomogeneous Helmholtz equation in the rectangular Cartesian system of coordinates has the form

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \lambda w = -\Phi(x, y).$$

#### 7.3.2-1. Particular solutions and some relations.

1°. Particular solutions of the homogeneous equation ( $\Phi \equiv 0$ ):

$$\begin{aligned} w &= (Ax + B)(C \cos \mu y + D \sin \mu y), & \lambda &= \mu^2, \\ w &= (Ax + B)(C \cosh \mu y + D \sinh \mu y), & \lambda &= -\mu^2, \\ w &= (A \cos \mu x + B \sin \mu x)(Cy + D), & \lambda &= \mu^2, \\ w &= (A \cosh \mu x + B \sinh \mu x)(Cy + D), & \lambda &= -\mu^2, \\ w &= (A \cos \mu_1 x + B \sin \mu_1 x)(C \cos \mu_2 y + D \sin \mu_2 y), & \lambda &= \mu_1^2 + \mu_2^2, \\ w &= (A \cos \mu_1 x + B \sin \mu_1 x)(C \cosh \mu_2 y + D \sinh \mu_2 y), & \lambda &= \mu_1^2 - \mu_2^2, \\ w &= (A \cosh \mu_1 x + B \sinh \mu_1 x)(C \cos \mu_2 y + D \sin \mu_2 y), & \lambda &= -\mu_1^2 + \mu_2^2, \\ w &= (A \cosh \mu_1 x + B \sinh \mu_1 x)(C \cosh \mu_2 y + D \sinh \mu_2 y), & \lambda &= -\mu_1^2 - \mu_2^2, \end{aligned}$$

where  $A$ ,  $B$ ,  $C$ , and  $D$  are arbitrary constants.

2°. Fundamental solutions:

$$\begin{aligned} \mathcal{E}(x, y) &= \frac{1}{2\pi} K_0(sr) & \text{if } \lambda &= -s^2 < 0, \\ \mathcal{E}(x, y) &= \frac{i}{4} H_0^{(1)}(kr) & \text{if } \lambda &= k^2 > 0, \\ \mathcal{E}(x, y) &= -\frac{i}{4} H_0^{(2)}(kr) & \text{if } \lambda &= k^2 > 0, \end{aligned}$$

where  $r = \sqrt{x^2 + y^2}$ ,  $K_0(z)$  is the modified Bessel function of the second kind,  $H_0^{(1)}(z)$  and  $H_0^{(2)}(z)$  are the Hankel functions of the first and second kind of order 0,  $x_0$  and  $y_0$  are arbitrary constants, and  $i^2 = -1$ . The leading term of the asymptotic expansion of the fundamental solutions, as  $r \rightarrow 0$ , is given by  $\frac{1}{2\pi} \ln \frac{1}{r}$ .

3°. Suppose  $w = w(x, y)$  is a solution of the homogeneous Helmholtz equation. Then the functions

$$\begin{aligned} w_1 &= w(x + C_1, \pm y + C_2), \\ w_2 &= w(-x + C_1, \pm y + C_2), \\ w_3 &= w(x \cos \theta + y \sin \theta + C_1, -x \sin \theta + y \cos \theta + C_2), \end{aligned}$$

where  $C_1$ ,  $C_2$ , and  $\theta$  are arbitrary constants, are also solutions of the equation.

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

#### 7.3.2-2. Domain: $-\infty < x < \infty$ , $-\infty < y < \infty$ .

1°. Solution for  $\lambda = -s^2 < 0$ :

$$w(x, y) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) K_0(s\varrho) d\xi d\eta, \quad \varrho = \sqrt{(x - \xi)^2 + (y - \eta)^2}.$$

2°. Solution for  $\lambda = k^2 > 0$ :

$$w(x, y) = -\frac{i}{4} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) H_0^{(2)}(k\varrho) d\xi d\eta, \quad \varrho = \sqrt{(x - \xi)^2 + (y - \eta)^2}.$$

The radiation conditions (Sommerfeld conditions) at infinity were used to obtain this solution (see Paragraph 7.3.1-5, Item 2°).

⊙ *References:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), A. N. Tikhonov and A. A. Samarskii (1990).

**7.3.2-3. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.**

A half-plane is considered. A boundary condition is prescribed:

$$w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} f(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi + \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta.$$

1°. The Green's function for  $\lambda = -s^2 < 0$ :

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} [K_0(s\varrho_1) - K_0(s\varrho_2)],$$

$$\varrho_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2}, \quad \varrho_2 = \sqrt{(x-\xi)^2 + (y+\eta)^2}.$$

2°. The Green's function for  $\lambda = k^2 > 0$ :

$$G(x, y, \xi, \eta) = -\frac{i}{4} [H_0^{(2)}(k\varrho_1) - H_0^{(2)}(k\varrho_2)].$$

The radiation conditions at infinity were used to obtain this relation (see Paragraph 7.3.1-5, Item 2°).

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.2-4. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ . Second boundary value problem.**

A half-plane is considered. A boundary condition is prescribed:

$$\partial_y w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = - \int_{-\infty}^{\infty} f(\xi) G(x, y, \xi, 0) d\xi + \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta.$$

1°. The Green's function for  $\lambda = -s^2 < 0$ :

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} [K_0(s\varrho_1) + K_0(s\varrho_2)],$$

$$\varrho_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2}, \quad \varrho_2 = \sqrt{(x-\xi)^2 + (y+\eta)^2}.$$

2°. The Green's function for  $\lambda = k^2 > 0$ :

$$G(x, y, \xi, \eta) = -\frac{i}{4} [H_0^{(2)}(k\varrho_1) + H_0^{(2)}(k\varrho_2)].$$

The radiation conditions at infinity were used to obtain this relation (see Paragraph 7.3.1-5, Item 2°).

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.2-5. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . First boundary value problem.**

A quadrant of the plane is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at} \quad x = 0, \quad w = f_2(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \int_0^\infty f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta + \int_0^\infty f_2(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi \\ + \int_0^\infty \int_0^\infty \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta.$$

1°. The Green's function for  $\lambda = -s^2 < 0$ :

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} [K_0(s\rho_1) - K_0(s\rho_2) - K_0(s\rho_3) + K_0(s\rho_4)], \\ \rho_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2}, \quad \rho_2 = \sqrt{(x-\xi)^2 + (y+\eta)^2}, \\ \rho_3 = \sqrt{(x+\xi)^2 + (y-\eta)^2}, \quad \rho_4 = \sqrt{(x+\xi)^2 + (y+\eta)^2}.$$

2°. The Green's function for  $\lambda = k^2 > 0$ :

$$G(x, y, \xi, \eta) = -\frac{i}{4} [H_0^{(2)}(k\rho_1) - H_0^{(2)}(k\rho_2) - H_0^{(2)}(k\rho_3) + H_0^{(2)}(k\rho_4)].$$

**7.3.2-6. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ . Second boundary value problem.**

A quadrant of the plane is considered. Boundary conditions are prescribed:

$$\partial_x w = f_1(y) \quad \text{at} \quad x = 0, \quad \partial_y w = f_2(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = - \int_0^\infty f_1(\eta) G(x, y, 0, \eta) d\eta - \int_0^\infty f_2(\xi) G(x, y, \xi, 0) d\xi \\ + \int_0^\infty \int_0^\infty \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta.$$

1°. The Green's function for  $\lambda = -s^2 < 0$ :

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} [K_0(s\rho_1) + K_0(s\rho_2) + K_0(s\rho_3) + K_0(s\rho_4)], \\ \rho_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2}, \quad \rho_2 = \sqrt{(x-\xi)^2 + (y+\eta)^2}, \\ \rho_3 = \sqrt{(x+\xi)^2 + (y-\eta)^2}, \quad \rho_4 = \sqrt{(x+\xi)^2 + (y+\eta)^2}.$$

2°. The Green's function for  $\lambda = k^2 > 0$ :

$$G(x, y, \xi, \eta) = -\frac{i}{4} [H_0^{(2)}(k\rho_1) + H_0^{(2)}(k\rho_2) + H_0^{(2)}(k\rho_3) + H_0^{(2)}(k\rho_4)].$$



**7.3.2-7. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . First boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(x) \quad \text{at} \quad y = 0, \quad w = f_2(x) \quad \text{at} \quad y = a.$$

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} f_1(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi - \int_{-\infty}^{\infty} f_2(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=a} d\xi \\ + \int_0^a \int_{-\infty}^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta.$$

Green's function:

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=1}^{\infty} \frac{1}{\beta_n} \exp(-\beta_n |x - \xi|) \sin(q_n y) \sin(q_n \eta), \quad q_n = \frac{\pi n}{a}, \quad \beta_n = \sqrt{q_n^2 - \lambda}.$$

Alternatively, the Green's function for  $\lambda = -s^2 < 0$  can be represented as

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} [K_0(s \varrho_{1n}) - K_0(s \varrho_{2n})], \\ \varrho_{n1} = \sqrt{(x - \xi)^2 + (y - \eta - 2na)^2}, \quad \varrho_{n2} = \sqrt{(x - \xi)^2 + (y + \eta + 2na)^2}.$$

**7.3.2-8. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . Second boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$\partial_y w = f_1(x) \quad \text{at} \quad y = 0, \quad \partial_y w = f_2(x) \quad \text{at} \quad y = a.$$

Solution:

$$w(x, y) = - \int_{-\infty}^{\infty} f_1(\xi) G(x, y, \xi, 0) d\xi + \int_{-\infty}^{\infty} f_2(\xi) G(x, y, \xi, a) d\xi \\ + \int_0^a \int_{-\infty}^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta.$$

Green's function:

$$G(x, y, \xi, \eta) = \frac{1}{2a} \sum_{n=0}^{\infty} \frac{\varepsilon_n}{\beta_n} \exp(-\beta_n |x - \xi|) \cos(q_n y) \cos(q_n \eta), \\ q_n = \frac{\pi n}{a}, \quad \beta_n = \sqrt{q_n^2 - \lambda}, \quad \varepsilon = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}$$

Alternatively, the Green's function for  $\lambda = -s^2 < 0$  can be represented as

$$G(x, y, \xi, \eta) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} [K_0(s \varrho_{1n}) + K_0(s \varrho_{2n})], \\ \varrho_{n1} = \sqrt{(x - \xi)^2 + (y - \eta_{n1})^2}, \quad \eta_{n1} = 2na + \eta, \\ \varrho_{n2} = \sqrt{(x - \xi)^2 + (y - \eta_{n2})^2}, \quad \eta_{n2} = 2na - \eta.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.2-9. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . Third boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$\partial_y w - k_1 w = f_1(x) \quad \text{at } y = 0, \quad \partial_y w + k_2 w = f_2(x) \quad \text{at } y = a.$$

The solution  $w(x, y)$  is determined by the formula in Paragraph 7.3.2-8 where

$$G(x, y, \xi, \eta) = \frac{1}{2} \sum_{n=1}^{\infty} \frac{\varphi_n(y)\varphi_n(\eta)}{\|\varphi_n\|^2 \beta_n} \exp(-\beta_n |x - \xi|), \quad \beta_n = \sqrt{\mu_n^2 - \lambda},$$

$$\varphi_n(y) = \mu_n \cos(\mu_n y) + k_1 \sin(\mu_n y), \quad \|\varphi_n\|^2 = \frac{1}{2}(\mu_n^2 + k_1^2) \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right].$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}$ .

**7.3.2-10. Domain:  $-\infty < x < \infty$ ,  $0 \leq y \leq a$ . Mixed boundary value problem.**

An infinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(x) \quad \text{at } y = 0, \quad \partial_y w = f_2(x) \quad \text{at } y = a.$$

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} f_1(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi + \int_{-\infty}^{\infty} f_2(\xi) G(x, y, \xi, a) d\xi + \int_0^a \int_{-\infty}^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta,$$

where

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=0}^{\infty} \frac{1}{\beta_n} \exp(-\beta_n |x - \xi|) \sin(q_n y) \sin(q_n \eta), \quad q_n = \frac{\pi(2n+1)}{2a}, \quad \beta_n = \sqrt{q_n^2 - \lambda}.$$

**7.3.2-11. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . First boundary value problem.**

A semiinfinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at } x = 0, \quad w = f_2(x) \quad \text{at } y = 0, \quad w = f_3(x) \quad \text{at } y = a.$$

Solution:

$$w(x, y) = \int_0^a \int_0^{\infty} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta + \int_0^a f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta + \int_0^{\infty} f_2(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi - \int_0^{\infty} f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=a} d\xi,$$

where

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=1}^{\infty} \frac{1}{\beta_n} [\exp(-\beta_n |x - \xi|) - \exp(-\beta_n |x + \xi|)] \sin(q_n y) \sin(q_n \eta),$$

$$q_n = \frac{\pi n}{a}, \quad \beta_n = \sqrt{q_n^2 - \lambda}.$$

**7.3.2-12. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . Second boundary value problem.**

A semiinfinite strip is considered. Boundary conditions are prescribed:

$$\partial_x w = f_1(y) \quad \text{at } x = 0, \quad \partial_y w = f_2(x) \quad \text{at } y = 0, \quad \partial_y w = f_3(x) \quad \text{at } y = a.$$

Solution:

$$w(x, y) = \int_0^a \int_0^\infty \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta - \int_0^a f_1(\eta) G(x, y, 0, \eta) d\eta \\ - \int_0^\infty f_2(\xi) G(x, y, \xi, 0) d\xi + \int_0^\infty f_3(\xi) G(x, y, \xi, a) d\xi,$$

where

$$G(x, y, \xi, \eta) = \frac{1}{2a} \sum_{n=0}^{\infty} \frac{\varepsilon_n}{\beta_n} [\exp(-\beta_n|x - \xi|) + \exp(-\beta_n|x + \xi|)] \cos(q_n y) \cos(q_n \eta), \\ q_n = \frac{\pi n}{a}, \quad \beta_n = \sqrt{q_n^2 - \lambda}, \quad \varepsilon = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}$$

**7.3.2-13. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . Third boundary value problem.**

A semiinfinite strip is considered. Boundary conditions are prescribed:

$$\partial_x w - k_1 w = f_1(y) \quad \text{at } x = 0, \quad \partial_y w - k_2 w = f_2(x) \quad \text{at } y = 0, \quad \partial_y w + k_3 w = f_3(x) \quad \text{at } y = a.$$

The solution  $w(x, y)$  is determined by the formula in Paragraph 7.3.2-12 where

$$G(x, y, \xi, \eta) = \sum_{n=1}^{\infty} \frac{\varphi_n(y)\varphi_n(\eta)}{\|\varphi_n\|^2 \beta_n(\beta_n + k_1)} H_n(x, \xi), \quad \beta_n = \sqrt{\mu_n^2 - \lambda}, \\ \varphi_n(y) = \mu_n \cos(\mu_n y) + k_2 \sin(\mu_n y), \quad \|\varphi_n\|^2 = \frac{1}{2}(\mu_n^2 + k_2^2) \left[ a + \frac{(k_2 + k_3)(\mu_n^2 + k_2 k_3)}{(\mu_n^2 + k_2^2)(\mu_n^2 + k_3^2)} \right], \\ H_n(x, \xi) = \begin{cases} \exp(-\beta_n x) [\beta_n \cosh(\beta_n \xi) + k_1 \sinh(\beta_n \xi)] & \text{for } x > \xi, \\ \exp(-\beta_n \xi) [\beta_n \cosh(\beta_n x) + k_1 \sinh(\beta_n x)] & \text{for } \xi > x. \end{cases}$$

Here, the  $\mu_n$  are positive roots of the transcendental equation  $\tan(\mu a) = \frac{(k_2 + k_3)\mu}{\mu^2 - k_2 k_3}$ .

**7.3.2-14. Domain:  $0 \leq x < \infty$ ,  $0 \leq y \leq a$ . Mixed boundary value problems.**

1°. A semiinfinite strip is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at } x = 0, \quad \partial_y w = f_2(x) \quad \text{at } y = 0, \quad \partial_y w = f_3(x) \quad \text{at } y = a.$$

Solution:

$$w(x, y) = \int_0^a f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta - \int_0^\infty f_2(\xi) G(x, y, \xi, 0) d\xi \\ + \int_0^\infty f_3(\xi) G(x, y, \xi, a) d\xi + \int_0^a \int_0^\infty \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta,$$

where

$$G(x, y, \xi, \eta) = \frac{1}{2a} \sum_{n=0}^{\infty} \frac{\varepsilon_n}{\beta_n} [\exp(-\beta_n|x - \xi|) - \exp(-\beta_n|x + \xi|)] \cos(q_n y) \cos(q_n \eta), \\ q_n = \frac{\pi n}{a}, \quad \beta_n = \sqrt{q_n^2 - \lambda}, \quad \varepsilon = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}$$

2°. A semiinfinite strip is considered. Boundary conditions are prescribed:

$$\partial_x w = f_1(y) \quad \text{at } x = 0, \quad w = f_2(x) \quad \text{at } y = 0, \quad w = f_3(x) \quad \text{at } y = a.$$

Solution:

$$w(x, y) = - \int_0^a f_1(\eta) G(x, y, 0, \eta) d\eta + \int_0^\infty f_2(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi \\ - \int_0^\infty f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=a} d\xi + \int_0^a \int_0^\infty \Phi(\xi, \eta) G(x, y, \xi, \eta) d\xi d\eta,$$

where

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=1}^{\infty} \frac{1}{\beta_n} [\exp(-\beta_n|x - \xi|) + \exp(-\beta_n|x + \xi|)] \sin(q_n y) \sin(q_n \eta), \\ q_n = \frac{\pi n}{a}, \quad \beta_n = \sqrt{q_n^2 - \lambda}.$$

7.3.2-15. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . First boundary value problem.

A rectangle is considered. Boundary conditions are prescribed:

$$w = f_1(y) \quad \text{at } x = 0, \quad w = f_2(y) \quad \text{at } x = a, \\ w = f_3(x) \quad \text{at } y = 0, \quad w = f_4(x) \quad \text{at } y = b.$$

1°. Eigenvalues of the one-dimensional problem (it is convenient to label them with a double subscript):

$$\lambda_{nm} = \pi^2 \left( \frac{n^2}{a^2} + \frac{m^2}{b^2} \right); \quad n = 1, 2, \dots; \quad m = 1, 2, \dots$$

Eigenfunctions and the norm squared:

$$w_{nm} = \sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{m\pi y}{b}\right), \quad \|w_{nm}\|^2 = \frac{ab}{4}.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

2°. Solution for  $\lambda \neq \lambda_{nm}$ :

$$w(x, y) = \int_0^a \int_0^b \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi \\ + \int_0^b f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=0} d\eta - \int_0^b f_2(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=a} d\eta \\ + \int_0^a f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi - \int_0^a f_4(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=b} d\xi.$$

Two forms of representation of the Green's function:

$$G(x, y, \xi, \eta) = \frac{2}{a} \sum_{n=1}^{\infty} \frac{\sin(p_n x) \sin(p_n \xi)}{\beta_n \sinh(\beta_n b)} H_n(y, \eta) = \frac{2}{b} \sum_{m=1}^{\infty} \frac{\sin(q_m y) \sin(q_m \eta)}{\mu_m \sinh(\mu_m a)} Q_m(x, \xi),$$

where

$$p_n = \frac{\pi n}{a}, \quad \beta_n = \sqrt{p_n^2 - \lambda}, \quad H_n(y, \eta) = \begin{cases} \sinh(\beta_n \eta) \sinh[\beta_n(b - y)] & \text{for } b \geq y > \eta \geq 0, \\ \sinh(\beta_n y) \sinh[\beta_n(b - \eta)] & \text{for } b \geq \eta > y \geq 0, \end{cases} \\ q_m = \frac{\pi m}{b}, \quad \mu_m = \sqrt{q_m^2 - \lambda}, \quad Q_m(x, \xi) = \begin{cases} \sinh(\mu_m \xi) \sinh[\mu_m(a - x)] & \text{for } a \geq x > \xi \geq 0, \\ \sinh(\mu_m x) \sinh[\mu_m(a - \xi)] & \text{for } a \geq \xi > x \geq 0. \end{cases}$$

Alternatively, the Green's function can be written as the double series

$$G(x, y, \xi, \eta) = \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta)}{p_n^2 + q_m^2 - \lambda}, \quad p_n = \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}.$$

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w &= f_1(y) & \text{at } x=0, & & \partial_x w &= f_2(y) & \text{at } x=a, \\ \partial_y w &= f_3(x) & \text{at } y=0, & & \partial_y w &= f_4(x) & \text{at } y=b. \end{aligned}$$

1°. Eigenvalues of the homogeneous problem:

$$\lambda_{nm} = \pi^2 \left( \frac{n^2}{a^2} + \frac{m^2}{b^2} \right); \quad n = 0, 1, 2, \dots; \quad m = 0, 1, 2, \dots$$

Eigenfunctions and the norm squared:

$$w_{nm} = \cos\left(\frac{n\pi x}{a}\right) \cos\left(\frac{m\pi y}{b}\right), \quad \|w_{nm}\|^2 = \frac{ab}{4}(1 + \delta_{n0})(1 + \delta_{m0}), \quad \delta_{n0} = \begin{cases} 1 & \text{for } n=0, \\ 0 & \text{for } n \neq 0. \end{cases}$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

2°. Solution for  $\lambda \neq \lambda_{nm}$ :

$$\begin{aligned} w(x, y) &= \int_0^a \int_0^b \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi \\ &\quad - \int_0^b f_1(\eta) G(x, y, 0, \eta) d\eta + \int_0^b f_2(\eta) G(x, y, a, \eta) d\eta \\ &\quad - \int_0^a f_3(\xi) G(x, y, \xi, 0) d\xi + \int_0^a f_4(\xi) G(x, y, \xi, b) d\xi. \end{aligned}$$

Two forms of representation of the Green's function:

$$G(x, y, \xi, \eta) = \frac{1}{a} \sum_{n=0}^{\infty} \frac{\varepsilon_n \cos(p_n x) \cos(p_n \xi)}{\beta_n \sinh(\beta_n b)} H_n(y, \eta) = \frac{1}{b} \sum_{m=0}^{\infty} \frac{\varepsilon_m \cos(q_m y) \cos(q_m \eta)}{\mu_m \sinh(\mu_m a)} Q_m(x, \xi),$$

where

$$\begin{aligned} p_n &= \frac{\pi n}{a}, \quad H_n(y, \eta) = \begin{cases} \cosh(\beta_n \eta) \cosh[\beta_n(b-y)] & \text{for } y > \eta, \\ \cosh(\beta_n y) \cosh[\beta_n(b-\eta)] & \text{for } \eta > y, \end{cases} \\ q_m &= \frac{\pi m}{b}, \quad Q_m(x, \xi) = \begin{cases} \cosh(\mu_m \xi) \cosh[\mu_m(a-x)] & \text{for } x > \xi, \\ \cosh(\mu_m x) \cosh[\mu_m(a-\xi)] & \text{for } \xi > x, \end{cases} \\ \beta_n &= \sqrt{p_n^2 - \lambda}, \quad \mu_m = \sqrt{q_m^2 - \lambda}, \quad \varepsilon_n = \begin{cases} 1 & \text{for } n=0, \\ 2 & \text{for } n \neq 0. \end{cases} \end{aligned}$$

The Green's function can also be written as the double series

$$G(x, y, \xi, \eta) = \frac{1}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{\varepsilon_n \varepsilon_m \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta)}{p_n^2 + q_m^2 - \lambda}, \quad p_n = \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}.$$

► In Paragraphs 7.3.2-17 through 7.3.2-20, only the eigenvalues and eigenfunctions of homogeneous boundary value problems for the homogeneous Helmholtz equation (with  $\Phi \equiv 0$ ) are given. The solutions of the corresponding nonhomogeneous problems can be constructed using formulas presented in Paragraphs 7.3.1-3 and 7.3.1-4.

**7.3.2-17. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . Third boundary value problem.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w - k_1 w &= 0 & \text{at } x = 0, & \quad \partial_x w + k_2 w = 0 & \text{at } x = a, \\ \partial_y w - k_3 w &= 0 & \text{at } y = 0, & \quad \partial_y w + k_4 w = 0 & \text{at } y = b. \end{aligned}$$

Eigenvalues:

$$\lambda_{nm} = \mu_n^2 + \nu_m^2,$$

where the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}, \quad \tan(\nu b) = \frac{(k_3 + k_4)\nu}{\nu^2 - k_3 k_4}.$$

Eigenfunctions:

$$w_{nm} = (\mu_n \cos \mu_n x + k_1 \sin \mu_n x)(\nu_m \cos \nu_m y + k_3 \sin \nu_m y).$$

The square of the norm of an eigenfunction:

$$\|w_{nm}\|^2 = \frac{1}{4}(\mu_n^2 + k_1^2)(\nu_m^2 + k_3^2) \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right] \left[ b + \frac{(k_3 + k_4)(\nu_m^2 + k_3 k_4)}{(\nu_m^2 + k_3^2)(\nu_m^2 + k_4^2)} \right].$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.2-18. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ . Mixed boundary value problems.**

1°. A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= 0 & \text{at } x = 0, & \quad w = 0 & \text{at } x = a, \\ \partial_y w &= 0 & \text{at } y = 0, & \quad \partial_y w = 0 & \text{at } y = b. \end{aligned}$$

Eigenvalues:

$$\lambda_{nm} = \pi^2 \left( \frac{n^2}{a^2} + \frac{m^2}{b^2} \right); \quad n = 1, 2, 3, \dots; \quad m = 0, 1, 2, \dots$$

Eigenfunctions and the norm squared:

$$w_{nm} = \sin\left(\frac{n\pi x}{a}\right) \cos\left(\frac{m\pi y}{b}\right), \quad \|w_{nm}\|^2 = \frac{ab}{4}(1 + \delta_{m0}), \quad \delta_{m0} = \begin{cases} 1 & \text{for } m = 0, \\ 0 & \text{for } m \neq 0. \end{cases}$$

2°. A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= 0 & \text{at } x = 0, & \quad \partial_x w = 0 & \text{at } x = a, \\ w &= 0 & \text{at } y = 0, & \quad \partial_y w = 0 & \text{at } y = b. \end{aligned}$$

Eigenvalues:

$$\lambda_{nm} = \frac{\pi^2}{4} \left[ \frac{(2n+1)^2}{a^2} + \frac{(2m+1)^2}{b^2} \right]; \quad n = 0, 1, 2, \dots; \quad m = 0, 1, 2, \dots$$

Eigenfunctions and the norm squared:

$$w_{nm} = \sin\left[\frac{\pi(2n+1)x}{2a}\right] \sin\left[\frac{\pi(2m+1)y}{2b}\right], \quad \|w_{nm}\|^2 = \frac{ab}{4}.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 7.3.2-19. First boundary value problem for a triangular domain.

The sides of the triangle are defined by the equations

$$x = 0, \quad y = 0, \quad y = a - x.$$

The unknown quantity is zero for these sides.

Eigenvalues:

$$\lambda_{nm} = \frac{\pi^2}{a^2} [(n+m)^2 + m^2]; \quad n = 1, 2, \dots; \quad m = 1, 2, \dots$$

Eigenfunctions:

$$w_{nm} = \sin \left[ \frac{\pi}{a}(n+m)x \right] \sin \left( \frac{\pi}{a}my \right) - (-1)^n \sin \left( \frac{\pi}{a}mx \right) \sin \left[ \frac{\pi}{a}(n+m)y \right].$$

☉ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

### 7.3.2-20. Second boundary value problem for a triangular domain.

The sides of the triangle are defined by the equations

$$x = 0, \quad y = 0, \quad y = a - x.$$

The normal derivative of the unknown quantity for these sides is zero.

Eigenvalues:

$$\lambda_{nm} = \frac{\pi^2}{a^2} [(n+m)^2 + m^2]; \quad n = 0, 1, \dots; \quad m = 0, 1, \dots$$

Eigenfunctions:

$$w_{nm} = \cos \left[ \frac{\pi}{a}(n+m)x \right] \cos \left( \frac{\pi}{a}my \right) - (-1)^n \cos \left( \frac{\pi}{a}mx \right) \cos \left[ \frac{\pi}{a}(n+m)y \right].$$

## 7.3.3. Problems in Polar Coordinate System

A two-dimensional nonhomogeneous Helmholtz equation in the polar coordinate system is written as

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \lambda w = -\Phi(r, \varphi, z), \quad r = \sqrt{x^2 + y^2}.$$

### 7.3.3-1. Particular solutions of the homogeneous equation ( $\Phi \equiv 0$ ):

$$w = [AJ_0(\mu r) + BY_0(\mu r)](C\varphi + D), \quad \lambda = \mu^2,$$

$$w = [AI_0(\mu r) + BK_0(\mu r)](C\varphi + D), \quad \lambda = -\mu^2,$$

$$w = [AJ_m(\mu r) + BY_m(\mu r)](C \cos m\varphi + D \sin m\varphi), \quad \lambda = \mu^2,$$

$$w = [AI_m(\mu r) + BK_m(\mu r)](C \cos m\varphi + D \sin m\varphi), \quad \lambda = -\mu^2,$$

where  $m = 1, 2, \dots$ ;  $A, B, C, D$  are arbitrary constants; the  $J_m(\mu)$  and  $Y_m(\mu)$  are the Bessel functions; and the  $I_m(\mu)$  and  $K_m(\mu)$  are the modified Bessel functions.

► In Paragraphs 7.3.3-2 through 7.3.3-11, only the eigenvalues and eigenfunctions of homogeneous boundary value problems for the homogeneous Helmholtz equation (with  $\Phi \equiv 0$ ) are given. The solutions of the corresponding nonhomogeneous problems can be constructed using formulas presented in Paragraphs 7.3.1-3 and 7.3.1-4.

### 7.3.3-2. Domain: $0 \leq r \leq R$ . First boundary value problem.

A circle is considered. A boundary condition is prescribed:

$$w = 0 \quad \text{at} \quad r = R.$$

Eigenvalues:

$$\lambda_{nm} = \frac{\mu_{nm}^2}{R^2}; \quad n = 0, 1, 2, \dots; \quad m = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are positive zeros of the Bessel functions,  $J_n(\mu) = 0$ .

Eigenfunctions:

$$w_{nm}^{(1)} = J_n(r\sqrt{\lambda_{nm}}) \cos n\varphi, \quad w_{nm}^{(2)} = J_n(r\sqrt{\lambda_{nm}}) \sin n\varphi.$$

Eigenfunctions possessing the axial symmetry property:  $w_{0m}^{(1)} = J_0(r\sqrt{\lambda_{0m}})$ .

The square of the norm of an eigenfunction is given by

$$\|w_{nm}^{(k)}\|^2 = \frac{1}{2}\pi R^2(1 + \delta_{n0})[J'_n(\mu_{nm})]^2, \quad k = 1, 2; \quad \delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 7.3.3-3. Domain: $0 \leq r \leq R$ . Second boundary value problem.

A circle is considered. A boundary condition is prescribed:

$$\partial_r w = 0 \quad \text{at} \quad r = R.$$

Eigenvalues:

$$\lambda_{nm} = \frac{\mu_{nm}^2}{R^2},$$

where the  $\mu_{nm}$  are roots of the transcendental equation  $J'_n(\mu) = 0$ .

Eigenfunctions:

$$w_{nm}^{(1)} = J_n(r\sqrt{\lambda_{nm}}) \cos n\varphi, \quad w_{nm}^{(2)} = J_n(r\sqrt{\lambda_{nm}}) \sin n\varphi.$$

Here,  $n = 0, 1, 2, \dots$ ; for  $n \neq 0$ , the parameter  $m$  assumes the values  $m = 1, 2, 3, \dots$ ; for  $n = 0$ , a root  $\mu_{00} = 0$  (the corresponding eigenfunction is  $w_{00} = 1$ ).

Eigenfunctions possessing the axial symmetry property:  $w_{0m}^{(1)} = J_0(r\sqrt{\lambda_{0m}})$ .

The square of the norm of an eigenfunction is given by

$$\|w_{nm}^{(k)}\|^2 = \frac{\pi^2 R^2(1 + \delta_{n0})}{2\mu_{nm}^2}(\mu_{nm}^2 - n^2)[J_n(\mu_{nm})]^2, \quad \|w_{00}\|^2 = \pi R^2,$$

where  $k = 1, 2$ ;  $\delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).



### 7.3.3-4. Domain: $0 \leq r \leq R$ . Third boundary value problem.

A circle is considered. A boundary condition is prescribed:

$$\partial_r w + kw = 0 \quad \text{at} \quad r = R.$$

Eigenvalues:

$$\lambda_{nm} = \frac{\mu_{nm}^2}{R^2}; \quad n = 0, 1, 2, \dots; \quad m = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  is the  $m$ th root of the transcendental equation  $\mu J'_n(\mu) + kR J_n(\mu) = 0$ .

Eigenfunctions:

$$w_{nm}^{(1)} = J_n(r\sqrt{\lambda_{nm}}) \cos n\varphi, \quad w_{nm}^{(2)} = J_n(r\sqrt{\lambda_{nm}}) \sin n\varphi.$$

The square of the norm of an eigenfunction is given by

$$\|w_{nm}^{(1)}\|^2 = \|w_{nm}^{(2)}\|^2 = \frac{\pi R^2(1 + \delta_{n0})}{2\mu_{nm}^2} (k^2 R^2 + \mu_{nm}^2 - n^2) [J_n(\mu_{nm})]^2, \quad \delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 7.3.3-5. Domain: $R_1 \leq r \leq R_2$ . First boundary value problem.

An annular domain is considered. Boundary conditions are prescribed:

$$w = 0 \quad \text{at} \quad r = R_1, \quad w = 0 \quad \text{at} \quad r = R_2.$$

Eigenvalues:

$$\lambda_{nm} = \mu_{nm}^2; \quad n = 0, 1, 2, \dots; \quad m = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - J_n(\mu R_2) Y_n(\mu R_1) = 0.$$

Eigenfunctions:

$$\begin{aligned} w_{nm}^{(1)} &= [J_n(\mu_{nm} R_1) Y_n(\mu_{nm} R_2) - J_n(\mu_{nm} R_2) Y_n(\mu_{nm} R_1)] \cos n\varphi, \\ w_{nm}^{(2)} &= [J_n(\mu_{nm} R_1) Y_n(\mu_{nm} R_2) - J_n(\mu_{nm} R_2) Y_n(\mu_{nm} R_1)] \sin n\varphi. \end{aligned}$$

The square of the norm of an eigenfunction is given by

$$\|w_{nm}^{(1)}\|^2 = \|w_{nm}^{(2)}\|^2 = \frac{2(1 + \delta_{n0})}{\pi \mu_{nm}^2} \frac{J_n^2(\mu_{nm} R_1) - J_n^2(\mu_{nm} R_2)}{J_n^2(\mu_{nm} R_2)}, \quad \delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 7.3.3-6. Domain: $R_1 \leq r \leq R_2$ . Second boundary value problem.

An annular domain is considered. Boundary conditions are prescribed:

$$\partial_r w = 0 \quad \text{at} \quad r = R_1, \quad \partial_r w = 0 \quad \text{at} \quad r = R_2.$$

Eigenvalues:

$$\lambda_{nm} = \mu_{nm}^2; \quad n = 0, 1, 2, \dots; \quad m = 0, 1, 2, \dots$$

Here, the  $\mu_{nm}$  are roots of the transcendental equation

$$J'_n(\mu R_1)Y'_n(\mu R_2) - J'_n(\mu R_2)Y'_n(\mu R_1) = 0.$$

If  $n = 0$ , there is a root  $\mu_{00} = 0$  and the corresponding eigenfunction is  $w_{00}^{(1)} = 1$ .

Eigenfunctions:

$$w_{nm}^{(1)} = [J_n(\mu_{nm}r)Y'_n(\mu_{nm}R_1) - J'_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \cos n\varphi,$$

$$w_{nm}^{(2)} = [J_n(\mu_{nm}r)Y'_n(\mu_{nm}R_1) - J'_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \sin n\varphi.$$

The square of the norm of an eigenfunction is given by

$$\|w_{nm}^{(1)}\|^2 = \|w_{nm}^{(2)}\|^2 = \frac{2(1 + \delta_{n0})}{\pi \mu_{nm}^2} \left\{ \left( 1 - \frac{n^2}{R_2^2 \mu_{nm}^2} \right) \left[ \frac{J'_n(\mu_{nm}R_1)}{J'_n(\mu_{nm}R_2)} \right]^2 - \left( 1 - \frac{n^2}{R_1^2 \mu_{nm}^2} \right) \right\},$$

$$\|w_{00}^{(1)}\|^2 = \pi(R_2^2 - R_1^2); \quad \delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 7.3.3-7. Domain: $R_1 \leq r \leq R_2$ . Third boundary value problem.

An annular domain is considered. Boundary conditions are prescribed:

$$\partial_r w - kw = 0 \quad \text{at } r = R_1, \quad \partial_r w + kw = 0 \quad \text{at } r = R_2.$$

Eigenvalues:

$$\lambda_{nm} = \mu_{nm}^2; \quad n = 0, 1, 2, \dots; \quad m = 1, 2, 3, \dots;$$

where the  $\mu_{nm}$  are positive roots of the transcendental equation

$$A_1(\mu R_1)B_2(\mu R_2) - A_2(\mu R_2)B_1(\mu R_1) = 0.$$

Here, we use the notation

$$A_1(\mu R) = J'_n(\mu R) - \frac{k}{\mu} J_n(\mu R), \quad B_1(\mu R) = Y'_n(\mu R) - \frac{k}{\mu} Y_n(\mu R),$$

$$A_2(\mu R) = J'_n(\mu R) + \frac{k}{\mu} J_n(\mu R), \quad B_2(\mu R) = Y'_n(\mu R) + \frac{k}{\mu} Y_n(\mu R).$$

Eigenfunctions:

$$w_{nm}^{(1)} = [B_1(\mu_{nm}R_1)J_n(\mu_{nm}r) - A_1(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \cos n\varphi,$$

$$w_{nm}^{(2)} = [B_1(\mu_{nm}R_1)J_n(\mu_{nm}r) - A_1(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \sin n\varphi.$$

The square of the norm of an eigenfunction is given by ( $s = 1, 2$ )

$$\|w_{nm}^{(s)}\|^2 = \frac{1}{2} \pi \varepsilon_n R_2^2 \left\{ [F'_{nm}(R_2)]^2 + \left( 1 - \frac{n^2}{R_2^2 \mu_{nm}^2} \right) F_{nm}^2(R_2) \right\}$$

$$- \frac{1}{2} \pi \varepsilon_n R_1^2 \left\{ [F'_{nm}(R_1)]^2 + \left( 1 - \frac{n^2}{R_1^2 \mu_{nm}^2} \right) F_{nm}^2(R_1) \right\},$$

$$F_{nm}(r) = B_1(\mu_{nm}R_1)J_n(\mu_{nm}r) - A_1(\mu_{nm}R_1)Y_n(\mu_{nm}r), \quad \varepsilon_{ij} = \begin{cases} 2 & \text{for } i = j, \\ 1 & \text{for } i \neq j. \end{cases}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.3-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \alpha$ . First boundary value problem.**

A circular sector is considered. Boundary conditions are prescribed:

$$w = 0 \quad \text{at} \quad r = R, \quad w = 0 \quad \text{at} \quad \varphi = 0, \quad w = 0 \quad \text{at} \quad \varphi = \alpha.$$

Eigenvalues:

$$\lambda_{nm} = \frac{\mu_{nm}^2}{R^2}; \quad n = 1, 2, 3, \dots; \quad m = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are positive zeros of the Bessel functions,  $J_{\frac{n\pi}{\alpha}}(\mu) = 0$ .

Eigenfunctions:

$$w_{nm} = J_{\frac{n\pi}{\alpha}}\left(\mu_{nm} \frac{r}{R}\right) \sin\left(\frac{n\pi}{\alpha}\varphi\right).$$

The square of the norm of an eigenfunction is given by

$$\|w_{nm}\|^2 = \frac{\alpha R^2}{4} \left[ J'_{\frac{n\pi}{\alpha}}(\mu_{nm}) \right]^2.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.3-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \alpha$ . Second boundary value problem.**

A circular sector is considered. Boundary conditions are prescribed:

$$\partial_r w = 0 \quad \text{at} \quad r = R, \quad \partial_\varphi w = 0 \quad \text{at} \quad \varphi = 0, \quad \partial_\varphi w = 0 \quad \text{at} \quad \varphi = \alpha.$$

Eigenvalues:

$$\lambda_{nm} = \frac{\mu_{nm}^2}{R^2}; \quad n = 0, 1, 2, \dots; \quad m = 0, 1, 2, \dots$$

Here, the  $\mu_{nm}$  are roots of the transcendental equation  $J'_{\frac{n\pi}{\alpha}}(\mu) = 0$ .

Eigenfunctions:

$$w_{nm} = J_{\frac{n\pi}{\alpha}}\left(\mu_{nm} \frac{r}{R}\right) \cos\left(\frac{n\pi}{\alpha}\varphi\right), \quad w_{00} = 1.$$

The square of the norm of an eigenfunction is given by

$$\|w_{nm}\|^2 = \frac{\alpha R^2}{4} (1 + \delta_{n0}) \left(1 - \frac{n^2}{\mu_{nm}^2}\right) \left[ J_{\frac{n\pi}{\alpha}}(\mu_{nm}) \right]^2, \quad \|w_{00}\|^2 = \frac{\alpha R^2}{2}.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.3-10. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \alpha$ . Third boundary value problem.**

A circular sector is considered. Boundary conditions are prescribed:

$$\partial_r w + k_1 w = 0 \quad \text{at} \quad r = R, \quad \partial_\varphi w - k_2 w = 0 \quad \text{at} \quad \varphi = 0, \quad \partial_\varphi w + k_3 w = 0 \quad \text{at} \quad \varphi = \alpha.$$

Eigenvalues:

$$\lambda_{nm} = \frac{\mu_{nm}^2}{R^2}; \quad n = 1, 2, 3, \dots; \quad m = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are positive roots of the transcendental equation  $\mu J'_{\nu_n}(\mu) + k_1 R J_{\nu_n}(\mu) = 0$ ; the  $\nu_n$  are positive roots of the transcendental equation  $\tan(\alpha\nu) = \frac{(k_2 + k_3)\nu}{\nu^2 - k_2 k_3}$ .

Eigenfunctions:

$$w_{nm} = J_{\nu_n}\left(\mu_{nm} \frac{r}{R}\right) \frac{\nu_n \cos(\nu_n \varphi) + k_2 \sin(\nu_n \varphi)}{\sqrt{\nu_n^2 + k_2^2}}.$$

The square of the norm of an eigenfunction is given by

$$\|w_{nm}\|^2 = \frac{R^2}{4} \left[ \alpha + \frac{(k_2 + k_3)(\nu_n^2 + k_2 k_3)}{(\nu_n^2 + k_2^2)(\nu_n^2 + k_3^2)} \right] \left( 1 + \frac{k_1^2 R^2 - \nu_n^2}{\mu_{nm}^2} \right) J_{\nu_n}^2(\mu_{nm}).$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**7.3.3-11. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq \alpha$ . First boundary value problem.**

Boundary conditions are prescribed:

$$\begin{aligned} w = 0 & \quad \text{at} \quad r = R_1, & w = 0 & \quad \text{at} \quad r = R_2, \\ w = 0 & \quad \text{at} \quad \varphi = 0, & w = 0 & \quad \text{at} \quad \varphi = \alpha. \end{aligned}$$

Eigenvalues:

$$\lambda_{nm} = \mu_{nm}^2,$$

where the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_{\nu_n}(\mu R_1)Y_{\nu_n}(\mu R_2) - J_{\nu_n}(\mu R_2)Y_{\nu_n}(\mu R_1) = 0, \quad \nu_n = \frac{n\pi}{\alpha}.$$

Eigenfunctions:

$$w_{nm} = [J_{\nu_n}(\mu_{nm}r)Y_{\nu_n}(\mu_{nm}R_1) - J_{\nu_n}(\mu_{nm}R_1)Y_{\nu_n}(\mu_{nm}r)] \sin(\nu_n\varphi).$$

The square of the norm of an eigenfunction is given by

$$\|w_{nm}\|^2 = \frac{\alpha}{\pi^2 \mu_{nm}^2} \frac{[J_{\nu_n}(\mu_{nm}R_1)]^2 - [J_{\nu_n}(\mu_{nm}R_2)]^2}{[J_{\nu_n}(\mu_{nm}R_2)]^2}.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 7.3.4. Other Orthogonal Coordinate Systems. Elliptic Domain

In Paragraphs 7.3.4-1 and 7.3.4-2, two other orthogonal systems of coordinates are described in which the homogeneous Helmholtz equation admits separation of variables.

**7.3.4-1. Parabolic coordinate system.**

In the parabolic coordinates that are introduced by the relations

$$x = \frac{1}{2}(\xi^2 - \eta^2), \quad y = \xi\eta \quad (0 \leq \xi < \infty, -\infty < \eta < \infty),$$

the Helmholtz equation has the form

$$\frac{\partial^2 w}{\partial \xi^2} + \frac{\partial^2 w}{\partial \eta^2} + \lambda(\xi^2 + \eta^2)w = 0.$$

Setting  $w = f(\xi)g(\eta)$ , we arrive at the following linear ordinary differential equations for  $f = f(\xi)$  and  $g = g(\eta)$ :

$$f'' + (\lambda\xi^2 + k)f = 0, \quad g'' + (\lambda\eta^2 - k)g = 0,$$

where  $k$  is the separation constant. The general solutions of these equations are given by

$$\begin{aligned} f(\xi) &= A_1 D_{\mu-1/2}(\sigma\xi) + A_2 D_{\mu-1/2}(-\sigma\xi), & g(\eta) &= B_1 D_{-\mu-1/2}(\sigma\eta) + B_2 D_{-\mu-1/2}(-\sigma\eta), \\ \mu &= \frac{1}{2}k(-\lambda)^{-1/2}, & \sigma &= (-4\lambda)^{1/4}. \end{aligned}$$

Here,  $A_1$ ,  $B_1$ ,  $A_2$ , and  $B_2$  are arbitrary constants, and  $D_\nu(z)$  is the parabolic cylinder function,

$$D_\nu(z) = 2^{1/2} \exp(-\frac{1}{4}z^2) \left[ \frac{\Gamma(\frac{1}{2})}{\Gamma(\frac{1}{2} - \frac{\nu}{2})} \Phi(-\frac{\nu}{2}, \frac{1}{2}; \frac{1}{2}z^2) + 2^{-1/2} \frac{\Gamma(-\frac{1}{2})}{\Gamma(-\frac{\nu}{2})} z \Phi(\frac{1}{2} - \frac{\nu}{2}, \frac{3}{2}; \frac{1}{2}z^2) \right].$$

For  $\nu = n = 0, 1, 2, \dots$ , we have

$$D_n(z) = 2^{-n/2} \exp(-\frac{1}{4}z^2) H_n(2^{-1/2}z), \quad \text{where} \quad H_n(z) = (-1)^n \exp(z^2) \frac{d^n}{dz^n} \exp(-z^2).$$

⊙ *References:* M. Abramowitz and I. Stegun (1964), W. Miller, Jr. (1977).

### 7.3.4-2. Elliptic coordinate system.

In the elliptic coordinates that are introduced by the relations

$$x = a \cosh u \cos v, \quad y = a \sinh u \sin v \quad (0 \leq u < \infty, 0 \leq v < 2\pi, a > 0),$$

the Helmholtz equation is expressed as

$$\frac{\partial^2 w}{\partial \xi^2} + \frac{\partial^2 w}{\partial \eta^2} + a^2 \lambda (\cosh^2 u - \cos^2 v) w = 0.$$

Setting  $w = F(u)G(v)$ , we arrive at the following linear ordinary differential equations for  $F = F(u)$  and  $G = G(v)$ :

$$F'' + \left(\frac{1}{2}a^2 \lambda \cosh 2u - k\right) F = 0, \quad G'' - \left(\frac{1}{2}a^2 \lambda \cos 2v - k\right) G = 0,$$

where  $k$  is the separation constant. The solutions of these equations periodic in  $v$  are given by

$$F(u) = \begin{cases} \text{Ce}_n(u, q), \\ \text{Se}_n(u, q), \end{cases} \quad G(v) = \begin{cases} \text{ce}_n(v, q), \\ \text{se}_n(v, q), \end{cases} \quad q = \frac{1}{4}a^2 \lambda,$$

where  $\text{Ce}_n(u, q)$  and  $\text{Se}_n(u, q)$  are the modified Mathieu functions, and  $\text{ce}_n(v, q)$  and  $\text{se}_n(v, q)$  are the Mathieu functions; to each value of  $q$  there is a corresponding  $k = k_n(q)$ .

⊙ *References:* M. Abramowitz and I. Stegun (1964), W. Miller, Jr. (1977).

### 7.3.4-3. Domain: $(x/a)^2 + (y/b)^2 \leq 1$ . First boundary value problem.

The unknown quantity is zero at the boundary of the elliptic domain:

$$w = 0 \quad \text{if} \quad (x/a)^2 + (y/b)^2 = 1 \quad (a \geq b).$$

The first three eigenvalues and eigenfunctions are given by the approximate relations

$$\begin{aligned} \lambda_1 &= \frac{\gamma_{10}^2}{2} \left( \frac{1}{a^2} + \frac{1}{b^2} \right), & w_1(\mathcal{R}) &= J_0(\gamma_{10}\mathcal{R}), \\ \lambda_2^{(c)} &= \frac{\gamma_{11}^2}{4} \left( \frac{3}{a^2} + \frac{1}{b^2} \right), & w_2^{(c)}(\mathcal{R}, \varphi) &= J_1(\gamma_{11}\mathcal{R}) \cos \varphi, \\ \lambda_2^{(s)} &= \frac{\gamma_{11}^2}{4} \left( \frac{1}{a^2} + \frac{3}{b^2} \right), & w_2^{(s)}(\mathcal{R}, \varphi) &= J_1(\gamma_{11}\mathcal{R}) \sin \varphi, \end{aligned}$$

where  $\gamma_{10} = 2.4048$  and  $\gamma_{11} = 3.8317$  are the first roots of the Bessel functions  $J_0$  and  $J_1$ , i.e.,  $J_0(\gamma_{10}) = 0$  and  $J_1(\gamma_{11}) = 0$ ;  $\mathcal{R} = \sqrt{(x/a)^2 + (y/b)^2}$ .

The above relations were obtained using the generalized (nonorthogonal) polar coordinates  $\mathcal{R}, \varphi$  defined by

$$x = a\mathcal{R} \cos \varphi, \quad y = b\mathcal{R} \sin \varphi \quad (0 \leq \mathcal{R} \leq 1, 0 \leq \varphi \leq 2\pi)$$

and the variational method.

For  $\varepsilon = \sqrt{1 - (b/a)^2} \leq 0.9$ , the above formulas provide an accuracy of 1% for  $\lambda_1$  and 2% for  $\lambda_2^{(c)}$  and  $\lambda_2^{(s)}$ . For  $\varepsilon \leq 0.5$ , the errors in calculating  $\lambda_1$  and  $\lambda_2^{(c)}$  do not exceed 0.01%, and the maximum error in determining  $\lambda_2^{(s)}$  is 0.12%. In the limit case  $\varepsilon = 0$  that corresponds to a circular domain, the above formulas are exact.

⊙ *Reference:* L. D. Akulenko and S. V. Nesterov (2000).

TABLE 24

Transformations reducing equation 7.4.1.3 to the Helmholtz equation  $\frac{\partial^2 w}{\partial \xi^2} + \frac{\partial^2 w}{\partial \eta^2} = bw$

| No | Exponent $k$                  | Transformation                                                                                                                                  | Factor $b$ |
|----|-------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------|------------|
| 1  | $k = 1$                       | $\xi = \frac{1}{2}(x^2 - y^2), \eta = xy$                                                                                                       | $b = a$    |
| 2  | $k = 2$                       | $\xi = \frac{1}{3}x^3 - xy^2, \eta = x^2y - \frac{1}{3}y^3$                                                                                     | $b = a$    |
| 3  | $k = -1$                      | $\xi = \frac{1}{2} \ln(x^2 + y^2), \eta = \arctan \frac{y}{x}$                                                                                  | $b = a$    |
| 4  | $k = -2$                      | $\xi = -\frac{x}{x^2 + y^2}, \eta = \frac{y}{x^2 + y^2}$                                                                                        | $b = a$    |
| 5  | $k = -\frac{1}{2}$            | $x = \frac{1}{2}(\xi^2 - \eta^2), y = \xi\eta$                                                                                                  | $b = 2a$   |
| 6  | $k = \pm 3, \pm 4, \dots$     | $\xi = \frac{(x+iy)^{k+1} + (x-iy)^{k+1}}{2(k+1)}, \eta = \frac{(x+iy)^{k+1} - (x-iy)^{k+1}}{2(k+1)i}$                                          | $b = a$    |
| 7  | $k$ is any<br>( $k \neq -1$ ) | $\xi = \frac{\rho^{k+1} \cos[(k+1)\varphi]}{x^{k+1} \rho \cos \varphi}, \eta = \frac{\rho^{k+1} \sin[(k+1)\varphi]}{y^{k+1} \rho \sin \varphi}$ | $b = a$    |

## 7.4. Other Equations

### 7.4.1. Stationary Schrödinger Equation $\Delta_2 w = f(x, y)w$

1.  $\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = a(x^2 + y^2)w.$

The transformation

$$z = \frac{1}{2}(x^2 - y^2), \quad \zeta = xy$$

leads to the Helmholtz equation

$$\frac{\partial^2 w}{\partial z^2} + \frac{\partial^2 w}{\partial \zeta^2} - aw = 0,$$

which is discussed in Subsection 7.3.2.

2.  $\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = a(x^2 + y^2)^2 w.$

The transformation

$$z = \frac{1}{3}x^3 - xy^2, \quad \zeta = x^2y - \frac{1}{3}y^3$$

leads to the Helmholtz equation

$$\frac{\partial^2 w}{\partial z^2} + \frac{\partial^2 w}{\partial \zeta^2} - aw = 0,$$

which is discussed in Subsection 7.3.2.

3.  $\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = a(x^2 + y^2)^k w.$

This is a special case of equation 7.4.1.7 for  $f(u) = au^k$ . Table 24 presents transformations that reduce this equation to the Helmholtz equation that is discussed in Subsection 7.3.2; the sixth row involves the imaginary unit,  $i^2 = -1$ .

$$4. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = a e^{\beta x} w.$$

The transformation

$$u(x, y) = \exp\left(\frac{1}{2}\beta x\right) \cos\left(\frac{1}{2}\beta y\right), \quad v(x, y) = \exp\left(\frac{1}{2}\beta x\right) \sin\left(\frac{1}{2}\beta y\right)$$

leads to the Helmholtz equation

$$\frac{\partial^2 w}{\partial u^2} + \frac{\partial^2 w}{\partial v^2} = 4a\beta^{-2}w,$$

which is discussed in Subsection 7.3.2.

$$5. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = k e^{ax+by} w.$$

The transformation

$$\xi = ax + by, \quad \eta = bx - ay$$

leads to an equation of the form 7.4.1.4:

$$\frac{\partial^2 w}{\partial \xi^2} + \frac{\partial^2 w}{\partial \eta^2} = \frac{k}{a^2 + b^2} e^{\xi} w.$$

$$6. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = f(ax + by)w.$$

This is a special case of equation 7.4.1.9 for  $g(u) = 0$ . Particular solutions:

$$w(x, y) = \{C_1 \cos[k(bx - ay)] + C_2 \sin[k(bx - ay)]\} \varphi(ax + by),$$

where  $C_1$ ,  $C_2$ , and  $k$  are arbitrary constants, and the function  $\varphi = \varphi(\xi)$  is determined by the ordinary differential equation

$$\varphi''_{\xi\xi} - \left[ \frac{1}{a^2 + b^2} f(\xi) + k^2 \right] \varphi = 0.$$

$$7. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = f(x^2 + y^2)w.$$

1°. This equation admits separation of variables in the polar coordinates  $\rho$ ,  $\varphi$  ( $x = \rho \cos \varphi$ ,  $y = \rho \sin \varphi$ ). Particular solution:

$$w(x, y) = [C_1 \cos(k\varphi) + C_2 \sin(k\varphi)] U(\rho),$$

where  $C_1$ ,  $C_2$ , and  $k$  are arbitrary constants, and the function  $U = U(\rho)$  is determined by the ordinary differential equation

$$\rho(\rho U'_\rho)' - [k^2 + \rho^2 f(\rho^2)] U = 0.$$

2°. The transformation

$$z = \frac{1}{2}(x^2 - y^2), \quad \zeta = xy$$

leads to a similar equation

$$\frac{\partial^2 w}{\partial z^2} + \frac{\partial^2 w}{\partial \zeta^2} = F(z^2 + \zeta^2)w, \quad F(u) = \frac{f(2\sqrt{u})}{2\sqrt{u}}.$$

In the special case  $f(u) = 2a$ , we have  $F(u) = a/\sqrt{u}$ . For  $f(u) = bu^3$ , we obtain an equation of the form 7.4.1.1 with  $F(u) = 4bu$ .

$$8. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = [f(x) + g(y)]w.$$

A particular separable solution:

$$w(x, y) = \varphi(x)\psi(y),$$

where the functions  $\varphi(x)$  and  $\psi(y)$  are determined by the second-order ordinary differential equations

$$\varphi''_{xx} - [f(x) - C]\varphi = 0, \quad \psi''_{yy} - [g(y) + C]\psi = 0,$$

where  $C$  is an arbitrary constant.

$$9. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = [f(ax + by) + g(bx - ay)]w.$$

The transformation

$$\xi = ax + by, \quad \eta = bx - ay$$

leads to an equation of the form 7.4.1.8:

$$\frac{\partial^2 w}{\partial \xi^2} + \frac{\partial^2 w}{\partial \eta^2} = \left[ \frac{f(\xi)}{a^2 + b^2} + \frac{g(\eta)}{a^2 + b^2} \right] w.$$

$$10. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = (x^2 + y^2)[f(x^2 - y^2) + g(xy)]w.$$

The transformation

$$z = \frac{1}{2}(x^2 - y^2), \quad \zeta = xy$$

leads to an equation of the form 7.4.1.8:

$$\frac{\partial^2 w}{\partial z^2} + \frac{\partial^2 w}{\partial \zeta^2} = [f(2z) + g(\zeta)]w.$$

## 7.4.2. Convective Heat and Mass Transfer Equations

$$1. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = \alpha \frac{\partial w}{\partial x}.$$

This is a convective heat and mass transfer equation. It describes a stationary temperature (concentration) field in a continuous medium moving with a constant velocity along the  $x$ -axis. In particular, it models convective-molecular heat transfer from a heated flat plate in a flow of a thermal-transfer ideal fluid moving along the plate. This occurs, for example, if a liquid-metal coolant flows past a flat plate or if a plate is in a seepage flow through a granular medium.

In the sequel, it is assumed that the equation is written in dimensionless variables  $x, y$  related to the characteristic length (for a flat plate of length  $2h$ , the characteristic length is taken to be  $h$ ).

1°. The substitution  $w(x, y) = \exp(\frac{1}{2}\alpha x)U(x, y)$  brings the original equation to the Helmholtz equation

$$\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} = \frac{1}{4}\alpha^2 U.$$

Particular solutions of this equation in Cartesian and polar coordinates can be found in Subsections 7.3.2 and 7.3.3.



2°. In the elliptic coordinates

$$x = \cosh \zeta \cos \eta, \quad y = \sinh \zeta \sin \eta$$

a wide class of particular solutions (vanishing as  $\zeta \rightarrow \infty$ ) can be indicated; this class of solutions of the original equation is represented in series form as

$$w = \exp\left(\frac{1}{2}\alpha x\right) \sum_{m=0}^{\infty} A_m \operatorname{ce}_m(\eta, -q) \operatorname{Fek}_m(\zeta, -q), \quad q = -\frac{1}{16}\alpha^2,$$

where the  $A_m$  are arbitrary constants, the  $\operatorname{ce}_m(\eta, -q)$  are the Mathieu functions, and the  $\operatorname{Fek}_m(\zeta, -q)$  are the modified Mathieu functions [e.g., see McLachlan (1947) and Bateman and Erdélyi (1955)].

3°. Consider the first boundary value problem in the upper half-plane ( $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ). We assume that the surface of a plate of finite length is maintained at a constant temperature  $w_0$  and the medium has a temperature  $w_\infty = \text{const}$  far away from the plate:

$$\begin{aligned} w &= w_0 & \text{for } y = 0, \quad |x| < 1, \\ \partial_y w &= 0 & \text{for } y = 0, \quad |x| > 1, \\ w &\rightarrow w_\infty & \text{for } x^2 + y^2 \rightarrow \infty. \end{aligned}$$

The solution of this problem in the elliptic coordinates  $\zeta, \eta$  (see Item 2°) has the form

$$w(\eta, \zeta) = w_\infty + (w_0 - w_\infty) \exp\left(\frac{1}{2}\alpha \cos \eta \cosh \zeta\right) \sum_{m=0}^{\infty} D_m \operatorname{ce}_m(\eta, -q) \frac{\operatorname{Fek}_m(\zeta, -q)}{\operatorname{Fer}_m(0, -q)},$$

where

$$D_{2n} = 2 \frac{\operatorname{ce}_{2n}(0, -q)}{\operatorname{ce}_{2n}(0, q)} A_0^{(2n)}, \quad D_{2n+1} = -\frac{1}{2} \frac{\operatorname{ce}_{2n+1}(0, -q)}{\operatorname{ce}_{2n+1}(0, q)} \alpha B_1^{(2n+1)}, \quad q = -\frac{1}{16}\alpha^2.$$

Here, the  $A_0^{(2n)}$  and  $B_1^{(2n+1)}$  are the coefficients in the series expansions of the Mathieu functions; these can be found in McLachlan (1947).

4°. Consider the second boundary value problem in the upper half-plane ( $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ). We assume that a thermal flux is prescribed on the surface of a plate of finite length and the medium has a constant temperature far away from the plate:

$$\begin{aligned} \partial_y w &= f(x) & \text{for } y = 0, \quad |x| < 1, \\ \partial_y w &= 0 & \text{for } y = 0, \quad |x| > 1, \\ w &\rightarrow w_\infty & \text{as } x^2 + y^2 \rightarrow \infty. \end{aligned}$$

The solution of this problem in the Cartesian coordinates has the form

$$w(x, y) = w_\infty - \frac{1}{\pi} \int_{-1}^1 f(\xi) \exp\left[\frac{1}{2}\alpha(x - \xi)\right] K_0\left(\frac{1}{2}\alpha\sqrt{(x - \xi)^2 + y^2}\right) d\xi,$$

where  $K_0(z)$  is the modified Bessel function of the second kind.

⊙ *References:* P. V. Cherpakov (1975), A. A. Borzykh and G. P. Cherepanov (1978).

$$2. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = \alpha \frac{\partial w}{\partial x} + \beta \frac{\partial w}{\partial y} + \gamma w.$$

This equation describes a stationary temperature field in a medium moving with a constant velocity, provided there is volume release heat (or absorption) proportional to temperature.

The substitution

$$w(x, y) = \exp\left[\frac{1}{2}(\alpha x + \beta y)\right] U(x, y)$$

brings the original equation to the Helmholtz equation

$$\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} = \left(\gamma + \frac{1}{4}\alpha^2 + \frac{1}{4}\beta^2\right) U,$$

which is discussed in Subsections 7.3.1 through 7.3.3.

$$3. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = \text{Pe} (1 - y^2) \frac{\partial w}{\partial x}.$$

*The Graetz–Nusselt equation.* It governs steady-state heat exchange in a laminar fluid flow with a parabolic velocity profile in a plane channel. The equation is written in terms of the dimensionless Cartesian coordinates  $x, y$  related to the channel half-width  $h$ ;  $\text{Pe} = Uh/a$  is the Peclet number and  $U$  is the fluid velocity at the channel axis ( $y = 0$ ). The walls of the channel correspond to  $y = \pm 1$ .

1°. Particular solutions:

$$\begin{aligned} w(y) &= A + By, \\ w(x, y) &= 12Ax + A \text{Pe} (6y^2 - y^4) + B, \\ w(x, y) &= \sum_{n=1}^m A_n \exp\left(-\frac{\lambda_n^2}{\text{Pe}} x\right) f_n(y). \end{aligned}$$

Here,  $A, B, A_n$ , and  $\lambda_n$  are arbitrary constants, and the functions  $f_n$  are defined by

$$f_n(y) = \exp\left(-\frac{1}{2}\lambda_n y^2\right) \Phi\left(\alpha_n, \frac{1}{2}; \lambda_n y^2\right), \quad \alpha_n = \frac{1}{4} - \frac{1}{4}\lambda_n - \frac{1}{4}\lambda_n^3 \text{Pe}^{-2}, \quad (1)$$

where  $\Phi(\alpha, \beta; \xi) = 1 + \sum_{k=1}^{\infty} \frac{\alpha(\alpha+1)\dots(\alpha+k-1)}{\beta(\beta+1)\dots(\beta+k-1)} \frac{\xi^k}{k!}$  is the degenerate hypergeometric function.

2°. Let the walls of the channel be maintained at a constant temperature,  $w = 0$  for  $x < 0$  and  $w = w_0$  for  $x > 0$ . Due to the symmetry of the problem about the  $x$ -axis, it suffices to consider only half of the domain,  $0 \leq y \leq 1$ . The boundary conditions are written as

$$\begin{aligned} y = 0, \quad \frac{\partial w}{\partial y} &= 0; \quad y = 1, \quad w = \begin{cases} 0 & \text{for } x < 0, \\ w_0 & \text{for } x > 0; \end{cases} \\ x \rightarrow -\infty, \quad w &\rightarrow 0; \quad x \rightarrow \infty, \quad w \rightarrow w_0. \end{aligned}$$

The solution of the original equation under these boundary conditions is sought in the form

$$\begin{aligned} w(x, y) &= w_0 \sum_{n=1}^{\infty} B_n \exp\left(\frac{\mu_n^2}{\text{Pe}} x\right) g_n(y) \quad \text{for } x < 0, \\ w(x, y) &= w_0 \left[ 1 - \sum_{n=1}^{\infty} A_n \exp\left(-\frac{\lambda_n^2}{\text{Pe}} x\right) f_n(y) \right] \quad \text{for } x > 0. \end{aligned}$$

The series coefficients must satisfy the matching conditions at the boundary:

$$\begin{aligned} w(x, y)|_{x \rightarrow 0, x < 0} - w(x, y)|_{x \rightarrow 0, x > 0} &= 0, \\ \partial_x w(x, y)|_{x \rightarrow 0, x < 0} - \partial_x w(x, y)|_{x \rightarrow 0, x > 0} &= 0. \end{aligned}$$

For  $x > 0$ , the function  $f_n(y)$  is defined by relation (1), where the eigenvalues  $\lambda_n$  are roots of the transcendental equation

$$\Phi\left(\alpha_n, \frac{1}{2}; \lambda_n\right) = 0, \quad \text{where } \alpha_n = \frac{1}{4} - \frac{1}{4}\lambda_n - \frac{1}{4}\lambda_n^3 \text{Pe}^{-2}.$$

For  $\text{Pe} \rightarrow \infty$ , it is convenient to use the following approximate relation to identify the  $\lambda_n$ :

$$\lambda_n = 4(n-1) + 1.68 \quad (n = 1, 2, 3, \dots). \quad (2)$$

The error of this formula does not exceed 0.2%. The corresponding numerical values of the coefficients  $A_n$  are rather well approximated by the relations

$$A_1 = 1.2, \quad A_n = 2.27 (-1)^{n-1} \lambda_n^{-7/6} \quad \text{for } n = 2, 3, 4, \dots,$$

whose maximum error is less than 0.1%, provided that the  $\lambda_n$  are calculated by (2).

For  $\text{Pe} \rightarrow 0$ , the following asymptotic relations hold:

$$\lambda_n = \sqrt{\pi\left(n - \frac{1}{2}\right) \text{Pe}}, \quad A_n = \frac{4(-1)^{n-1}}{\pi^2(2n-1)^2}, \quad f_n(y) = \cos\left[\pi\left(n - \frac{1}{2}\right)y\right] \quad (n = 1, 2, 3, \dots).$$

No results for  $x < 0$  are given here, because they are of secondary importance in applications.

3°. Let a constant thermal flux be prescribed at the walls for  $x > 0$  and let, for  $x < 0$ , the walls be insulated from heat and the temperature vanishes as  $x \rightarrow -\infty$ . Then the boundary conditions have the form

$$y = 0, \quad \frac{\partial w}{\partial y} = 0; \quad y = 1, \quad \frac{\partial w}{\partial y} = \begin{cases} 0 & \text{for } x < 0, \\ q & \text{for } x > 0; \end{cases} \quad x \rightarrow -\infty, \quad w \rightarrow 0.$$

In the domain of thermal stabilization, the asymptotic behavior of the solution (as  $x \rightarrow \infty$ ) is as follows:

$$w(x, y) = q \left( \frac{3}{2} \frac{x}{\text{Pe}} + \frac{3}{4} y^2 - \frac{1}{8} y^4 + \frac{9}{4 \text{Pe}^2} - \frac{39}{280} \right).$$

⊙ *References:* L. Graetz (1883), W. Nusselt (1910), C. A. Deavours (1974), A. D. Polyanin, A. M. Kutepov, A. V. Vyazmin, and D. A. Kazenin (2001).

$$4. \quad \frac{\partial^2 w}{\partial r^2} + \frac{1}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} = \text{Pe} (1 - r^2) \frac{\partial w}{\partial z}.$$

This equation governs steady-state heat exchange in a laminar fluid flow with parabolic (Poiseuille's) velocity profile in a circular tube. The equation is written in terms of the dimensionless cylindrical coordinates  $x, y$  related to the tube radius  $R$ ;  $\text{Pe} = UR/a$  is the Peclet number and  $U$  is the fluid velocity at the tube axis (at  $r = 0$ ). The walls of the tube correspond to  $r = 1$ .

1°. Particular solutions:

$$\begin{aligned} w(r) &= A + B \ln r, \\ w(r, z) &= 16Az + A \text{Pe} (4r^2 - r^4) + B, \\ w(r, z) &= \sum_{n=1}^m A_n \exp\left(-\frac{\lambda_n^2}{\text{Pe}} z\right) f_n(r). \end{aligned}$$

Here,  $A, B, A_n$ , and  $\lambda_n$  are arbitrary constants, and the functions  $f_n$  are defined by

$$f_n(r) = \exp\left(-\frac{1}{2} \lambda_n r^2\right) \Phi(\alpha_n, 1; \lambda_n r^2), \quad \alpha_n = \frac{1}{2} - \frac{1}{4} \lambda_n - \frac{1}{4} \lambda_n^3 \text{Pe}^{-2}, \quad (1)$$

where  $\Phi(\alpha, \beta; \xi)$  is the degenerate hypergeometric function (see equation 7.4.2.3, Item 1°).

2°. Let the tube wall be maintained at a constant temperature such that  $w = 0$  for  $z < 0$  and  $w = w_0$  for  $z > 0$ . The boundary conditions are written as

$$\begin{aligned} r = 0, \quad \frac{\partial w}{\partial r} &= 0; \quad r = 1, \quad w = \begin{cases} 0 & \text{for } z < 0, \\ w_0 & \text{for } z > 0; \end{cases} \\ z \rightarrow -\infty, \quad w &\rightarrow 0; \quad z \rightarrow \infty, \quad w \rightarrow w_0. \end{aligned}$$

The solution of the original equation under these boundary conditions is sought in the form

$$\begin{aligned} w(r, z) &= w_0 \sum_{n=1}^{\infty} B_n \exp\left(\frac{\mu_n^2}{\text{Pe}} z\right) g_n(r) \quad \text{for } z < 0, \\ w(r, z) &= w_0 \left[ 1 - \sum_{n=1}^{\infty} A_n \exp\left(-\frac{\lambda_n^2}{\text{Pe}} z\right) f_n(r) \right] \quad \text{for } z > 0. \end{aligned}$$

The series coefficients must satisfy the matching conditions at the boundary,

$$\begin{aligned} w(r, z) \Big|_{z \rightarrow 0, z < 0} - w(r, z) \Big|_{z \rightarrow 0, z > 0} &= 0, \\ \partial_z w(r, z) \Big|_{z \rightarrow 0, z < 0} - \partial_z w(r, z) \Big|_{z \rightarrow 0, z > 0} &= 0. \end{aligned}$$

For  $z > 0$ , the functions  $f_n(r)$  are defined by relations (1), where the eigenvalues  $\lambda_n$  are roots of the transcendental equation

$$\Phi(\alpha_n, 1; \lambda_n) = 0, \quad \text{where} \quad \alpha_n = \frac{1}{2} - \frac{1}{4} \lambda_n - \frac{1}{4} \lambda_n^3 \text{Pe}^{-2}.$$

For  $Pe \rightarrow \infty$ , it is convenient to use the following approximate relation to identify the  $\lambda_n$ :

$$\lambda_n = 4(n-1) + 2.7 \quad (n = 1, 2, 3, \dots). \quad (2)$$

The error of this formula does not exceed 0.3%. The corresponding numerical values of the coefficients  $A_n$  are rather well approximated by the relations

$$A_n = 2.85 (-1)^{n-1} \lambda_n^{-2/3} \quad \text{for } n = 1, 2, 3, \dots,$$

whose maximum error is 0.5%,

No results for  $z < 0$  are given here, since they are of secondary importance in applications.

3°. Let a constant thermal flux be prescribed at the wall for  $z > 0$  and let, for  $z < 0$ , the tube surface be insulated from heat and the temperature vanishes as  $z \rightarrow -\infty$ . Then the boundary conditions have the form

$$r = 0, \quad \frac{\partial w}{\partial r} = 0; \quad r = 1, \quad \frac{\partial w}{\partial r} = \begin{cases} 0 & \text{for } z < 0, \\ q & \text{for } z > 0; \end{cases} \quad z \rightarrow -\infty, \quad w \rightarrow 0.$$

In the domain of thermal stabilization, the asymptotic behavior of the solution (as  $z \rightarrow \infty$ ) is as follows:

$$w(r, z) = q \left( 4 \frac{z}{Pe} + r^2 - \frac{1}{4} r^4 + \frac{8}{Pe^2} - \frac{7}{24} \right).$$

⊙ *References:* C. A. Deavours (1974), A. D. Polyaniin, A. M. Kutepov, A. V. Vyazmin, and D. A. Kazenin (2001).

$$5. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = f(y) \frac{\partial w}{\partial x}.$$

This equation describes steady-state heat exchange in a laminar fluid flow with an arbitrary velocity profile  $f = f(y)$  in a plane channel.

1°. Particular solutions:

$$w(x, y) = Ax + A \int_{y_0}^y (y - \xi) f(\xi) d\xi + By + C, \quad (1)$$

$$w(x, y) = B + \sum_{n=1}^m A_n \exp(-\beta_n x) u_n(y). \quad (2)$$

Here,  $A, B, C, y_0, A_n$ , and  $\beta_n$  are arbitrary constants, and the functions  $u_n = u_n(y)$  are determined by the second-order linear ordinary differential equation

$$\frac{d^2 u_n}{dy^2} + [\beta_n f(y) + \beta_n^2] u_n = 0.$$

2°. Solution (1) describes the temperature distribution far away from the inlet section of the tube, in the domain of thermal stabilization, provided that a constant thermal flux is prescribed at the channel walls.

$$6. \quad a \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) = v_1(x, y) \frac{\partial w}{\partial x} + v_2(x, y) \frac{\partial w}{\partial y}.$$

This is an equation of steady-state convective heat and mass transfer in the Cartesian coordinate system. Here,  $v_1 = v_1(x, y)$  and  $v_2 = v_2(x, y)$  are the components of the fluid velocity that are assumed to be known from the solution of the hydrodynamic problem.

1°. In plane problems of convective heat exchange in liquid metals modeled by an ideal fluid, as well as in describing seepage (filtration) streams employing the model of potential flows, the fluid velocity components  $v_1(x, y)$  and  $v_2(x, y)$  can be expressed in terms of the potential  $\varphi = \varphi(x, y)$  and stream function  $\psi = \psi(x, y)$  as follows:

$$v_1 = \frac{\partial \varphi}{\partial x} = -\frac{\partial \psi}{\partial y}, \quad v_2 = \frac{\partial \varphi}{\partial y} = \frac{\partial \psi}{\partial x}. \quad (1)$$

The function  $\varphi$  is determined by solving the Laplace equation  $\Delta \varphi = 0$ . In specific problems, the potential  $\varphi$  and stream function  $\psi$  may be identified by invoking the complex variable theory [e.g., see Lavrent'ev and Shabat (1973) and Sedov (1980)].

By passing in the convective heat exchange equation from  $x, y$  to the new variables  $\varphi, \psi$  (Boussinesq transformation) and taking into account (1), we arrive a simpler equation with constant coefficients of the form 7.4.2.1:

$$\frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial \psi^2} = \frac{1}{a} \frac{\partial w}{\partial \varphi}. \quad (2)$$

The Boussinesq transformation brings any plane contour in a potential flow to a cut in the  $\varphi$ -axis, simultaneously with the reduction of the original equation to the form (2). Consequently, the heat transfer problem of a potential flow about this contour is reduced to the heat exchange problem of a longitudinal flow of an ideal fluid past a flat plate (see equation 7.4.2.1, Items 3° and 4°).

2°. Asymptotic analyses of plane problems on heat/mass exchange of bodies of various shape with laminar translational and shear flows of a viscous (and ideal) incompressible fluid for large and small Peclet numbers were carried out in the references cited below. In the thermal boundary layer approximation, the solution of the heat exchange problem for a flat plate in a longitudinal translational flow of a viscous incompressible fluid at large Reynolds numbers is presented in 1.9.1.4, Item 3°.

⊙ *References:* V. G. Levich (1962), P. V. Cherpakov (1975), A. A. Borzykh and G. P. Cherepanov (1978), Yu. P. Gupalo, A. D. Polyaniin, and Yu. S. Ryazantsev (1985), A. D. Polyaniin, A. M. Kutepov, A. V. Vyazmin, and D. A. Kazenin (2001).

$$7. \quad \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) = \cos \theta \frac{\partial w}{\partial r} - \frac{\sin \theta}{r} \frac{\partial w}{\partial \theta}.$$

This is a special case of equation 7.4.1.8 with  $a = 1$ ,  $v_r = \cos \theta$ , and  $v_\theta = -\sin \theta$ . This equation is obtained from the equation  $\partial_{xx} w + \partial_{yy} w + \partial_{zz} w = \partial_x w$  by the passage to the spherical coordinate system in the axisymmetric case.

The general solution satisfying the decay condition ( $w \rightarrow 0$  as  $r \rightarrow \infty$ ) is expressed as

$$w(r, \theta) = \left( \frac{\pi}{r} \right)^{1/2} \exp \left( \frac{r \cos \theta}{2} \right) \sum_{n=0}^{\infty} A_n K_{n+\frac{1}{2}} \left( \frac{r}{2} \right) P_n(\cos \theta),$$

where the  $A_n$  are arbitrary constants. The Legendre polynomials  $P_n(\xi)$  and the modified Bessel functions  $K_{n+\frac{1}{2}}(z)$  are given by

$$P_n(\xi) = \frac{1}{n! 2^n} \frac{d^n}{d\xi^n} (\xi^2 - 1)^n, \quad K_{n+\frac{1}{2}} \left( \frac{r}{2} \right) = \left( \frac{\pi}{r} \right)^{1/2} \exp \left( -\frac{r}{2} \right) \sum_{m=0}^n \frac{(n+m)!}{(n-m)! m! r^m}.$$

⊙ *Reference:* P. L. Rimmer (1968).

$$8. \quad a \left[ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) \right] = v_r \frac{\partial w}{\partial r} + \frac{v_\theta}{r} \frac{\partial w}{\partial \theta}.$$

This equation is often encountered in axisymmetric problems of convective heat and mass exchange of solid particles, drops, and bubbles with a flow of a viscous incompressible fluid. The fluid velocity components  $v_r = v_r(r, \theta)$  and  $v_\theta = v_\theta(r, \theta)$  can be expressed in terms of the stream function  $\psi = \psi(r, \theta)$  as

$$v_r = \frac{1}{r^2 \sin \theta} \frac{\partial \psi}{\partial \theta}, \quad v_\theta = -\frac{1}{r \sin \theta} \frac{\partial \psi}{\partial r}. \quad (1)$$

Asymptotic analyses for a wide class of axisymmetric problems on heat/mass exchange of solid particles, drops, and bubbles of various shape with a laminar translational or straining flow of a viscous incompressible fluid at large and small Peclet numbers  $Pe = UR/a$  are performed in the books cited below. The Peclet number is written in terms of the characteristic velocity  $U$  (e.g., the unperturbed fluid velocity far away from the particle in the case of translation flow), the characteristic size of the particle  $R$  (e.g., the radius for a spherical particle), and the thermal conductivity or diffusion coefficient  $a$ .

The following boundary conditions are usually specified:

$$w = w_0 \quad \text{at} \quad r = R, \quad w \rightarrow w_\infty \quad \text{as} \quad r \rightarrow \infty, \quad (2)$$

where  $R$  is the particle radius,  $w_0$  the temperature at the particle surface, and  $w_\infty$  the temperature far away from the particle ( $w_0$  and  $w_\infty$  are constant).

Convective mass transfer problems are characterized by large Peclet numbers. To solve such problems, the diffusion boundary layer approximation is often used; in this case, the left-hand side of the equation takes into account only the diffusion mass transfer in the normal direction to the particle surface (the tangential mass transfer is neglected). The convective terms on the right-hand side are partially preserved—the fluid velocity components are approximated by their leading terms of the asymptotic expansion near the phase surface. Presented below are some important results obtained by solving the original equation under the boundary conditions (2) in the diffusion boundary layer approximation.

**Example 1.** For the translational Stokes flow of a viscous incompressible fluid about a spherical bubble, the stream function is expressed as

$$\psi(r, \theta) = \frac{1}{2} U r (r - R) \sin^2 \theta.$$

Here,  $U$  is the unperturbed fluid velocity in the incident flow,  $R$  the bubble radius (the value  $\theta = \pi$  corresponds to the front critical point at the bubble surface).

In this case, the solution of the convective heat/mass transfer equation with the boundary conditions (2) for  $Pe = UR/a \gg 1$  in the diffusion boundary layer approximation is given by

$$w(r, \theta) = w_0 + (w_\infty - w_0) \operatorname{erf} \xi, \quad \xi = \sqrt{\frac{3}{8}} Pe \left( \frac{r}{R} - 1 \right) \frac{1 - \cos \theta}{\sqrt{2 - \cos \theta}},$$

where  $\operatorname{erf} \xi$  is the error function.

**Example 2.** For the translational Stokes flow of a viscous incompressible fluid about a solid spherical particle, the stream function is expressed as

$$\psi(r, \theta) = \frac{1}{4} U (r - R)^2 \left( 2 + \frac{R}{r} \right) \sin^2 \theta.$$

Here, the notation is the same as in the case of a bubble above.

For a solid particle, the solution of the convective heat/mass transfer equation with the boundary conditions (2) for  $Pe = UR/a \gg 1$  in the diffusion boundary layer approximation is given by

$$w(r, \theta) = w_0 + (w_\infty - w_0) \left[ \Gamma\left(\frac{1}{3}\right) \right]^{-1} \gamma\left(\frac{1}{3}, \xi\right), \quad \xi = \frac{Pe (r - R)^3 \sin^3 \theta}{3R^3 \left( \pi - \theta + \frac{1}{2} \sin 2\theta \right)},$$

where  $\Gamma(\beta)$  is the gamma function and  $\gamma(\beta, \xi) = \int_0^\xi e^{-z} z^{\beta-1} dz$  is the incomplete gamma function.

⊙ *References:* V. G. Levich (1962), Yu. P. Gupalo, A. D. Polyaniin, and Yu. S. Ryazantsev (1985), A. D. Polyaniin, A. M. Kutepov, A. V. Vyazmin, and D. A. Kazenin (2001).

### 7.4.3. Equations of Heat and Mass Transfer in Anisotropic Media

$$1. \quad \frac{\partial}{\partial x} \left( a x^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( b y^m \frac{\partial w}{\partial y} \right) = 0.$$

This is a two-dimensional equation of the heat and mass transfer theory in a inhomogeneous anisotropic medium. Here,  $a_1(x) = a x^n$  and  $a_2(y) = b y^m$  are the principal thermal diffusivities.

1°. Particular solutions ( $A, B, C$  are arbitrary constants):

$$\begin{aligned}w(x, y) &= Ax^{1-n} + By^{1-m} + C, \\w(x, y) &= A \left[ \frac{x^{2-n}}{a(2-n)} - \frac{y^{2-m}}{b(2-m)} \right] + B, \\w(x, y) &= Ax^{1-n}y^{1-m} + B.\end{aligned}$$

2°. For  $n \neq 2$  and  $m \neq 2$ , there are particular solutions of the form

$$w = w(\xi), \quad \xi = [b(2-m)^2x^{2-n} + a(2-n)^2y^{2-m}]^{1/2}.$$

The function  $w = w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{A}{\xi}w'_{\xi} = 0, \quad A = \frac{4-nm}{(2-n)(2-m)}. \quad (1)$$

The general solution of equation (1) is given by

$$w(\xi) = \begin{cases} C_1\xi^{1-A} + C_2 & \text{for } A \neq 1, \\ C_1 \ln \xi + C_2 & \text{for } A = 1, \end{cases}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

3°. There are multiplicatively separable particular solutions in the form

$$w(x, y) = \varphi(x)\psi(y), \quad (2)$$

where  $\varphi(x)$  and  $\psi(y)$  are determined by the following second-order linear ordinary differential equations ( $A_1$  is an arbitrary constant):

$$(ax^n\varphi'_x)'_x = -A_1\varphi, \quad (3)$$

$$(by^m\psi'_y)'_y = A_1\psi. \quad (4)$$

The solution of equation (3) is given by

$$\begin{aligned}\varphi(x) &= \begin{cases} x^{\frac{1-n}{2}} \left[ C_1 J_{\nu} \left( \beta x^{\frac{2-n}{2}} \right) + C_2 Y_{\nu} \left( \beta x^{\frac{2-n}{2}} \right) \right] & \text{for } A_1 > 0, \\ x^{\frac{1-n}{2}} \left[ C_1 I_{\nu} \left( \beta x^{\frac{2-n}{2}} \right) + C_2 K_{\nu} \left( \beta x^{\frac{2-n}{2}} \right) \right] & \text{for } A_1 < 0, \end{cases} \\ \nu &= \frac{|1-n|}{2-n}, \quad \beta = \frac{2}{2-n} \sqrt{\frac{|A_1|}{a}},\end{aligned}$$

where  $C_1$  and  $C_2$  are arbitrary constants,  $J_{\nu}(z)$  and  $Y_{\nu}(z)$  are the Bessel functions, and  $I_{\nu}(z)$  and  $K_{\nu}(z)$  are the modified Bessel functions.

The solution of equation (4) is expressed as

$$\begin{aligned}\psi(y) &= \begin{cases} y^{\frac{1-m}{2}} \left[ C_1 J_{\sigma} \left( \mu y^{\frac{2-m}{2}} \right) + C_2 Y_{\sigma} \left( \mu y^{\frac{2-m}{2}} \right) \right] & \text{for } A_1 < 0, \\ y^{\frac{1-m}{2}} \left[ C_1 I_{\sigma} \left( \mu y^{\frac{2-m}{2}} \right) + C_2 K_{\sigma} \left( \mu y^{\frac{2-m}{2}} \right) \right] & \text{for } A_1 > 0, \end{cases} \\ \sigma &= \frac{|1-m|}{2-m}, \quad \mu = \frac{2}{2-m} \sqrt{\frac{|A_1|}{b}},\end{aligned}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

The sum of solutions of the form (2) corresponding to different values of the parameter  $A_1$  is also a solution of the original equation; the solutions of some boundary value problems may be obtained by separation of variables.

4°. See equation 7.4.3.3, Item 4°, for  $c = 0$ .

$$2. \quad \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( by^m \frac{\partial w}{\partial y} \right) = c.$$

This is a two-dimensional equation of the heat and mass transfer theory with constant volume release of heat in an inhomogeneous anisotropic medium. Here,  $a_1(x) = ax^n$  and  $a_2(y) = by^m$  are the principal thermal diffusivities.

1°. For  $n \neq 2$  and  $m \neq 2$ , there are particular solutions of the form

$$w = w(\xi), \quad \xi = [b(2-m)^2 x^{2-n} + a(2-n)^2 y^{2-m}]^{1/2}. \quad (1)$$

The function  $w = w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{A}{\xi} w'_{\xi} = B, \quad (2)$$

where

$$A = \frac{4-nm}{(2-n)(2-m)}, \quad B = \frac{4c}{ab(2-n)^2(2-m)^2}. \quad (3)$$

The general solution of equation (2) is given by

$$w(\xi) = \begin{cases} C_1 \xi^{1-A} + C_2 + \frac{B}{2(A+1)} \xi^2 & \text{for } A \neq \pm 1, \\ C_1 \ln \xi + C_2 + \frac{1}{4} B \xi^2 & \text{for } A = 1, \\ C_1 \xi^2 + C_2 + \frac{1}{2} B \xi^2 \ln \xi & \text{for } A = -1, \end{cases}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

2°. The substitution

$$w(x, y) = U(x, y) + \frac{c}{a(2-n)} x^{2-n}$$

leads to a homogeneous equation of the form 7.4.3.1:

$$\frac{\partial}{\partial x} \left( ax^n \frac{\partial U}{\partial x} \right) + \frac{\partial}{\partial y} \left( by^m \frac{\partial U}{\partial y} \right) = 0.$$

$$3. \quad \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( by^m \frac{\partial w}{\partial y} \right) = cw.$$

This is a two-dimensional equation of the heat and mass transfer theory with a linear source in an inhomogeneous anisotropic medium.

1°. For  $n \neq 2$  and  $m \neq 2$ , there are particular solutions of the form

$$w = w(\xi), \quad \xi = [b(2-m)^2 x^{2-n} + a(2-n)^2 y^{2-m}]^{1/2}.$$

The function  $w = w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{A}{\xi} w'_{\xi} = Bw, \quad (1)$$

where

$$A = \frac{4-nm}{(2-n)(2-m)}, \quad B = \frac{4c}{ab(2-n)^2(2-m)^2}.$$

The general solution of equation (1) is given by

$$w(\xi) = \xi^{\frac{1-A}{2}} \left[ C_1 J_{\nu}(\xi \sqrt{|B|}) + C_2 Y_{\nu}(\xi \sqrt{|B|}) \right] \quad \text{for } B < 0,$$

$$w(\xi) = \xi^{\frac{1-A}{2}} \left[ C_1 I_{\nu}(\xi \sqrt{B}) + C_2 K_{\nu}(\xi \sqrt{B}) \right] \quad \text{for } B > 0,$$

where  $\nu = \frac{1}{2}|1-A|$ ;  $C_1$  and  $C_2$  are arbitrary constants;  $J_{\nu}(z)$  and  $Y_{\nu}(z)$  are the Bessel functions; and  $I_{\nu}(z)$  and  $K_{\nu}(z)$  are the modified Bessel functions.



2°. There are multiplicatively separable particular solutions of the form

$$w(x, y) = \varphi(x)\psi(y),$$

where  $\varphi(x)$  and  $\psi(y)$  are determined by the following second-order linear ordinary differential equations ( $A_1$  is an arbitrary constant):

$$(ax^n \varphi'_x)'_x = A_1 \varphi, \quad (by^m \psi'_y)'_y = (c - A_1) \psi. \quad (2)$$

The solutions of equations (2) are expressed in terms of the Bessel functions (or modified Bessel functions); see equation 7.4.3.1, Item 3°.

3°. There are additively separable particular solutions of the form

$$w(x, y) = f(x) + g(y),$$

where  $f(x)$  and  $g(y)$  are determined by the following second-order linear ordinary differential equations ( $A_2$  is an arbitrary constant):

$$(ax^n f'_x)'_x - cf = A_2, \quad (by^m g'_y)'_y - cg = -A_2. \quad (3)$$

The solutions of equations (3) are expressed in terms of the Bessel functions (or modified Bessel functions).

4°. The transformation (specified by A. I. Zhurov, 2001)

$$x^{\frac{2-n}{2}} = Ar \cos \theta, \quad y^{\frac{2-m}{2}} = Br \sin \theta,$$

where  $A^2 = a(2-n)^2$  and  $B^2 = b(2-m)^2$ , leads to the equation

$$\frac{\partial^2 w}{\partial r^2} + \frac{4-nm}{(2-n)(2-m)} \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2} - \frac{2}{r^2} \frac{(nm-n-m) \cos 2\theta + (n-m)}{(2-n)(2-m) \sin 2\theta} \frac{\partial w}{\partial \theta} = 4cw,$$

which admits separable solutions of the form  $w(r, \theta) = F_1(r)F_2(\theta)$ .

$$4. \quad \frac{\partial}{\partial x} \left[ a(x+k)^n \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ b(y+s)^m \frac{\partial w}{\partial y} \right] = c.$$

The transformation  $\zeta = x+k$ ,  $\eta = y+s$  leads to an equation of the form 7.4.3.2:

$$\frac{\partial}{\partial \zeta} \left( a \zeta^n \frac{\partial w}{\partial \zeta} \right) + \frac{\partial}{\partial \eta} \left( b \eta^m \frac{\partial w}{\partial \eta} \right) = c.$$

$$5. \quad \frac{\partial}{\partial x} \left[ a(x+k)^n \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ b(y+s)^m \frac{\partial w}{\partial y} \right] = cw.$$

The transformation  $\zeta = x+k$ ,  $\eta = y+s$  leads to an equation of the form 7.4.3.3:

$$\frac{\partial}{\partial \zeta} \left( a \zeta^n \frac{\partial w}{\partial \zeta} \right) + \frac{\partial}{\partial \eta} \left( b \eta^m \frac{\partial w}{\partial \eta} \right) = cw.$$

$$6. \quad \frac{\partial}{\partial x} \left( a e^{\beta x} \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( b e^{\mu y} \frac{\partial w}{\partial y} \right) = 0.$$

This is a two-dimensional equation of the heat and mass transfer theory in an inhomogeneous anisotropic medium. Here,  $a_1(x) = a e^{\beta x}$  and  $a_2(y) = b e^{\mu y}$  are the principal thermal diffusivities.

1°. Particular solutions ( $A, B, C$  are arbitrary constants):

$$w(x, y) = Ae^{-\beta x} + Be^{-\mu y} + C,$$

$$w(x, y) = \frac{A}{a\beta^2}(\beta x + 1)e^{-\beta x} - \frac{A}{b\mu^2}(\mu y + 1)e^{-\mu y} + B,$$

$$w(x, y) = Ae^{-\beta x - \mu y} + B.$$

2°. There are multiplicatively separable particular solutions of the form

$$w(x, y) = \varphi(x)\psi(y), \quad (1)$$

where  $\varphi(x)$  and  $\psi(y)$  are determined by the following second-order linear ordinary differential equations ( $A_1$  is an arbitrary constant):

$$(ae^{\beta x}\varphi'_x)'_x = -A_1\varphi, \quad (2)$$

$$(be^{\mu y}\psi'_y)'_y = A_1\psi. \quad (3)$$

The solution of equation (2) is given by

$$\varphi(x) = \begin{cases} e^{-\beta x/2} [C_1 J_1(ke^{-\beta x/2}) + C_2 Y_1(ke^{-\beta x/2})] & \text{for } A_1 > 0, \\ e^{-\beta x/2} [C_1 I_1(ke^{-\beta x/2}) + C_2 K_1(ke^{-\beta x/2})] & \text{for } A_1 < 0, \end{cases}$$

where  $k = -(2/\beta)\sqrt{|A_1|/a}$ ;  $C_1$  and  $C_2$  are arbitrary constants;  $J_1(z)$  and  $Y_1(z)$  are the Bessel functions; and  $I_1(z)$  and  $K_1(z)$  are the modified Bessel functions.

The solution of equation (3) is given by

$$\psi(y) = \begin{cases} e^{-\mu y/2} [C_1 J_1(se^{-\mu y/2}) + C_2 Y_1(se^{-\mu y/2})] & \text{for } A_1 < 0, \\ e^{-\mu y/2} [C_1 I_1(se^{-\mu y/2}) + C_2 K_1(se^{-\mu y/2})] & \text{for } A_1 > 0, \end{cases}$$

where  $s = -(2/\mu)\sqrt{|A_1|/b}$ ;  $C_1$  and  $C_2$  are arbitrary constants.

The sum of solutions of the form (1) corresponding to different values of the parameter  $A_1$  is also a solution of the original equation.

3°. See equation 7.4.3.8, Item 3°, for  $c = 0$ .

$$7. \quad \frac{\partial}{\partial x} \left( ae^{\beta x} \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( be^{\mu y} \frac{\partial w}{\partial y} \right) = c.$$

This is a two-dimensional equation of the heat and mass transfer theory with constant volume release of heat in an inhomogeneous anisotropic medium. Here,  $a_1(x) = ae^{\beta x}$  and  $a_2(y) = be^{\mu y}$  are the principal thermal diffusivities.

The substitution

$$w(x, y) = U(x, y) - \frac{c}{a\beta^2}(\beta x + 1)e^{-\beta x}$$

leads to a homogeneous equation of the form 7.4.3.6:

$$\frac{\partial}{\partial x} \left( ae^{\beta x} \frac{\partial U}{\partial x} \right) + \frac{\partial}{\partial y} \left( be^{\mu y} \frac{\partial U}{\partial y} \right) = 0.$$

$$8. \quad \frac{\partial}{\partial x} \left( ae^{\beta x} \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( be^{\mu y} \frac{\partial w}{\partial y} \right) = cw.$$

This is a two-dimensional equation of the heat and mass transfer theory with a linear source in an inhomogeneous anisotropic medium.

1°. For  $\beta\mu \neq 0$ , there are particular solutions of the form

$$w = w(\xi), \quad \xi = (b\mu^2 e^{-\beta x} + a\beta^2 e^{-\mu y})^{1/2}.$$

The function  $w = w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} - \frac{1}{\xi} w'_{\xi} = Bw, \quad B = \frac{4c}{ab\beta^2\mu^2}.$$

For the solution of this equation, see 7.4.3.3 (Item 1° for  $A = -1$ ).

2°. The original equation admits multiplicatively (and additively) separable solutions. See equation 7.4.3.12 with  $f(x) = ae^{\beta x}$  and  $g(y) = be^{\mu y}$ .

3°. The transformation (specified by A. I. Zhurov, 2001)

$$e^{-\beta x/2} = Ar \cos \theta, \quad e^{-\mu y/2} = Br \sin \theta,$$

where  $A^2 = a\beta^2$  and  $B^2 = b\mu^2$ , leads to the equation

$$\frac{\partial^2 w}{\partial r^2} - \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2} - \frac{2}{r^2} \cot 2\theta \frac{\partial w}{\partial \theta} = 4cw,$$

which admits separable solutions of the form  $w(r, \theta) = F_1(r)F_2(\theta)$ .

$$9. \quad \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( be^{\beta y} \frac{\partial w}{\partial x} \right) = cw.$$

1°. For  $n \neq 2$  and  $\beta \neq 0$ , there are particular solutions of the form

$$w = w(r), \quad r^2 = \frac{x^{2-n}}{a(2-n)^2} + \frac{e^{-\beta y}}{b\beta^2}.$$

The function  $w = w(r)$  is determined by the ordinary differential equation

$$\frac{\partial^2 w}{\partial r^2} + \frac{n}{2-n} \frac{1}{r} \frac{\partial w}{\partial r} = 4cw.$$

For the solution of this equation, see 7.4.3.3 (Item 1°).

2°. The original equation admits multiplicatively (and additively) separable solutions. See equation 7.4.3.12 with  $f(x) = ax^n$  and  $g(y) = be^{\beta y}$ .

3°. The transformation (specified by A. I. Zhurov, 2001)

$$x^{1-\frac{1}{2}n} = Ar \cos \theta, \quad e^{-\frac{1}{2}\beta y} = Br \sin \theta,$$

where  $A^2 = a(2-n)^2$  and  $B^2 = b\beta^2$ , leads to the equation

$$\frac{\partial^2 w}{\partial r^2} + \frac{n}{2-n} \frac{1}{r} \frac{\partial w}{\partial r} + \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2} - \frac{2}{r^2} \frac{(1-n) \cos 2\theta + 1}{(2-n) \sin 2\theta} \frac{\partial w}{\partial \theta} = 4cw,$$

which admits separable solutions of the form  $w(r, \theta) = F_1(r)F_2(\theta)$ .

$$10. \quad \frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + \frac{\partial^2 w}{\partial y^2} = 0.$$

1°. Particular solutions:

$$w = C_1 y^2 + C_2 y - 2 \int \frac{C_1 x + C_3}{f(x)} dx + C_4,$$

$$w = C_1 y^3 + C_2 y - 6y \int \frac{C_1 x + C_3}{f(x)} dx + C_4,$$

$$w = [C_1 \Phi(x) + C_2]y + C_3 \Phi(x) + C_4, \quad \Phi(x) = \int \frac{dx}{f(x)},$$

$$w = [C_1 \Phi(x) + C_2]y^2 + C_3 \Phi(x) + C_4 - 2 \int \left\{ \frac{1}{f(x)} \int [C_1 \Phi(x) + C_2] dx \right\} dx,$$

where  $C_1, C_2, C_3, C_4$ , and  $C_5$  are arbitrary constants.

2°. Separable particular solution:

$$w = (C_1 e^{\lambda y} + C_2 e^{-\lambda y}) H(x),$$

where  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants, and the function  $H = H(x)$  is determined by the ordinary differential equation  $[f(x)H'_x]'_x + \lambda^2 H = 0$ .

3°. Separable particular solution:

$$w = [C_1 \sin(\lambda y) + C_2 \cos(\lambda y)] Z(x),$$

where  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants, and the function  $Z = Z(x)$  is determined by the ordinary differential equation  $[f(x)Z'_x]'_x - \lambda^2 Z = 0$ .

4°. Particular solutions with even powers of  $y$ :

$$w = \sum_{k=0}^n \zeta_k(x) y^{2k},$$

where the functions  $\zeta_k = \zeta_k(x)$  are defined by the recurrence relations

$$\begin{aligned} \zeta_n(x) &= A_n \Phi(x) + B_n, & \Phi(x) &= \int \frac{dx}{f(x)}, \\ \zeta_{k-1}(x) &= A_k \Phi(x) + B_k - 2k(2k-1) \int \frac{1}{f(x)} \left\{ \int \zeta_k(x) dx \right\} dx, \end{aligned}$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

5°. Particular solutions with odd powers of  $y$ :

$$w = \sum_{k=0}^n \eta_k(x) y^{2k+1},$$

where the functions  $\eta_k = \eta_k(x)$  are defined by the recurrence relations

$$\begin{aligned} \eta_n(x) &= A_n \Phi(x) + B_n, & \Phi(x) &= \int \frac{dx}{f(x)}, \\ \eta_{k-1}(x) &= A_k \Phi(x) + B_k - 2k(2k+1) \int \frac{1}{f(x)} \left\{ \int \eta_k(x) dx \right\} dx, \end{aligned}$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

$$11. \quad \frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(y) \frac{\partial w}{\partial y} \right] = 0.$$

This is a two-dimensional sourceless equation of the heat and mass transfer theory in an inhomogeneous anisotropic medium. The functions  $f = f(x)$  and  $g = g(y)$  are the principal thermal diffusivities.

1°. Particular solutions:

$$\begin{aligned} w(x, y) &= A_1 \int \frac{dx}{f(x)} + B_1 \int \frac{dy}{g(y)} + C_1, \\ w(x, y) &= A_2 \int \frac{x dx}{f(x)} - A_2 \int \frac{y dy}{g(y)} + B_2, \\ w(x, y) &= A_3 \int \frac{dx}{f(x)} \int \frac{dy}{g(y)} + B_3, \end{aligned}$$

where the  $A_k$ ,  $B_k$ , and  $C_1$  are arbitrary constants. A linear combination of these solutions is also a solution of the original equation.

2°. There are multiplicatively separable particular solutions of the form

$$w(x, y) = \varphi(x)\psi(y), \quad (1)$$

where  $\varphi(x)$  and  $\psi(y)$  are determined by the following second-order linear ordinary differential equations ( $A$  is an arbitrary constant):

$$\begin{aligned} (f\varphi'_x)'_x &= A\varphi, & f &= f(x), \\ (g\psi'_y)'_y &= -A\psi, & g &= g(y). \end{aligned} \quad (2)$$

The sum of solutions of the form (1) corresponding to different values of the parameter  $A$  in (2) is also a solution of the original equation (the solutions of some boundary value problems may be obtained by separation of variables).

12. 
$$\frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(y) \frac{\partial w}{\partial y} \right] = \beta w.$$

This is a two-dimensional equation of the heat and mass transfer theory with a linear source in an inhomogeneous anisotropic medium. The functions  $f = f(x)$  and  $g = g(y)$  are the principal thermal diffusivities.

1°. There are multiplicatively separable particular solutions of the form

$$w(x, y) = \varphi(x)\psi(y), \quad (1)$$

where  $\varphi(x)$  and  $\psi(y)$  are determined by the following second-order linear ordinary differential equations ( $A$  is an arbitrary constant):

$$\begin{aligned} (f\varphi'_x)'_x &= A\varphi, & f &= f(x), \\ (g\psi'_y)'_y &= (\beta - A)\psi, & g &= g(y). \end{aligned} \quad (2)$$

The sum of solutions of the form (1) corresponding to different values of the parameter  $A$  in (2) is also a solution of the original equation; the solutions of some boundary value problems may be obtained by separation of variables.

2°. There are additively separable particular solutions of the form

$$w(x, y) = \Phi(x) + \Psi(y),$$

where  $\Phi(x)$  and  $\Psi(y)$  are determined by the following second-order linear ordinary differential equations ( $C$  is an arbitrary constant):

$$\begin{aligned} (f\Phi'_x)'_x - \beta\Phi &= C, & f &= f(x), \\ (g\Psi'_y)'_y - \beta\Psi &= -C, & g &= g(y). \end{aligned}$$

In the special case  $\beta = 0$ , the solutions of these equations can be represented as

$$\begin{aligned} \Phi(x) &= C \int \frac{x dx}{f(x)} + A_1 \int \frac{dx}{f(x)} + B_1, \\ \Psi(y) &= -C \int \frac{y dy}{g(y)} + A_2 \int \frac{dy}{g(y)} + B_2, \end{aligned}$$

where  $A_1$ ,  $A_2$ ,  $B_1$ , and  $B_2$  are arbitrary constants.

## 7.4.4. Other Equations Arising in Applications

1.  $y \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = 0.$

*Tricomi equation.* It is used to describe near-sonic flows of gas.

1°. Particular solutions:

$$\begin{aligned} w &= Axy + Bx + Cy + D, \\ w &= A(3x^2 - y^3) + B(x^3 - xy^3) + C(6yx^2 - y^4), \end{aligned}$$

where  $A$ ,  $B$ ,  $C$ , and  $D$  are arbitrary constants.

2°. Particular solutions with even powers of  $x$ :

$$w = \sum_{k=0}^n \varphi_k(y) x^{2k},$$

where the functions  $\varphi_k = \varphi_k(y)$  are defined by the recurrence relations

$$\varphi_n(y) = A_n y + B_n, \quad \varphi_{k-1}(y) = A_k y + B_k - 2k(2k-1) \int_0^y (y-t)t\varphi_k(t) dt,$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

3°. Particular solutions with odd powers of  $x$ :

$$w = \sum_{k=0}^n \psi_k(y) x^{2k+1},$$

where the functions  $\psi_k = \psi_k(y)$  are defined by the recurrence relations

$$\psi_n(y) = A_n y + B_n, \quad \psi_{k-1}(y) = A_k y + B_k - 2k(2k+1) \int_0^y (y-t)t\psi_k(t) dt,$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ).

4°. Separable particular solutions:

$$\begin{aligned} w(x, y) &= [A \sinh(3\lambda x) + B \cosh(3\lambda x)] \sqrt{y} [C J_{1/3}(2\lambda y^{3/2}) + D Y_{1/3}(2\lambda y^{3/2})], \\ w(x, y) &= [A \sin(3\lambda x) + B \cos(3\lambda x)] \sqrt{y} [C I_{1/3}(2\lambda y^{3/2}) + D K_{1/3}(2\lambda y^{3/2})], \end{aligned}$$

where  $A$ ,  $B$ ,  $C$ ,  $D$ , and  $\lambda$  are arbitrary constants,  $J_{1/3}(z)$  and  $Y_{1/3}(z)$  are the Bessel functions, and  $I_{1/3}(z)$  and  $K_{1/3}(z)$  are the modified Bessel functions.

5°. For  $y > 0$ , see also equation 7.4.4.2 with  $n = 1$ . For  $y < 0$ , the change of variable  $y = -t$  leads to an equation of the form 4.3.3.11 with  $n = 1$ .

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

2.  $y^n \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = 0.$

1°. Particular solutions:

$$\begin{aligned} w &= Axy + Bx + Cy + D, \\ w &= Ax^2 - \frac{2A}{(n+1)(n+2)} y^{n+2}, \\ w &= Ax^3 - \frac{6A}{(n+1)(n+2)} xy^{n+2}, \\ w &= Ayx^2 - \frac{2A}{(n+2)(n+3)} y^{n+3}, \end{aligned}$$

where  $A$ ,  $B$ ,  $C$ , and  $D$  are arbitrary constants.

2°. Particular solutions with even powers of  $x$ :

$$w = \sum_{k=0}^m \varphi_k(y) x^{2k},$$

where the functions  $\varphi_k = \varphi_k(y)$  are defined by the recurrence relations

$$\varphi_m(y) = A_m y + B_m, \quad \varphi_{k-1}(y) = A_k y + B_k - 2k(2k-1) \int_a^y (y-t)^n \varphi_k(t) dt,$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = m, \dots, 1$ ),  $a$  is any number.

3°. Particular solutions with odd powers of  $x$ :

$$w = \sum_{k=0}^m \psi_k(y) x^{2k+1},$$

where the functions  $\psi_k = \psi_k(y)$  are defined by the recurrence relations

$$\psi_m(y) = A_m y + B_m, \quad \psi_{k-1}(y) = A_k y + B_k - 2k(2k+1) \int_a^y (y-t)^n \psi_k(t) dt,$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = m, \dots, 1$ ),  $a$  is any number.

4°. Separable particular solutions:

$$w(x, y) = [A \sinh(\lambda q x) + B \cosh(\lambda q x)] \sqrt{y} [C J_{\frac{1}{2q}}(\lambda y^q) + D Y_{\frac{1}{2q}}(\lambda y^q)], \quad q = \frac{1}{2}(n+2),$$

$$w(x, y) = [A \sin(\lambda q x) + B \cos(\lambda q x)] \sqrt{y} [C I_{\frac{1}{2q}}(\lambda y^q) + D K_{\frac{1}{2q}}(\lambda y^q)],$$

where  $A, B, C, D$ , and  $\lambda$  are arbitrary constants,  $J_\nu(z)$  and  $Y_\nu(z)$  are the Bessel functions, and  $I_\nu(z)$  and  $K_\nu(z)$  are the modified Bessel functions.

5°. Fundamental solutions (for  $y > 0$ ):

$$w_1(x, y, x_0, y_0) = k_1 (r_1^2)^{-\beta} F(\beta, \beta, 2\beta; 1 - \xi), \quad \beta = \frac{n}{2(n+2)}, \quad \xi = \frac{r_2^2}{r_1^2},$$

$$w_2(x, y, x_0, y_0) = k_2 (r_1^2)^{-\beta} (1 - \xi)^{1-2\beta} F(1 - \beta, 1 - \beta, 2 - 2\beta; 1 - \xi).$$

Here,  $F(a, b, c; \xi)$  is the hypergeometric function and

$$r_1^2 = (x - x_0)^2 + \frac{4}{(n+2)^2} \left( y^{\frac{n+2}{2}} + y_0^{\frac{n+2}{2}} \right), \quad k_1 = \frac{1}{4\pi} \left( \frac{4}{n+2} \right)^{2\beta} \frac{\Gamma^2(\beta)}{\Gamma(2\beta)},$$

$$r_2^2 = (x - x_0)^2 + \frac{4}{(n+2)^2} \left( y^{\frac{n+2}{2}} - y_0^{\frac{n+2}{2}} \right), \quad k_2 = \frac{1}{4\pi} \left( \frac{4}{n+2} \right)^{2\beta} \frac{\Gamma^2(1-\beta)}{\Gamma(2-2\beta)},$$

where  $\Gamma(\beta)$  is the gamma function;  $x_0$  and  $y_0$  are arbitrary constants.

The fundamental solutions satisfy the conditions

$$\partial_y w_1|_{y=0} = 0, \quad w_2|_{y=0} = 0 \quad (x \text{ and } x_0 \text{ are any, } y_0 > 0).$$

The solutions of some boundary value problems can be found in the first book cited below.

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), A. D. Polyanin (2001a).

$$3. \quad \frac{\partial^2 w}{\partial r^2} + \frac{\alpha}{r} \frac{\partial w}{\partial r} + \frac{\partial^2 w}{\partial z^2} = 0.$$

*Elliptic analogue of the Euler–Poisson–Darboux equation.*

1°. For  $\alpha = 1$ , see Subsections 8.1.2 and 8.2.3 with  $w = w(r, z)$ . For  $\alpha \neq 1$ , the transformation  $x = (1 - \alpha)z$ ,  $y = r^{1-\alpha}$  leads to an equation of the form 7.4.4.1:

$$y^{\frac{2\alpha}{1-\alpha}} \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = 0.$$

2°. Suppose  $w_\alpha = w_\alpha(r, z)$  is a solution of the equation in question for a fixed value of the parameter  $\alpha$ . Then the functions  $\tilde{w}_\alpha$  defined by the relations

$$\begin{aligned}\tilde{w}_\alpha &= \frac{\partial w_\alpha}{\partial z}, \\ \tilde{w}_\alpha &= r \frac{\partial w_\alpha}{\partial r} + z \frac{\partial w_\alpha}{\partial z}, \\ \tilde{w}_\alpha &= 2rz \frac{\partial w_\alpha}{\partial r} + (z^2 - r^2) \frac{\partial w_\alpha}{\partial z} + \alpha z w_\alpha\end{aligned}$$

are also solutions of this equation.

3°. Suppose  $w_\alpha = w_\alpha(r, z)$  is a solution of the equation in question for a fixed value of the parameter  $\alpha$ . Using this  $w_\alpha$ , one can construct solutions of the equation with other values of the parameter by the formulas

$$\begin{aligned}w_{2-\alpha} &= r^{\alpha-1} w_\alpha, \\ w_{\alpha-2} &= r \frac{\partial w_\alpha}{\partial r} + (\alpha - 1) w_\alpha, \\ w_{\alpha-2} &= rz \frac{\partial w_\alpha}{\partial r} - r^2 \frac{\partial w_\alpha}{\partial z} + (\alpha - 1) z w_\alpha, \\ w_{\alpha-2} &= r(r^2 - z^2) \frac{\partial w_\alpha}{\partial r} + 2r^2 z \frac{\partial w_\alpha}{\partial z} + [r^2 - (\alpha - 1)z^2] w_\alpha, \\ w_{\alpha+2} &= \frac{1}{r} \frac{\partial w_\alpha}{\partial r}, \\ w_{\alpha+2} &= \frac{z}{r} \frac{\partial w_\alpha}{\partial r} - \frac{\partial w_\alpha}{\partial z}, \\ w_{\alpha+2} &= \frac{r^2 - z^2}{r} \frac{\partial w_\alpha}{\partial r} + 2z \frac{\partial w_\alpha}{\partial z} + \alpha w_\alpha.\end{aligned}$$

⊙ *Reference:* A. V. Aksenov (2001).

$$4. \quad \frac{\partial^2 w}{\partial x^2} + f(x) \frac{\partial^2 w}{\partial y^2} = 0.$$

1°. Particular solutions:

$$\begin{aligned}w &= C_1 xy + C_2 y + C_3 x + C_4, \\ w &= C_1 y^2 + C_2 xy + C_3 y + C_4 x - 2C_1 \int_a^x (x-t)f(t) dt + C_5, \\ w &= C_1 y^3 + C_2 xy + C_3 y + C_4 x - 6C_1 y \int_a^x (x-t)f(t) dt + C_5, \\ w &= (C_1 x + C_2) y^2 + C_3 xy + C_4 y + C_5 x - 2 \int_a^x (x-t)(C_1 t + C_2) f(t) dt + C_6,\end{aligned}$$

where  $C_1, C_2, C_3, C_4, C_5$ , and  $C_6$  are arbitrary constants,  $a$  is any number.



2°. Separable particular solution:

$$w = (C_1 e^{\lambda y} + C_2 e^{-\lambda y})H(x),$$

where  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants, and the function  $H = H(x)$  is determined by the ordinary differential equation  $H''_{xx} + \lambda^2 f(x)H = 0$ .

3°. Separable particular solution:

$$w = [C_1 \sin(\lambda y) + C_2 \cos(\lambda y)]Z(x),$$

where  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants, and the function  $Z = Z(x)$  is determined by the ordinary differential equation  $Z''_{xx} - \lambda^2 f(x)Z = 0$ .

4°. Particular solutions with even powers of  $y$ :

$$w = \sum_{k=0}^n \varphi_k(x) y^{2k},$$

where the functions  $\varphi_k = \varphi_k(x)$  are defined by the recurrence relations

$$\varphi_n(x) = A_n x + B_n, \quad \varphi_{k-1}(x) = A_k x + B_k - 2k(2k-1) \int_a^x (x-t)f(t)\varphi_k(t) dt,$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ),  $a$  is any number.

5°. Particular solutions with odd powers of  $y$ :

$$w = \sum_{k=0}^n \psi_k(x) y^{2k+1},$$

where the functions  $\psi_k = \psi_k(x)$  are defined by the recurrence relations

$$\psi_n(x) = A_n x + B_n, \quad \psi_{k-1}(x) = A_k x + B_k - 2k(2k+1) \int_a^x (x-t)f(t)\psi_k(t) dt,$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = n, \dots, 1$ ), and  $a$  is any number.

$$5. \quad \frac{\partial}{\partial x} \left[ f_1(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ f_2(y) \frac{\partial w}{\partial y} \right] + \lambda [g_1(x) + g_2(y)] w = 0.$$

This equation is encountered in the theory of vibration of inhomogeneous membranes. Its separable solutions are sought in the form  $w(x, y) = \varphi(x)\psi(y)$ .

The article cited below presents an algorithm for accelerated convergence of solutions to eigenvalue boundary value problems for this equation.

⊙ *Reference:* L. D. Akulenko and S. V. Nesterov (1999).

## 7.4.5. Equations of the Form

$$a(x) \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + b(x) \frac{\partial w}{\partial x} + c(x)w = -\Phi(x, y)$$

7.4.5-1. Statements of boundary value problems. Relations for the Green's function.

Consider two-dimensional boundary value problems for the equation

$$a(x) \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + b(x) \frac{\partial w}{\partial x} + c(x)w = -\Phi(x, y) \quad (1)$$

with general boundary conditions in  $x$ ,

$$\begin{aligned} s_1 \partial_x w - k_1 w &= f_1(y) \quad \text{at } x = x_1, \\ s_2 \partial_x w + k_2 w &= f_2(y) \quad \text{at } x = x_2, \end{aligned} \quad (2)$$

and different boundary conditions in  $y$ . We assume that the coefficients of equation (1) and the boundary conditions (2) meet the requirement

$a(x)$ ,  $b(x)$ ,  $c(x)$  are continuous functions ( $x_1 \leq x \leq x_2$ );  $a > 0$ ,  $|s_1| + |k_1| > 0$ ,  $|s_2| + |k_2| > 0$ .

In the general case, the Green's function can be represented as

$$G(x, y, \xi, \eta) = \rho(\xi) \sum_{n=1}^{\infty} \frac{u_n(x)u_n(\xi)}{\|u_n\|^2} \Psi_n(y, \eta; \lambda_n). \quad (3)$$

Here,

$$\rho(x) = \frac{1}{a(x)} \exp \left[ \int \frac{b(x)}{a(x)} dx \right], \quad \|u_n\|^2 = \int_{x_1}^{x_2} \rho(x) u_n^2(x) dx, \quad (4)$$

and the  $\lambda_n$  and  $u_n(x)$  are the eigenvalues and eigenfunctions of the homogeneous boundary value problem for the ordinary differential equation

$$a(x)u''_{xx} + b(x)u'_x + [\lambda + c(x)]u = 0, \quad (5)$$

$$s_1 u'_x - k_1 u = 0 \quad \text{at } x = x_1, \quad (6)$$

$$s_2 u'_x + k_2 u = 0 \quad \text{at } x = x_2. \quad (7)$$

The functions  $\Psi_n$  for various boundary conditions in  $y$  are specified in [Table 25](#).

Equation (5) can be rewritten in self-adjoint form as

$$[p(x)u'_x]'_x + [\lambda \rho(x) - q(x)]u = 0, \quad (8)$$

where the functions  $p(x)$  and  $q(x)$  are given by

$$p(x) = \exp \left[ \int \frac{b(x)}{a(x)} dx \right], \quad q(x) = -\frac{c(x)}{a(x)} \exp \left[ \int \frac{b(x)}{a(x)} dx \right],$$

and  $\rho(x)$  is defined in (4).

The eigenvalue problem (8), (6), (7) possesses the following properties:

1°. All eigenvalues  $\lambda_1, \lambda_2, \dots$  are real and  $\lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ .

2°. The system of eigenfunctions  $\{u_1(x), u_2(x), \dots\}$  is orthogonal on the interval  $x_1 \leq x \leq x_2$  with weight  $\rho(x)$ , that is,

$$\int_{x_1}^{x_2} \rho(x) u_n(x) u_m(x) dx = 0 \quad \text{for } n \neq m.$$

3°. If the conditions

$$q(x) \geq 0, \quad s_1 k_1 \geq 0, \quad s_2 k_2 \geq 0 \quad (9)$$

are satisfied, there are no negative eigenvalues. If  $q \equiv 0$  and  $k_1 = k_2 = 0$ , then the least eigenvalue is  $\lambda_0 = 0$  and the corresponding eigenfunction is  $u_0 = \text{const}$ ; in this case, the summation in (3) must start with  $n = 0$ . In the other cases, if conditions (9) are satisfied, all eigenvalues are positive; for example, the first inequality in (9) holds if  $c(x) \leq 0$ .

TABLE 25

The functions  $\Psi_n$  in (3) for various boundary conditions.\* Notation:  $\beta_n = \sqrt{\lambda_n}$ 

| Domain                 | Boundary conditions                                          | Function $\Psi_n(y, \eta; \lambda_n)$                                                                                                                                                                                                                     |
|------------------------|--------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $-\infty < y < \infty$ | $ w  < \infty$ for $y \rightarrow \pm\infty$                 | $\frac{1}{2\beta_n} e^{-\beta_n y-\eta }$                                                                                                                                                                                                                 |
| $0 \leq y < \infty$    | $w=0$ for $y=0$                                              | $\frac{1}{\beta_n} \begin{cases} e^{-\beta_n y} \sinh(\beta_n \eta) & \text{for } y > \eta, \\ e^{-\beta_n \eta} \sinh(\beta_n y) & \text{for } \eta > y \end{cases}$                                                                                     |
| $0 \leq y < \infty$    | $\partial_y w = 0$ for $y=0$                                 | $\frac{1}{\beta_n} \begin{cases} e^{-\beta_n y} \cosh(\beta_n \eta) & \text{for } y > \eta, \\ e^{-\beta_n \eta} \cosh(\beta_n y) & \text{for } \eta > y \end{cases}$                                                                                     |
| $0 \leq y < \infty$    | $\partial_y w - k_3 w = 0$ for $y=0$                         | $\frac{1}{\beta_n(\beta_n + k_3)} \begin{cases} e^{-\beta_n y} [\beta_n \cosh(\beta_n \eta) + k_3 \sinh(\beta_n \eta)] & \text{for } y > \eta, \\ e^{-\beta_n \eta} [\beta_n \cosh(\beta_n y) + k_3 \sinh(\beta_n y)] & \text{for } \eta > y \end{cases}$ |
| $0 \leq y \leq h$      | $w=0$ at $y=0$ ,<br>$w=0$ at $y=h$                           | $\frac{1}{\beta_n \sinh(\beta_n h)} \begin{cases} \sinh(\beta_n \eta) \sinh[\beta_n(h-y)] & \text{for } y > \eta, \\ \sinh(\beta_n y) \sinh[\beta_n(h-\eta)] & \text{for } \eta > y \end{cases}$                                                          |
| $0 \leq y \leq h$      | $\partial_y w = 0$ at $y=0$ ,<br>$\partial_y w = 0$ at $y=h$ | $\frac{1}{\beta_n \sinh(\beta_n h)} \begin{cases} \cosh(\beta_n \eta) \cosh[\beta_n(h-y)] & \text{for } y > \eta, \\ \cosh(\beta_n y) \cosh[\beta_n(h-\eta)] & \text{for } \eta > y \end{cases}$                                                          |
| $0 \leq y \leq h$      | $w=0$ at $y=0$ ,<br>$\partial_y w = 0$ at $y=h$              | $\frac{1}{\beta_n \cosh(\beta_n h)} \begin{cases} \sinh(\beta_n \eta) \cosh[\beta_n(h-y)] & \text{for } y > \eta, \\ \sinh(\beta_n y) \cosh[\beta_n(h-\eta)] & \text{for } \eta > y \end{cases}$                                                          |

Subsection 1.8.9 presents some relations for estimating the eigenvalues  $\lambda_n$  and eigenfunctions  $u_n(x)$ .

The Green's function of the two-dimensional third boundary value problem (1)–(2) augmented by the boundary conditions

$$\frac{\partial w}{\partial y} - k_3 w = 0 \quad \text{at } y = 0, \quad \frac{\partial w}{\partial y} + k_4 w = 0 \quad \text{at } y = h$$

is given by relation (3) with

$$\Psi_n(y, \eta; \lambda_n) = \begin{cases} \frac{[\beta_n \cosh(\beta_n \eta) + k_3 \sinh(\beta_n \eta)] \{ \beta_n \cosh[\beta_n(h-y)] + k_4 \sinh[\beta_n(h-y)] \}}{\beta_n [\beta_n(k_3 + k_4) \cosh(\beta_n h) + (\beta_n^2 + k_3 k_4) \sinh(\beta_n h)]} & \text{for } y > \eta, \\ \frac{[\beta_n \cosh(\beta_n y) + k_3 \sinh(\beta_n y)] \{ \beta_n \cosh[\beta_n(h-\eta)] + k_4 \sinh[\beta_n(h-\eta)] \}}{\beta_n [\beta_n(k_3 + k_4) \cosh(\beta_n h) + (\beta_n^2 + k_3 k_4) \sinh(\beta_n h)]} & \text{for } y < \eta. \end{cases}$$

#### 7.4.5-2. Representation of solutions to boundary value problems using the Green's function.

1°. The solution of the first boundary value problem for equation (1) with the boundary conditions

$$\begin{aligned} w &= f_1(y) \quad \text{at } x = x_1, & w &= f_2(y) \quad \text{at } x = x_2, \\ w &= f_3(x) \quad \text{at } y = 0, & w &= f_4(x) \quad \text{at } y = h \end{aligned}$$

\* For unbounded domains, the condition of boundedness of the solution as  $y \rightarrow \pm\infty$  is set; in Table 25, this condition is omitted.

is expressed in terms of the Green's function as

$$\begin{aligned}
 w(x, y) = & a(x_1) \int_0^h f_1(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=x_1} d\eta - a(x_2) \int_0^h f_2(\eta) \left[ \frac{\partial}{\partial \xi} G(x, y, \xi, \eta) \right]_{\xi=x_2} d\eta \\
 & + \int_{x_1}^{x_2} f_3(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=0} d\xi - \int_{x_1}^{x_2} f_4(\xi) \left[ \frac{\partial}{\partial \eta} G(x, y, \xi, \eta) \right]_{\eta=h} d\xi \\
 & + \int_{x_1}^{x_2} \int_0^h \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi.
 \end{aligned}$$

2°. The solution of the second boundary value problem for equation (1) with boundary conditions

$$\begin{aligned}
 \partial_x w = f_1(y) \quad \text{at} \quad x = x_1, \quad \partial_x w = f_2(y) \quad \text{at} \quad x = x_2, \\
 \partial_y w = f_3(x) \quad \text{at} \quad y = 0, \quad \partial_y w = f_4(x) \quad \text{at} \quad y = h
 \end{aligned}$$

is expressed in terms of the Green's function as

$$\begin{aligned}
 w(x, y) = & -a(x_1) \int_0^h f_1(\eta) G(x, y, x_1, \eta) d\eta + a(x_2) \int_0^h f_2(\eta) G(x, y, x_2, \eta) d\eta \\
 & - \int_{x_1}^{x_2} f_3(\xi) G(x, y, \xi, 0) d\xi + \int_{x_1}^{x_2} f_4(\xi) G(x, y, \xi, h) d\xi \\
 & + \int_{x_1}^{x_2} \int_0^h \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi.
 \end{aligned}$$

3°. The solution of the third boundary value problem for equation (1) in terms of the Green's function is represented in the same way as the solution of the second boundary value problem (the Green's function is now different).

## Chapter 8

# Elliptic Equations with Three or More Space Variables

---

### 8.1. Laplace Equation $\Delta_3 w = 0$

The three-dimensional Laplace equation is often encountered in heat and mass transfer theory, fluid mechanics, elasticity, electrostatics, and other areas of mechanics and physics. For example, in heat and mass transfer theory, this equation describes stationary temperature distribution in the absence of heat sources and sinks in the domain under study.

A regular solution of the Laplace equation is called a harmonic function. The first boundary value problem for the Laplace equation is often referred to as the Dirichlet problem, and the second boundary value problem, as the Neumann problem.

*Extremum principle:* Given a domain  $D$ , a harmonic function  $w$  in  $D$  that is not identically constant in  $D$  cannot attain its maximum or minimum value at any interior point of  $D$ .

#### 8.1.1. Problems in Cartesian Coordinates

The three-dimensional Laplace equation in the rectangular Cartesian system of coordinates is written as

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} = 0.$$

**8.1.1-1. Particular solutions and some relations.**

1°. Particular solutions:

$$\begin{aligned}w(x, y, z) &= Ax + By + Cz + D, \\w(x, y, z) &= Ax^2 + By^2 - (A + B)z^2 + Cxy + Dxz + Eyz, \\w(x, y, z) &= \cos(\mu_1 x + \mu_2 y) \exp(\pm \mu z), \\w(x, y, z) &= \sin(\mu_1 x + \mu_2 y) \exp(\pm \mu z), \\w(x, y, z) &= \exp(\mu_1 x + \mu_2 y) \cos(\mu z + A), \\w(x, y, z) &= \exp(\pm \mu x) \cos(\mu_1 y + A) \cos(\mu_2 z + B), \\w(x, y, z) &= \cosh(\mu_1 x) \cosh(\mu_2 y) \cos(\mu z + B), \\w(x, y, z) &= \cosh(\mu_1 x) \sinh(\mu_2 y) \cos(\mu z + B), \\w(x, y, z) &= \cosh(\mu x) \cos(\mu_1 y + A) \cos(\mu_2 z + B), \\w(x, y, z) &= \sinh(\mu_1 x) \sinh(\mu_2 y) \sin(\mu z + B), \\w(x, y, z) &= \sinh(\mu x) \sin(\mu_1 y + A) \sin(\mu_2 z + B),\end{aligned}$$

where  $A, B, C, D, E, \mu_1$ , and  $\mu_2$  are arbitrary constants, and  $\mu = \sqrt{\mu_1^2 + \mu_2^2}$ .

2°. Fundamental solution:

$$\mathcal{E}(x, y, z) = \frac{1}{4\pi\sqrt{x^2 + y^2 + z^2}}.$$

3°. Suppose  $w = w(x, y, z)$  is a solution of the Laplace equation. Then the functions

$$w_1 = Aw(\pm\lambda x + C_1, \pm\lambda y + C_2, \pm\lambda x + C_3),$$

$$w_2 = \frac{A}{r} w\left(\frac{x}{r^2}, \frac{y}{r^2}, \frac{z}{r^2}\right), \quad r = \sqrt{x^2 + y^2 + z^2},$$

$$w_3 = \frac{A}{\sqrt{\Xi}} w\left(\frac{x - ar^2}{\Xi}, \frac{y - br^2}{\Xi}, \frac{z - cr^2}{\Xi}\right), \quad \Xi = 1 - 2(ax + by + cz) + (a^2 + b^2 + c^2)r^2,$$

where  $A$ ,  $C_n$ ,  $a$ ,  $b$ ,  $c$ , and  $\lambda$  are arbitrary constants, are also solutions of this equation. The signs at  $\lambda$  in the expression of  $w_1$  can be taken independently of one another.

⊙ *References:* W. Miller, Jr. (1977), R. Courant and D. Hilbert (1989).

8.1.1-2. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z < \infty$ . First boundary value problem.

A half-space is considered. A boundary condition is prescribed:

$$w = f(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$w(x, y, z) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{zf(\xi, \eta) d\xi d\eta}{[(x - \xi)^2 + (y - \eta)^2 + z^2]^{3/2}}.$$

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

8.1.1-3. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z < \infty$ . Second boundary value problem.

A half-space is considered. A boundary condition is prescribed:

$$\partial_z w = f(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$w(x, y, z) = -\frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{f(\xi, \eta) d\xi d\eta}{\sqrt{(x - \xi)^2 + (y - \eta)^2 + z^2}} + C,$$

where  $C$  is an arbitrary constant.

⊙ *Reference:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

8.1.1-4. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . First boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$w = f_1(y, z) \quad \text{at} \quad x = 0, \quad w = f_2(y, z) \quad \text{at} \quad x = a,$$

$$w = f_3(x, z) \quad \text{at} \quad y = 0, \quad w = f_4(x, z) \quad \text{at} \quad y = b,$$

$$w = f_5(x, y) \quad \text{at} \quad z = 0, \quad w = f_6(x, y) \quad \text{at} \quad z = c.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{f_{nm}^2 \sinh(\lambda_{nm}^1 x) + f_{nm}^1 \sinh[\lambda_{nm}^1 (a - x)]}{\sinh(\lambda_{nm}^1 a)} \sin\left(\frac{\pi n y}{b}\right) \sin\left(\frac{\pi m z}{c}\right) \\ & + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{f_{nm}^4 \sinh(\lambda_{nm}^2 y) + f_{nm}^3 \sinh[\lambda_{nm}^2 (b - y)]}{\sinh(\lambda_{nm}^2 b)} \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m z}{c}\right) \\ & + \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{f_{nm}^6 \sinh(\lambda_{nm}^3 z) + f_{nm}^5 \sinh[\lambda_{nm}^3 (c - z)]}{\sinh(\lambda_{nm}^3 c)} \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m y}{b}\right), \end{aligned}$$

where the constant coefficients are given by

$$\lambda_{nm}^1 = \pi \sqrt{\frac{n^2}{b^2} + \frac{m^2}{c^2}}, \quad \lambda_{nm}^2 = \pi \sqrt{\frac{n^2}{a^2} + \frac{m^2}{c^2}}, \quad \lambda_{nm}^3 = \pi \sqrt{\frac{n^2}{a^2} + \frac{m^2}{b^2}},$$

$$f_{nm}^i = \begin{cases} \frac{4}{bc} \int_0^b \int_0^c f_i(y, z) \sin\left(\frac{\pi n y}{b}\right) \sin\left(\frac{\pi m z}{c}\right) dy dz & \text{for } i = 1, 2; \\ \frac{4}{ac} \int_0^a \int_0^c f_i(x, z) \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m z}{c}\right) dx dz & \text{for } i = 3, 4; \\ \frac{4}{ab} \int_0^a \int_0^b f_i(x, y) \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m y}{b}\right) dx dy & \text{for } i = 5, 6. \end{cases}$$

**Example.** The planes  $x = 0$  and  $x = a$  have constant temperatures  $w_1$  and  $w_2$ , respectively. The other planes are maintained at zero temperature ( $f_3 = f_4 = f_5 = f_6 = 0$ ).

Solution:

$$w = \frac{16}{\pi^2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{w_2 \sinh(\mu_{nm} x) + w_1 \sinh[\mu_{nm}(a-x)]}{(2n+1)(2m+1) \sinh(\mu_{nm} a)} \sin(p_n y) \sin(q_m z),$$

$$p_n = \frac{\pi(2n+1)}{b}, \quad q_m = \frac{\pi(2m+1)}{c}, \quad \mu_{nm} = \sqrt{p_n^2 + q_m^2}.$$

☉ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), H. S. Carslaw and J. C. Jaeger (1984).

► *For the solution of other boundary value problems for the three-dimensional Laplace equation in the Cartesian coordinate system, see Subsection 8.2.2 for  $\Phi \equiv 0$ .*

## 8.1.2. Problems in Cylindrical Coordinates

The three-dimensional Laplace equation in the cylindrical coordinate system is written as

$$\Delta_3 w \equiv \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} = 0, \quad r = \sqrt{x^2 + y^2}.$$

### 8.1.2-1. Particular solutions:

$$w(r, \varphi, z) = \left( Ar^m + \frac{B}{r^m} \right) (C \cos m\varphi + D \sin m\varphi)(\alpha + \beta z),$$

$$w(r, \varphi, z) = J_m(\mu r)(A \cos m\varphi + B \sin m\varphi)(C \cosh \mu z + D \sinh \mu z),$$

$$w(r, \varphi, z) = Y_m(\mu r)(A \cos m\varphi + B \sin m\varphi)(C \cosh \mu z + D \sinh \mu z),$$

$$w(r, \varphi, z) = I_m(\mu r)(A \cos m\varphi + B \sin m\varphi)(C \cos \mu z + D \sin \mu z),$$

$$w(r, \varphi, z) = K_m(\mu r)(A \cos m\varphi + B \sin m\varphi)(C \cos \mu z + D \sin \mu z),$$

where  $m = 0, 1, 2, \dots$ ;  $A, B, C, D, \alpha, \beta$ , and  $\mu$  are arbitrary constants; the  $J_m(\xi)$  and  $Y_m(\xi)$  are the Bessel functions; and the  $I_m(\xi)$  and  $K_m(\xi)$  are the modified Bessel functions.

### 8.1.2-2. Domain: $0 \leq r \leq a$ , $0 \leq \varphi \leq 2\pi$ , $-\infty < z < \infty$ . First boundary value problem.

An infinite circular cylinder is considered. A boundary condition is prescribed:

$$w = f(\varphi, z) \quad \text{at} \quad r = a.$$

Solution:

$$w(r, \varphi, z) = -\frac{1}{a} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{J_n(\lambda_{nm} r)}{J'_n(\lambda_{nm} a)} \int_{-\infty}^{\infty} [A_n(\xi) \cos n\varphi + B_n(\xi) \sin n\varphi] \exp(-\lambda_{nm} |\xi - z|) d\xi,$$

where the  $J_n(r)$  are the Bessel functions and  $\lambda_{mn}$  are positive roots of the transcendental equation  $J_n(a\lambda) = 0$ . The functions  $A_n(z)$  and  $B_n(z)$  are the coefficients of the Fourier series expansion of  $f(\varphi, z)$ ,

$$A_n(z) = \frac{\varepsilon_n}{\pi} \int_0^{2\pi} f(\varphi, z) \cos(n\varphi) d\varphi, \quad B_n(z) = \frac{1}{\pi} \int_0^{2\pi} f(\varphi, z) \sin(n\varphi) d\varphi,$$

where  $\varepsilon_0 = 1/2$  and  $\varepsilon_n = 1$  for  $n = 1, 2, \dots$

If the surface temperature is independent of  $\varphi$ , i.e.,  $f(z, \varphi) = f(z)$ , then the solution takes the form

$$w(r, \varphi, z) = \frac{1}{a} \sum_{m=1}^{\infty} \frac{J_0(\lambda_m r)}{J_1(\lambda_m a)} \int_0^{\infty} [f(z + \zeta) + f(z - \zeta)] \exp(-\lambda_m \zeta) d\zeta,$$

where the  $\lambda_m$  are positive roots of the transcendental equation  $J_0(a\lambda) = 0$ .

⊙ *Reference:* H. S. Carslaw and J. C. Jaeger (1984).

8.1.2-3. Domain:  $0 \leq r \leq a$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq b$ . First boundary value problem.

A circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$w = f(\varphi, z) \quad \text{at} \quad r = a, \quad w = g_1(r, \varphi) \quad \text{at} \quad z = 0, \quad w = g_2(r, \varphi) \quad \text{at} \quad z = b.$$

**Solution:**

$$\begin{aligned} w(r, \varphi, z) = & \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{I_n\left(\frac{\pi m r}{b}\right)}{I_n\left(\frac{\pi m a}{b}\right)} (A_{nm} \cos n\varphi + B_{nm} \sin n\varphi) \sin\left(\frac{\pi m z}{b}\right) \\ & + \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} J_n\left(\frac{\mu_{mn} r}{a}\right) (C_{nm}^{(1)} \cos n\varphi + D_{nm}^{(1)} \sin n\varphi) \frac{\sinh\left(\frac{\mu_{mn}(b-z)}{a}\right)}{\sinh\left(\frac{\mu_{mn} b}{a}\right)} \\ & + \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} J_n\left(\frac{\mu_{mn} r}{a}\right) (C_{nm}^{(2)} \cos n\varphi + D_{nm}^{(2)} \sin n\varphi) \frac{\sinh\left(\frac{\mu_{mn} z}{a}\right)}{\sinh\left(\frac{\mu_{mn} b}{a}\right)}, \end{aligned}$$

where the  $J_n(r)$  are the Bessel functions, the  $I_n(r)$  are the modified Bessel functions, and  $\mu_{mn}$  is the  $m$ th root of the equation  $J_n(\mu) = 0$ . The coefficients  $A_{nm}$ ,  $B_{nm}$ ,  $C_{nm}^{(i)}$ , and  $D_{nm}^{(i)}$  are defined by

$$\begin{aligned} A_{nm} &= \frac{\varepsilon_n}{\pi b} \int_0^{2\pi} \int_0^b f(\varphi, z) \cos(n\varphi) \sin\left(\frac{\pi m z}{b}\right) d\varphi dz, \\ B_{nm} &= \frac{2}{\pi b} \int_0^{2\pi} \int_0^b f(\varphi, z) \sin(n\varphi) \sin\left(\frac{\pi m z}{b}\right) d\varphi dz, \\ C_{nm}^{(i)} &= \frac{\varepsilon_n}{\pi a^2 [J'_n(\mu_{mn})]^2} \int_0^{2\pi} \int_0^a g_i(r, \varphi) \cos(n\varphi) J_n\left(\frac{\mu_{mn} r}{a}\right) r dr d\varphi, \\ D_{nm}^{(i)} &= \frac{2}{\pi a^2 [J'_n(\mu_{mn})]^2} \int_0^{2\pi} \int_0^a g_i(r, \varphi) \sin(n\varphi) J_n\left(\frac{\mu_{mn} r}{a}\right) r dr d\varphi, \\ \varepsilon_n &= \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad i = 1, 2. \end{aligned}$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

► *For the solution of other boundary value problems for the three-dimensional Laplace equation in the cylindrical coordinate system, see Subsection 8.2.3 for  $\Phi \equiv 0$ .*



### 8.1.3. Problems in Spherical Coordinates

The three-dimensional Laplace equation in the spherical coordinate system is written as

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} = 0, \quad r = \sqrt{x^2 + y^2 + z^2}.$$

8.1.3-1. Particular solutions:

$$\begin{aligned} w(r) &= A + \frac{B}{r}, \\ w(r, \theta) &= \left( Ar^n + \frac{B}{r^{n+1}} \right) P_n(\cos \theta), \\ w(r, \theta, \varphi) &= \left( Ar^n + \frac{B}{r^{n+1}} \right) P_n^m(\cos \theta) (C \cos m\varphi + D \sin m\varphi), \end{aligned}$$

where  $n = 0, 1, 2, \dots$ ;  $m = 0, 1, 2, \dots, n$ ;  $A, B, C, D$  are arbitrary constants; the  $P_n(\xi)$  are the Legendre polynomials; and the  $P_n^m(\xi)$  are the associated Legendre functions that are expressed as

$$P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n, \quad P_n^m(x) = (1 - x^2)^{m/2} \frac{d^m}{dx^m} P_n(x).$$

8.1.3-2. Domain:  $0 \leq r \leq R$  or  $R \leq r < \infty$ . First boundary value problem.

A boundary condition at the sphere surface is prescribed:

$$w = f(\theta, \varphi) \quad \text{at} \quad r = R.$$

1°. Solution of the inner problem (for  $r \leq R$ ):

$$\begin{aligned} w(r, \theta, \varphi) &= \frac{R}{4\pi} \int_0^{2\pi} \int_0^\pi f(\theta_0, \varphi_0) \frac{R^2 - r^2}{(r^2 - 2Rr \cos \gamma + R^2)^{3/2}} \sin \theta_0 \, d\theta_0 \, d\varphi_0, \\ \cos \gamma &= \cos \theta \cos \theta_0 + \sin \theta \sin \theta_0 \cos(\varphi - \varphi_0). \end{aligned}$$

This formula is conventionally called the Poisson integral for a sphere.

Series solution:

$$w(r, \theta, \varphi) = \sum_{n=0}^{\infty} \left( \frac{r}{R} \right)^n Y_n(\theta, \varphi), \quad Y_n(\theta, \varphi) = \sum_{m=0}^n (A_{nm} \cos m\varphi + B_{nm} \sin m\varphi) P_n^m(\cos \theta),$$

where

$$\begin{aligned} A_{00} &= \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi f(\theta, \varphi) \sin \theta \, d\theta \, d\varphi, \\ A_{nm} &= \frac{(2n+1)(n-m)!}{2\pi(n+m)!} \int_0^{2\pi} \int_0^\pi f(\theta, \varphi) P_n^m(\cos \theta) \cos m\varphi \sin \theta \, d\theta \, d\varphi, \\ B_{nm} &= \frac{(2n+1)(n-m)!}{2\pi(n+m)!} \int_0^{2\pi} \int_0^\pi f(\theta, \varphi) P_n^m(\cos \theta) \sin m\varphi \sin \theta \, d\theta \, d\varphi. \end{aligned}$$

2°. Solution of the outer problem (for  $r \geq R$ ):

$$w(r, \theta, \varphi) = \frac{R}{4\pi} \int_0^{2\pi} \int_0^\pi f(\theta_0, \varphi_0) \frac{r^2 - R^2}{(r^2 - 2Rr \cos \gamma + R^2)^{3/2}} \sin \theta_0 \, d\theta_0 \, d\varphi_0,$$

where  $\cos \gamma$  is expressed in the same way as in the inner problem.

Series solution:

$$w(r, \theta, \varphi) = \sum_{n=0}^{\infty} \left( \frac{R}{r} \right)^{n+1} Y_n(\theta, \varphi),$$

$$Y_n(\theta, \varphi) = \sum_{m=0}^n (A_{nm} \cos m\varphi + B_{nm} \sin m\varphi) P_n^m(\cos \theta),$$

where the coefficients  $A_{nm}$  and  $B_{nm}$  are defined by the same relations as in the inner problem.

⊙ *References:* G. N. Polozhii (1964), V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), A. N. Tikhonov and A. A. Samarskii (1990).

8.1.3-3. Domain:  $0 \leq r \leq R$  or  $R \leq r < \infty$ . Second boundary value problem.

A boundary condition at the sphere surface is prescribed:

$$\frac{\partial w}{\partial r} = f(\theta, \varphi) \quad \text{at} \quad r = R.$$

The function  $f(\theta, \varphi)$  must satisfy the solvability condition

$$\int_0^{2\pi} \int_0^\pi f(\theta, \varphi) \sin \theta \, d\theta \, d\varphi = 0.$$

1°. Solution of the inner problem (for  $r \leq R$ ):

$$w(r, \theta, \varphi) = \frac{R}{4\pi} \int_0^{2\pi} \int_0^\pi f(\theta_0, \varphi_0) \left[ \frac{1}{R} \ln(R + r_1 - r \cos \gamma) - \frac{2}{r_1} \right] \sin \theta_0 \, d\theta_0 \, d\varphi_0,$$

$$r_1 = \sqrt{r^2 - 2Rr \cos \gamma + R^2}, \quad \cos \gamma = \cos \theta \cos \theta_0 + \sin \theta \sin \theta_0 \cos(\varphi - \varphi_0).$$

Series solution:

$$w(r, \theta, \varphi) = \sum_{n=1}^{\infty} \sum_{m=0}^n \frac{R}{n} \left( \frac{r}{R} \right)^n (A_{nm} \cos m\varphi + B_{nm} \sin m\varphi) P_n^m(\cos \theta) + C,$$

where the coefficients  $A_{nm}$  and  $B_{nm}$  are expressed in the same way as in the inner first boundary value problem (see Paragraph 8.1.3-2), and  $C$  is an arbitrary constant.

2°. Solution of the outer problem (for  $r \geq R$ ):

$$w(r, \theta, \varphi) = -\frac{R}{4\pi} \int_0^{2\pi} \int_0^\pi f(\theta_0, \varphi_0) \left[ \frac{1}{R} \ln \frac{R + r_1 - r \cos \gamma}{r(1 - \cos \gamma)} - \frac{2}{r_1} \right] \sin \theta_0 \, d\theta_0 \, d\varphi_0,$$

$$r_1 = \sqrt{r^2 - 2Rr \cos \gamma + R^2}, \quad \cos \gamma = \cos \theta \cos \theta_0 + \sin \theta \sin \theta_0 \cos(\varphi - \varphi_0).$$

Series solution:

$$w(r, \theta, \varphi) = -\sum_{n=0}^{\infty} \sum_{m=0}^n \frac{R}{n+1} \left( \frac{R}{r} \right)^{n+1} (A_{nm} \cos m\varphi + B_{nm} \sin m\varphi) P_n^m(\cos \theta) + C,$$

where the coefficients  $A_{nm}$  and  $B_{nm}$  are expressed in the same way as in the inner first boundary value problem, and  $C$  is an arbitrary constant.

3°. Outer boundary value problems where unbounded solutions as  $r \rightarrow \infty$  are sought are also encountered in applications.

**Example.** A potential translational flow of an ideal incompressible fluid about a sphere of radius  $R$  is governed by the Laplace equation with the boundary conditions:

$$\partial_r w = 0 \quad \text{at} \quad r = R, \quad |w - Ur \cos \theta| \rightarrow 0 \quad \text{as} \quad r \rightarrow \infty,$$

where  $w$  is the potential,  $U$  the unperturbed flow velocity at infinity; the fluid velocity is expressed in terms of the potential as  $\mathbf{v} = \nabla \varphi$ .

Solution:

$$w = Ur \left( 1 + \frac{R^3}{2r^3} \right) \cos \theta.$$

This solution is a special case of the second formula from Paragraph for 8.1.3-1 for  $n = 1$ .

⊙ *References:* G. N. Polozhii (1964), V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), L. G. Loitsyanskii (1996).

► *For the solution of other boundary value problems for the three-dimensional Laplace equation in the spherical coordinate system, see Subsection 8.2.4 for  $\Phi \equiv 0$ .*

## 8.1.4. Other Orthogonal Curvilinear Systems of Coordinates

The three-dimensional Laplace equation admits separation of variables in the eleven orthogonal coordinate systems that are listed in Table 26.

For the general ellipsoidal and conical coordinate systems, the functions  $f$ ,  $g$ , and  $h$  are determined by Lamé equations that involve the Jacobian elliptic function  $\text{sn } z = \text{sn}(z, k)$ . The solutions of these equations under some conditions can be represented in the form of finite series called Lamé polynomials. For details about the Lamé equation and its solutions, see Whittaker and Watson (1963), Bateman and Erdélyi (1955), Arscott (1964), and Miller, Jr. (1977).

There are also coordinate systems that allow the so-called  $\mathcal{R}$ -separation of variables of the three-dimensional Laplace equation. Such solutions in the new coordinate system,  $\mu, \nu, \rho$ , can be represented in the form  $w = \sqrt{\mathcal{R}(\mu, \nu, \rho)} f(\mu)g(\nu)h(\rho)$ . Coordinates that allow the  $\mathcal{R}$ -separation of variables are listed in Table 27.

Only the bicylindrical and toroidal coordinate systems are fairly widely used in applications. In three subsequent coordinate systems, the functions  $f = f(\mu)$  and  $g = g(\rho)$  are determined by identical equations. With the change of variables  $\mu = \text{sn}^2(\alpha, k)$ ,  $\rho = \text{sn}^2(\beta, k)$ , where  $k = a^{-1/2}$ , these equations are reduced to Lamé equations ( $\alpha$  and  $\beta$  are the new independent variables).

⊙ *References for Subsection 8.1.4:* M. Bôcher (1894), F. M. Morse and H. Feshbach (1953, Vols. 1–2), N. N. Lebedev, I. P. Skal'skaya, and Ya. S. Uflyand (1955), P. Moon and D. Spencer (1961), A. Makarov, J. Smorodinsky, K. Valiev, and P. Winternitz (1967), W. Miller, Jr. (1977).

## 8.2. Poisson Equation $\Delta_3 w + \Phi(\mathbf{x}) = 0$

### 8.2.1. Preliminary Remarks. Solution Structure

Like the three-dimensional Laplace equation, the three-dimensional Poisson equation is often encountered in heat and mass transfer theory, fluid mechanics, elasticity, electrostatics, and other areas of mechanics and physics. In particular, the Poisson equation describes stationary temperature distribution in the presence of thermal sources or sinks in the domain under consideration.

The Laplace equation is a special case of the Poisson equation with  $\Phi \equiv 0$ .

Throughout this section, we consider a three-dimensional bounded domain  $V$  with a sufficiently smooth boundary  $S$ . We assume that  $\mathbf{r} \in V$  and  $\boldsymbol{\rho} \in V$ , where  $\mathbf{r} = \{x, y, z\}$  and  $\boldsymbol{\rho} = \{\xi, \eta, \zeta\}$ .

TABLE 26

Orthogonal coordinates  $\bar{x}, \bar{y}, \bar{z}$  that allow separable solutions of the form  $w = f(\bar{x})g(\bar{y})h(\bar{z})$  for the three-dimensional Laplace equation  $\Delta_3 w = 0$

| Coordinates                                | Transformations                                                                                                                              | Particular solutions (or equations for $f, g, h$ )                                                                                                                                                                                                                                                                                                                     |
|--------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cartesian<br>$x, y, z$                     | $x = x,$<br>$y = y,$<br>$z = z$                                                                                                              | $w = \cos(k_1 x + s_1) \cos(k_2 y + s_2) \cosh(k_3 z + s_3),$<br>where $k_1^2 + k_2^2 = k_3^2;$<br>see also Paragraph 8.1.1-1                                                                                                                                                                                                                                          |
| Cylindrical<br>$r, \varphi, z$             | $x = r \cos \varphi,$<br>$y = r \sin \varphi,$<br>$z = z$                                                                                    | $w = [AJ_n(kr) + BY_n(kr)] \cos(n\varphi + c) \exp(\pm kz),$<br>$J_n(z)$ and $Y_n(z)$ are the Bessel functions;<br>see also Paragraph 8.1.2-1                                                                                                                                                                                                                          |
| Parabolic<br>cylindrical<br>$\xi, \eta, z$ | $x = \frac{1}{2}(\xi^2 - \eta^2),$<br>$y = \xi\eta,$<br>$z = z$                                                                              | $w = D_{\mu-1/2}(\pm\sigma\xi)D_{-\mu-1/2}(\pm\sigma\eta) \cos(kz + s),$<br>where $\sigma = \sqrt{2k}, D_\mu(z)$ is the parabolic<br>cylinder function                                                                                                                                                                                                                 |
| Elliptic<br>cylindrical<br>$u, v, z$       | $x = a \cosh u \cos v,$<br>$y = a \sinh u \sin v,$<br>$z = z$                                                                                | $w = \begin{cases} \text{Ce}_n(u, -q) \text{ce}_n(v, -q) \cos(kz + s), \\ \text{Se}_n(u, -q) \text{se}_n(v, -q) \cos(kz + s), \end{cases}$<br>$\text{Ce}_n$ and $\text{Se}_n$ are the modified Mathieu functions,<br>$\text{ce}_n$ and $\text{se}_n$ are the Mathieu functions, $q = \frac{1}{4}a^2 k^2$                                                               |
| Spherical<br>$r, \theta, \varphi$          | $x = r \sin \theta \cos \varphi,$<br>$y = r \sin \theta \sin \varphi,$<br>$z = r \cos \theta$                                                | $w = (Ar^n + Br^{-n-1})P_n^k(\cos \theta) \cos(k\varphi + s),$<br>$P_n^k(\xi)$ are the associated Legendre functions,<br>see also Paragraph 8.1.3-1                                                                                                                                                                                                                    |
| Prolate<br>spheroidal<br>$u, v, \varphi$   | $x = a \sinh u \sin v \cos \varphi,$<br>$y = a \sinh u \sin v \sin \varphi,$<br>$z = a \cosh u \cos v$                                       | $w = P_n^k(\cosh u)P_n^k(\cos v) \cos(k\varphi + s),$<br>$P_n^k(\xi)$ are the associated Legendre functions                                                                                                                                                                                                                                                            |
| Oblate<br>spheroidal<br>$u, v, \varphi$    | $x = a \cosh u \sin v \cos \varphi,$<br>$y = a \cosh u \sin v \sin \varphi,$<br>$z = a \sinh u \cos v$                                       | $w = P_n^k(-i \sinh u)P_n^k(\cos v) \cos(k\varphi + s),$<br>$P_n^k(\xi)$ are the associated Legendre functions                                                                                                                                                                                                                                                         |
| Parabolic<br>$\xi, \eta, \varphi$          | $x = a\xi\eta \cos \varphi,$<br>$y = a\xi\eta \sin \varphi,$<br>$z = \frac{1}{2}a(\xi^2 - \eta^2)$                                           | $w = I_{\pm k}(\beta\xi)J_{\pm k}(\beta\eta) \cos(k\varphi + s),$<br>$J_k(z)$ are the Bessel functions,<br>$I_k(z)$ are the modified Bessel functions                                                                                                                                                                                                                  |
| Paraboloidal<br>$u, v, \varphi$            | $x = 2a \cosh u \cos v \sinh \varphi,$<br>$y = 2a \sinh u \sin v \cosh \varphi,$<br>$z = \frac{1}{2}a(\cosh 2u + \cos 2v - \cosh 2\varphi)$  | $w = \begin{cases} \text{Ce}_n(u, -b) \text{ce}_n(v, -b) \text{Ce}_n(\varphi + i\pi/2, -b), \\ \text{Se}_n(u, -b) \text{se}_n(v, -b) \text{Se}_n(\varphi + i\pi/2, -b), \end{cases}$<br>$b = \frac{1}{2}a\beta;$ $\text{ce}_n$ and $\text{se}_n$ are the Mathieu functions,<br>$\text{Ce}_n$ and $\text{Se}_n$ are the modified Mathieu functions                      |
| General<br>ellipsoidal<br>$\mu, \nu, \rho$ | $x = \sqrt{\frac{(\mu-a)(\nu-a)(\rho-a)}{a(a-1)}},$<br>$y = \sqrt{\frac{(\mu-1)(\nu-1)(\rho-1)}{1-a}},$<br>$z = \sqrt{\frac{\mu\nu\rho}{a}}$ | $f''_{\xi\xi} + (\beta_2 + \beta_1 \text{sn}^2 \xi)f = 0$ (Lamé equation),<br>$g''_{\eta\eta} + (\beta_2 + \beta_1 \text{sn}^2 \eta)g = 0$ (Lamé equation),<br>$h''_{\zeta\zeta} + (\beta_2 + \beta_1 \text{sn}^2 \zeta)h = 0$ (Lamé equation),<br>$\mu = \text{sn}^2(\xi, k), \nu = \text{sn}^2(\eta, k), \rho = \text{sn}^2(\zeta, k), k = a^{-1/2}$                 |
| Conical<br>$\rho, \mu, \nu$                | $x = \rho \sqrt{\frac{a(\mu-1)(\nu-1)}{1-a}},$<br>$y = \rho \sqrt{\frac{a(\mu-1)(\nu-1)}{a-1}},$<br>$z = \rho \sqrt{a\mu\nu}$                | $f(\rho) = A\rho^n + B\rho^{-n-1}, \quad n = 0, 1, \dots,$<br>$g''_{\xi\xi} + [\beta - n(n+1)k^2 \text{sn}^2 \xi]g = 0$ (Lamé equation),<br>$h''_{\eta\eta} + [\beta - n(n+1)k^2 \text{sn}^2 \eta]h = 0$ (Lamé equation),<br>where $\mu = \text{sn}^2(\xi, k), \nu = \text{sn}^2(\eta, k), k = a^{1/2},$<br>$g$ and $h$ are expressed in terms of the Lamé polynomials |

TABLE 27

Coordinates  $\bar{x}, \bar{y}, \bar{z}$  that allow  $\mathcal{R}$ -separated solutions of the form  
 $w = \sqrt{\mathcal{R}(\bar{x}, \bar{y}, \bar{z})} f(\bar{x})g(\bar{y})h(\bar{z})$  for the three-dimensional Laplace equation  $\Delta_3 w = 0$

| New coordinates,<br>function $\mathcal{R}$                                                                                                                                                          | Transformations<br>of coordinates                                                                                                                                                                                                                                           | Functions $f, g, h$<br>(equations for $f, g, h$ )                                                                                                                                                                                                                                           |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Bicylindrical<br>coordinates $\alpha, \beta, \varphi$ ,<br>$\mathcal{R} = \cosh \beta - \cos \alpha$                                                                                                | $x = c\mathcal{R}^{-1} \sin \alpha \cos \varphi$ ,<br>$y = c\mathcal{R}^{-1} \sin \alpha \sin \varphi$ ,<br>$z = c\mathcal{R}^{-1} \sinh \beta$ ;<br>$0 \leq \alpha \leq \pi$ , $\beta$ is any,<br>$0 \leq \varphi < 2\pi$                                                  | $f(\alpha) = A_1 P_n^m(\cos \alpha) + A_2 Q_n^m(\cos \alpha)$ ,<br>$g(\beta) = B_1 \cosh[(n + \frac{1}{2})\beta] + B_2 \sinh[(n + \frac{1}{2})\beta]$ ,<br>$h(\varphi) = C_1 \cos(m\varphi) + C_2 \sin(m\varphi)$ ,<br>$n = 0, 1, 2, \dots$ ; $m = 0, 1, 2, \dots$                          |
| Toroidal<br>coordinates $\alpha, \beta, \varphi$ ,<br>$\mathcal{R} = \cosh \alpha - \cos \beta$                                                                                                     | $x = c\mathcal{R}^{-1} \sinh \alpha \cos \varphi$ ,<br>$y = c\mathcal{R}^{-1} \sinh \alpha \sin \varphi$ ,<br>$z = c\mathcal{R}^{-1} \sin \beta$ ;<br>$\alpha \geq 0$ , $-\pi \leq \beta \leq \pi$ , $0 \leq \varphi < 2\pi$                                                | $f(\alpha) = A_1 P_{n-1/2}^m(\cosh \alpha) + A_2 Q_{n-1/2}^m(\cosh \alpha)$ ,<br>$g(\beta) = B_1 \cos(n\beta) + B_2 \sin(n\beta)$ ,<br>$h(\varphi) = C_1 \cos(m\varphi) + C_2 \sin(m\varphi)$ ,<br>$n = 0, 1, 2, \dots$ ; $m = 0, 1, 2, \dots$                                              |
| Coordinates $\mu, \rho, \varphi$ ,<br>$\mathcal{R} = \sqrt{\frac{(\mu-a)(a-\rho)}{a(a-1)}}$<br>$-\sqrt{\frac{(\mu-1)(1-\rho)}{a-1}}$                                                                | $x = \mathcal{R}^{-1} \cos \varphi$ ,<br>$y = \mathcal{R}^{-1} \sin \varphi$ ,<br>$z = \mathcal{R}^{-1} \sqrt{-\mu\rho/a}$ ;<br>$\mu > a > 1$ , $\rho < 0$ , $0 \leq \varphi < 2\pi$                                                                                        | $\sqrt{U(\mu)} [\sqrt{U(\mu)} f']' + [(\frac{1}{4} - n^2)\mu - \lambda] f = 0$ ,<br>$\sqrt{U(\rho)} [\sqrt{U(\rho)} g']' + [(\frac{1}{4} - n^2)\rho - \lambda] g = 0$ ,<br>$h(\varphi) = C_1 \cos(n\varphi) + C_2 \sin(n\varphi)$ ,<br>$U(t) = 4t(t-1)(t-a)$                                |
| Coordinates $\mu, \rho, \varphi$ ,<br>$\mathcal{R} = \sqrt{\frac{\mu\rho}{a}}$<br>$+\sqrt{\frac{(\mu-1)(\rho-1)}{a-1}}$                                                                             | $x = \mathcal{R}^{-1} \cos \varphi$ ,<br>$y = \mathcal{R}^{-1} \sin \varphi$ ,<br>$z = \mathcal{R}^{-1} \sqrt{\frac{(\mu-a)(a-\rho)}{a(a-1)}}$ ;<br>$1 < \rho < a < \mu$ , $0 \leq \varphi < 2\pi$                                                                          | $\sqrt{U(\mu)} [\sqrt{U(\mu)} f']' + [(\frac{1}{4} - n^2)\mu - \lambda] f = 0$ ,<br>$\sqrt{U(\rho)} [\sqrt{U(\rho)} g']' + [(\frac{1}{4} - n^2)\rho - \lambda] g = 0$ ,<br>$h(\varphi) = C_1 \cos(n\varphi) + C_2 \sin(n\varphi)$ ,<br>$U(t) = 4t(t-1)(t-a)$                                |
| Coordinates $\mu, \rho, \varphi$ ,<br>$\mathcal{R} = 2 \operatorname{Re} \sqrt{\frac{i(\mu-a)(\rho-a)}{a(a-b)}}$ ,<br>$a = \bar{b} = \alpha + i\beta$ ,<br>$\alpha, \beta$ are real numbers         | $x = \mathcal{R}^{-1} \cos \varphi$ ,<br>$y = \mathcal{R}^{-1} \sin \varphi$ ,<br>$z = \mathcal{R}^{-1} \sqrt{-\mu\rho/(ab)}$ ;<br>$\mu > 0$ , $\rho < 0$ , $0 \leq \varphi < 2\pi$                                                                                         | $\sqrt{U(\mu)} [\sqrt{U(\mu)} f']' + [(\frac{1}{4} - n^2)\mu - \lambda] f = 0$ ,<br>$\sqrt{U(\rho)} [\sqrt{U(\rho)} g']' + [(\frac{1}{4} - n^2)\rho - \lambda] g = 0$ ,<br>$h(\varphi) = C_1 \cos(n\varphi) + C_2 \sin(n\varphi)$ ,<br>$U(t) = 4t(t-a)(t-b)$                                |
| Coordinates $\mu, \nu, \rho$ ,<br>$\mathcal{R} = 1 + \sqrt{\frac{\mu\nu\rho}{ab}}$                                                                                                                  | $x = \mathcal{R}^{-1} \sqrt{\frac{(\mu-a)(\nu-a)(\rho-a)}{(b-a)(a-1)a}}$ ,<br>$y = \mathcal{R}^{-1} \sqrt{\frac{(\mu-b)(\nu-b)(\rho-b)}{(a-b)(b-1)b}}$ ,<br>$z = \mathcal{R}^{-1} \sqrt{\frac{(\mu-1)(\nu-1)(\rho-1)}{(a-1)(b-1)}}$ ;<br>$0 < \rho < 1 < \nu < b < \mu < a$ | $\sqrt{U(\mu)} [\sqrt{U(\mu)} f']' - (3\mu^2 + \lambda_1\mu + \lambda_2)f = 0$ ,<br>$\sqrt{U(\nu)} [\sqrt{U(\nu)} g']' - (3\nu^2 + \lambda_1\nu + \lambda_2)g = 0$ ,<br>$\sqrt{U(\rho)} [\sqrt{U(\rho)} h']' - (3\rho^2 + \lambda_1\rho + \lambda_2)h = 0$ ,<br>$U(t) = 16t(t-1)(t-a)(t-b)$ |
| Coordinates $\mu, \nu, \rho$ ,<br>$\mathcal{R} = 2 \operatorname{Re} \sqrt{\frac{(\mu-a)(\nu-a)(\rho-a)}{ia(a-1)(a-b)}}$ ,<br>$a = \bar{b} = \alpha + i\beta$ ,<br>$\alpha, \beta$ are real numbers | $x = \mathcal{R}^{-1} \sqrt{\frac{(\mu-1)(\nu-1)(\rho-1)}{(a-1)(b-1)}}$ ,<br>$y = \mathcal{R}^{-1} \sqrt{-\frac{\mu\nu\rho}{ab}}$ ,<br>$z = \mathcal{R}^{-1}$ ;<br>$\rho < 0 < \mu < 1 < \nu$                                                                               | $\sqrt{U(\mu)} [\sqrt{U(\mu)} f']' - (3\mu^2 + \lambda_1\mu + \lambda_2)f = 0$ ,<br>$\sqrt{U(\nu)} [\sqrt{U(\nu)} g']' - (3\nu^2 + \lambda_1\nu + \lambda_2)g = 0$ ,<br>$\sqrt{U(\rho)} [\sqrt{U(\rho)} h']' - (3\rho^2 + \lambda_1\rho + \lambda_2)h = 0$ ,<br>$U(t) = 16t(t-1)(t-a)(t-b)$ |

### 8.2.1-1. First boundary value problem.

The solution of the first boundary value problem for the Poisson equation

$$\Delta_3 w + \Phi(\mathbf{r}) = 0 \quad (1)$$

in a domain  $V$  with the nonhomogeneous boundary condition

$$w = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in S$$

TABLE 28

The volume elements and distances occurring in relations (2) and (5) in some coordinate systems. In all cases,  $\rho = \{\xi, \eta, \zeta\}$

| Coordinate system                                  | Volume element, $dV_\rho$           | Gradient, $\nabla_\rho u$<br>( $ \mathbf{i}_\xi  =  \mathbf{i}_\eta  =  \mathbf{i}_\zeta  = 1$ )                                                                                               | Distance,<br>$d =  \mathbf{r} - \rho $                                                                                                      |
|----------------------------------------------------|-------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------|
| Cartesian<br>$\mathbf{r} = \{x, y, z\}$            | $d\xi d\eta d\zeta$                 | $\mathbf{i}_\xi \frac{\partial u}{\partial \xi} + \mathbf{i}_\eta \frac{\partial u}{\partial \eta} + \mathbf{i}_\zeta \frac{\partial u}{\partial \zeta}$                                       | $d = \sqrt{(x - \xi)^2 + (y - \eta)^2 + (z - \zeta)^2}$                                                                                     |
| Cylindrical<br>$\mathbf{r} = \{r, \varphi, z\}$    | $\xi d\xi d\eta d\zeta$             | $\mathbf{i}_\xi \frac{\partial u}{\partial \xi} + \mathbf{i}_\eta \frac{1}{\xi} \frac{\partial u}{\partial \eta} + \mathbf{i}_\zeta \frac{\partial u}{\partial \zeta}$                         | $d = \sqrt{r^2 + \xi^2 - 2r\xi \cos(\varphi - \eta) + (z - \zeta)^2}$                                                                       |
| Spherical<br>$\mathbf{r} = \{r, \theta, \varphi\}$ | $\xi^2 \sin \eta d\xi d\eta d\zeta$ | $\mathbf{i}_\xi \frac{\partial u}{\partial \xi} + \mathbf{i}_\eta \frac{1}{\xi} \frac{\partial u}{\partial \eta} + \mathbf{i}_\zeta \frac{1}{\xi \sin \eta} \frac{\partial u}{\partial \zeta}$ | $d = \sqrt{r^2 + \xi^2 - 2r\xi \cos \gamma}$ , where<br>$\cos \gamma = \cos \theta \cos \eta + \sin \theta \sin \eta \cos(\varphi - \zeta)$ |

can be represented in the form

$$w(\mathbf{r}) = \int_V \Phi(\rho) G(\mathbf{r}, \rho) dV_\rho - \int_S f(\rho) \frac{\partial G}{\partial N_\rho} dS_\rho. \quad (2)$$

Here,  $G(\mathbf{r}, \rho)$  is the Green's function of the first boundary value problem,  $\frac{\partial G}{\partial N_\rho}$  is the derivative of the Green's function with respect to  $\xi, \eta, \zeta$  along the outward normal  $\mathbf{N}$  the boundary  $S$  of the domain  $V$ . Integration is everywhere with respect to  $\xi, \eta, \zeta$ .

The volume elements in solution (2) for basic coordinate systems are presented in Table 28. In addition, the expressions of the gradients are given, which enable one to find the derivative along the normal in accordance with the formula  $\frac{\partial G}{\partial N_\rho} = (\mathbf{N} \cdot \nabla_\rho G)$ .

The Green's function  $G = G(\mathbf{r}, \rho)$  of the first boundary value problem is determined by the following conditions:

- 1°. The function  $G$  satisfies the Laplace equation with respect to  $x, y, z$  in the domain  $V$  everywhere except for the point  $(\xi, \eta, \zeta)$ , at which it can have a singularity of the form  $\frac{1}{4\pi} \frac{1}{|\mathbf{r} - \rho|}$ .
- 2°. The function  $G$ , with respect to  $x, y, z$ , satisfies the homogeneous boundary condition of the first kind at the boundary, i.e., the condition  $G|_S = 0$ .

The Green's function can be represented as

$$G(\mathbf{r}, \rho) = \frac{1}{4\pi} \frac{1}{|\mathbf{r} - \rho|} + u, \quad (3)$$

where the auxiliary function  $u = u(\mathbf{r}, \rho)$  is determined by solving the first boundary value problem for the Laplace equation  $\Delta_3 u = 0$  with the boundary condition  $u|_S = -\frac{1}{4\pi} \frac{1}{|\mathbf{r} - \rho|}$ ; the vector quantity  $\rho$  in this problem is treated as a three-dimensional free parameter.

The Green's function possesses the symmetry property with respect to their arguments:  $G(\mathbf{r}, \rho) = G(\rho, \mathbf{r})$ .

The construction of Green's functions is discussed in Paragraphs 8.3.1-4 and 8.3.1-6 through 8.3.1-8 for  $\lambda = 0$ .

*Remark 1.* For outer first boundary value problems for the Laplace equation, the following condition is usually set at infinity:  $|w| < A/|\mathbf{r}|$  ( $|\mathbf{r}| \rightarrow \infty$ ,  $A = \text{const}$ ).

#### 8.2.1-2. Second boundary value problem.

The second boundary value problem for the Poisson equation (1) is characterized by the boundary condition

$$\frac{\partial w}{\partial N} = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in S.$$

Necessary condition solvability of the inner problem:

$$\int_V \Phi(\mathbf{r}) d\mathbf{r} + \int_S f(\mathbf{r}) dS = 0. \quad (4)$$

The solution of the second boundary value problem can be written as

$$w(\mathbf{r}) = \int_V \Phi(\rho) G(\mathbf{r}, \rho) dV_\rho + \int_S f(\rho) G(\mathbf{r}, \rho) dS_\rho + C, \quad (5)$$

where  $C$  is an arbitrary constant, provided that the solvability condition is met.

The Green's function  $G = G(\mathbf{r}, \rho)$  of the second boundary value problem is determined by the following conditions:

1°. The function  $G$  satisfies the Laplace equation with respect to  $x, y, z$  in the domain  $V$  everywhere except for the point  $(\xi, \eta, \zeta)$  at which it has a singularity of the form  $\frac{1}{4\pi} \frac{1}{|\mathbf{r}-\rho|}$ .

2°. The function  $G$ , with respect to  $x, y, z$ , satisfies the homogeneous condition of the second kind at the boundary, i.e., the condition

$$\left. \frac{\partial G}{\partial N} \right|_S = \frac{1}{S_0},$$

where  $S_0$  is the area of the surface  $S$ .

The Green's function is unique up to an additive constant.

*Remark 2.* The Green's function cannot be identified with condition 1° and the homogeneous boundary condition  $\left. \frac{\partial G}{\partial N} \right|_S = 0$ ; this problem for  $G$  has no solution, because, on representing  $G$  in the form (3), for  $u$  we obtain a problem with a nonhomogeneous boundary condition of the second kind, for which the solvability condition (2) is not met.

*Remark 3.* Condition (4) is not extended to the outer second boundary value problem (for infinite domain).

### 8.2.1-3. Third boundary value problem.

The solution of the third boundary value problem for the Poisson equation (1) in a bounded domain  $V$  with the nonhomogeneous boundary condition

$$\frac{\partial w}{\partial N} + kw = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in S$$

is given by relation (5) with  $C = 0$ , where  $G = G(\mathbf{r}, \rho)$  is the Green's function of the third boundary value problem; the Green's function is determined by the following conditions:

1°. The function  $G$  satisfies the Laplace equation with respect to  $x, y, z$  in  $V$  everywhere except for the point  $(\xi, \eta, \zeta)$  at which it has a singularity of the form  $\frac{1}{4\pi} \frac{1}{|\mathbf{r}-\rho|}$ .

2°. The function  $G$ , with respect to  $x, y, z$ , satisfies the homogeneous boundary condition of the third kind at the boundary, i.e., the condition  $\left[ \frac{\partial G}{\partial N} + kG \right]_S = 0$ .

The Green's function can be represented in the form (3), where the auxiliary function  $u$  is determined by solving the corresponding third boundary value problem for the Laplace equation  $\Delta_3 u = 0$ .

The construction of Green's functions is discussed in Paragraphs 8.3.1-4 and 8.3.1-6 through 8.3.1-8 for  $\lambda = 0$ .

⊙ *References for Subsection 8.2.1:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970).

## 8.2.2. Problems in Cartesian Coordinates

The three-dimensional Poisson equation in the rectangular Cartesian system of coordinates has the form

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} + \Phi(x, y, z) = 0.$$

8.2.2-1. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ .

Solution:

$$w(x, y, z) = \frac{1}{4\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{\Phi(\xi, \eta, \zeta) d\xi d\eta d\zeta}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}}.$$

⊙ *Reference:* R. Courant and D. Hilbert (1989).

8.2.2-2. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z < \infty$ . First boundary value problem.

A half-space is considered. A boundary condition is prescribed:

$$w = f(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{z f(\xi, \eta) d\xi d\eta}{[(x-\xi)^2 + (y-\eta)^2 + z^2]^{3/2}} \\ & + \frac{1}{4\pi} \int_0^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left( \frac{1}{R_-} - \frac{1}{R_+} \right) \Phi(\xi, \eta, \zeta) d\xi d\eta d\zeta, \end{aligned}$$

where

$$R_- = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}, \quad R_+ = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2}.$$

⊙ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

8.2.2-3. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z < \infty$ . Third boundary value problem.

A half-space is considered. A boundary condition is prescribed:

$$\partial_z w - kw = f(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(x, y, z) = & - \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta) G(x, y, z, \xi, \eta, 0) d\xi d\eta \\ & + \int_0^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) = & \frac{1}{4\pi} \left[ \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}} + \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2}} \right. \\ & \left. - 2k \int_0^{\infty} \frac{\exp(-ks) ds}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta+s)^2}} \right]. \end{aligned}$$



**8.2.2-4. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . First boundary value problem.**

A dihedral angle is considered. Boundary conditions are prescribed:

$$w = f_1(x, z) \quad \text{at} \quad y = 0, \quad w = f_2(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_0^\infty \int_{-\infty}^\infty f_1(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\xi d\zeta \\ & + \int_0^\infty \int_{-\infty}^\infty f_2(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta \\ & + \int_0^\infty \int_0^\infty \int_{-\infty}^\infty \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) = & \frac{1}{4\pi} \left[ \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}} - \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2}} \right. \\ & \left. - \frac{1}{\sqrt{(x-\xi)^2 + (y+\eta)^2 + (z-\zeta)^2}} + \frac{1}{\sqrt{(x-\xi)^2 + (y+\eta)^2 + (z+\zeta)^2}} \right]. \end{aligned}$$

⊙ *References:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), A. G. Butkovskiy (1979).

**8.2.2-5. Domain:  $0 \leq x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . First boundary value problem.**

An octant is considered. Boundary conditions are prescribed:

$$w = f_1(y, z) \quad \text{at} \quad x = 0, \quad w = f_2(x, z) \quad \text{at} \quad y = 0, \quad w = f_3(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_0^\infty \int_0^\infty f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\eta d\zeta \\ & + \int_0^\infty \int_0^\infty f_2(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\xi d\zeta \\ & + \int_0^\infty \int_0^\infty f_3(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta \\ & + \int_0^\infty \int_0^\infty \int_0^\infty \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) = & \frac{1}{4\pi} \left[ \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}} - \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2}} \right. \\ & - \frac{1}{\sqrt{(x-\xi)^2 + (y+\eta)^2 + (z-\zeta)^2}} + \frac{1}{\sqrt{(x-\xi)^2 + (y+\eta)^2 + (z+\zeta)^2}} \\ & - \frac{1}{\sqrt{(x+\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}} + \frac{1}{\sqrt{(x+\xi)^2 + (y-\eta)^2 + (z+\zeta)^2}} \\ & \left. + \frac{1}{\sqrt{(x+\xi)^2 + (y+\eta)^2 + (z-\zeta)^2}} - \frac{1}{\sqrt{(x+\xi)^2 + (y+\eta)^2 + (z+\zeta)^2}} \right]. \end{aligned}$$

⊙ *References:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), A. G. Butkovskiy (1979).

**8.2.2-6. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq a$ . First boundary value problem.**

An infinite layer is considered. Boundary conditions are prescribed:

$$w = f_1(x, y) \quad \text{at} \quad z = 0, \quad w = f_2(x, y) \quad \text{at} \quad z = a.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_1(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta \\ & - \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_2(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=a} d\xi d\eta \\ & + \int_0^a \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \left( \frac{1}{r_{n1}} - \frac{1}{r_{n2}} \right),$$

where

$$\begin{aligned} r_{n1} &= \sqrt{(x - \xi)^2 + (y - \eta)^2 + (z - \zeta - 2na)^2}, \\ r_{n2} &= \sqrt{(x - \xi)^2 + (y - \eta)^2 + (z + \zeta - 2na)^2}. \end{aligned}$$

⊙ *References:* A. G. Butkovskiy (1979), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.2.2-7. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq a$ . Mixed boundary value problem.**

An infinite layer is considered. Boundary conditions are prescribed:

$$w = f_1(x, y) \quad \text{at} \quad z = 0, \quad \partial_z w = f_2(x, y) \quad \text{at} \quad z = a.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_1(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta \\ & + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_2(\xi, \eta) G(x, y, z, \xi, \eta, a) d\xi d\eta \\ & + \int_0^a \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \left( \frac{1}{r_{n1}} - \frac{1}{r_{n2}} \right),$$

where

$$\begin{aligned} r_{n1} &= \sqrt{(x - \xi)^2 + (y - \eta)^2 + [z - (-1)^n \zeta - 2na]^2}, \\ r_{n2} &= \sqrt{(x - \xi)^2 + (y - \eta)^2 + [z + (-1)^n \zeta - 2na]^2}. \end{aligned}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.2.2-8. Domain:  $0 \leq x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq a$ . First boundary value problem.**

A semiinfinite layer is considered. Boundary conditions are prescribed:

$$w = f_1(y, z) \quad \text{at} \quad x = 0, \quad w = f_2(x, y) \quad \text{at} \quad z = 0, \quad w = f_3(x, y) \quad \text{at} \quad z = a.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_0^a \int_{-\infty}^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\eta d\zeta \\ & + \int_{-\infty}^{\infty} \int_0^{\infty} f_2(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta \\ & - \int_{-\infty}^{\infty} \int_0^{\infty} f_3(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=a} d\xi d\eta \\ & + \int_0^a \int_{-\infty}^{\infty} \int_0^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \left( \frac{1}{r_{n1}} - \frac{1}{r_{n2}} - \frac{1}{r_{n3}} + \frac{1}{r_{n4}} \right),$$

where

$$\begin{aligned} r_{n1} &= \sqrt{(x - \xi)^2 + (y - \eta)^2 + (z - \zeta - 2na)^2}, \\ r_{n2} &= \sqrt{(x - \xi)^2 + (y - \eta)^2 + (z + \zeta - 2na)^2}, \\ r_{n3} &= \sqrt{(x + \xi)^2 + (y - \eta)^2 + (z - \zeta - 2na)^2}, \\ r_{n4} &= \sqrt{(x + \xi)^2 + (y - \eta)^2 + (z + \zeta - 2na)^2}. \end{aligned}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.2.2-9. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . First boundary value problem.**

An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y, z) \quad \text{at} \quad x = 0, & w &= f_2(y, z) \quad \text{at} \quad x = a, \\ w &= f_3(x, z) \quad \text{at} \quad y = 0, & w &= f_4(x, z) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\ & - \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=a} d\zeta d\eta \\ & + \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\zeta d\xi \\ & - \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=b} d\zeta d\xi \\ & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) = & \frac{2}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \exp(-\beta_{nm} |z - \zeta|), \\ & p_n = \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2}. \end{aligned}$$

Alternatively, the Green's function can be represented as

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \left( \frac{1}{r_{nm}^{(1)}} - \frac{1}{r_{nm}^{(2)}} - \frac{1}{r_{nm}^{(3)}} + \frac{1}{r_{nm}^{(4)}} \right),$$

where

$$\begin{aligned} r_{nm}^{(1)} &= \sqrt{(x - \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z - \zeta)^2}, \\ r_{nm}^{(2)} &= \sqrt{(x + \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z - \zeta)^2}, \\ r_{nm}^{(3)} &= \sqrt{(x - \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z - \zeta)^2}, \\ r_{nm}^{(4)} &= \sqrt{(x + \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z - \zeta)^2}. \end{aligned}$$

8.2.2-10. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . Third boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w - k_1 w &= f_1(y, z) \quad \text{at} \quad x = 0, & \partial_x w + k_2 w &= f_2(y, z) \quad \text{at} \quad x = a, \\ \partial_y w - k_3 w &= f_3(x, z) \quad \text{at} \quad y = 0, & \partial_y w + k_4 w &= f_4(x, z) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z) = & - \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) G(x, y, z, 0, \eta, \zeta) d\zeta d\eta + \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) G(x, y, z, a, \eta, \zeta) d\zeta d\eta \\ & - \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) G(x, y, z, \xi, 0, \zeta) d\zeta d\xi + \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\ & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{u_{nm}(x, y) u_{nm}(\xi, \eta)}{\|u_{nm}\|^2 \beta_{nm}} \exp(-\beta_{nm}|z - \zeta|),$$

where

$$\begin{aligned} w_{nm}(x, y) &= (\mu_n \cos \mu_n x + k_1 \sin \mu_n x)(\nu_m \cos \nu_m y + k_3 \sin \nu_m y), \quad \beta_{nm} = \sqrt{\mu_n^2 + \nu_m^2}, \\ \|w_{nm}\|^2 &= \frac{1}{4}(\mu_n^2 + k_1^2)(\nu_m^2 + k_3^2) \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right] \left[ b + \frac{(k_3 + k_4)(\nu_m^2 + k_3 k_4)}{(\nu_m^2 + k_3^2)(\nu_m^2 + k_4^2)} \right]. \end{aligned}$$

Here, the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}, \quad \tan(\nu b) = \frac{(k_3 + k_4)\nu}{\nu^2 - k_3 k_4}.$$

8.2.2-11. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . Mixed boundary value problems.

1°. An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y, z) \quad \text{at} \quad x = 0, & \partial_x w &= f_2(y, z) \quad \text{at} \quad x = a, \\ w &= f_3(x, z) \quad \text{at} \quad y = 0, & \partial_y w &= f_4(x, z) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z) = & \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\
 & + \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) G(x, y, z, a, \eta, \zeta) d\zeta d\eta \\
 & + \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\zeta d\xi \\
 & + \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\
 & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi.
 \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{2}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \exp(-\beta_{nm}|z - \zeta|),$$

where

$$p_n = \frac{(2n+1)\pi}{2a}, \quad q_m = \frac{(2m+1)\pi}{2b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2}.$$

2°. An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned}
 w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad w = f_2(y, z) \quad \text{at} \quad x = a, \\
 \partial_y w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad \partial_y w = f_4(x, z) \quad \text{at} \quad y = b.
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z) = & \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\
 & - \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=a} d\zeta d\eta \\
 & - \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) G(x, y, z, \xi, 0, \zeta) d\zeta d\xi \\
 & + \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\
 & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi.
 \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{ab} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{\beta_{nm}} \sin(p_n x) \cos(q_m y) \sin(p_n \xi) \cos(q_m \eta) \exp(-\beta_{nm}|z - \zeta|),$$

where

$$p_n = \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases}$$

► Paragraphs 8.2.2-12 through 8.2.2-17 present only Green's functions; the complete solution is constructed with the formulas given in Paragraphs 8.2.1-1 through 8.2.1-3.

8.2.2-12. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . First boundary value problem.

An infinite cylindrical domain of triangular cross-section is considered. Boundary conditions are prescribed:

$$w = f_1(y, z) \quad \text{at} \quad x = 0, \quad w = f_2(x, z) \quad \text{at} \quad y = 0, \quad w = f_3(x, z) \quad \text{at} \quad y = x.$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = H(x, y, z, \xi, \eta, \zeta) - H(x, y, z, \eta, \xi, \zeta),$$

where

$$H(x, y, z, \xi, \eta, \zeta) = \frac{2}{a^2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(p_m y) \sin(p_n \xi) \sin(p_m \eta) \exp(-\beta_{nm} |z - \zeta|),$$

$$p_n = \frac{n\pi}{a}, \quad p_m = \frac{m\pi}{a}, \quad \beta_{nm} = \sqrt{p_n^2 + p_m^2}.$$

An alternative representation of the Green's function can be obtained by setting

$$H(x, y, z, \xi, \eta, \zeta) = \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \left( \frac{1}{r_{nm}^{(1)}} - \frac{1}{r_{nm}^{(2)}} - \frac{1}{r_{nm}^{(3)}} + \frac{1}{r_{nm}^{(4)}} \right),$$

where the functions  $r_{nm}^{(k)}$  ( $k = 1, 2, 3, 4$ ) are specified in Paragraph 8.2.2-9 for  $a = b$ .

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

8.2.2-13. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z < \infty$ . First boundary value problem.

A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y, z) \quad \text{at} \quad x = 0, & w &= f_2(y, z) \quad \text{at} \quad x = a, \\ w &= f_3(x, z) \quad \text{at} \quad y = 0, & w &= f_4(x, z) \quad \text{at} \quad y = b, \\ w &= f_5(x, y) \quad \text{at} \quad z = 0. \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) H_{nm}(z, \zeta),$$

$$p_n = \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2},$$

$$H_{nm}(z, \zeta) = \begin{cases} \exp(-\beta_{nm} z) \sinh(\beta_{nm} \zeta) & \text{for } z > \zeta \geq 0, \\ \exp(-\beta_{nm} \zeta) \sinh(\beta_{nm} z) & \text{for } \zeta > z \geq 0. \end{cases}$$

An alternative representation of the Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \left( \frac{1}{r_{nm}^{(1)}} - \frac{1}{r_{nm}^{(2)}} - \frac{1}{r_{nm}^{(3)}} + \frac{1}{r_{nm}^{(4)}} - \frac{1}{r_{nm}^{(5)}} + \frac{1}{r_{nm}^{(6)}} + \frac{1}{r_{nm}^{(7)}} - \frac{1}{r_{nm}^{(8)}} \right),$$

where

$$\begin{aligned} r_{nm}^{(1)} &= \sqrt{(x - \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z - \zeta)^2}, \\ r_{nm}^{(2)} &= \sqrt{(x + \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z - \zeta)^2}, \\ r_{nm}^{(3)} &= \sqrt{(x - \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z - \zeta)^2}, \\ r_{nm}^{(4)} &= \sqrt{(x + \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z - \zeta)^2}, \\ r_{nm}^{(5)} &= \sqrt{(x - \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z + \zeta)^2}, \\ r_{nm}^{(6)} &= \sqrt{(x + \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z + \zeta)^2}, \\ r_{nm}^{(7)} &= \sqrt{(x - \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z + \zeta)^2}, \\ r_{nm}^{(8)} &= \sqrt{(x + \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z + \zeta)^2}. \end{aligned}$$

**8.2.2-14. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z < \infty$ . Third boundary value problem.**

A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w - k_1 w &= f_1(y, z) & \text{at } x = 0, & & \partial_x w + k_2 w &= f_2(y, z) & \text{at } x = a, \\ \partial_y w - k_3 w &= f_3(x, z) & \text{at } y = 0, & & \partial_y w + k_4 w &= f_4(x, z) & \text{at } y = b, \\ \partial_z w - k_5 w &= f_5(x, y) & \text{at } z = 0. & & & & \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{u_{nm}(x, y) u_{nm}(\xi, \eta)}{\|u_{nm}\|^2} H_{nm}(z, \zeta),$$

where

$$\begin{aligned} w_{nm}(x, y) &= (\mu_n \cos \mu_n x + k_1 \sin \mu_n x)(\nu_m \cos \nu_m y + k_3 \sin \nu_m y), \\ \|w_{nm}\|^2 &= \frac{1}{4}(\mu_n^2 + k_1^2)(\nu_m^2 + k_3^2) \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right] \left[ b + \frac{(k_3 + k_4)(\nu_m^2 + k_3 k_4)}{(\nu_m^2 + k_3^2)(\nu_m^2 + k_4^2)} \right], \\ H_{nm}(z, \zeta) &= \begin{cases} \frac{\exp(-\beta_{nm} z) [\beta_{nm} \cosh(\beta_{nm} \zeta) + k_5 \sinh(\beta_{nm} \zeta)]}{\beta_{nm}(\beta_{nm} + k_5)} & \text{for } z > \zeta, \\ \frac{\exp(-\beta_{nm} \zeta) [\beta_{nm} \cosh(\beta_{nm} z) + k_5 \sinh(\beta_{nm} z)]}{\beta_{nm}(\beta_{nm} + k_5)} & \text{for } \zeta > z, \end{cases} & \beta_{nm} = \sqrt{\mu_n^2 + \nu_m^2}. \end{aligned}$$

Here, the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}, \quad \tan(\nu b) = \frac{(k_3 + k_4)\nu}{\nu^2 - k_3 k_4}.$$

**8.2.2-15. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z < \infty$ . Mixed boundary value problems.**

1°. A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y, z) & \text{at } x = 0, & & w &= f_2(y, z) & \text{at } x = a, \\ w &= f_3(x, z) & \text{at } y = 0, & & w &= f_4(x, z) & \text{at } y = b, \\ \partial_z w &= f_5(x, y) & \text{at } z = 0. & & & & \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) &= \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) H_{nm}(z, \zeta), \\ p_n &= \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2}, \\ H_{nm}(z, \zeta) &= \begin{cases} \exp(-\beta_{nm} z) \cosh(\beta_{nm} \zeta) & \text{for } z > \zeta \geq 0, \\ \exp(-\beta_{nm} \zeta) \cosh(\beta_{nm} z) & \text{for } \zeta > z \geq 0. \end{cases} \end{aligned}$$

2°. A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w &= f_1(y, z) & \text{at } x = 0, & & \partial_x w &= f_2(y, z) & \text{at } x = a, \\ \partial_y w &= f_3(x, z) & \text{at } y = 0, & & \partial_y w &= f_4(x, z) & \text{at } y = b, \\ w &= f_5(x, y) & \text{at } z = 0. & & & & \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_n A_m}{\beta_{nm}} \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) H_{nm}(z, \zeta),$$

$$p_n = \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

$$H_{nm}(z, \zeta) = \begin{cases} \exp(-\beta_{nm} z) \sinh(\beta_{nm} \zeta) & \text{for } z > \zeta \geq 0, \\ \exp(-\beta_{nm} \zeta) \sinh(\beta_{nm} z) & \text{for } \zeta > z \geq 0. \end{cases}$$

8.2.2-16. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . First boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y, z) & \text{at } x = 0, & & w &= f_2(y, z) & \text{at } x = a, \\ w &= f_3(x, z) & \text{at } y = 0, & & w &= f_4(x, z) & \text{at } y = b, \\ w &= f_5(x, y) & \text{at } z = 0, & & w &= f_6(x, y) & \text{at } z = c. \end{aligned}$$

1°. Representation of the Green's function in the form of a double series:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) F_{nm}(z, \zeta),$$

$$F_{nm}(z, \zeta) = \begin{cases} \frac{\sinh(\beta_{nm} \zeta) \sinh[\beta_{nm}(c - z)]}{\beta_{nm} \sinh(\beta_{nm} c)} & \text{for } c \geq z > \zeta \geq 0, \\ \frac{\sinh(\beta_{nm} z) \sinh[\beta_{nm}(c - \zeta)]}{\beta_{nm} \sinh(\beta_{nm} c)} & \text{for } c \geq \zeta > z \geq 0, \end{cases}$$

$$p_n = \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2}.$$

This relation can be used to obtain two other representations of the Green's function by means of the following cyclic permutations:

$$\begin{array}{ccc} & (x, \xi, a) & \\ \nearrow & & \searrow \\ (z, \zeta, c) & \leftarrow & (y, \eta, b) \end{array}$$

2°. Representation of the Green's function in the form of a triple series:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{8}{abc} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{\sin(p_n x) \sin(q_m y) \sin(s_k z) \sin(p_n \xi) \sin(q_m \eta) \sin(s_k \zeta)}{p_n^2 + q_m^2 + s_k^2},$$

$$p_n = \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}, \quad s_k = \frac{\pi k}{c}.$$

3°. An alternative representation of the Green's function in the form of a triple series:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{4\pi} \sum_{n=-\infty}^{\infty} \sum_{m=-\infty}^{\infty} \sum_{k=-\infty}^{\infty} \left( \frac{1}{r_{nmk}^{(1)}} - \frac{1}{r_{nmk}^{(2)}} - \frac{1}{r_{nmk}^{(3)}} + \frac{1}{r_{nmk}^{(4)}} \right. \\ \left. - \frac{1}{r_{nmk}^{(5)}} + \frac{1}{r_{nmk}^{(6)}} + \frac{1}{r_{nmk}^{(7)}} - \frac{1}{r_{nmk}^{(8)}} \right),$$



where

$$\begin{aligned}
 r_{nmk}^{(1)} &= \sqrt{(x - \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z - \zeta - 2kc)^2}, \\
 r_{nmk}^{(2)} &= \sqrt{(x + \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z - \zeta - 2kc)^2}, \\
 r_{nmk}^{(3)} &= \sqrt{(x - \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z - \zeta - 2kc)^2}, \\
 r_{nmk}^{(4)} &= \sqrt{(x + \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z - \zeta - 2kc)^2}, \\
 r_{nmk}^{(5)} &= \sqrt{(x - \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z + \zeta - 2kc)^2}, \\
 r_{nmk}^{(6)} &= \sqrt{(x + \xi - 2na)^2 + (y - \eta - 2mb)^2 + (z + \zeta - 2kc)^2}, \\
 r_{nmk}^{(7)} &= \sqrt{(x - \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z + \zeta - 2kc)^2}, \\
 r_{nmk}^{(8)} &= \sqrt{(x + \xi - 2na)^2 + (y + \eta - 2mb)^2 + (z + \zeta - 2kc)^2}.
 \end{aligned}$$

8.2.2-17. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . Third boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned}
 \partial_x w - k_1 w &= f_1(y, z) & \text{at } x = 0, & & \partial_x w + k_2 w &= f_2(y, z) & \text{at } x = a, \\
 \partial_y w - k_3 w &= f_3(x, z) & \text{at } y = 0, & & \partial_y w + k_4 w &= f_4(x, z) & \text{at } y = b, \\
 \partial_z w - k_5 w &= f_5(x, y) & \text{at } z = 0, & & \partial_z w + k_6 w &= f_6(x, y) & \text{at } z = c.
 \end{aligned}$$

Green's function:

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta) &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi) \psi_m(y) \psi_m(\eta) \chi_s(z) \chi_s(\zeta)}{\|\varphi_n\|^2 \|\psi_m\|^2 \|\chi_s\|^2 (\mu_n^2 + \lambda_m^2 + \nu_s^2)}, \\
 \varphi_n(x) &= \cos(\mu_n x) + \frac{k_1}{\mu_n} \sin(\mu_n x), & \|\varphi_n\|^2 &= \frac{k_2}{2\mu_n^2} \frac{\mu_n^2 + k_1^2}{\mu_n^2 + k_2^2} + \frac{k_1}{2\mu_n^2} + \frac{a}{2} \left( 1 + \frac{k_1^2}{\mu_n^2} \right), \\
 \psi_m(y) &= \cos(\lambda_m y) + \frac{k_3}{\lambda_m} \sin(\lambda_m y), & \|\psi_m\|^2 &= \frac{k_4}{2\lambda_m^2} \frac{\lambda_m^2 + k_3^2}{\lambda_m^2 + k_4^2} + \frac{k_3}{2\lambda_m^2} + \frac{b}{2} \left( 1 + \frac{k_3^2}{\lambda_m^2} \right), \\
 \chi_s(z) &= \cos(\nu_s z) + \frac{k_5}{\nu_s} \sin(\nu_s z), & \|\chi_s\|^2 &= \frac{k_6}{2\nu_s^2} \frac{\nu_s^2 + k_5^2}{\nu_s^2 + k_6^2} + \frac{k_5}{2\nu_s^2} + \frac{c}{2} \left( 1 + \frac{k_5^2}{\nu_s^2} \right),
 \end{aligned}$$

where the  $\mu_n$ ,  $\lambda_m$ , and  $\nu_s$  are positive roots of the transcendental equations

$$\frac{\tan(\mu a)}{\mu} = \frac{k_1 + k_2}{\mu^2 - k_1 k_2}, \quad \frac{\tan(\lambda b)}{\lambda} = \frac{k_3 + k_4}{\lambda^2 - k_3 k_4}, \quad \frac{\tan(\nu c)}{\nu} = \frac{k_5 + k_6}{\nu^2 - k_5 k_6}.$$

8.2.2-18. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . Mixed boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned}
 w &= f_1(y, z) & \text{at } x = 0, & & \partial_x w &= f_2(y, z) & \text{at } x = a, \\
 w &= f_3(x, z) & \text{at } y = 0, & & \partial_y w &= f_4(x, z) & \text{at } y = b, \\
 w &= f_5(x, y) & \text{at } z = 0, & & \partial_z w &= f_6(x, y) & \text{at } z = c.
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z) = & \int_0^a \int_0^b \int_0^c \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi \\
 & + \int_0^b \int_0^c f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta + \int_0^b \int_0^c f_2(\eta, \zeta) G(x, y, z, a, \eta, \zeta) d\zeta d\eta \\
 & + \int_0^a \int_0^c f_3(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\zeta d\xi + \int_0^a \int_0^c f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\
 & + \int_0^a \int_0^b f_5(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\eta d\xi + \int_0^a \int_0^b f_6(\xi, \eta) G(x, y, z, \xi, \eta, c) d\eta d\xi.
 \end{aligned}$$

1°. A double-series representation of the Green's function:

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta) &= \frac{4}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) F_{nm}(z, \zeta), \\
 F_{nm}(z, \zeta) &= \begin{cases} \frac{\sinh(\beta_{nm} \zeta) \cosh[\beta_{nm}(c-z)]}{\beta_{nm} \cosh(\beta_{nm} c)} & \text{for } c \geq z > \zeta \geq 0, \\ \frac{\sinh(\beta_{nm} z) \cosh[\beta_{nm}(c-\zeta)]}{\beta_{nm} \cosh(\beta_{nm} c)} & \text{for } c \geq \zeta > z \geq 0, \end{cases} \\
 p_n &= \frac{\pi(2n+1)}{2a}, \quad q_m = \frac{\pi(2m+1)}{2b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2}.
 \end{aligned}$$

This relation can be used to obtain two other representations of the Green's function by means of the following cyclic permutations:

$$\begin{array}{ccc}
 & (x, \xi, a) & \\
 \nearrow & & \searrow \\
 (z, \zeta, c) & \longleftarrow & (y, \eta, b)
 \end{array}$$

2°. A triple series representation of the Green's function:

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta) &= \frac{8}{abc} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{\sin(p_n x) \sin(q_m y) \sin(s_k z) \sin(p_n \xi) \sin(q_m \eta) \sin(s_k \zeta)}{p_n^2 + q_m^2 + s_k^2}, \\
 p_n &= \frac{\pi(2n+1)}{2a}, \quad q_m = \frac{\pi(2m+1)}{2b}, \quad s_k = \frac{\pi(2k+1)}{2c}.
 \end{aligned}$$

### 8.2.3. Problems in Cylindrical Coordinates

The three-dimensional Poisson equation in the cylindrical coordinate system is written as

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} = -\Phi(r, \varphi, z), \quad r = \sqrt{x^2 + y^2}.$$

8.2.3-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . First boundary value problem.

An infinite circular cylinder is considered. A boundary condition is prescribed:

$$w = f(\varphi, z) \quad \text{at} \quad r = R.$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z) = & -R \int_0^{2\pi} \int_{-\infty}^{\infty} f(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\xi=R} d\zeta d\eta \\
 & + \int_0^R \int_0^{2\pi} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi.
 \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{2\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{[J'_n(\mu_{nm} R)]^2 \mu_{nm}} \cos[n(\varphi - \eta)] \exp(-\mu_{nm}|z - \zeta|),$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n \neq 0$ ; the  $J_n(\xi)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

8.2.3-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Third boundary value problem.

An infinite circular cylinder is considered. A boundary condition is prescribed:

$$\partial_r w + kw = f(\varphi, z) \quad \text{at} \quad r = R.$$

Solution:

$$\begin{aligned} w(r, \varphi, z) = & R \int_0^{2\pi} \int_{-\infty}^{\infty} f(\eta, \zeta) G(r, \varphi, z, R, \eta, \zeta) d\zeta d\eta \\ & + \int_0^R \int_0^{2\pi} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{2\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)]}{(\mu_{nm}^2 R^2 + k^2 R^2 - n^2) J_n^2(\mu_{nm} R)} \exp(-\mu_{nm}|z - \zeta|),$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n \neq 0$ ; the  $J_n(\xi)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k J_n(\mu R) = 0.$$

8.2.3-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . First boundary value problem.

A semiinfinite circular cylinder is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi, z) \quad \text{at} \quad r = R, \quad w = f_2(r, \varphi) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(r, \varphi, z) = & -R \int_0^{2\pi} \int_0^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\xi=R} d\zeta d\eta \\ & + \int_0^{2\pi} \int_0^R f_2(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\zeta=0} \xi d\xi d\eta \\ & + \int_0^R \int_0^{2\pi} \int_0^{\infty} \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{2\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{[J'_n(\mu_{nm} R)]^2 \mu_{nm}} \cos[n(\varphi - \eta)] F_{nm}(z, \zeta),$$

$$F_{nm}(z, \zeta) = \exp(-\mu_{nm}|z - \zeta|) - \exp(-\mu_{nm}|z + \zeta|), \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

**8.2.3-4. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Third boundary value problem.**

A semiinfinite circular cylinder is considered. Boundary conditions are prescribed:

$$\partial_r w + k_1 w = f_1(\varphi, z) \quad \text{at} \quad r = R, \quad \partial_z w - k_2 w = f_2(r, \varphi) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(r, \varphi, z) = & R \int_0^{2\pi} \int_0^\infty f_1(\eta, \zeta) G(r, \varphi, z, R, \eta, \zeta) d\zeta d\eta \\ & - \int_0^{2\pi} \int_0^R f_2(\xi, \eta) G(r, \varphi, z, R, \eta, 0) \xi d\xi d\eta \\ & + \int_0^R \int_0^{2\pi} \int_0^\infty \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)]}{(\mu_{nm}^2 R^2 + k_1^2 R^2 - n^2) J_n^2(\mu_{nm} R)} F_{nm}(z, \zeta),$$

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad F_{nm}(z, \zeta) = \begin{cases} \frac{\exp(-\mu_{nm} z) [\mu_{nm} \cosh(\mu_{nm} \zeta) + k_2 \sinh(\mu_{nm} \zeta)]}{\mu_{nm} (\mu_{nm} + k_2)} & \text{for } z > \zeta, \\ \frac{\exp(-\mu_{nm} \zeta) [\mu_{nm} \cosh(\mu_{nm} z) + k_2 \sinh(\mu_{nm} z)]}{\mu_{nm} (\mu_{nm} + k_2)} & \text{for } \zeta > z, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J_n'(\mu R) + k_1 J_n(\mu R) = 0.$$

**8.2.3-5. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Mixed boundary value problem.**

A semiinfinite circular cylinder is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi, z) \quad \text{at} \quad r = R, \quad \partial_z w = f_2(r, \varphi) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(r, \varphi, z) = & -R \int_0^{2\pi} \int_0^\infty f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\xi=R} d\zeta d\eta \\ & - \int_0^{2\pi} \int_0^R f_2(\xi, \eta) G(r, \varphi, z, \xi, \eta, 0) \xi d\xi d\eta \\ & + \int_0^R \int_0^{2\pi} \int_0^\infty \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{2\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)]}{[J_n'(\mu_{nm} R)]^2 \mu_{nm}} F_{nm}(z, \zeta),$$

$$F_{nm}(z, \zeta) = \exp(-\mu_{nm}|z - \zeta|) + \exp(-\mu_{nm}|z + \zeta|), \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are roots of the transcendental equation  $J_n(\mu R) = 0$ .

► Paragraphs 8.2.3-6 through 8.3.3-10 present only Green's functions; the complete solution is constructed with the formulas given in Subsection 8.2.1. See also Paragraphs 8.3.1-4 and 8.3.1-8 for  $\lambda = 0$ .

**8.2.3-6. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . First boundary value problem.**

A circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi, z) \quad \text{at} \quad r = R, \quad w = f_2(r, \varphi) \quad \text{at} \quad z = 0, \quad w = f_3(r, \varphi) \quad \text{at} \quad z = a.$$

A double series representation of the Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{[J'_n(\mu_{nm} R)]^2 \mu_{nm} \sinh(\mu_{nm} a)} \cos[n(\varphi - \eta)] F_{nm}(z, \zeta),$$

$$F_{nm}(z, \zeta) = \begin{cases} \sinh(\mu_{nm} \zeta) \sinh[\mu_{nm}(a - z)] & \text{for } a \geq z > \zeta \geq 0, \\ \sinh(\mu_{nm} z) \sinh[\mu_{nm}(a - \zeta)] & \text{for } a \geq \zeta > z \geq 0, \end{cases} \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument) and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

A triple series representation of the Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{2a}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n}{[J'_n(\mu_{nm} R)]^2 [(a\mu_{nm})^2 + (\pi k)^2]} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\ \times \cos[n(\varphi - \eta)] \sin\left(\frac{k\pi z}{a}\right) \sin\left(\frac{k\pi \zeta}{a}\right).$$

**8.2.3-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . Third boundary value problem.**

A circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_r w + k_1 w &= f_1(\varphi, z) \quad \text{at} \quad r = R, \\ \partial_z w - k_2 w &= f_2(r, \varphi) \quad \text{at} \quad z = 0, \\ \partial_z w + k_3 w &= f_3(r, \varphi) \quad \text{at} \quad z = a. \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)] h_s(z) h_s(\zeta)}{(\mu_{nm}^2 R^2 + k_1^2 R^2 - n^2)(\mu_{nm}^2 + \lambda_s^2) [J_n(\mu_{nm} R)]^2 \|h_s\|^2},$$

$$h_s(z) = \cos(\lambda_s z) + \frac{k_2}{\lambda_s} \sin(\lambda_s z), \quad \|h_s\|^2 = \frac{k_3}{2\lambda_s^2} \frac{\lambda_s^2 + k_2^2}{\lambda_s^2 + k_3^2} + \frac{k_2}{2\lambda_s^2} + \frac{a}{2} \left(1 + \frac{k_2^2}{\lambda_s^2}\right).$$

Here,  $A_0 = 1$  and  $A_n = 2$  for  $n \neq 0$ ; the  $J_n(\xi)$  are the Bessel functions; and the  $\mu_{nm}$  and  $\lambda_s$  are positive roots of the transcendental equations

$$\mu J'_n(\mu R) + k_1 J_n(\mu R) = 0, \quad \frac{\tan(\lambda a)}{\lambda} = \frac{k_2 + k_3}{\lambda^2 - k_2 k_3}.$$

**8.2.3-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . Mixed boundary value problem.**

A circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi, z) \quad \text{at} \quad r = R, \quad \partial_z w = f_2(r, \varphi) \quad \text{at} \quad z = 0, \quad \partial_z w = f_3(r, \varphi) \quad \text{at} \quad z = a.$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{a}{\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{A_n A_k}{[J'_n(\mu_{nm} R)]^2 [(a\mu_{nm})^2 + (\pi k)^2]} J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \\ \times \cos[n(\varphi - \eta)] \cos\left(\frac{k\pi z}{a}\right) \cos\left(\frac{k\pi \zeta}{a}\right),$$

where  $A_0 = 1$  and  $A_n = 2$  for  $n \neq 0$ ; the  $J_n(\xi)$  are the Bessel functions (the prime denotes the derivative with respect to the argument); and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

**8.2.3-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq a$ . First boundary value problem.**

A cylindrical sector of finite thickness is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(r, z) \quad \text{at} \quad \varphi = 0, \quad w = f_2(r, z) \quad \text{at} \quad \varphi = \varphi_0, \quad w = f_3(\varphi, z) \quad \text{at} \quad r = R, \\ w = f_4(r, \varphi) \quad \text{at} \quad z = 0, \quad w = f_5(r, \varphi) \quad \text{at} \quad z = a. \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{8a}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{J_{n\pi/\varphi_0}(\mu_{nm}r) J_{n\pi/\varphi_0}(\mu_{nm}\xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm}R)]^2 [(a\mu_{nm})^2 + (\pi k)^2]} \\ \times \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \sin\left(\frac{k\pi z}{a}\right) \sin\left(\frac{k\pi\zeta}{a}\right),$$

where the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

**8.2.3-10. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq a$ . Mixed boundary value problem.**

A cylindrical sector of finite thickness is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(r, z) \quad \text{at} \quad \varphi = 0, \quad w = f_2(r, z) \quad \text{at} \quad \varphi = \varphi_0, \quad w = f_3(\varphi, z) \quad \text{at} \quad r = R, \\ \partial_z w = f_4(r, \varphi) \quad \text{at} \quad z = 0, \quad \partial_z w = f_5(r, \varphi) \quad \text{at} \quad z = a. \end{aligned}$$

Green's function:

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{4a}{R^2 \varphi_0} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^{\infty} \frac{A_k J_{n\pi/\varphi_0}(\mu_{nm}r) J_{n\pi/\varphi_0}(\mu_{nm}\xi)}{[J'_{n\pi/\varphi_0}(\mu_{nm}R)]^2 [(a\mu_{nm})^2 + (\pi k)^2]} \\ \times \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{n\pi\eta}{\varphi_0}\right) \cos\left(\frac{k\pi z}{a}\right) \cos\left(\frac{k\pi\zeta}{a}\right),$$

where  $A_0 = 1$  and  $A_k = 2$  for  $k \neq 0$ ; the  $J_{n\pi/\varphi_0}(r)$  are the Bessel functions; and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu R) = 0$ .

## 8.2.4. Problems in Spherical Coordinates

The three-dimensional Poisson equation in the spherical coordinate system is written as

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} = -\Phi(r, \theta, \varphi), \quad r = \sqrt{x^2 + y^2 + z^2}.$$

► Only Green's functions are presented below; the complete solutions can be constructed with the formulas given in Subsection 8.2.1.

**8.2.4-1. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

A spherical domain is considered. A boundary condition is prescribed:

$$w = f(\varphi, \theta) \quad \text{at} \quad r = R.$$

Green's function:

$$G(r, \theta, \varphi, \xi, \eta, \zeta) = \frac{1}{4\pi \sqrt{r^2 - 2r\xi \cos \gamma + \xi^2}} - \frac{1}{4\pi \sqrt{r^2 \xi^2 - 2R^2 r \xi \cos \gamma + R^4}}, \\ \cos \gamma = \cos \theta \cos \eta + \sin \theta \sin \eta \cos(\varphi - \zeta).$$

An alternative representation of the Green's function:

$$G(\mathbf{r}, \mathbf{r}_0) = \frac{1}{4\pi} \frac{1}{|\mathbf{r} - \mathbf{r}_0|} - \frac{1}{4\pi} \frac{R}{r_0|(R/r_0)^2 \mathbf{r}_0 - \mathbf{r}|}, \quad r_0 = |\mathbf{r}_0|,$$

where

$$\mathbf{r} = \{x, y, z\}, \quad x = r \sin \theta \cos \varphi, \quad y = r \sin \theta \sin \varphi, \quad z = r \cos \theta$$

$$\mathbf{r}_0 = \{x_0, y_0, z_0\}, \quad x_0 = \xi \sin \eta \cos \zeta, \quad y_0 = \xi \sin \eta \sin \zeta, \quad z_0 = \xi \cos \eta.$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

8.2.4-2. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.

A spherical domain is considered. A boundary condition is prescribed:

$$\partial_r w = f(\varphi, \theta) \quad \text{at} \quad r = R.$$

Green's function:

$$G(r, \theta, \varphi, \xi, \eta, \zeta) = \frac{1}{4\pi} \left\{ \frac{1}{|\mathbf{r} - \mathbf{r}_0|} + \frac{R}{|\mathbf{r}_0| |\mathbf{r}_1|} + \frac{1}{R} \ln \frac{2R^2}{R^2 + |\mathbf{r}_0| |\mathbf{r}_1| - (\mathbf{r} \cdot \mathbf{r}_0)} \right\},$$

where

$$|\mathbf{r} - \mathbf{r}_0| = \sqrt{r^2 - 2r\xi \cos \gamma + \xi^2}, \quad |\mathbf{r}_0| |\mathbf{r}_1| = \sqrt{r^2 \xi^2 - 2R^2 r \xi \cos \gamma + R^4},$$

$$|\mathbf{r}_0| = \xi, \quad (\mathbf{r} \cdot \mathbf{r}_0) = r\xi \cos \gamma, \quad \cos \gamma = \cos \theta \cos \eta + \sin \theta \sin \eta \cos(\varphi - \zeta).$$

For a solution of the second boundary value problem to exist the solvability condition must be satisfied (see Paragraph 8.2.1-2).

⊙ *Reference:* N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970).

8.2.4-3. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Third boundary value problem.

A spherical domain is considered. A boundary condition is prescribed:

$$\partial_r w + kw = f(\theta, \varphi) \quad \text{at} \quad r = R.$$

Green's function:

$$G(r, \theta, \varphi, \xi, \eta, \zeta) = \frac{1}{2\pi\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{s=0}^n A_s B_{nms} J_{n+1/2}(\lambda_{nm} r) J_{n+1/2}(\lambda_{nm} \xi) \times P_n^s(\cos \theta) P_n^s(\cos \eta) \cos[s(\varphi - \zeta)],$$

$$A_s = \begin{cases} 1 & \text{for } s = 0, \\ 2 & \text{for } s \neq 0, \end{cases} \quad B_{nms} = \frac{(2n+1)(n-s)!}{(n+s)! [R^2 \lambda_{nm}^2 + (kR+n)(kR-n-1)] [J_{n+1/2}(\lambda_{nm} R)]^2}.$$

Here, the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^s(\mu)$  are the associated Legendre functions that are expressed in terms of the Legendre polynomials  $P_n(\mu)$  as

$$P_n^s(\mu) = (1 - \mu^2)^{s/2} \frac{d^s}{d\mu^s} P_n(\mu), \quad P_n(\mu) = \frac{1}{n! 2^n} \frac{d^n}{d\mu^n} (\mu^2 - 1)^n,$$

and the  $\lambda_{nm}$  are positive roots of the transcendental equation

$$\lambda R J'_{n+1/2}(\lambda R) + \left(kR - \frac{1}{2}\right) J_{n+1/2}(\lambda R) = 0.$$

**8.2.4-4. Domain:  $R \leq r < \infty$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

Three-dimensional space with a spherical cavity is considered. A boundary condition is prescribed:

$$w = f(\varphi, \theta) \quad \text{at} \quad r = R.$$

The Green's function of the outer first boundary value problem is given by the same relation as that for the inner first boundary value problem (see Paragraph 8.2.4-1), except that  $r \geq R$  and  $\xi \geq R$ .

**8.2.4-5. Domain:  $R \leq r < \infty$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . Second boundary value problem.**

Three-dimensional space with a spherical cavity is considered. A boundary condition is prescribed:

$$\partial_r w = f(\varphi, \theta) \quad \text{at} \quad r = R.$$

Green's function:

$$G(r, \theta, \varphi, \xi, \eta, \zeta) = \frac{1}{4\pi} \left\{ \frac{1}{|\mathbf{r} - \mathbf{r}_0|} + \frac{R}{|\mathbf{r}_0| |\mathbf{r}_1|} + \frac{1}{R} \ln \frac{(1 - \cos \gamma) |\mathbf{r}| |\mathbf{r}_0|}{R^2 + |\mathbf{r}_0| |\mathbf{r}_1| - (\mathbf{r} \cdot \mathbf{r}_0)} \right\},$$

where

$$|\mathbf{r}| = r, \quad |\mathbf{r}_0| = \xi, \quad |\mathbf{r} - \mathbf{r}_0| = \sqrt{r^2 - 2r\xi \cos \gamma + \xi^2}, \quad |\mathbf{r}_0| |\mathbf{r}_1| = \sqrt{r^2 \xi^2 - 2R^2 r \xi \cos \gamma + R^4},$$

$$(\mathbf{r} \cdot \mathbf{r}_0) = r\xi \cos \gamma, \quad \cos \gamma = \cos \theta \cos \eta + \sin \theta \sin \eta \cos(\varphi - \zeta).$$

⊙ *Reference:* N. S. Koshlyakov, E. B. Gliner, and M. M. Smirnov (1970).

**8.2.4-6. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \theta \leq \pi$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

A spherical layer is considered. Boundary conditions are prescribed:

$$w = f_1(\theta, \varphi) \quad \text{at} \quad r = R_1, \quad w = f_2(\theta, \varphi) \quad \text{at} \quad r = R_2.$$

Green's function:

$$G(r, \theta, \varphi, \xi, \eta, \zeta) = \frac{\pi}{8\sqrt{r\xi}} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \sum_{k=0}^n A_k B_{nmk} Z_{n+1/2}(\lambda_{nm} r) Z_{n+1/2}(\lambda_{nm} \xi)$$

$$\times P_n^k(\cos \theta) P_n^k(\cos \eta) \cos[k(\varphi - \zeta)],$$

where

$$Z_{n+1/2}(\lambda_{nm} r) = J_{n+1/2}(\lambda_{nm} R_1) Y_{n+1/2}(\lambda_{nm} r) - Y_{n+1/2}(\lambda_{nm} R_1) J_{n+1/2}(\lambda_{nm} r),$$

$$A_k = \begin{cases} 1 & \text{for } k = 0, \\ 2 & \text{for } k \neq 0, \end{cases} \quad B_{nmk} = \frac{(2n+1)(n-k)! J_{n+1/2}^2(\lambda_{nm} R_2)}{(n+k)! [J_{n+1/2}^2(\lambda_{nm} R_1) - J_{n+1/2}^2(\lambda_{nm} R_2)]};$$

the  $J_{n+1/2}(r)$  are the Bessel functions, the  $P_n^k(\mu)$  are the associated Legendre functions (see Paragraph 8.2.4-3), and the  $\lambda_{nm}$  are positive roots of the transcendental equation  $Z_{n+1/2}(\lambda R_2) = 0$ .

**8.2.4-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \theta \leq \pi/2$ ,  $0 \leq \varphi \leq 2\pi$ . First boundary value problem.**

A hemisphere is considered. Boundary conditions are prescribed:

$$w = f_1(\theta, \varphi) \quad \text{at} \quad r = R, \quad w = f_2(r, \varphi) \quad \text{at} \quad \theta = \pi/2.$$

Green's function in the spherical coordinate system:

$$G(r, \theta, \varphi, \xi, \eta, \zeta) = G_s(r, \theta, \varphi, \xi, \eta, \zeta) - G_s(r, \theta, \varphi, \xi, \pi - \eta, \zeta),$$

where  $G_s(r, \theta, \varphi, \xi, \eta, \zeta)$  is the Green's functions for a sphere; see Paragraph 8.2.4-1, where  $G$  must be replaced by  $G_s$ .

Green's function in the Cartesian coordinate system:

$$G(x, y, z, x_0, y_0, z_0) = \frac{1}{4\pi} \left( \frac{1}{|\mathbf{r} - \mathbf{r}_0|} - \frac{R}{|\mathbf{r}_0| |\mathbf{r} - \mathbf{r}_0^*|} \right) - \frac{1}{4\pi} \left( \frac{1}{|\mathbf{r} - \mathbf{r}_1|} - \frac{R}{|\mathbf{r}_0| |\mathbf{r} - \mathbf{r}_1^*|} \right),$$

$$\mathbf{r} = \{x, y, z\}, \quad \mathbf{r}_0 = \{x_0, y_0, z_0\}, \quad \mathbf{r}_1 = \{x_0, y_0, -z_0\}, \quad \mathbf{r}_k^* = (R/r_0)^2 \mathbf{r}_k, \quad k = 0, 1.$$

⊙ *References:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).



A quarter of a sphere is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(\varphi, \theta) & \text{at } r &= R, & w &= f_2(r, \varphi) & \text{at } \theta &= \pi/2, \\ w &= f_3(r, \theta) & \text{at } \varphi &= 0, & w &= f_4(r, \theta) & \text{at } \varphi &= \pi. \end{aligned}$$

Green's function in the spherical coordinate system:

$$\begin{aligned} G(r, \theta, \varphi, \xi, \eta, \zeta) &= G_s(r, \theta, \varphi, \xi, \eta, \zeta) - G_s(r, \theta, \varphi, \xi, \pi - \eta, \zeta) \\ &\quad + G_s(r, \theta, \varphi, \xi, \pi - \eta, 2\pi - \zeta) - G_s(r, \theta, \varphi, \xi, \eta, 2\pi - \zeta), \end{aligned}$$

where  $G_s(r, \theta, \varphi, \xi, \eta, \zeta)$  is the Green's function for a sphere; see Paragraph 8.2.4-1, where  $G$  must be replaced by  $G_s$ .

Green's function in the Cartesian coordinate system:

$$G(x, y, z, x_0, y_0, z_0) = \frac{1}{4\pi} \sum_{n,k=0}^1 (-1)^{n+k} \left( \frac{1}{|\mathbf{r} - \mathbf{r}_{nk}|} - \frac{R}{|\mathbf{r}_0| |\mathbf{r} - \mathbf{r}_{nk}^*|} \right),$$

$$\mathbf{r} = \{x, y, z\}, \quad \mathbf{r}_0 = \{x_0, y_0, z_0\}, \quad \mathbf{r}_{nk} = \{x_0, (-1)^n y_0, (-1)^k z_0\}, \quad \mathbf{r}_{nk}^* = (R/r_0)^2 \mathbf{r}_{nk},$$

where  $r_0 = |\mathbf{r}_0|$ ;  $n = 0, 1$ ;  $k = 0, 1$ .

⊙ *References:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

## 8.3. Helmholtz Equation $\Delta_3 w + \lambda w = -\Phi(\mathbf{x})$

A variety of problems related to steady-state oscillations (mechanical, acoustic, thermal, electromagnetic, etc.) lead to the three-dimensional Helmholtz equation with  $\lambda > 0$ . This equation governs mass transfer phenomena with volume chemical reaction of the first order for  $\lambda < 0$ . Any elliptic equation with constant coefficients can be reduced to the Helmholtz equation.

### 8.3.1. General Remarks, Results, and Formulas

#### 8.3.1-1. Some definitions.

The Helmholtz equation is called homogeneous if  $\Phi = 0$  and nonhomogeneous if  $\Phi \neq 0$ . A homogeneous boundary value problem is a boundary value problem for a homogeneous equation with homogeneous boundary conditions;  $w = 0$  is a particular solution of a homogeneous boundary value problem.

The values  $\lambda_n$  of the parameter  $\lambda$  for which there are nontrivial solutions (i.e., not identically zero solutions) of a homogeneous boundary value problem are called eigenvalues. The corresponding solutions,  $w = w_n$ , are called eigenfunctions of this boundary value problem.

In what follows, we consider simultaneously the first, second, and third boundary value problems for the three-dimensional Helmholtz equation in a finite three-dimensional domain  $V$  with a sufficiently smooth surface  $S$ . It is assumed that  $k > 0$  for the third boundary value problem with the boundary condition

$$\frac{\partial w}{\partial N} + kw = 0 \quad \text{for } \mathbf{r} \in S,$$

where  $\frac{\partial w}{\partial N}$  is the derivative along the outward normal to the surface  $S$ , and  $\mathbf{r} = \{x, y, z\}$ .

### 8.3.1-2. Properties of eigenvalues and eigenfunctions.

1°. There are infinitely many eigenvalues  $\{\lambda_n\}$ ; they form a discrete spectrum of the boundary value problem.

2°. All eigenvalues are positive, except for one eigenvalue  $\lambda_0 = 0$  of the second boundary value problem (the corresponding eigenfunction is  $w_0 = \text{const}$ ). The eigenvalues are assumed to be ordered so that  $\lambda_1 < \lambda_2 < \lambda_3 < \dots$ .

3°. The eigenvalues tend to infinity as the number  $n$  increases. The following asymptotic estimate holds:

$$\lim_{n \rightarrow \infty} \frac{n}{\lambda_n^{3/2}} = \frac{V_3}{6\pi^2},$$

where  $V_3$  is the volume of the domain under consideration.

4°. The eigenfunctions are defined up to a constant multiplier. Any two eigenfunctions,  $w_n$  and  $w_m$ , that correspond to different eigenvalues  $\lambda_n \neq \lambda_m$  are orthogonal, that is,

$$\int_V w_n w_m dV = 0.$$

5°. Any twice continuously differentiable function  $f = f(\mathbf{r})$  that satisfies the boundary conditions of a boundary value problem can be expanded into a uniformly convergent series in the eigenfunctions of this boundary value problem, specifically,

$$f = \sum_{n=1}^{\infty} a_n w_n, \quad \text{where} \quad a_n = \frac{1}{\|w_n\|^2} \int_V f w_n dV, \quad \|w_n\|^2 = \int_V w_n^2 dV.$$

If  $f$  is square summable, then the series is convergent in mean.

6°. The eigenvalues of the first boundary value problem do not increase if the domain is extended.

*Remark 1.* In a three-dimensional problem, to each eigenvalue  $\lambda_n$  finitely many linearly independent eigenfunctions  $w_n^{(1)}, \dots, w_n^{(p)}$  generally correspond. These functions can always be replaced by their linear combinations

$$\bar{w}_n^{(s)} = c_{s,1} w_n^{(1)} + \dots + c_{s,s-1} w_n^{(s-1)} + w_n^{(s)}, \quad s = 1, 2, \dots, p,$$

such that  $\bar{w}_n^{(1)}, \dots, \bar{w}_n^{(p)}$  are now pairwise orthogonal. Therefore, without loss of generality, we assume that all eigenfunctions are orthogonal.

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1984).

### 8.3.1-3. Nonhomogeneous Helmholtz equation with homogeneous boundary conditions.

Three cases are possible.

1°. If  $\lambda$  is not equal to any one of the eigenvalues, then the solution of the problem is given by

$$w = \sum_{n=1}^{\infty} \frac{A_n}{\lambda_n - \lambda} w_n, \quad \text{where} \quad A_n = \frac{1}{\|w_n\|^2} \int_V \Phi w_n dV, \quad \|w_n\|^2 = \int_V w_n^2 dV.$$

2°. If  $\lambda$  coincides with one of the eigenvalues,  $\lambda = \lambda_m$ , then the condition of the orthogonality of the function  $\Phi$  to the eigenfunction  $w_m$ ,

$$\int_V \Phi w_m dV = 0,$$

is a necessary condition for a solution of the nonhomogeneous problem to exist. The solution is then given by

$$w = \sum_{n=1}^{m-1} \frac{A_n}{\lambda_n - \lambda_m} w_n + \sum_{n=m+1}^{\infty} \frac{A_n}{\lambda_n - \lambda_m} w_n + C w_m, \quad A_n = \frac{1}{\|w_n\|^2} \int_V \Phi w_n dV,$$

where  $C$  is an arbitrary constant and  $\|w_n\|^2 = \int_V w_n^2 dV$ .

3°. If  $\lambda = \lambda_m$  and  $\int_V \Phi w_m dV \neq 0$ , then the boundary value problem for the nonhomogeneous equation has no solution.

*Remark 2.* If to each eigenvalue  $\lambda_n$  there are corresponding  $p_n$  mutually orthogonal eigenfunctions  $w_n^{(s)}$  ( $s = 1, \dots, p_n$ ), then the solution is written as

$$w = \sum_{n=1}^{\infty} \sum_{s=1}^{p_n} \frac{A_n^{(s)}}{\lambda_n - \lambda} w_n^{(s)}, \quad \text{where} \quad A_n^{(s)} = \frac{1}{\|w_n^{(s)}\|^2} \int_V \Phi w_n^{(s)} dV, \quad \|w_n^{(s)}\|^2 = \int_V [w_n^{(s)}]^2 dV,$$

provided that  $\lambda \neq \lambda_n$ .

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1984).

### 8.3.1-4. Solution of nonhomogeneous boundary value problems of general form.

1°. The solution of the first boundary value problem for the Helmholtz equation with the boundary condition

$$w = f(\mathbf{r}) \quad \text{for} \quad \mathbf{r} \in S$$

can be represented in the form

$$w(\mathbf{r}) = \int_V \Phi(\rho) G(\mathbf{r}, \rho) dV_\rho - \int_S f(\rho) \frac{\partial}{\partial N_\rho} G(\mathbf{r}, \rho) dS_\rho. \quad (1)$$

Here,  $\mathbf{r} = \{x, y, z\}$ ,  $\rho = \{\xi, \eta, \zeta\}$  ( $\mathbf{r} \in V$ ,  $\rho \in V$ );  $\frac{\partial}{\partial N_\rho}$  denotes the derivative along the outward normal to the surface  $S$  with respect to  $\xi, \eta, \zeta$ . The Green's function is given by the series

$$G(\mathbf{r}, \rho) = \sum_{n=1}^{\infty} \frac{w_n(\mathbf{r}) w_n(\rho)}{\|w_n\|^2 (\lambda_n - \lambda)}, \quad \lambda \neq \lambda_n, \quad (2)$$

where the  $w_n$  and  $\lambda_n$  are the eigenfunctions and eigenvalues of the homogeneous first boundary value problem.

2°. The solution of the second boundary value problem with the boundary condition

$$\frac{\partial w}{\partial N} = f(\mathbf{r}) \quad \text{for} \quad \mathbf{r} \in S$$

can be represented in the form

$$w(\mathbf{r}) = \int_V \Phi(\rho) G(\mathbf{r}, \rho) dV_\rho + \int_S f(\rho) G(\mathbf{r}, \rho) dS_\rho. \quad (3)$$

Here, the Green's function is given by the series

$$G(\mathbf{r}, \rho) = -\frac{1}{V_3 \lambda} + \sum_{n=1}^{\infty} \frac{w_n(\mathbf{r}) w_n(\rho)}{\|w_n\|^2 (\lambda_n - \lambda)}, \quad (4)$$

where  $V_3$  is the volume of the three-dimensional domain under consideration, and the  $\lambda_n$  and  $w_n$  are the positive eigenvalues and corresponding eigenfunctions of the homogeneous second boundary value problem. For clarity, the term corresponding to the zero eigenvalue  $\lambda_0 = 0$  ( $w_0 = \text{const}$ ) is singled out in (4). It is assumed that  $\lambda \neq 0$  and  $\lambda \neq \lambda_n$ .

3°. The solution of the third boundary value problem for the Helmholtz equation with the boundary condition

$$\frac{\partial w}{\partial N} + kw = f(\mathbf{r}) \quad \text{for} \quad \mathbf{r} \in S$$

is given by relation (3) in which the Green's function is defined by series (2) with the eigenfunctions  $w_n$  and eigenvalues  $\lambda_n$  of the homogeneous third boundary value problem.

4°. Let nonhomogeneous boundary conditions of various types be set on different portions  $S_i$  of the surface  $S = \sum_{i=1}^m S_i$ ,

$$\Gamma_i[w] = f_i(\mathbf{r}) \quad \text{for } \mathbf{r} \in S_i.$$

Then the solution of the corresponding mixed boundary value problem can be written as

$$w(\mathbf{r}) = \int_V \Phi(\rho) G(\mathbf{r}, \rho) dV_\rho + \sum_{i=1}^m \int_{S_i} f_i(\rho) \Lambda_i(\mathbf{r}, \rho) dS_\rho^{(i)},$$

where

$$\Lambda_i(\mathbf{r}, \rho) = \begin{cases} -\frac{\partial}{\partial N_\rho} G(\mathbf{r}, \rho) & \text{if a first-kind boundary condition is set on } S_i, \\ G(\mathbf{r}, \rho) & \text{if a second- or third-kind boundary condition is set on } S_i. \end{cases}$$

The Green's function is expressed by series (2) that involves the eigenfunctions  $w_n$  and eigenvalues  $\lambda_n$  of the homogeneous mixed boundary value problem.

#### 8.3.1-5. Boundary conditions at infinity in the case of an unbounded domain.

Below it is assumed that the function  $\Phi$  is finite or sufficiently rapidly decaying as  $r \rightarrow \infty$ .

1°. If  $\lambda < 0$  and the domain is unbounded, the additional condition that the solution must vanish at infinity is set:

$$w \rightarrow 0 \quad \text{as } r \rightarrow \infty.$$

2°. If  $\lambda > 0$ , the radiation conditions (Sommerfeld conditions) are often used at infinity. In three-dimensional problems, these conditions are expressed as

$$\lim_{r \rightarrow \infty} r w = \text{const}, \quad \lim_{r \rightarrow \infty} r \left( \frac{\partial w}{\partial r} + i\sqrt{\lambda} w \right) = 0,$$

where  $i^2 = -1$ .

The principle of limit absorption and the principle of limit amplitude are also employed to separate a single solution.

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

#### 8.3.1-6. Green's function for an infinite cylindrical domain of arbitrary cross-section.

Consider the three-dimensional Helmholtz equation

$$\Delta_3 w + \lambda w = -\Phi(\mathbf{r}) \quad (5)$$

inside an infinite cylindrical domain  $V = \{(x, y) \in D, -\infty < z < \infty\}$  with arbitrary cross-section  $D$ . On the surface of this domain, let  $S = \{(x, y) \in L, -\infty < z < \infty\}$ , where  $L$  is the boundary of  $D$ , the homogeneous boundary condition of general form

$$s \frac{\partial w}{\partial N} + k w = 0 \quad \text{for } \mathbf{r} \in S \quad (6)$$

be set, with  $sk \geq 0$ . By appropriately choosing the constants  $s$  and  $k$  in (6), one can obtain boundary conditions of the first ( $s = 0, k = 1$ ), second ( $s = 1, k = 0$ ), and third ( $sk \neq 0$ ) kind.

The Green's function of the first or third boundary value problem can be represented in the form\*

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{2} \sum_{n=1}^{\infty} \frac{u_n(x, y) u_n(\xi, \eta)}{\|u_n\|^2 \sqrt{\mu_n - \lambda}} e^{-\sqrt{\mu_n - \lambda} |z - \zeta|}, \quad \|u_n\|^2 = \int_D u_n^2(x, y) dx dy, \quad (7)$$

\* In Paragraphs 8.3.1-6 through 8.3.1-8, the cross-section  $D$  is assumed to have finite dimensions.

where the  $\mu_n$  and  $u_n$  are the eigenvalues and eigenfunctions of the corresponding two-dimensional boundary value problem in  $D$ ,

$$\begin{aligned}\Delta_2 u + \mu u &= 0 & \text{for } (x, y) \in D, \\ s \frac{\partial u}{\partial N} + k u &= 0 & \text{for } (x, y) \in L.\end{aligned}\quad (8)$$

Recall that all  $\mu_n$  are positive.

In the second boundary value problem, the zero eigenvalue  $\mu_0 = 0$  appears, and hence the summation in (7) must start with  $n = 0$ . In this case,  $u_0 = 1$  and  $\|u_0\|^2 = D_2$ , where  $D_2$  is the area of the cross-section  $D$ .

⊙ *References:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980), A. N. Tikhonov and A. A. Samarskii (1990).

### 8.3.1-7. Green's function for a semiinfinite cylindrical domain.

1°. The Green's function of the three-dimensional first boundary value problem for equation (5) in a semiinfinite cylindrical domain  $V = \{(x, y) \in D, 0 \leq z < \infty\}$  with arbitrary cross-section  $D$  is given by

$$G(x, y, z, \xi, \eta, \zeta) = \sum_{n=1}^{\infty} \frac{u_n(x, y)u_n(\xi, \eta)}{\|u_n\|^2} H_n(z, \zeta), \quad (9)$$

where

$$\begin{aligned}H_n(z, \zeta) &= \frac{1}{2\beta_n} [\exp(-\beta_n|z - \zeta|) - \exp(-\beta_n|z + \zeta|)] \\ &= \begin{cases} \frac{1}{\beta_n} \exp(-\beta_n z) \sinh(\beta_n \zeta) & \text{for } z > \zeta \geq 0, \\ \frac{1}{\beta_n} \exp(-\beta_n \zeta) \sinh(\beta_n z) & \text{for } \zeta > z \geq 0, \end{cases} \quad \beta_n = \sqrt{\mu_n - \lambda}.\end{aligned}\quad (10)$$

Relations (9) and (10) involve the eigenfunctions  $u_n$  and eigenvalues  $\mu_n$  of the two-dimensional first boundary value problem (8) with  $s = 0$  and  $k = 1$ .

2°. The Green's function of the three-dimensional second boundary value problem for equation (5) in a semiinfinite cylindrical domain  $V = \{(x, y) \in D, 0 \leq z < \infty\}$  with arbitrary cross-section  $D$  is given by

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{D_2} H_0(z, \zeta) + \sum_{n=1}^{\infty} \frac{u_n(x, y)u_n(\xi, \eta)}{\|u_n\|^2} H_n(z, \zeta), \quad (11)$$

where

$$\begin{aligned}H_n(z, \zeta) &= \frac{1}{2\beta_n} [\exp(-\beta_n|z - \zeta|) + \exp(-\beta_n|z + \zeta|)] \\ &= \begin{cases} \frac{1}{\beta_n} \exp(-\beta_n z) \cosh(\beta_n \zeta) & \text{for } z > \zeta \geq 0, \\ \frac{1}{\beta_n} \exp(-\beta_n \zeta) \cosh(\beta_n z) & \text{for } \zeta > z \geq 0, \end{cases} \quad \beta_n = \sqrt{\mu_n - \lambda}.\end{aligned}\quad (12)$$

Relations (11) and (12) involve the eigenfunctions  $u_n$  and eigenvalues  $\mu_n$  of the two-dimensional second boundary value problem (8) with  $s = 1$  and  $k = 0$ . Note that in (11) the term corresponding to the zero eigenvalue  $\mu_0 = 0$  is specially singled out;  $D_2$  is the area of the cross-section  $D$ .

3°. The Green's function of the three-dimensional third boundary value problem for equation (5) with the boundary conditions

$$\frac{\partial w}{\partial z} - k_1 w = 0 \quad \text{for } z = 0, \quad \frac{\partial w}{\partial N} + k_2 w = 0 \quad \text{for } \mathbf{r} \in S$$

in a semiinfinite cylindrical domain  $V = \{(x, y) \in D, 0 \leq z < \infty\}$  with arbitrary cross-section  $D$  and lateral surface  $S$  is given by relation (9) with

$$H_n(z, \zeta) = \begin{cases} \frac{\exp(-\beta_n z) [\beta_n \cosh(\beta_n \zeta) + k_1 \sinh(\beta_n \zeta)]}{\beta_n (\beta_n + k_1)} & \text{for } z > \zeta \geq 0, \\ \frac{\exp(-\beta_n \zeta) [\beta_n \cosh(\beta_n z) + k_1 \sinh(\beta_n z)]}{\beta_n (\beta_n + k_1)} & \text{for } \zeta > z \geq 0, \end{cases} \quad \beta_n = \sqrt{\mu_n - \lambda}. \quad (13)$$

Relations (9) and (13) involve the eigenfunctions  $u_n$  and eigenvalues  $\mu_n$  of the two-dimensional third boundary value problem (8) with  $s = 1$  and  $k = k_2$ .

4°. The Green's function of the three-dimensional mixed boundary value problem for equation (5) with a second-kind boundary condition at the end face and a first-kind boundary condition at the lateral surface is given by relations (9) and (12), where the  $\mu_n$  and  $u_n$  are the eigenvalues and eigenfunctions of the two-dimensional first boundary value problem (8) with  $s = 0$  and  $k = 1$ .

The Green's functions of other mixed boundary value problems can be constructed likewise.

### 8.3.1-8. Green's function for a cylindrical domain of finite dimensions.

1°. The Green's function of the three-dimensional first boundary value problem for equation (5) in a cylindrical domain of finite dimensions  $V = \{(x, y) \in D, 0 \leq z \leq a\}$  with arbitrary cross-section  $D$  is given by relation (9) with

$$H_n(z, \zeta) = \begin{cases} \frac{\sinh(\beta_n \zeta) \sinh[\beta_n(a - z)]}{\beta_n \sinh(\beta_n a)} & \text{for } a \geq z > \zeta \geq 0, \\ \frac{\sinh(\beta_n z) \sinh[\beta_n(a - \zeta)]}{\beta_n \sinh(\beta_n a)} & \text{for } a \geq \zeta > z \geq 0, \end{cases} \quad \beta_n = \sqrt{\mu_n - \lambda}. \quad (14)$$

Relations (9) and (14) involve the eigenfunctions  $u_n$  and eigenvalues  $\mu_n$  of the two-dimensional first boundary value problem (8) with  $s = 0$  and  $k = 1$ .

Another representation of the Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{2}{a} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{u_n(x, y) u_n(\xi, \eta) \sin(q_m z) \sin(q_m \zeta)}{\|u_n\|^2 (\mu_n + q_m^2 - \lambda)}, \quad q_m = \frac{\pi m}{a}.$$

It is a consequence of formula (2).

2°. The Green's function of the three-dimensional second boundary value problem for equation (5) in a cylindrical domain of finite dimensions  $V = \{(x, y) \in D, 0 \leq z \leq a\}$  with arbitrary cross-section  $D$  is given by relation (11) with

$$H_n(z, \zeta) = \begin{cases} \frac{\cosh(\beta_n \zeta) \cosh[\beta_n(a - z)]}{\beta_n \sinh(\beta_n a)} & \text{for } a \geq z > \zeta \geq 0, \\ \frac{\cosh(\beta_n z) \cosh[\beta_n(a - \zeta)]}{\beta_n \sinh(\beta_n a)} & \text{for } a \geq \zeta > z \geq 0, \end{cases} \quad \beta_n = \sqrt{\mu_n - \lambda}. \quad (15)$$

Relations (11) and (15) involve the eigenfunctions  $u_n$  and eigenvalues  $\mu_n$  of the two-dimensional second boundary value problem (8) with  $s = 1$  and  $k = 0$ .

Another representation of the Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{a} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{\varepsilon_m u_n(x, y) u_n(\xi, \eta) \cos(q_m z) \cos(q_m \zeta)}{\|u_n\|^2 (\mu_n + q_m^2 - \lambda)},$$

$$q_m = \frac{\pi m}{a}, \quad \varepsilon_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0, \end{cases} \quad \mu_0 = 0, \quad u_0 = 1.$$

It is a consequence of formula (4).

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

3°. The Green's function of the three-dimensional third boundary value problem for equation (5) with the boundary conditions

$$\frac{\partial w}{\partial z} - k_1 w = 0 \quad \text{at } z = 0, \quad \frac{\partial w}{\partial z} + k_2 w = 0 \quad \text{at } z = a, \quad \frac{\partial w}{\partial N} + k_3 w = 0 \quad \text{for } \mathbf{r} \in S$$

in a cylindrical domain of finite dimensions  $V = \{(x, y) \in D, 0 \leq z \leq a\}$  with arbitrary cross-section  $D$  and lateral surface  $S$  is given by relation (9) with

$$H_n(z, \zeta) = \begin{cases} \frac{[\beta_n \cosh(\beta_n \zeta) + k_1 \sinh(\beta_n \zeta)] \{ \beta_n \cosh[\beta_n(a-z)] + k_2 \sinh[\beta_n(a-z)] \}}{\beta_n [\beta_n (k_1 + k_2) \cosh(\beta_n a) + (\beta_n^2 + k_1 k_2) \sinh(\beta_n a)]} & \text{for } z > \zeta, \\ \frac{[\beta_n \cosh(\beta_n z) + k_1 \sinh(\beta_n z)] \{ \beta_n \cosh[\beta_n(a-\zeta)] + k_2 \sinh[\beta_n(a-\zeta)] \}}{\beta_n [\beta_n (k_1 + k_2) \cosh(\beta_n a) + (\beta_n^2 + k_1 k_2) \sinh(\beta_n a)]} & \text{for } z < \zeta, \end{cases} \quad (16)$$

$$\beta_n = \sqrt{\mu_n - \lambda} \quad (0 \leq z \leq a, 0 \leq \zeta \leq a).$$

Relations (9) and (16) involve the eigenfunctions  $u_n$  and eigenvalues  $\mu_n$  of the two-dimensional third boundary value problem (8) with  $s = 1$  and  $k = k_3$ .

4°. The Green's function of the three-dimensional mixed boundary value problem for equation (5) with second-kind boundary conditions at the end faces and a first-kind boundary condition at the lateral surface is given by relations (9) and (15), where the  $\mu_n$  and  $u_n$  are the eigenvalues and eigenfunctions of the two-dimensional first boundary value problem (8) with  $s = 0$  and  $k = 1$ .

The Green's function of the three-dimensional mixed boundary value problem for equation (5) with the boundary conditions

$$w = 0 \quad \text{for } z = 0, \quad \partial_z w = 0 \quad \text{for } z = a, \quad w = 0 \quad \text{for } \mathbf{r} \in S$$

in a cylindrical domain of finite dimensions  $V = \{(x, y) \in D, 0 \leq z \leq a\}$  with arbitrary cross-section  $D$  and lateral surface  $S$  is given by relation (9) with

$$H_n(z, \zeta) = \begin{cases} \frac{\sinh(\beta_n \zeta) \cosh[\beta_n(a-z)]}{\beta_n \cosh(\beta_n a)} & \text{for } a \geq z > \zeta \geq 0, \\ \frac{\sinh(\beta_n z) \cosh[\beta_n(a-\zeta)]}{\beta_n \cosh(\beta_n a)} & \text{for } a \geq \zeta > z \geq 0, \end{cases} \quad \beta_n = \sqrt{\mu_n - \lambda}. \quad (17)$$

Relations (9) and (17) involve the eigenfunctions  $u_n$  and eigenvalues  $\mu_n$  of the two-dimensional first boundary value problem (8) with  $s = 0$  and  $k = 1$ .

The Green's functions of other mixed boundary value problems can be constructed likewise.

### 8.3.2. Problems in Cartesian Coordinates

The three-dimensional nonhomogeneous Helmholtz equation in the rectangular Cartesian system of coordinates has the form

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} + \lambda w = -\Phi(x, y, z).$$

### 8.3.2-1. Particular solutions of the homogeneous equation ( $\Phi \equiv 0$ ):

$$w = (A_1 \cos kx + A_2 \sin kx)(B_1 \cos my + B_2 \sin my)(C_1 z + C_2), \quad \lambda = k^2 + m^2;$$

$$w = (A_1 \cos kx + A_2 \sin kx)(B_1 \cosh my + B_2 \sinh my)(C_1 z + C_2), \quad \lambda = k^2 - m^2;$$

$$w = (A_1 \cos kx + A_2 \sin kx)(B_1 \cos my + B_2 \sin my)(C_1 \cos nz + C_2 \sin nz), \quad \lambda = k^2 + m^2 + n^2;$$

$$w = (A_1 \cosh kx + A_2 \sinh kx)(B_1 \cos my + B_2 \sin my)(C_1 \cos nz + C_2 \sin nz), \quad \lambda = -k^2 + m^2 + n^2;$$

$$w = (A_1 \cosh kx + A_2 \sinh kx)(B_1 \cosh my + B_2 \sinh my)(C_1 \cos nz + C_2 \sin nz), \quad \lambda = -k^2 - m^2 + n^2;$$

$$w = (A_1 \cosh kx + A_2 \sinh kx)(B_1 \cosh my + B_2 \sinh my)(C_1 \cosh nz + C_2 \sinh nz), \quad \lambda = -k^2 - m^2 - n^2,$$

where  $A_1, A_2, B_1, B_2, C_1$ , and  $C_2$  are arbitrary constants.

Fundamental solutions:

$$\mathcal{E}(x, y, z) = \frac{1}{4\pi r} \exp(-kr), \quad \lambda = -k^2 < 0,$$

$$\mathcal{E}(x, y, z) = \frac{1}{4\pi r} \exp(\mp ikr), \quad \lambda = k^2 > 0,$$

where  $r = \sqrt{x^2 + y^2 + z^2}$ ,  $k > 0$ ,  $i^2 = -1$ .

### 8.3.2-2. Domain: $-\infty < x < \infty$ , $-\infty < y < \infty$ , $-\infty < z < \infty$ .

1°. Solution for  $\lambda = -k^2 < 0$ :

$$w(x, y, z) = \frac{1}{4\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) \frac{\exp[-k\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}]}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}} d\xi d\eta d\zeta.$$

2°. Solution for  $\lambda = k^2 > 0$ :

$$w(x, y, z) = \frac{1}{4\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) \frac{\exp[-ik\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}]}{\sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}} d\xi d\eta d\zeta.$$

This solution was obtained taking into account the radiation condition at infinity (see Paragraph 8.3.1-5, Item 2°).

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

### 8.3.2-3. Domain: $-\infty < x < \infty$ , $-\infty < y < \infty$ , $0 \leq z < \infty$ . First boundary value problem.

A half-space is considered. A boundary condition is prescribed:

$$w = f(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$w(x, y, z) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta + \int_0^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta.$$

Green's function for  $\lambda = -k^2 < 0$ :

$$G(x, y, z, \xi, \eta, \zeta) = \frac{\exp(-k\mathcal{R}_1)}{4\pi\mathcal{R}_1} - \frac{\exp(-k\mathcal{R}_2)}{4\pi\mathcal{R}_2},$$

$$\mathcal{R}_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}, \quad \mathcal{R}_2 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2}.$$

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).



**8.3.2-4. Domain:**  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z < \infty$ . **Second boundary value problem.**

A half-space is considered. A boundary condition is prescribed:

$$\partial_z w = f(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(x, y, z) = & - \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(\xi, \eta) G(x, y, z, \xi, \eta, 0) d\xi d\eta \\ & + \int_0^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function for  $\lambda = -k^2 < 0$ :

$$G(x, y, z, \xi, \eta, \zeta) = \frac{\exp(-k\mathcal{R}_1)}{4\pi\mathcal{R}_1} + \frac{\exp(-k\mathcal{R}_2)}{4\pi\mathcal{R}_2},$$

$$\mathcal{R}_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}, \quad \mathcal{R}_2 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2}.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.3.2-5. Domain:**  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . **First boundary value problem.**

A dihedral angle is considered. Boundary conditions are prescribed:

$$w = f_1(x, z) \quad \text{at} \quad y = 0, \quad w = f_2(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_0^{\infty} \int_{-\infty}^{\infty} f_1(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\xi d\zeta \\ & + \int_0^{\infty} \int_{-\infty}^{\infty} f_2(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta \\ & + \int_0^{\infty} \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function for  $\lambda = -k^2 < 0$ :

$$G(x, y, z, \xi, \eta, \zeta) = \frac{\exp(-k\mathcal{R}_1)}{4\pi\mathcal{R}_1} - \frac{\exp(-k\mathcal{R}_2)}{4\pi\mathcal{R}_2} - \frac{\exp(-k\mathcal{R}_3)}{4\pi\mathcal{R}_3} + \frac{\exp(-k\mathcal{R}_4)}{4\pi\mathcal{R}_4},$$

$$\mathcal{R}_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}, \quad \mathcal{R}_2 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2},$$

$$\mathcal{R}_3 = \sqrt{(x-\xi)^2 + (y+\eta)^2 + (z-\zeta)^2}, \quad \mathcal{R}_4 = \sqrt{(x-\xi)^2 + (y+\eta)^2 + (z+\zeta)^2}.$$

**8.3.2-6. Domain:**  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ ,  $0 \leq z < \infty$ . **Second boundary value problem.**

A dihedral angle is considered. Boundary conditions are prescribed:

$$\partial_y w = f_1(x, z) \quad \text{at} \quad y = 0, \quad \partial_z w = f_2(x, y) \quad \text{at} \quad z = 0.$$

Solution:

$$\begin{aligned} w(x, y, z) = & - \int_0^{\infty} \int_{-\infty}^{\infty} f_1(\xi, \zeta) G(x, y, z, \xi, 0, \zeta) d\xi d\zeta \\ & - \int_0^{\infty} \int_{-\infty}^{\infty} f_2(\xi, \eta) G(x, y, z, \xi, \eta, 0) d\xi d\eta \\ & + \int_0^{\infty} \int_0^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function for  $\lambda = -k^2 < 0$ :

$$G(x, y, z, \xi, \eta, \zeta) = \frac{\exp(-k\mathcal{R}_1)}{4\pi\mathcal{R}_1} + \frac{\exp(-k\mathcal{R}_2)}{4\pi\mathcal{R}_2} + \frac{\exp(-k\mathcal{R}_3)}{4\pi\mathcal{R}_3} + \frac{\exp(-k\mathcal{R}_4)}{4\pi\mathcal{R}_4},$$

$$\mathcal{R}_1 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta)^2}, \quad \mathcal{R}_2 = \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta)^2},$$

$$\mathcal{R}_3 = \sqrt{(x-\xi)^2 + (y+\eta)^2 + (z-\zeta)^2}, \quad \mathcal{R}_4 = \sqrt{(x-\xi)^2 + (y+\eta)^2 + (z+\zeta)^2}.$$

8.3.2-7. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq a$ . First boundary value problem.

An infinite layer is considered. Boundary conditions are prescribed:

$$w = f_1(x, y) \quad \text{at} \quad z = 0, \quad w = f_2(x, y) \quad \text{at} \quad z = a.$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_1(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\xi d\eta \\ & - \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_2(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=a} d\xi d\eta \\ & + \int_0^a \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function for  $\lambda = -k^2 < 0$ :

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) = & \sum_{n=-\infty}^{\infty} \left[ \frac{\exp(-k\mathcal{R}_{1n})}{4\pi\mathcal{R}_{1n}} - \frac{\exp(-k\mathcal{R}_{2n})}{4\pi\mathcal{R}_{2n}} \right], \\ \mathcal{R}_{1n} = & \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta-2na)^2}, \\ \mathcal{R}_{2n} = & \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta-2na)^2}. \end{aligned}$$

8.3.2-8. Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $0 \leq z \leq a$ . Second boundary value problem.

An infinite layer is considered. Boundary conditions are prescribed:

$$\partial_z w = f_1(x, y) \quad \text{at} \quad z = 0, \quad \partial_z w = f_2(x, y) \quad \text{at} \quad z = a.$$

Solution:

$$\begin{aligned} w(x, y, z) = & - \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_1(\xi, \eta) G(x, y, z, \xi, \eta, 0) d\xi d\eta \\ & + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f_2(\xi, \eta) G(x, y, z, \xi, \eta, a) d\xi d\eta \\ & + \int_0^a \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\xi d\eta d\zeta. \end{aligned}$$

Green's function for  $\lambda = -k^2 < 0$ :

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) = & \sum_{n=-\infty}^{\infty} \left[ \frac{\exp(-k\mathcal{R}_{1n})}{4\pi\mathcal{R}_{1n}} + \frac{\exp(-k\mathcal{R}_{2n})}{4\pi\mathcal{R}_{2n}} \right], \\ \mathcal{R}_{1n} = & \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z-\zeta-2na)^2}, \\ \mathcal{R}_{2n} = & \sqrt{(x-\xi)^2 + (y-\eta)^2 + (z+\zeta-2na)^2}. \end{aligned}$$

☉ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

8.3.2-9. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . First boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad w = f_2(y, z) \quad \text{at} \quad x = a, \\ w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad w = f_4(x, z) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z) = & \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\
 & - \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=a} d\zeta d\eta \\
 & + \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\zeta d\xi \\
 & - \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=b} d\zeta d\xi \\
 & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi.
 \end{aligned}$$

Green's function:

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta) = & \frac{2}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \exp(-\beta_{nm}|z - \zeta|), \\
 p_n = & \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}.
 \end{aligned}$$

⊙ Reference: A. N. Tikhonov and A. A. Samarskii (1990).

8.3.2-10. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . Second boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned}
 \partial_x w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad \partial_x w = f_2(y, z) \quad \text{at} \quad x = a, \\
 \partial_y w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad \partial_y w = f_4(x, z) \quad \text{at} \quad y = b.
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z) = & - \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) G(x, y, z, 0, \eta, \zeta) d\zeta d\eta + \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) G(x, y, z, a, \eta, \zeta) d\zeta d\eta \\
 & - \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) G(x, y, z, \xi, 0, \zeta) d\zeta d\xi + \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\
 & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi.
 \end{aligned}$$

Green's function:

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta) = & \frac{1}{2ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_n A_m}{\beta_{nm}} \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) \exp(-\beta_{nm}|z - \zeta|), \\
 p_n = & \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}
 \end{aligned}$$

⊙ Reference: A. N. Tikhonov and A. A. Samarskii (1990).

8.3.2-11. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . Third boundary value problem.

An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned}
 \partial_x w - k_1 w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad \partial_x w + k_2 w = f_2(y, z) \quad \text{at} \quad x = a, \\
 \partial_y w - k_3 w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad \partial_y w + k_4 w = f_4(x, z) \quad \text{at} \quad y = b.
 \end{aligned}$$

The solution  $w(x, y, z)$  is determined by the formula in Paragraph 8.3.2-10 where

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{u_{nm}(x, y) u_{nm}(\xi, \eta)}{\|u_{nm}\|^2 \beta_{nm}} \exp(-\beta_{nm}|z - \zeta|).$$

Here,

$$w_{nm}(x, y) = (\mu_n \cos \mu_n x + k_1 \sin \mu_n x)(\nu_m \cos \nu_m y + k_3 \sin \nu_m y), \quad \beta_{nm} = \sqrt{\mu_n^2 + \nu_m^2} - \lambda,$$

$$\|w_{nm}\|^2 = \frac{1}{4}(\mu_n^2 + k_1^2)(\nu_m^2 + k_3^2) \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right] \left[ b + \frac{(k_3 + k_4)(\nu_m^2 + k_3 k_4)}{(\nu_m^2 + k_3^2)(\nu_m^2 + k_4^2)} \right],$$

where the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}, \quad \tan(\nu b) = \frac{(k_3 + k_4)\nu}{\nu^2 - k_3 k_4}.$$

8.3.2-12. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $-\infty < z < \infty$ . Mixed boundary value problems.

1°. An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad \partial_x w = f_2(y, z) \quad \text{at} \quad x = a, \\ w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad \partial_y w = f_4(x, z) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z) = & \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\ & + \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) G(x, y, z, a, \eta, \zeta) d\zeta d\eta \\ & + \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\zeta d\xi \\ & + \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\ & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{2}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \exp(-\beta_{nm}|z - \zeta|),$$

$$p_n = \frac{(2n+1)\pi}{2a}, \quad q_m = \frac{(2m+1)\pi}{2b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2} - \lambda.$$

2°. An infinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad w = f_2(y, z) \quad \text{at} \quad x = a, \\ \partial_y w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad \partial_y w = f_4(x, z) \quad \text{at} \quad y = b. \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z) = & \int_0^b \int_{-\infty}^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\
 & - \int_0^b \int_{-\infty}^{\infty} f_2(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=a} d\zeta d\eta \\
 & - \int_0^a \int_{-\infty}^{\infty} f_3(\xi, \zeta) G(x, y, z, \xi, 0, \zeta) d\zeta d\xi \\
 & + \int_0^a \int_{-\infty}^{\infty} f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\
 & + \int_0^a \int_0^b \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi.
 \end{aligned}$$

Green's function:

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta) = & \frac{1}{ab} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \frac{A_m}{\beta_{nm}} \sin(p_n x) \cos(q_m y) \sin(p_n \xi) \cos(q_m \eta) \exp(-\beta_{nm}|z - \zeta|), \\
 p_n = & \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}, \quad A_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases}
 \end{aligned}$$

8.3.2-13. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z < \infty$ . First boundary value problem.

A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned}
 w = f_1(y, z) & \quad \text{at } x = 0, & w = f_2(y, z) & \quad \text{at } x = a, \\
 w = f_3(x, z) & \quad \text{at } y = 0, & w = f_4(x, z) & \quad \text{at } y = b, \\
 w = f_5(x, y) & \quad \text{at } z = 0.
 \end{aligned}$$

Solution:

$$\begin{aligned}
 w(x, y, z) = & \int_0^b \int_0^{\infty} f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\
 & - \int_0^b \int_0^{\infty} f_2(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=a} d\zeta d\eta \\
 & + \int_0^a \int_0^{\infty} f_3(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\zeta d\xi \\
 & - \int_0^a \int_0^{\infty} f_4(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=b} d\zeta d\xi \\
 & + \int_0^a \int_0^b f_5(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\eta d\xi \\
 & + \int_0^a \int_0^b \int_0^{\infty} \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi.
 \end{aligned}$$

Green's function:

$$\begin{aligned}
 G(x, y, z, \xi, \eta, \zeta) = & \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) H_{nm}(z, \zeta), \\
 p_n = & \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}, \\
 H_{nm}(z, \zeta) = & \begin{cases} \exp(-\beta_{nm}z) \sinh(\beta_{nm}\zeta) & \text{for } z > \zeta \geq 0, \\ \exp(-\beta_{nm}\zeta) \sinh(\beta_{nm}z) & \text{for } \zeta > z \geq 0. \end{cases}
 \end{aligned}$$

**8.3.2-14. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z < \infty$ . Second boundary value problem.**

A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned}\partial_x w &= f_1(y, z) & \text{at } x = 0, & & \partial_x w &= f_2(y, z) & \text{at } x = a, \\ \partial_y w &= f_3(x, z) & \text{at } y = 0, & & \partial_y w &= f_4(x, z) & \text{at } y = b, \\ \partial_z w &= f_5(x, y) & \text{at } z = 0.\end{aligned}$$

Solution:

$$\begin{aligned}w(x, y, z) &= \int_0^a \int_0^b \int_0^\infty \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi \\ &\quad - \int_0^b \int_0^\infty f_1(\eta, \zeta) G(x, y, z, 0, \eta, \zeta) d\zeta d\eta + \int_0^b \int_0^\infty f_2(\eta, \zeta) G(x, y, z, a, \eta, \zeta) d\zeta d\eta \\ &\quad - \int_0^a \int_0^\infty f_3(\xi, \zeta) G(x, y, z, \xi, 0, \zeta) d\zeta d\xi + \int_0^a \int_0^\infty f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\ &\quad - \int_0^a \int_0^b f_5(\xi, \eta) G(x, y, z, \xi, \eta, 0) d\eta d\xi.\end{aligned}$$

Green's function:

$$\begin{aligned}G(x, y, z, \xi, \eta, \zeta) &= \frac{1}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_n A_m}{\beta_{nm}} \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) H_{nm}(z, \zeta), \\ p_n &= \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \\ H_{nm}(z, \zeta) &= \begin{cases} \exp(-\beta_{nm} z) \cosh(\beta_{nm} \zeta) & \text{for } z > \zeta \geq 0, \\ \exp(-\beta_{nm} \zeta) \cosh(\beta_{nm} z) & \text{for } \zeta > z \geq 0. \end{cases}\end{aligned}$$

**8.3.2-15. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z < \infty$ . Third boundary value problem.**

A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned}\partial_x w - k_1 w &= f_1(y, z) & \text{at } x = 0, & & \partial_x w + k_2 w &= f_2(y, z) & \text{at } x = a, \\ \partial_y w - k_3 w &= f_3(x, z) & \text{at } y = 0, & & \partial_y w + k_4 w &= f_4(x, z) & \text{at } y = b, \\ \partial_z w - k_5 w &= f_5(x, y) & \text{at } z = 0.\end{aligned}$$

The solution  $w(x, y, z)$  is determined by the formula in Paragraph 8.3.2-14 where

$$G(x, y, z, \xi, \eta, \zeta) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{u_{nm}(x, y) u_{nm}(\xi, \eta)}{\|u_{nm}\|^2} H_{nm}(z, \zeta).$$

Here,

$$\begin{aligned}w_{nm}(x, y) &= (\mu_n \cos \mu_n x + k_1 \sin \mu_n x)(\nu_m \cos \nu_m y + k_3 \sin \nu_m y), \\ \|w_{nm}\|^2 &= \frac{1}{4} (\mu_n^2 + k_1^2)(\nu_m^2 + k_3^2) \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right] \left[ b + \frac{(k_3 + k_4)(\nu_m^2 + k_3 k_4)}{(\nu_m^2 + k_3^2)(\nu_m^2 + k_4^2)} \right], \\ H_{nm}(z, \zeta) &= \begin{cases} \frac{\exp(-\beta_{nm} z) [\beta_{nm} \cosh(\beta_{nm} \zeta) + k_5 \sinh(\beta_{nm} \zeta)]}{\beta_{nm}(\beta_{nm} + k_5)} & \text{for } z > \zeta, \\ \frac{\exp(-\beta_{nm} \zeta) [\beta_{nm} \cosh(\beta_{nm} z) + k_5 \sinh(\beta_{nm} z)]}{\beta_{nm}(\beta_{nm} + k_5)} & \text{for } \zeta > z, \end{cases} \quad \beta_{nm} = \sqrt{\mu_n^2 + \nu_m^2 - \lambda},\end{aligned}$$

where the  $\mu_n$  and  $\nu_m$  are positive roots of the transcendental equations

$$\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}, \quad \tan(\nu b) = \frac{(k_3 + k_4)\nu}{\nu^2 - k_3 k_4}.$$

1°. A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y, z) & \text{at } x &= 0, & w &= f_2(y, z) & \text{at } x &= a, \\ w &= f_3(x, z) & \text{at } y &= 0, & w &= f_4(x, z) & \text{at } y &= b, \\ \partial_z w &= f_5(x, y) & \text{at } z &= 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z) &= \int_0^b \int_0^\infty f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=0} d\zeta d\eta \\ &\quad - \int_0^b \int_0^\infty f_2(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(x, y, z, \xi, \eta, \zeta) \right]_{\xi=a} d\zeta d\eta \\ &\quad + \int_0^a \int_0^\infty f_3(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=0} d\zeta d\xi \\ &\quad - \int_0^a \int_0^\infty f_4(\xi, \zeta) \left[ \frac{\partial}{\partial \eta} G(x, y, z, \xi, \eta, \zeta) \right]_{\eta=b} d\zeta d\xi \\ &\quad - \int_0^a \int_0^b f_5(\xi, \eta) G(x, y, z, \xi, \eta, 0) d\eta d\xi \\ &\quad + \int_0^a \int_0^b \int_0^\infty \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta) &= \frac{4}{ab} \sum_{n=1}^\infty \sum_{m=1}^\infty \frac{1}{\beta_{nm}} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) H_{nm}(z, \zeta), \\ p_n &= \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2} - \lambda, \\ H_{nm}(z, \zeta) &= \begin{cases} \exp(-\beta_{nm} z) \cosh(\beta_{nm} \zeta) & \text{for } z > \zeta \geq 0, \\ \exp(-\beta_{nm} \zeta) \cosh(\beta_{nm} z) & \text{for } \zeta > z \geq 0. \end{cases} \end{aligned}$$

2°. A semiinfinite cylindrical domain of a rectangular cross-section is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w &= f_1(y, z) & \text{at } x &= 0, & \partial_x w &= f_2(y, z) & \text{at } x &= a, \\ \partial_y w &= f_3(x, z) & \text{at } y &= 0, & \partial_y w &= f_4(x, z) & \text{at } y &= b, \\ w &= f_5(x, y) & \text{at } z &= 0. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z) &= \int_0^a \int_0^b \int_0^\infty \Phi(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta) d\zeta d\eta d\xi \\ &\quad - \int_0^b \int_0^\infty f_1(\eta, \zeta) G(x, y, z, 0, \eta, \zeta) d\zeta d\eta + \int_0^b \int_0^\infty f_2(\eta, \zeta) G(x, y, z, a, \eta, \zeta) d\zeta d\eta \\ &\quad - \int_0^a \int_0^\infty f_3(\xi, \zeta) G(x, y, z, \xi, 0, \zeta) d\zeta d\xi + \int_0^a \int_0^\infty f_4(\xi, \zeta) G(x, y, z, \xi, b, \zeta) d\zeta d\xi \\ &\quad + \int_0^a \int_0^b f_5(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(x, y, z, \xi, \eta, \zeta) \right]_{\zeta=0} d\eta d\xi. \end{aligned}$$

Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{1}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{A_n A_m}{\beta_{nm}} \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) H_{nm}(z, \zeta).$$

Here,

$$p_n = \frac{n\pi}{a}, \quad q_m = \frac{m\pi}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

$$H_{nm}(z, \zeta) = \begin{cases} \exp(-\beta_{nm} z) \sinh(\beta_{nm} \zeta) & \text{for } z > \zeta \geq 0, \\ \exp(-\beta_{nm} \zeta) \sinh(\beta_{nm} z) & \text{for } \zeta > z \geq 0. \end{cases}$$

► Paragraphs 8.3.2-17 through 8.3.2-23 present only the eigenvalues and eigenfunctions of homogeneous boundary value problems for the homogeneous Helmholtz equation (with  $\Phi \equiv 0$ ). The solutions of the corresponding nonhomogeneous boundary value problems (with  $\Phi \not\equiv 0$ ) can be constructed by the relations specified in Paragraphs 8.3.1-4 and 8.3.1-8.

8.3.2-17. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . First boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_1(y, z) & \text{at } x = 0, & & w &= f_2(y, z) & \text{at } x = a, \\ w &= f_3(x, z) & \text{at } y = 0, & & w &= f_4(x, z) & \text{at } y = b, \\ w &= f_5(x, y) & \text{at } z = 0, & & w &= f_6(x, y) & \text{at } z = c. \end{aligned}$$

1°. Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \pi^2 \left( \frac{n^2}{a^2} + \frac{m^2}{b^2} + \frac{k^2}{c^2} \right); \quad n, m, k = 1, 2, 3, \dots$$

Eigenfunctions and the norm squared:

$$w_{nmk} = \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m y}{b}\right) \sin\left(\frac{\pi k z}{c}\right), \quad \|w_{nmk}\|^2 = \frac{abc}{8}.$$

2°. A double-series representation of the Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{4}{ab} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) H_{nm}(z, \zeta),$$

$$H_{nm}(z, \zeta) = \begin{cases} \frac{\sinh(\beta_{nm} \zeta) \sinh[\beta_{nm}(c - z)]}{\beta_{nm} \sinh(\beta_{nm} c)} & \text{for } c \geq z > \zeta \geq 0, \\ \frac{\sinh(\beta_{nm} z) \sinh[\beta_{nm}(c - \zeta)]}{\beta_{nm} \sinh(\beta_{nm} c)} & \text{for } c \geq \zeta > z \geq 0, \end{cases}$$

$$p_n = \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}.$$

This relation can be used to obtain two other representations of the Green's function with the aid of the cyclic permutations of triples:

$$\begin{array}{ccc} & (x, \xi, a) & \\ \nearrow & & \searrow \\ (z, \zeta, c) & \longleftarrow & (y, \eta, b) \end{array}$$

A triple series representation of the Green's function:

$$G(x, y, z, \xi, \eta, \zeta) = \frac{8}{abc} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{\sin(p_n x) \sin(q_m y) \sin(s_k z) \sin(p_n \xi) \sin(q_m \eta) \sin(s_k \zeta)}{p_n^2 + q_m^2 + s_k^2 - \lambda},$$

$$p_n = \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}, \quad s_k = \frac{\pi k}{c}.$$

© References: V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).



**8.3.2-18. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . Second boundary value problem.**

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned}\partial_x w &= f_1(y, z) & \text{at } x = 0, & \quad \partial_x w = f_2(y, z) & \text{at } x = a, \\ \partial_y w &= f_3(x, z) & \text{at } y = 0, & \quad \partial_y w = f_4(x, z) & \text{at } y = b, \\ \partial_z w &= f_5(x, y) & \text{at } z = 0, & \quad \partial_z w = f_6(x, y) & \text{at } z = c.\end{aligned}$$

1°. Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \pi^2 \left( \frac{n^2}{a^2} + \frac{m^2}{b^2} + \frac{k^2}{c^2} \right); \quad n, m, k = 0, 1, 2, \dots$$

Eigenfunctions:

$$w_{nmk} = \cos\left(\frac{\pi n x}{a}\right) \cos\left(\frac{\pi m y}{b}\right) \cos\left(\frac{\pi k z}{c}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}\|^2 = \frac{abc}{8} (1 + \delta_{n0})(1 + \delta_{m0})(1 + \delta_{k0}), \quad \delta_{n0} = \begin{cases} 1 & \text{for } n = 0, \\ 0 & \text{for } n \neq 0. \end{cases}$$

2°. A double series representation of the Green's function:

$$\begin{aligned}G(x, y, z, \xi, \eta, \zeta) &= \frac{1}{ab} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \varepsilon_n \varepsilon_m \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) H_{nm}(z, \zeta), \\ H_{nm}(z, \zeta) &= \begin{cases} \frac{\cosh(\beta_{nm} \zeta) \cosh[\beta_{nm}(c - z)]}{\beta_{nm} \sinh(\beta_{nm} c)} & \text{for } c \geq z > \zeta \geq 0, \\ \frac{\cosh(\beta_{nm} z) \cosh[\beta_{nm}(c - \zeta)]}{\beta_{nm} \sinh(\beta_{nm} c)} & \text{for } c \geq \zeta > z \geq 0, \end{cases} \\ p_n &= \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}, \quad \beta_{nm} = \sqrt{p_n^2 + q_m^2 - \lambda}, \quad \varepsilon_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}\end{aligned}$$

This relation can be used to obtain two other representations of the Green's function with the aid of the cyclic permutations:

$$\begin{array}{ccc} & (x, \xi, a) & \\ \nearrow & & \searrow \\ (z, \zeta, c) & \longleftarrow & (y, \eta, b) \end{array}$$

A triple series representation of the Green's function:

$$\begin{aligned}G(x, y, z, \xi, \eta, \zeta) &= \frac{1}{abc} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sum_{k=0}^{\infty} \frac{\varepsilon_n \varepsilon_m \varepsilon_k \cos(p_n x) \cos(q_m y) \cos(s_k z) \cos(p_n \xi) \cos(q_m \eta) \cos(s_k \zeta)}{p_n^2 + q_m^2 + s_k^2 - \lambda}, \\ p_n &= \frac{\pi n}{a}, \quad q_m = \frac{\pi m}{b}, \quad s_k = \frac{\pi k}{c}.\end{aligned}$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.3.2-19. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . Third boundary value problem.**

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned}\partial_x w - k_1 w &= f_1(y, z) & \text{at } x = 0, & \quad \partial_x w + k_2 w = f_2(y, z) & \text{at } x = a, \\ \partial_y w - k_3 w &= f_3(x, z) & \text{at } y = 0, & \quad \partial_y w + k_4 w = f_4(x, z) & \text{at } y = b, \\ \partial_z w - k_5 w &= f_5(x, y) & \text{at } z = 0, & \quad \partial_z w + k_6 w = f_6(x, y) & \text{at } z = c.\end{aligned}$$

Eigenvalues of the homogeneous problem:

$$\lambda_{nml} = \mu_n^2 + \nu_m^2 + \sigma_l^2; \quad n, m, l = 1, 2, 3, \dots$$

Here, the  $\mu_n$ ,  $\nu_m$ , and  $\sigma_l$  are positive roots of the transcendental equations

$$\tan(\mu a) = \frac{(k_1 + k_2)\mu}{\mu^2 - k_1 k_2}, \quad \tan(\nu b) = \frac{(k_3 + k_4)\nu}{\nu^2 - k_3 k_4}, \quad \tan(\sigma c) = \frac{(k_5 + k_6)\sigma}{\sigma^2 - k_5 k_6}.$$

Eigenfunctions:

$$w_{nml} = \frac{1}{A_n B_m C_l} (\mu_n \cos \mu_n x + k_1 \sin \mu_n x) (\nu_m \cos \nu_m y + k_3 \sin \nu_m y) (\sigma_l \cos \sigma_l z + k_5 \sin \sigma_l z),$$

$$A_n = \sqrt{\mu_n^2 + k_1^2}, \quad B_m = \sqrt{\nu_m^2 + k_3^2}, \quad C_l = \sqrt{\sigma_l^2 + k_5^2}.$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nml}\|^2 = \frac{1}{8} \left[ a + \frac{(k_1 + k_2)(\mu_n^2 + k_1 k_2)}{(\mu_n^2 + k_1^2)(\mu_n^2 + k_2^2)} \right] \left[ b + \frac{(k_3 + k_4)(\nu_m^2 + k_3 k_4)}{(\nu_m^2 + k_3^2)(\nu_m^2 + k_4^2)} \right] \left[ c + \frac{(k_5 + k_6)(\sigma_l^2 + k_5 k_6)}{(\sigma_l^2 + k_5^2)(\sigma_l^2 + k_6^2)} \right].$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

8.3.2-20. Domain:  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ . Mixed boundary value problems.

1°. A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad w = f_2(y, z) \quad \text{at} \quad x = a, \\ w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad w = f_4(x, z) \quad \text{at} \quad y = b, \\ \partial_z w = f_5(x, y) \quad \text{at} \quad z = 0, & \quad \partial_z w = f_6(x, y) \quad \text{at} \quad z = c. \end{aligned}$$

Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \pi^2 \left( \frac{n^2}{a^2} + \frac{m^2}{b^2} + \frac{k^2}{c^2} \right); \quad n, m = 1, 2, 3, \dots; \quad k = 0, 1, 2, \dots$$

Eigenfunctions:

$$w_{nmk} = \sin\left(\frac{\pi n x}{a}\right) \sin\left(\frac{\pi m y}{b}\right) \cos\left(\frac{\pi k z}{c}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}\|^2 = \frac{abc}{8} (1 + \delta_{k0}), \quad \delta_{k0} = \begin{cases} 1 & \text{for } k = 0, \\ 0 & \text{for } k \neq 0. \end{cases}$$

2°. A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y, z) \quad \text{at} \quad x = 0, & \quad w = f_2(y, z) \quad \text{at} \quad x = a, \\ \partial_y w = f_3(x, z) \quad \text{at} \quad y = 0, & \quad \partial_y w = f_4(x, z) \quad \text{at} \quad y = b, \\ \partial_z w = f_5(x, y) \quad \text{at} \quad z = 0, & \quad \partial_z w = f_6(x, y) \quad \text{at} \quad z = c. \end{aligned}$$

Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \pi^2 \left( \frac{n^2}{a^2} + \frac{m^2}{b^2} + \frac{k^2}{c^2} \right); \quad n = 1, 2, 3, \dots; \quad m, k = 0, 1, 2, \dots$$

Eigenfunctions:

$$w_{nmk} = \sin\left(\frac{\pi n x}{a}\right) \cos\left(\frac{\pi m y}{b}\right) \cos\left(\frac{\pi k z}{c}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}\|^2 = \frac{abc}{8} (1 + \delta_{m0})(1 + \delta_{k0}), \quad \delta_{m0} = \begin{cases} 1 & \text{for } m = 0, \\ 0 & \text{for } m \neq 0. \end{cases}$$

**8.3.2-21. Domain:  $0 \leq x \leq a, 0 \leq y \leq x, 0 \leq z \leq c$ . First boundary value problem.**

A right prism whose base is an isosceles right-angled triangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y, z) \quad \text{at} \quad x = 0, \quad w = f_2(x, z) \quad \text{at} \quad y = 0, \quad w = f_3(x, z) \quad \text{at} \quad y = x, \\ w = f_4(x, y) \quad \text{at} \quad z = 0, \quad w = f_5(x, y) \quad \text{at} \quad z = c. \end{aligned}$$

Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \frac{\pi^2}{a^2} [(n+m)^2 + m^2] + \frac{\pi^2 k^2}{c^2}; \quad n, m, k = 1, 2, 3, \dots$$

Eigenfunctions:

$$w_{nmk} = \left\{ \sin \left[ \frac{\pi}{a} (n+m)x \right] \sin \left( \frac{\pi}{a} my \right) - (-1)^n \sin \left( \frac{\pi}{a} mx \right) \sin \left[ \frac{\pi}{a} (n+m)y \right] \right\} \sin \left( \frac{\pi kz}{c} \right).$$

**8.3.2-22. Domain:  $0 \leq x \leq a, 0 \leq y \leq x, 0 \leq z \leq c$ . Second boundary value problem.**

A right prism whose base is an isosceles right-angled triangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w = f_1(y, z) \quad \text{at} \quad x = 0, \quad \partial_y w = f_2(x, z) \quad \text{at} \quad y = 0, \quad \partial_N w = f_3(x, z) \quad \text{at} \quad y = x, \\ \partial_z w = f_4(x, y) \quad \text{at} \quad z = 0, \quad \partial_z w = f_5(x, y) \quad \text{at} \quad z = c, \end{aligned}$$

where  $\partial_N w = \mathbf{N} \cdot \nabla w = \frac{1}{\sqrt{2}}(\partial_x w + \partial_y w)$ .

Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \frac{\pi^2}{a^2} [(n+m)^2 + m^2] + \frac{\pi^2 k^2}{c^2}; \quad n, m, k = 0, 1, 2, \dots$$

Eigenfunctions:

$$w_{nmk} = \left\{ \cos \left[ \frac{\pi}{a} (n+m)x \right] \cos \left( \frac{\pi}{a} my \right) - (-1)^n \cos \left( \frac{\pi}{a} mx \right) \cos \left[ \frac{\pi}{a} (n+m)y \right] \right\} \cos \left( \frac{\pi kz}{c} \right).$$

**8.3.2-23. Domain:  $0 \leq x \leq a, 0 \leq y \leq x, 0 \leq z \leq c$ . Mixed boundary value problems.**

1°. A right prism whose base is an isosceles right-angled triangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = f_1(y, z) \quad \text{at} \quad x = 0, \quad w = f_2(x, z) \quad \text{at} \quad y = 0, \quad w = f_3(x, z) \quad \text{at} \quad y = x, \\ \partial_z w = f_4(x, y) \quad \text{at} \quad z = 0, \quad \partial_z w = f_5(x, y) \quad \text{at} \quad z = c. \end{aligned}$$

Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \frac{\pi^2}{a^2} [(n+m)^2 + m^2] + \frac{\pi^2 k^2}{c^2}; \quad n, m = 1, 2, 3, \dots; \quad k = 0, 1, 2, \dots$$

Eigenfunctions:

$$w_{nmk} = \left\{ \sin \left[ \frac{\pi}{a} (n+m)x \right] \sin \left( \frac{\pi}{a} my \right) - (-1)^n \sin \left( \frac{\pi}{a} mx \right) \sin \left[ \frac{\pi}{a} (n+m)y \right] \right\} \cos \left( \frac{\pi kz}{c} \right).$$

2°. A right prism whose base is an isosceles right-angled triangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_x w = f_1(y, z) \quad \text{at} \quad x = 0, \quad \partial_y w = f_2(x, z) \quad \text{at} \quad y = 0, \quad \partial_N w = f_3(x, z) \quad \text{at} \quad y = x, \\ w = f_4(x, y) \quad \text{at} \quad z = 0, \quad w = f_5(x, y) \quad \text{at} \quad z = c. \end{aligned}$$

Eigenvalues of the homogeneous problem:

$$\lambda_{nmk} = \frac{\pi^2}{a^2} [(n+m)^2 + m^2] + \frac{\pi^2 k^2}{c^2}; \quad n, m = 0, 1, 2, \dots; \quad k = 1, 2, 3, \dots$$

Eigenfunctions:

$$w_{nmk} = \left\{ \cos \left[ \frac{\pi}{a} (n+m)x \right] \cos \left( \frac{\pi}{a} my \right) - (-1)^n \cos \left( \frac{\pi}{a} mx \right) \cos \left[ \frac{\pi}{a} (n+m)y \right] \right\} \sin \left( \frac{\pi kz}{c} \right).$$

### 8.3.3. Problems in Cylindrical Coordinates

The three-dimensional nonhomogeneous Helmholtz equation in the cylindrical coordinate system is written as

$$\frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \varphi^2} + \frac{\partial^2 w}{\partial z^2} + \lambda w = -\Phi(r, \varphi, z), \quad r = \sqrt{x^2 + y^2}.$$

#### 8.3.3-1. Particular solutions of the homogeneous equation ( $\Phi \equiv 0$ ):

$$\begin{aligned} w &= [AJ_0(r\sqrt{\lambda}) + BY_0(r\sqrt{\lambda})](C_1\varphi + D_1)(C_2z + D_2), \\ w &= J_m(r\sqrt{\lambda - \mu^2})(A \cos m\varphi + B \sin m\varphi)(C \cos \mu z + D \sin \mu z), \quad \lambda > \mu^2, \\ w &= Y_m(r\sqrt{\lambda - \mu^2})(A \cos m\varphi + B \sin m\varphi)(C \cos \mu z + D \sin \mu z), \quad \lambda > \mu^2, \\ w &= J_m(r\sqrt{\lambda + \mu^2})(A \cos m\varphi + B \sin m\varphi)(C \cosh \mu z + D \sinh \mu z), \quad \lambda > -\mu^2, \\ w &= Y_m(r\sqrt{\lambda + \mu^2})(A \cos m\varphi + B \sin m\varphi)(C \cosh \mu z + D \sinh \mu z), \quad \lambda > -\mu^2, \\ w &= I_m(r\sqrt{\mu^2 - \lambda})(A \cos m\varphi + B \sin m\varphi)(C \cos \mu z + D \sin \mu z), \quad \lambda < \mu^2, \\ w &= Y_m(r\sqrt{\mu^2 - \lambda})(A \cos m\varphi + B \sin m\varphi)(C \cos \mu z + D \sin \mu z), \quad \lambda < \mu^2, \end{aligned}$$

where  $m = 0, 1, 2, \dots$ ;  $A, B, C, D, C_1, C_2, D_1, D_2$ , and  $\mu$  are arbitrary constants; the  $J_m(\xi)$  and  $Y_m(\xi)$  are the Bessel functions; and the  $I_m(\xi)$  and  $K_m(\xi)$  are the modified Bessel functions.

#### 8.3.3-2. Domain: $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi, -\infty < z < \infty$ . First boundary value problem.

An infinite circular cylinder is considered. A boundary condition is prescribed:

$$w = f(\varphi, z) \quad \text{at} \quad r = R.$$

Solution:

$$\begin{aligned} w(r, \varphi, z) &= -R \int_0^{2\pi} \int_{-\infty}^{\infty} f(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\xi=R} d\zeta d\eta \\ &\quad + \int_0^R \int_0^{2\pi} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi. \end{aligned}$$

Here,

$$\begin{aligned} G(r, \varphi, z, \xi, \eta, \zeta) &= \frac{1}{2\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{[J'_n(\mu_{nm} R)]^2 \beta_{nm}} \cos[n(\varphi - \eta)] \exp(-\beta_{nm}|z - \zeta|), \\ \beta_{nm} &= \sqrt{\mu_{nm}^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \end{aligned}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

#### 8.3.3-3. Domain: $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi, -\infty < z < \infty$ . Second boundary value problem.

An infinite circular cylinder is considered. A boundary condition is prescribed:

$$\partial_r w = f(\varphi, z) \quad \text{at} \quad r = R.$$

Solution:

$$w(r, \varphi, z) = R \int_0^{2\pi} \int_{-\infty}^{\infty} f(\eta, \zeta) G(r, \varphi, z, R, \eta, \zeta) d\zeta d\eta \\ + \int_0^R \int_0^{2\pi} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi.$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{\exp(-\sqrt{-\lambda}|z - \zeta|)}{2\pi R^2 \sqrt{-\lambda}} \\ + \frac{1}{2\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)]}{(\mu_{nm}^2 R^2 - n^2) J_n^2(\mu_{nm} R) \beta_{nm}} \exp(-\beta_{nm}|z - \zeta|), \\ \beta_{nm} = \sqrt{\mu_{nm}^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

8.3.3-4. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $-\infty < z < \infty$ . Third boundary value problem.

An infinite circular cylinder is considered. A boundary condition is prescribed:

$$\partial_r w + kw = f(\varphi, z) \quad \text{at } r = R.$$

Solution:

$$w(r, \varphi, z) = R \int_0^{2\pi} \int_{-\infty}^{\infty} f(\eta, \zeta) G(r, \varphi, z, R, \eta, \zeta) d\zeta d\eta \\ + \int_0^R \int_0^{2\pi} \int_{-\infty}^{\infty} \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi.$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{2\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)]}{(\mu_{nm}^2 R^2 + k^2 R^2 - n^2) J_n^2(\mu_{nm} R) \beta_{nm}} \exp(-\beta_{nm}|z - \zeta|), \\ \beta_{nm} = \sqrt{\mu_{nm}^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k J_n(\mu R) = 0.$$

8.3.3-5. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . First boundary value problem.

A semiinfinite circular cylinder is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi, z) \quad \text{at } r = R, \quad w = f_2(r, \varphi) \quad \text{at } z = 0.$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z) = & -R \int_0^{2\pi} \int_0^\infty f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\xi=R} d\zeta d\eta \\
 & + \int_0^{2\pi} \int_0^R f_2(\xi, \eta) \left[ \frac{\partial}{\partial \zeta} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\zeta=0} \xi d\xi d\eta \\
 & + \int_0^R \int_0^{2\pi} \int_0^\infty \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi.
 \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{2\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{[J'_n(\mu_{nm} R)]^2 \beta_{nm}} \cos[n(\varphi - \eta)] F_{nm}(z, \zeta),$$

$$F_{nm}(z, \zeta) = \exp(-\beta_{nm}|z - \zeta|) - \exp(-\beta_{nm}|z + \zeta|), \quad \beta_{nm} = \sqrt{\mu_{nm}^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

8.3.3-6. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Second boundary value problem.

A semiinfinite circular cylinder is considered. Boundary conditions are prescribed:

$$\partial_r w = f_1(\varphi, z) \quad \text{at } r = R, \quad \partial_z w = f_2(r, \varphi) \quad \text{at } z = 0.$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z) = & R \int_0^{2\pi} \int_0^\infty f_1(\eta, \zeta) G(r, \varphi, z, R, \eta, \zeta) d\zeta d\eta \\
 & - \int_0^{2\pi} \int_0^R f_2(\xi, \eta) G(r, \varphi, z, R, \eta, 0) \xi d\xi d\eta \\
 & + \int_0^R \int_0^{2\pi} \int_0^\infty \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi.
 \end{aligned}$$

Here,

$$\begin{aligned}
 G(r, \varphi, z, \xi, \eta, \zeta) = & \frac{\exp(-\sqrt{-\lambda}|z - \zeta|) + \exp(-\sqrt{-\lambda}|z + \zeta|)}{2\pi R^2 \sqrt{-\lambda}} \\
 & + \frac{1}{2\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)]}{(\mu_{nm}^2 R^2 - n^2) J_n^2(\mu_{nm} R) \beta_{nm}} F_{nm}(z, \zeta),
 \end{aligned}$$

$$F_{nm}(z, \zeta) = \exp(-\beta_{nm}|z - \zeta|) + \exp(-\beta_{nm}|z + \zeta|), \quad \beta_{nm} = \sqrt{\mu_{nm}^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J'_n(\mu R) = 0$ .

8.3.3-7. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Third boundary value problem.

A semiinfinite circular cylinder is considered. Boundary conditions are prescribed:

$$\partial_r w + k_1 w = f(\varphi, z) \quad \text{at } r = R, \quad \partial_z w - k_2 w = f_2(r, \varphi) \quad \text{at } z = 0.$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z) = & R \int_0^{2\pi} \int_0^\infty f_1(\eta, \zeta) G(r, \varphi, z, R, \eta, \zeta) d\zeta d\eta \\
 & - \int_0^{2\pi} \int_0^R f_2(\xi, \eta) G(r, \varphi, z, \xi, \eta, 0) \xi d\xi d\eta \\
 & + \int_0^R \int_0^{2\pi} \int_0^\infty \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi.
 \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{\pi} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n \mu_{nm}^2 J_n(\mu_{nm} r) J_n(\mu_{nm} \xi) \cos[n(\varphi - \eta)]}{(\mu_{nm}^2 R^2 + k_1^2 R^2 - n^2) J_n^2(\mu_{nm} R)} F_{nm}(z, \zeta),$$

$$A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases} \quad F_{nm}(z, \zeta) = \begin{cases} \frac{\exp(-\beta_{nm} z) [\beta_{nm} \cosh(\beta_{nm} \zeta) + k_2 \sinh(\beta_{nm} \zeta)]}{\beta_{nm} (\beta_{nm} + k_2)} & \text{for } z > \zeta, \\ \frac{\exp(-\beta_{nm} \zeta) [\beta_{nm} \cosh(\beta_{nm} z) + k_2 \sinh(\beta_{nm} z)]}{\beta_{nm} (\beta_{nm} + k_2)} & \text{for } \zeta > z, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions,  $\beta_{nm} = \sqrt{\mu_{nm}^2 - \lambda}$ , and the  $\mu_{nm}$  are positive roots of the transcendental equation

$$\mu J'_n(\mu R) + k_1 J_n(\mu R) = 0.$$

8.3.3-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z < \infty$ . Mixed boundary value problem.

A semiinfinite circular cylinder is considered. Boundary conditions are prescribed:

$$w = f_1(\varphi, z) \quad \text{at } r = R, \quad \partial_z w = f_2(r, \varphi) \quad \text{at } z = 0.$$

Solution:

$$\begin{aligned}
 w(r, \varphi, z) = & -R \int_0^{2\pi} \int_0^\infty f_1(\eta, \zeta) \left[ \frac{\partial}{\partial \xi} G(r, \varphi, z, \xi, \eta, \zeta) \right]_{\xi=R} d\zeta d\eta \\
 & - \int_0^{2\pi} \int_0^R f_2(\xi, \eta) G(r, \varphi, z, \xi, \eta, 0) \xi d\xi d\eta \\
 & + \int_0^R \int_0^{2\pi} \int_0^\infty \Phi(\xi, \eta, \zeta) G(r, \varphi, z, \xi, \eta, \zeta) \xi d\zeta d\eta d\xi.
 \end{aligned}$$

Here,

$$G(r, \varphi, z, \xi, \eta, \zeta) = \frac{1}{2\pi R^2} \sum_{n=0}^{\infty} \sum_{m=1}^{\infty} \frac{A_n J_n(\mu_{nm} r) J_n(\mu_{nm} \xi)}{[J'_n(\mu_{nm} R)]^2 \beta_{nm}} \cos[n(\varphi - \eta)] F_{nm}(z, \zeta),$$

$$F_{nm}(z, \zeta) = \exp(-\beta_{nm}|z - \zeta|) + \exp(-\beta_{nm}|z + \zeta|), \quad \beta_{nm} = \sqrt{\mu_{nm}^2 - \lambda}, \quad A_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0, \end{cases}$$

where the  $J_n(\xi)$  are the Bessel functions and the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_n(\mu R) = 0$ .

► Paragraphs 8.3.3-9 through 8.3.3-16 present only the eigenvalues and eigenfunctions of homogeneous boundary value problems for the homogeneous Helmholtz equation (with  $\Phi \equiv 0$ ). The solutions of the corresponding nonhomogeneous boundary value problems ( $\Phi \not\equiv 0$ ) can be constructed by the relations specified in Paragraphs 8.3.1-4 and 8.3.1-8.

**8.3.3-9. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . First boundary value problem.**

A circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$w = 0 \quad \text{at} \quad r = R, \quad w = 0 \quad \text{at} \quad z = 0, \quad w = 0 \quad \text{at} \quad z = a.$$

Eigenvalues:

$$\lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \frac{\mu_{nm}^2}{R^2}; \quad n = 0, 1, \dots; \quad m, k = 1, 2, \dots$$

Here, the  $\mu_{nm}$  are positive zeros of the Bessel functions,  $J_n(\mu) = 0$ .

Eigenfunctions:

$$w_{nmk}^{(1)} = J_n\left(\mu_{nm} \frac{r}{R}\right) \cos(n\varphi) \sin\left(\frac{\pi k z}{a}\right),$$

$$w_{nmk}^{(2)} = J_n\left(\mu_{nm} \frac{r}{R}\right) \sin(n\varphi) \sin\left(\frac{\pi k z}{a}\right).$$

Eigenfunctions possessing the axial symmetry property:

$$w_{0mk}^{(1)} = J_0\left(\mu_{0m} \frac{r}{R}\right) \sin\left(\frac{\pi k z}{a}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2 = \frac{\pi R^2 a}{4} (1 + \delta_{n0}) [J'_n(\mu_{nm})]^2, \quad \delta_{nm} = \begin{cases} 1 & \text{for } n = m, \\ 0 & \text{for } n \neq m. \end{cases}$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.3.3-10. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . Second boundary value problem.**

A circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$\partial_r w = 0 \quad \text{at} \quad r = R, \quad \partial_z w = 0 \quad \text{at} \quad z = 0, \quad \partial_z w = 0 \quad \text{at} \quad z = a.$$

Eigenvalues:

$$\lambda_{000} = 0, \quad \lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \frac{\mu_{nm}^2}{R^2}; \quad n = 0, 1, \dots; \quad k, m = 0, 1, \dots$$

Here, the  $\mu_{nm}$  are roots of the transcendental equation  $J'_n(\mu) = 0$ .

Eigenfunctions:

$$w_{nmk}^{(1)} = J_n\left(\mu_{nm} \frac{r}{R}\right) \cos(n\varphi) \cos\left(\frac{\pi k z}{a}\right), \quad w_{000}^{(1)} = 1,$$

$$w_{nmk}^{(2)} = J_n\left(\mu_{nm} \frac{r}{R}\right) \sin(n\varphi) \cos\left(\frac{\pi k z}{a}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2 = \frac{\pi R^2 a}{4 \mu_{nm}^2} (1 + \delta_{n0}) (\mu_{nm}^2 - n^2) [J_n(\mu_{nm})]^2, \quad \|w_{000}^{(1)}\|^2 = \pi R^2 a,$$

where  $\delta_{n0}$  is the Kronecker delta.

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).



**8.3.3-11. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . Third boundary value problem.**

A circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$\partial_r w + k_1 w = 0 \quad \text{at} \quad r = R, \quad \partial_z w - k_2 w = 0 \quad \text{at} \quad z = 0, \quad \partial_z w + k_3 w = 0 \quad \text{at} \quad z = a.$$

Eigenvalues:

$$\lambda_{nml} = \nu_l^2 + \frac{\mu_{nm}^2}{R^2},$$

where the  $\nu_l$  and  $\mu_{nm}$  are positive roots of the transcendental equations

$$\tan(\nu a) = \frac{(k_2 + k_3)\nu}{\nu^2 - k_2 k_3}, \quad \mu J'_n(\mu) + R k_1 J_n(\mu) = 0.$$

Eigenfunctions:

$$w_{nml}^{(1)} = J_n\left(\mu_{nm} \frac{r}{R}\right) \cos(n\varphi) \frac{\nu_l \cos \nu_l z + k_2 \sin \nu_l z}{\sqrt{\nu_l^2 + k_2^2}},$$

$$w_{nml}^{(2)} = J_n\left(\mu_{nm} \frac{r}{R}\right) \sin(n\varphi) \frac{\nu_l \cos \nu_l z + k_2 \sin \nu_l z}{\sqrt{\nu_l^2 + k_2^2}}.$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nml}^{(i)}\|^2 = \frac{\pi R^2}{4\mu_{nm}^2} (1 + \delta_{n0}) (R^2 k_1^2 + \mu_{nm}^2 - n^2) [J_n(\mu_{nm})]^2 \left[ a + \frac{(k_2 + k_3)(\nu_l^2 + k_2 k_3)}{(\nu_l^2 + k_2^2)(\nu_l^2 + k_3^2)} \right],$$

where  $\delta_{n0}$  is the Kronecker delta.

**8.3.3-12. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . First boundary value problem.**

A hollow circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$w = 0 \quad \text{at} \quad r = R_1, \quad w = 0 \quad \text{at} \quad r = R_2,$$

$$w = 0 \quad \text{at} \quad z = 0, \quad w = 0 \quad \text{at} \quad z = a.$$

Eigenvalues:

$$\lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \mu_{nm}^2; \quad n = 0, 1, 2, \dots; \quad m, k = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are positive roots of the transcendental equation

$$J_n(\mu R_1) Y_n(\mu R_2) - J_n(\mu R_2) Y_n(\mu R_1) = 0.$$

Eigenfunctions:

$$w_{nmk}^{(1)} = [J_n(\mu_{nm} r) Y_n(\mu_{nm} R_1) - J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r)] \cos(n\varphi) \sin\left(\frac{\pi k z}{a}\right),$$

$$w_{nmk}^{(2)} = [J_n(\mu_{nm} r) Y_n(\mu_{nm} R_1) - J_n(\mu_{nm} R_1) Y_n(\mu_{nm} r)] \sin(n\varphi) \sin\left(\frac{\pi k z}{a}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2 = \frac{a}{\pi \mu_{nm}^2} (1 + \delta_{n0}) \frac{[J_n(\mu_{nm} R_1)]^2 - [J_n(\mu_{nm} R_2)]^2}{[J_n(\mu_{nm} R_2)]^2}, \quad \delta_{ij} = \begin{cases} 1 & \text{for } i = j, \\ 0 & \text{for } i \neq j. \end{cases}$$

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.3.3-13. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . Second boundary value problem.**

A hollow circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_r w &= 0 & \text{at } r &= R_1, & \partial_r w &= 0 & \text{at } r &= R_2, \\ \partial_z w &= 0 & \text{at } z &= 0, & \partial_z w &= 0 & \text{at } z &= a. \end{aligned}$$

Eigenvalues:

$$\lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \mu_{nm}^2; \quad n, m, k = 0, 1, 2, \dots$$

Here, the  $\mu_{nm}$  are roots of the transcendental equation

$$J'_n(\mu R_1)Y'_n(\mu R_2) - J'_n(\mu R_2)Y'_n(\mu R_1) = 0.$$

Eigenfunctions:

$$\begin{aligned} w_{nmk}^{(1)} &= [J_n(\mu_{nm}r)Y'_n(\mu_{nm}R_1) - J'_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \cos(n\varphi) \cos\left(\frac{\pi kz}{a}\right), \\ w_{nmk}^{(2)} &= [J_n(\mu_{nm}r)Y'_n(\mu_{nm}R_1) - J'_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \sin(n\varphi) \cos\left(\frac{\pi kz}{a}\right). \end{aligned}$$

To the zero eigenvalue  $\lambda_{000} = 0$  there is a corresponding eigenfunction  $w_{000}^{(1)} = 1$ .

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2 = \frac{a(1 + \delta_{n0})(1 + \delta_{k0})}{\pi \mu_{nm}^2} \left\{ \left(1 - \frac{n^2}{R_2^2 \mu_{nm}^2}\right) \left[ \frac{J'_n(\mu_{nm}R_1)}{J'_n(\mu_{nm}R_2)} \right]^2 - \left(1 - \frac{n^2}{R_1^2 \mu_{nm}^2}\right) \right\},$$

where  $\delta_{n0}$  is the Kronecker delta.

⊙ *References:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**8.3.3-14. Domain:  $R_1 \leq r \leq R_2$ ,  $0 \leq \varphi \leq 2\pi$ ,  $0 \leq z \leq a$ . Mixed boundary value problems.**

1°. A hollow circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= 0 & \text{at } r &= R_1, & w &= 0 & \text{at } r &= R_2, \\ \partial_z w &= 0 & \text{at } z &= 0, & \partial_z w &= 0 & \text{at } z &= a. \end{aligned}$$

Eigenvalues:

$$\lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \mu_{nm}^2; \quad n, k = 0, 1, 2, \dots; \quad m = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are roots of the transcendental equation

$$J_n(\mu R_1)Y_n(\mu R_2) - J_n(\mu R_2)Y_n(\mu R_1) = 0.$$

Eigenfunctions:

$$\begin{aligned} w_{nmk}^{(1)} &= [J_n(\mu_{nm}r)Y_n(\mu_{nm}R_1) - J_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \cos(n\varphi) \cos\left(\frac{\pi kz}{a}\right), \\ w_{nmk}^{(2)} &= [J_n(\mu_{nm}r)Y_n(\mu_{nm}R_1) - J_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \sin(n\varphi) \cos\left(\frac{\pi kz}{a}\right). \end{aligned}$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2 = \frac{a \varepsilon_n \varepsilon_k}{\pi \mu_{nm}^2} \frac{[J_n(\mu_{nm}R_1)]^2 - [J_n(\mu_{nm}R_2)]^2}{[J_n(\mu_{nm}R_2)]^2}, \quad \varepsilon_n = \begin{cases} 2 & \text{for } n = 0, \\ 1 & \text{for } n \neq 0. \end{cases}$$

2°. A hollow circular cylinder of finite length is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_r w &= 0 & \text{at } r &= R_1, & \partial_r w &= 0 & \text{at } r &= R_2, \\ w &= 0 & \text{at } z &= 0, & w &= 0 & \text{at } z &= a. \end{aligned}$$

Eigenvalues:

$$\lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \mu_{nm}^2; \quad n = 0, 1, 2, \dots; \quad m, k = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are roots of the transcendental equation

$$J'_n(\mu R_1)Y'_n(\mu R_2) - J'_n(\mu R_2)Y'_n(\mu R_1) = 0.$$

Eigenfunctions:

$$\begin{aligned} w_{nmk}^{(1)} &= [J_n(\mu_{nm}r)Y'_n(\mu_{nm}R_1) - J'_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \cos(n\varphi) \sin\left(\frac{\pi kz}{a}\right), \\ w_{nmk}^{(2)} &= [J_n(\mu_{nm}r)Y'_n(\mu_{nm}R_1) - J'_n(\mu_{nm}R_1)Y_n(\mu_{nm}r)] \sin(n\varphi) \sin\left(\frac{\pi kz}{a}\right). \end{aligned}$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2 = \frac{a\varepsilon_n}{\pi\mu_{nm}^2} \left\{ \left(1 - \frac{n^2}{R_2^2\mu_{nm}^2}\right) \left[\frac{J'_n(\mu_{nm}R_1)}{J'_n(\mu_{nm}R_2)}\right]^2 - \left(1 - \frac{n^2}{R_1^2\mu_{nm}^2}\right) \right\},$$

where  $\varepsilon_n$  is defined in Item 1°.

8.3.3-15. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq a$ . First boundary value problem.

A cylindrical sector of finite thickness is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= 0 & \text{at } \varphi &= 0, & w &= 0 & \text{at } \varphi &= \varphi_0, & w &= 0 & \text{at } r &= R, \\ w &= 0 & \text{at } z &= 0, & w &= 0 & \text{at } z &= a. \end{aligned}$$

Eigenvalues:

$$\lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \frac{\mu_{nm}^2}{R^2}; \quad n, m, k = 1, 2, 3, \dots$$

Here, the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu) = 0$ .

Eigenfunctions:

$$w_{nmk} = J_{n\pi/\varphi_0}\left(\frac{\mu_{nm}r}{R}\right) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \sin\left(\frac{k\pi z}{a}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}\|^2 = \frac{1}{8}aR^2\varphi_0 [J'_{n\pi/\varphi_0}(\mu_{nm})]^2.$$

8.3.3-16. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq \varphi_0$ ,  $0 \leq z \leq a$ . Mixed boundary value problem.

A cylindrical sector of finite thickness is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= 0 & \text{at } \varphi &= 0, & w &= 0 & \text{at } \varphi &= \varphi_0, & w &= 0 & \text{at } r &= R, \\ \partial_z w &= 0 & \text{at } z &= 0, & \partial_z w &= 0 & \text{at } z &= a. \end{aligned}$$

Eigenvalues:

$$\lambda_{nmk} = \frac{\pi^2 k^2}{a^2} + \frac{\mu_{nm}^2}{R^2}; \quad n, m = 1, 2, 3, \dots; \quad k = 0, 1, 2, \dots$$

Here, the  $\mu_{nm}$  are positive roots of the transcendental equation  $J_{n\pi/\varphi_0}(\mu) = 0$ .

Eigenfunctions:

$$w_{nmk} = J_{n\pi/\varphi_0}\left(\frac{\mu_{nm}r}{R}\right) \sin\left(\frac{n\pi\varphi}{\varphi_0}\right) \cos\left(\frac{k\pi z}{a}\right).$$

The square of the norm of an eigenfunction is defined as

$$\|w_{nmk}\|^2 = \frac{1}{8}aR^2\varphi_0(1 + \delta_{k0})[J'_{n\pi/\varphi_0}(\mu_{nm})]^2, \quad \delta_{k0} = \begin{cases} 1 & \text{for } k = 0, \\ 0 & \text{for } k \neq 0. \end{cases}$$

### 8.3.4. Problems in Spherical Coordinates

The three-dimensional homogeneous Helmholtz equation in the spherical coordinate system is written as

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial w}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial w}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 w}{\partial \varphi^2} + \lambda w = 0, \quad r = \sqrt{x^2 + y^2 + z^2}.$$

#### 8.3.4-1. Particular solutions:

$$w = \frac{1}{r} (A \sin \mu r + B \cos \mu r), \quad \lambda = \mu^2,$$

$$w = \frac{1}{r} (A \sinh \mu r + B \cosh \mu r), \quad \lambda = -\mu^2,$$

$$w = \frac{1}{\sqrt{r}} J_{n+1/2}(\mu r) P_n^m(\cos \theta) (A \cos m\varphi + B \sin m\varphi), \quad \lambda = \mu^2,$$

$$w = \frac{1}{\sqrt{r}} Y_{n+1/2}(\mu r) P_n^m(\cos \theta) (A \cos m\varphi + B \sin m\varphi), \quad \lambda = \mu^2,$$

$$w = \frac{1}{\sqrt{r}} I_{n+1/2}(\mu r) P_n^m(\cos \theta) (A \cos m\varphi + B \sin m\varphi), \quad \lambda = -\mu^2,$$

$$w = \frac{1}{\sqrt{r}} K_{n+1/2}(\mu r) P_n^m(\cos \theta) (A \cos m\varphi + B \sin m\varphi), \quad \lambda = -\mu^2,$$

where  $n, m = 0, 1, 2, \dots$ ;  $A$  and  $B$  are arbitrary constants;  $J_\nu(\xi)$  and  $Y_\nu(\xi)$  are the Bessel functions;  $I_\nu(\xi)$  and  $K_\nu(\xi)$  are the modified Bessel functions; and the  $P_n^m(\xi)$  are the associated Legendre functions that are expressed in terms of the Legendre polynomials  $P_n(\xi)$  as

$$P_n^m(\xi) = (1 - \xi^2)^{m/2} \frac{d^m}{d\xi^m} P_n(\xi), \quad P_n(\xi) = \frac{1}{n! 2^n} \frac{d^n}{d\xi^n} (\xi^2 - 1)^n.$$

#### 8.3.4-2. Domain: $0 \leq r \leq R$ . First boundary value problem.

1°. A spherical domain is considered. A homogeneous boundary condition is prescribed,

$$w = 0 \quad \text{at} \quad r = R.$$

Eigenvalues:

$$\lambda_{nk} = \frac{\mu_{nk}^2}{R^2}; \quad n = 0, 1, 2, \dots; \quad k = 1, 2, 3, \dots$$

Here, the  $\mu_{nk}$  are positive zeros of the Bessel functions,  $J_{n+1/2}(\mu) = 0$ . Note that the  $J_{n+1/2}(\mu)$  can be expressed in terms of elementary functions, see Bateman and Erdélyi (1953, Vol. 2).

Eigenfunctions:

$$w_{nmk}^{(1)} = \frac{1}{\sqrt{r}} J_{n+1/2} \left( \mu_{nk} \frac{r}{R} \right) P_n^m(\cos \theta) \cos m\varphi, \quad m = 0, 1, 2, \dots;$$

$$w_{nmk}^{(2)} = \frac{1}{\sqrt{r}} J_{n+1/2} \left( \mu_{nk} \frac{r}{R} \right) P_n^m(\cos \theta) \sin m\varphi, \quad m = 1, 2, 3, \dots$$

Here, the  $P_n^m(\xi)$  are the associated Legendre functions.

Eigenfunctions possessing central symmetry (i.e., independent of  $\theta$  and  $\varphi$ ):

$$w_{00k}^{(1)} = J_{1/2} \left( \mu_{0k} \frac{r}{R} \right).$$

Eigenfunctions possessing axial symmetry (i.e., independent of  $\varphi$ ):

$$w_{n0k}^{(1)} = J_{n+1/2} \left( \mu_{nk} \frac{r}{R} \right) P_n(\cos \theta).$$

The square of the norm of an eigenfunction:

$$\|w_{nmk}^{(1)}\|^2 = \frac{\pi R^2 (1 + \delta_{m0})(n+m)!}{(2n+1)(n-m)!} [J'_{n+1/2}(\mu_{nk})]^2, \quad \delta_{m0} = \begin{cases} 1 & \text{for } m=0, \\ 0 & \text{for } m \neq 0, \end{cases}$$

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2, \quad m = 1, 2, 3, \dots$$

2°. A spherical domain is considered. A nonhomogeneous boundary condition is prescribed,

$$w = f(\theta, \varphi) \quad \text{at } r = R.$$

Solution:

$$w(r, \theta, \varphi) = \sum_{n=0}^{\infty} \sum_{m=-n}^n f_{nm} \frac{\Psi_n(r\sqrt{\lambda})}{\Psi_n(R\sqrt{\lambda})} Y_n^m(\theta, \varphi), \quad \Psi_n(x) = \frac{1}{\sqrt{x}} J_{n+1/2}(x),$$

where

$$f_{nm} = \frac{1}{\|Y_n^m\|} \int_0^{2\pi} \int_0^\pi f(\theta, \varphi) Y_n^m(\theta, \varphi) \sin \theta \, d\theta \, d\varphi, \quad \|Y_n^m\| = \frac{2\pi \varepsilon_m (n+m)!}{2n+1 (n-m)!},$$

$$Y_n^m(\theta, \varphi) = \begin{cases} P_n(\cos \theta) & \text{for } m=0, \\ P_n^m(\cos \theta) \sin m\varphi & \text{for } m=1, 2, \dots, \\ P_n^{|m|}(\cos \theta) \cos m\varphi & \text{for } m=-1, -2, \dots, \end{cases} \quad \varepsilon_m = \begin{cases} 2 & \text{for } m=0, \\ 1 & \text{for } m \neq 0. \end{cases}$$

The solution was written out under the assumption that  $J_{n+1/2}(R\sqrt{\lambda}) \neq 0$  for  $n = 0, 1, 2, \dots$

⊙ *References:* M. M. Smirnov (1975), A. N. Tikhonov and A. A. Samarskii (1990).

► Paragraphs 8.3.4-3 through 8.3.4-6 present only the eigenvalues and eigenfunctions of homogeneous boundary value problems for the homogeneous Helmholtz equation (with  $\Phi \equiv 0$ ). The solutions of the corresponding nonhomogeneous boundary value problems ( $\Phi \not\equiv 0$ ) can be constructed by the relations specified in Paragraph 8.3.1-4.

### 8.3.4-3. Domain: $0 \leq r \leq R$ . Second boundary value problem.

A spherical domain is considered. A boundary condition is prescribed:

$$\partial_r w = 0 \quad \text{at } r = R.$$

Eigenvalues:

$$\lambda_{00} = 0, \quad \lambda_{nk} = \frac{\mu_{nk}^2}{R^2}; \quad n = 0, 1, 2, \dots; \quad k = 1, 2, 3, \dots$$

Here, the  $\mu_{nk}$  are roots of the transcendental equation

$$2\mu J'_{n+1/2}(\mu) - J_{n+1/2}(\mu) = 0.$$

Eigenfunctions:

$$w_{000}^{(1)} = 1, \quad w_{nmk}^{(1)} = \frac{1}{\sqrt{r}} J_{n+1/2} \left( \mu_{nk} \frac{r}{R} \right) P_n^m(\cos \theta) \cos m\varphi, \quad m = 0, 1, 2, \dots;$$

$$w_{nmk}^{(2)} = \frac{1}{\sqrt{r}} J_{n+1/2} \left( \mu_{nk} \frac{r}{R} \right) P_n^m(\cos \theta) \sin m\varphi, \quad m = 1, 2, 3, \dots$$

The square of the norm of an eigenfunction:

$$\|w_{000}^{(1)}\|^2 = \frac{4}{3} \pi R^3, \quad \|w_{nmk}^{(1)}\|^2 = \frac{\pi R^2 \varepsilon_m (n+m)!}{(2n+1)(n-m)!} \left[ 1 - \frac{n(n+1)}{\mu_{nk}^2} \right] J_{n+1/2}^2(\mu_{nk}),$$

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2, \quad m = 1, 2, 3, \dots,$$

where  $\varepsilon_m = \begin{cases} 2 & \text{for } m=0, \\ 1 & \text{for } m \neq 0. \end{cases}$

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

#### 8.3.4-4. Domain: $0 \leq r \leq R$ . Third boundary value problem.

A spherical domain is considered. A boundary condition is prescribed:

$$\partial_r w + sw = 0 \quad \text{at} \quad r = R.$$

Eigenvalues:

$$\lambda_{nk} = \frac{\mu_{nk}^2}{R^2}; \quad n = 0, 1, 2, \dots; \quad k = 1, 2, 3, \dots$$

Here, the  $\mu_{nk}$  are positive roots of the transcendental equation

$$2\mu J'_{n+1/2}(\mu) - (1 - 2Rs)J_{n+1/2}(\mu) = 0.$$

Eigenfunctions:

$$\begin{aligned} w_{nmk}^{(1)} &= \frac{1}{\sqrt{r}} J_{n+1/2} \left( \mu_{nk} \frac{r}{R} \right) P_n^m(\cos \theta) \cos m\varphi, \quad m = 0, 1, 2, \dots; \\ w_{nmk}^{(2)} &= \frac{1}{\sqrt{r}} J_{n+1/2} \left( \mu_{nk} \frac{r}{R} \right) P_n^m(\cos \theta) \sin m\varphi, \quad m = 1, 2, 3, \dots \end{aligned}$$

Here, the  $P_n^m(\xi)$  are the associated Legendre functions.

The square of the norm of an eigenfunction:

$$\begin{aligned} \|w_{nmk}^{(1)}\|^2 &= \frac{\pi R^2 \varepsilon_m (n+m)!}{(2n+1)(n-m)!} \left[ 1 + \frac{(Rs+n)(Rs-n-1)}{\mu_{nk}^2} \right] J_{n+1/2}^2(\mu_{nk}), \quad \varepsilon_m = \begin{cases} 2 & \text{for } m = 0, \\ 1 & \text{for } m \neq 0, \end{cases} \\ \|w_{nmk}^{(1)}\|^2 &= \|w_{nmk}^{(2)}\|^2, \quad m = 1, 2, 3, \dots \end{aligned}$$

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

#### 8.3.4-5. Domain: $R \leq r < \infty$ . First boundary value problem.

A spherical cavity is considered and the dependent variable is prescribed at its surface:

$$w = f(\theta, \varphi) \quad \text{at} \quad r = R,$$

and the radiation conditions are prescribed at infinity (see Paragraph 8.3.1-5, Item 2°).

Solution for  $\lambda = k^2 > 0$ :

$$w(r, \theta, \varphi) = \sum_{n=0}^{\infty} \sum_{m=-n}^n f_{nm} \frac{\Xi_n(kr)}{\Xi_n(kR)} Y_n^m(\theta, \varphi), \quad \Xi_n(\rho) = \frac{1}{\sqrt{\rho}} H_{n+1/2}^{(2)}(\rho),$$

where  $H_{n+1/2}^{(2)}(\rho)$  is the Hankel function of the second kind and the other quantities are defined just as in Paragraph 8.3.4-2, Item 2°.

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

#### 8.3.4-6. Domain: $R_1 \leq r \leq R_2$ . First boundary value problem.

A spherical layer is considered. Boundary conditions are prescribed:

$$w = 0 \quad \text{at} \quad r = R_1, \quad w = 0 \quad \text{at} \quad r = R_2.$$

Eigenvalues:

$$\lambda_{nk} = \mu_{nk}^2; \quad n = 0, 1, 2, \dots; \quad k = 1, 2, 3, \dots$$

Here, the  $\mu_{nk}$  are positive roots of the transcendental equation

$$J_{n+1/2}(\mu R_1) Y_{n+1/2}(\mu R_2) - J_{n+1/2}(\mu R_2) Y_{n+1/2}(\mu R_1) = 0.$$

Eigenfunctions:

$$w_{nmk}^{(1)} = \frac{1}{\sqrt{r}} Z_{n+1/2}(\mu_{nk}r) P_n^m(\cos \theta) \cos m\varphi, \quad m = 0, 1, 2, \dots;$$

$$w_{nmk}^{(2)} = \frac{1}{\sqrt{r}} Z_{n+1/2}(\mu_{nk}r) P_n^m(\cos \theta) \sin m\varphi, \quad m = 1, 2, 3, \dots$$

Here, the  $P_n^m(\xi)$  are the associated Legendre functions and

$$Z_{n+1/2}(\mu r) = J_{n+1/2}(\mu R_1) Y_{n+1/2}(\mu r) - Y_{n+1/2}(\mu R_1) J_{n+1/2}(\mu r).$$

The square of the norm of an eigenfunction:

$$\|w_{nmk}^{(1)}\|^2 = \frac{4\varepsilon_m(n+m)!}{\pi(2n+1)(n-m)!} \frac{J_{n+1/2}^2(\mu_{nk}R_1) - J_{n+1/2}^2(\mu_{nk}R_2)}{\mu_{nk}^2 J_{n+1/2}^2(\mu_{nk}R_2)}, \quad \varepsilon_m = \begin{cases} 2 & \text{for } m = 0, \\ 1 & \text{for } m \neq 0, \end{cases}$$

$$\|w_{nmk}^{(1)}\|^2 = \|w_{nmk}^{(2)}\|^2, \quad m = 1, 2, 3, \dots$$

### 8.3.5. Other Orthogonal Curvilinear Coordinates

The homogenous three-dimensional Helmholtz equation admits separation of variables in the eleven orthogonal systems of coordinates listed in [Table 29](#).

For the parabolic cylindrical system of coordinates, the multipliers  $f$  and  $g$  are expressed in terms of the parabolic cylinder functions as

$$f(\xi) = A_1 D_{\mu-1/2}(\sigma\xi) + A_2 D_{\mu-1/2}(-\sigma\xi), \quad g(\eta) = B_1 D_{-\mu-1/2}(\sigma\eta) + B_2 D_{-\mu-1/2}(-\sigma\eta),$$

$$\mu = \frac{1}{2}\beta(k^2 - \lambda)^{-1/2}, \quad \sigma = [4(k^2 - \lambda)]^{1/4},$$

where  $A_1$ ,  $B_1$ ,  $A_2$ , and  $B_2$  are arbitrary constants.

For the elliptic cylindrical system of coordinates, the functions  $f$  and  $g$  are determined by the modified Mathieu equation and Mathieu equation, respectively, so that

$$f(u) = \begin{cases} \text{Ce}_n(u, q), \\ \text{Se}_n(u, q), \end{cases} \quad g(v) = \begin{cases} \text{ce}_n(v, q), \\ \text{se}_n(v, q), \end{cases} \quad q = \frac{1}{4}a^2(\lambda - k^2),$$

where  $\text{Ce}_n(u, q)$  and  $\text{Se}_n(u, q)$  are the modified Mathieu functions, and  $\text{ce}_n(v, q)$  and  $\text{se}_n(v, q)$  are the Mathieu functions; to each value of the parameter  $q$  there are certain corresponding eigenvalues  $\beta = \beta_n(q)$  [see Abramowitz and Stegun (1964)].

In the prolate and oblate spheroidal systems of coordinates, the equations for  $f$  and  $g$  are different forms of the spheroidal wave equation, whose bounded solutions are given by

$$f(u) = \text{Ps}_n^{(k)}(\cosh u, a^2\lambda), \quad g(u) = \text{Ps}_n^{(k)}(\cos v, a^2\lambda) \quad \text{for prolate spheroid,}$$

$$f(u) = \text{Ps}_n^{(k)}(-i \sinh u, a^2\lambda), \quad g(u) = \text{Ps}_n^{(k)}(\cos v, -a^2\lambda) \quad \text{for oblate spheroid,}$$

$$k \text{ is an integer, } n = 0, 1, 2, \dots, \quad -n \leq k \leq n,$$

where  $\text{Ps}_n^k(z, a)$  are the spheroidal wave functions; see Bateman and Erdélyi (1955, Vol. 3), Arscott (1964), and Meixner and Schäfke (1965). The separation of variables for the Helmholtz equation in modified prolate and oblate spheroidal systems of coordinates, as well as the spheroidal wave functions, are discussed in Abramowitz and Stegun (1964).

In the parabolic coordinate system, the solutions of the equations for  $f$  and  $g$  are expressed in terms of the degenerate hypergeometric functions [see Miller, Jr. (1977)] as follows:

$$f(\xi) = \xi^k \exp(\pm \frac{1}{2}\omega\xi^2) \Phi\left(-\frac{\beta}{4\omega} + \frac{k+1}{2}, k+1; \mp\omega\xi^2\right), \quad \omega = \sqrt{-\lambda},$$

$$g(\eta) = \eta^k \exp(\pm \frac{1}{2}\omega\eta^2) \Phi\left(\frac{\beta}{4\omega} + \frac{k+1}{2}, k+1; \mp\omega\eta^2\right).$$

TABLE 29

Orthogonal coordinates  $\bar{x}, \bar{y}, \bar{z}$  that allow separable solutions of the form  $w = f(\bar{x})g(\bar{y})h(\bar{z})$  for the three-dimensional Helmholtz equation  $\Delta_3 w + \lambda w = 0$

| Coordinates                                | Transformations                                                                                                                              | Particular solutions (or equations for $f, g, h$ )                                                                                                                                                                                                                                                                                      |
|--------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Cartesian<br>$x, y, z$                     | $x = x,$<br>$y = y,$<br>$z = z$                                                                                                              | $w = \cos(k_1 x + s_1) \cos(k_2 y + s_2) \cos(k_3 z + s_3),$<br>where $k_1^2 + k_2^2 + k_3^2 = \lambda;$<br>see also Paragraph 8.3.2-1                                                                                                                                                                                                  |
| Cylindrical<br>$r, \varphi, z$             | $x = r \cos \varphi,$<br>$y = r \sin \varphi,$<br>$z = z$                                                                                    | $w = [AJ_n(\beta r) + BY_n(\beta r)] \cos(n\varphi + c) \cos(kz + s),$<br>where $k^2 + \beta^2 = \lambda$ , see also Paragraph 8.3.3-1<br>( $J_n$ and $Y_n$ are the Bessel functions)                                                                                                                                                   |
| Parabolic<br>cylindrical<br>$\xi, \eta, z$ | $x = \frac{1}{2}(\xi^2 - \eta^2),$<br>$y = \xi\eta,$<br>$z = z$                                                                              | $w = f(\xi)g(\eta) \cos(kz + s),$<br>$f'' + [(\lambda - k^2)\xi^2 + \beta]f = 0,$<br>$g'' + [(\lambda - k^2)\eta^2 - \beta]g = 0$                                                                                                                                                                                                       |
| Elliptic<br>cylindrical<br>$u, v, z$       | $x = a \cosh u \cos v,$<br>$y = a \sinh u \sin v,$<br>$z = z$                                                                                | $w = f(u)g(v) \cos(kz + s),$<br>$f'' + [\frac{1}{2}a^2(\lambda - k^2) \cosh 2u - \beta]f = 0,$<br>$g'' - [\frac{1}{2}a^2(\lambda - k^2) \cos 2v - \beta]g = 0$                                                                                                                                                                          |
| Spherical<br>$r, \theta, \varphi$          | $x = r \sin \theta \cos \varphi,$<br>$y = r \sin \theta \sin \varphi,$<br>$z = r \cos \theta$                                                | $w = r^{-1/2} J_{n+1/2}(\beta r) P_n^m(\cos \theta) \cos(m\varphi + s),$<br>$w = r^{-1/2} Y_{n+1/2}(\beta r) P_n^m(\cos \theta) \cos(m\varphi + s),$<br>where $\lambda = \beta^2$ ; see also Paragraph 8.3.4-1                                                                                                                          |
| Prolate<br>spheroidal<br>$u, v, \varphi$   | $x = a \sinh u \sin v \cos \varphi,$<br>$y = a \sinh u \sin v \sin \varphi,$<br>$z = a \cosh u \cos v$                                       | $w = f(u)g(v) \cos(k\varphi + s),$<br>$f'' + f' \coth u + (-\beta + a^2 \lambda \sinh^2 u - k^2 / \sinh^2 u) f = 0,$<br>$g'' + g' \cot v + (\beta + a^2 \lambda \sin^2 v - k^2 / \sin^2 v) g = 0$                                                                                                                                       |
| Oblate<br>spheroidal<br>$u, v, \varphi$    | $x = a \cosh u \sin v \cos \varphi,$<br>$y = a \cosh u \sin v \sin \varphi,$<br>$z = a \sinh u \cos v$                                       | $w = f(u)g(v) \cos(k\varphi + s),$<br>$f'' + f' \tanh u + (-\beta + a^2 \lambda \cosh^2 u + k^2 / \cosh^2 u) f = 0,$<br>$g'' + g' \cot v + (\beta - a^2 \lambda \sin^2 v - k^2 / \sin^2 v) g = 0$                                                                                                                                       |
| Parabolic<br>$\xi, \eta, \varphi$          | $x = \xi\eta \cos \varphi,$<br>$y = \xi\eta \sin \varphi,$<br>$z = \frac{1}{2}(\xi^2 - \eta^2)$                                              | $w = f(\xi)g(\eta) \cos(k\varphi + s),$<br>$\xi^2 f'' + \xi f' + (\lambda \xi^4 - \beta \xi^2 - k^2) f = 0,$<br>$\eta^2 g'' + \eta g' + (\lambda \eta^4 + \beta \eta^2 - k^2) g = 0$                                                                                                                                                    |
| Paraboloidal<br>$u, v, \varphi$            | $x = 2a \cosh u \cos v \sinh \varphi,$<br>$y = 2a \sinh u \sin v \cosh \varphi,$<br>$z = \frac{1}{2}a(\cosh 2u + \cos 2v - \cosh 2\varphi)$  | $f'' + (-k - a\beta \cosh 2u + \frac{1}{2}a^2 \lambda \cosh 4u) f = 0,$<br>$g'' + (k + a\beta \cos 2v - \frac{1}{2}a^2 \lambda \cos 4v) g = 0,$<br>$h'' + (-k + a\beta \cosh 2\varphi - \frac{1}{2}a^2 \lambda \cosh 4\varphi) h = 0$                                                                                                   |
| General<br>ellipsoidal<br>$\mu, \nu, \rho$ | $x = \sqrt{\frac{(\mu-a)(\nu-a)(\rho-a)}{a(a-1)}},$<br>$y = \sqrt{\frac{(\mu-1)(\nu-1)(\rho-1)}{1-a}},$<br>$z = \sqrt{\frac{\mu\nu\rho}{a}}$ | $4\sqrt{\varphi(\mu)} [\sqrt{\varphi(\mu)} f']' + (\lambda\mu^2 + \beta_1\mu + \beta_2) f = 0,$<br>$4\sqrt{\varphi(\nu)} [\sqrt{\varphi(\nu)} g']' + (\lambda\nu^2 + \beta_1\nu + \beta_2) g = 0,$<br>$4\sqrt{\varphi(\rho)} [\sqrt{\varphi(\rho)} h']' + (\lambda\rho^2 + \beta_1\rho + \beta_2) h = 0,$<br>$\varphi(t) = t(t-1)(t-a)$ |
| Conical<br>$\rho, \mu, \nu$                | $x = \rho \sqrt{\frac{(a\mu-1)(a\nu-1)}{1-a}},$<br>$y = \rho \sqrt{\frac{a(\mu-1)(\nu-1)}{a-1}},$<br>$z = \rho \sqrt{a\mu\nu}$               | $w = \rho^{-1/2} J_{\pm(n+1/2)}(\rho\sqrt{\lambda}) g(\xi) h(\eta),$<br>$g'' + [\beta - n(n+1)k^2 \operatorname{sn}^2 \xi] g = 0,$<br>$h'' + [\beta - n(n+1)k^2 \operatorname{sn}^2 \eta] h = 0,$<br>where $\mu = \operatorname{sn}^2(\xi, k), \nu = \operatorname{sn}^2(\eta, k), k = \sqrt{a}$                                        |



In the case of the paraboloidal coordinate system, the equations for  $f$ ,  $g$ , and  $h$  are reduced to the Whittaker–Hill equation

$$G''_{\theta\theta} + \left(\mu + \frac{1}{8}b^2 + bc \cos 2\theta - \frac{1}{8}b^2 \cos 4\theta\right)G = 0.$$

Denote by  $gc_n(\theta; b, c)$  and  $gs_n(\theta; b, c)$ , respectively, the even and odd  $2\pi$ -periodic solutions of the Whittaker–Hill equation, which is a generalization of the Mathieu equation. The subscript  $n = 0, 1, 2, \dots$  labels the discrete eigenvalues  $\mu = \mu_n$ . Each of the solutions  $gc_n$  and  $gs_n$  can be represented in the form of an infinite convergent trigonometric series in  $\cos n\theta$  and  $\sin n\theta$ , respectively; see Urvin and Arscott (1970). The functions  $f$ ,  $g$ , and  $h$  can be expressed in terms of the periodic solutions of the Whittaker–Hill equation as follows [Miller, Jr. (1977)]:

$$f(u) = \begin{cases} gc_n(iu; 2a\omega, \frac{1}{2}\beta/\omega), \\ gs_n(iu; 2a\omega, \frac{1}{2}\beta/\omega), \end{cases} \quad g(v) = \begin{cases} gc_n(v; 2a\omega, \frac{1}{2}\beta/\omega), \\ gs_n(v; 2a\omega, \frac{1}{2}\beta/\omega), \end{cases} \quad h(\varphi) = \begin{cases} gc_n(i\varphi + \frac{\pi}{2}; 2a\omega, \frac{1}{2}\beta/\omega), \\ gs_n(i\varphi + \frac{\pi}{2}; 2a\omega, \frac{1}{2}\beta/\omega), \end{cases}$$

where  $\omega = \sqrt{\lambda}$  and  $k = \mu_n - \frac{1}{2}a^2\lambda$ .

For the general ellipsoidal coordinates, the functions  $f$ ,  $g$ , and  $h$  are expressed in terms of the ellipsoidal wave functions; for details, see Arscott (1964) and Miller, Jr. (1977).

For the conical coordinate system, the functions  $g$  and  $h$  are determined by the Lamé equations that involve the Jacobian elliptic function  $\text{sn } z = \text{sn}(z, k)$ .

The unambiguity conditions for the transformation yield  $n = 0, 1, 2, \dots$ . It is known that, for any positive integer  $n$ , there exist exactly  $2n+1$  solutions corresponding to  $2n+1$  different eigenvalues  $\beta$ . These solutions can be represented the form of finite series known as Lamé polynomials. For more details about the Lamé equation and its solutions, see Whittaker and Watson (1963), Arscott (1964), Bateman and Erdélyi (1955), and Miller, Jr. (1977).

Unlike the Laplace equation, there are no nontrivial transformations for the three-dimensional Helmholtz equation that allow the  $\mathcal{R}$ -separation of variables.

© *References for Subsection 8.3.5:* F. M. Morse and H. Feshbach (1953, Vols. 1–2), P. Moon and D. Spencer (1961), A. Makarov, J. Smorodinsky, K. Valiev, and P. Winternitz (1967), W. Miller, Jr. (1977).

## 8.4. Other Equations with Three Space Variables

### 8.4.1. Equations Containing Arbitrary Functions

$$1. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} + \left(\lambda + \frac{a}{r}\right)w = 0, \quad r^2 = x^2 + y^2 + z^2.$$

*Schrödinger's equation.* It governs the motion of an electron in the Coulomb field of a nucleus ( $a > 0$ ).

The desired solutions must satisfy the normalizing condition

$$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} |w(x, y, z)|^2 dx dy dz = 1.$$

Eigenvalues:

$$\lambda_n = -\frac{a^2}{4n^2}; \quad n = 1, 2, 3, \dots$$

Normalized eigenfunctions (in the spherical coordinate system  $r, \theta, \varphi$ ):

$$w_{nmk} = \left(\frac{2}{n}\right)^{3/2} \sqrt{\frac{(2k+1)(k-m)!(n-k-1)!}{4\pi\epsilon_n n(n+k)!(m+k)!}} \left(\frac{ar}{n}\right)^k \exp\left(-\frac{ar}{2n}\right) L_{n-k-1}^{2k+1}\left(\frac{ar}{n}\right) Y_k^{(m)}(\theta, \varphi),$$

$$n = 1, 2, 3, \dots; \quad m = 0, \pm 1, \pm 2, \dots, \pm k; \quad k = 0, 1, 2, \dots, n-1;$$

where

$$\varepsilon_m = \begin{cases} 2 & \text{for } m = 0, \\ 1 & \text{for } m \neq 0, \end{cases} \quad Y_k^{(m)}(\theta, \varphi) = \begin{cases} P_k(\cos \theta) & \text{for } m = 0, \\ P_k^m(\cos \theta) \sin m\varphi & \text{for } m = 1, 2, \dots, \\ P_k^{|m|}(\cos \theta) \cos m\varphi & \text{for } m = -1, -2, \dots, \end{cases}$$

$$L_k^s(x) = \frac{1}{k!} x^{-s} e^x \frac{d^k}{dx^k} (x^{k+s} e^{-x}), \quad P_k^m(x) = (1-x^2)^{m/2} \frac{d^m}{dx^m} P_k(x), \quad P_k(x) = \frac{1}{k! 2^k} \frac{d^k}{dx^k} (x^2 - 1)^k.$$

These relations involve the generalized Laguerre polynomials  $L_k^s(x)$  and the associated Legendre functions  $P_n^m(\xi)$ ; the  $P_n(\xi)$  are the Legendre polynomials.

⊙ *References:* G. Korn and T. Korn (1968), A. N. Tikhonov and A. A. Samarskii (1990).

$$2. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} = a y \frac{\partial w}{\partial x}.$$

This equation is encountered in problems of convective heat and mass transfer in a simple shear flow.

Fundamental solution:

$$\mathcal{E}(x, y, z, \xi, \eta, \zeta) = \frac{1}{(4\pi)^{3/2}} \int_0^\infty \exp \left\{ -\frac{[x-\xi - \frac{1}{2}at(y+\eta)]^2}{4t(1+\frac{1}{12}a^2t^2)} - \frac{(y-\eta)^2 + (z-\zeta)^2}{4t} \right\} \frac{dt}{\sqrt{t^3(1+\frac{1}{12}a^2t^2)}}.$$

⊙ *References:* E. A. Novikov (1958), D. E. Elrick (1962).

$$3. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} + \frac{\partial^2 w}{\partial z^2} + a_1 x \frac{\partial w}{\partial x} + a_2 y \frac{\partial w}{\partial y} + a_3 z \frac{\partial w}{\partial z} = 0.$$

This equation is encountered in problems of convective heat and mass transfer in a straining flow.

Fundamental solution:

$$\mathcal{E}(x, y, z, \xi, \eta, \zeta) = \int_0^\infty F(x, \xi, t; a_1) F(y, \eta, t; a_2) F(z, \zeta, t; a_3) dt,$$

$$F(x, \xi, t; a) = \left[ \frac{2\pi}{a} (e^{2at} - 1) \right]^{-1/2} \exp \left[ -\frac{a(xe^{at} - \xi)^2}{2(e^{2at} - 1)} \right].$$

$$4. \quad \frac{\partial^2 w}{\partial x_1^2} + \frac{\partial^2 w}{\partial x_2^2} + \frac{\partial^2 w}{\partial x_3^2} = \sum_{n,k=1}^3 a_{nk} x_n \frac{\partial w}{\partial x_k}.$$

This equation is encountered in problems of convective heat and mass transfer in an arbitrary linear shear flow.

The solution that corresponds to a source of unit power at the origin of coordinates is given by

$$w(x_1, x_2, x_3) = \frac{1}{(4\pi)^{3/2}} \int_0^\infty \exp \left[ -\sum_{n,k=1}^3 \frac{b_{nk}(t) x_n x_k}{4D(t)} \right] \frac{dt}{\sqrt{D(t)}}.$$

Here,  $D = D(t)$  is the determinant of the matrix  $\mathbf{B} = \{B_{nk}\}$ ; the  $b_{nk} = b_{nk}(t)$  are the cofactors of the entries  $B_{nk} = B_{nk}(t)$ ; the  $B_{nk}$  are determined by solving the following system of ordinary differential equations with constant coefficients:

$$\frac{dB_{nk}}{dt} = \delta_{nk} + \sum_{m=1}^3 a_{nm} B_{km} + \sum_{m=1}^3 a_{km} B_{nm},$$

$$B_{nk} \rightarrow \delta_{nk} t \quad \text{as } t \rightarrow 0 \quad (\text{initial conditions}),$$

where  $\delta_{nn} = 1$  and  $\delta_{nk} = 0$  if  $n \neq k$ .

⊙ *Reference:* G. K. Batchelor (1979).

$$5. \quad \frac{\partial}{\partial x} \left[ f_1(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ f_2(y) \frac{\partial w}{\partial y} \right] + \frac{\partial}{\partial z} \left[ f_3(z) \frac{\partial w}{\partial z} \right] = \beta w.$$

This is a three-dimensional linear equation of heat and mass transfer theory with a source in an inhomogeneous anisotropic medium. Here,  $f_1 = f_1(x)$ ,  $f_2 = f_2(y)$ , and  $f_3 = f_3(z)$  are the principal thermal diffusivities.

1°. The equation admits multiplicatively separable solutions,  $w(x, y, z) = \varphi_1(x)\varphi_2(y)\varphi_3(z)$ .

2°. There are also additively separable solutions,  $w(x, y, z) = \psi_1(x) + \psi_2(y) + \psi_3(z)$ .

3°. If  $f_1 = ax^n$ ,  $f_2 = by^m$ , and  $f_3 = cz^k$  ( $n \neq 2$ ,  $m \neq 2$ ,  $k \neq 2$ ), there are particular solutions of the form

$$w = w(\xi), \quad \xi^2 = 4 \left[ \frac{x^{2-n}}{a(2-n)^2} + \frac{y^{2-m}}{b(2-m)^2} + \frac{z^{2-k}}{c(2-k)^2} \right],$$

where the function  $w(\xi)$  is determined by the ordinary differential equation

$$\frac{d^2 w}{d\xi^2} + \frac{A}{\xi} \frac{dw}{d\xi} = \beta w, \quad A = 2 \left( \frac{1}{2-n} + \frac{1}{2-m} + \frac{1}{2-k} \right) - 1,$$

whose solutions are expressed in terms of the Bessel functions.

### 8.4.2. Equations of the Form

$$\operatorname{div} [a(x, y, z) \nabla w] - q(x, y, z)w = -\Phi(x, y, z)$$

Equations of this sort are often encountered in heat and mass transfer theory. For brevity, the equation is written using the notation

$$\operatorname{div} [a(\mathbf{r}) \nabla w] = \frac{\partial}{\partial x} \left[ a(\mathbf{r}) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ a(\mathbf{r}) \frac{\partial w}{\partial y} \right] + \frac{\partial}{\partial z} \left[ a(\mathbf{r}) \frac{\partial w}{\partial z} \right], \quad \mathbf{r} = \{x, y, z\}.$$

In what follows, the problems for the equation in question will be considered in a bounded domain  $V$  with a sufficiently smooth surface  $S$ . It is assumed that  $a(\mathbf{r}) > 0$  and  $q(\mathbf{r}) \geq 0$ .

#### 8.4.2-1. First boundary value problem.

The following boundary condition of the first kind is imposed:

$$w = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in S.$$

Solution:

$$w(\mathbf{r}) = \int_V \Phi(\rho) G(\mathbf{r}, \rho) dV_\rho - \int_S f(\rho) a(\rho) \frac{\partial}{\partial N_\rho} G(\mathbf{r}, \rho) dS_\rho. \quad (1)$$

Here, the Green's function is given by

$$G(\mathbf{r}, \rho) = \sum_{n=1}^{\infty} \frac{u_n(\mathbf{r}) u_n(\rho)}{\|u_n\|^2 \lambda_n}, \quad \|u_n\|^2 = \int_V u_n^2(\mathbf{r}) dV, \quad \rho = \{\xi, \eta, \zeta\}, \quad (2)$$

where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and eigenfunctions of the Sturm–Liouville problem for the following second-order elliptic equation with a homogeneous boundary condition of the first kind:

$$\operatorname{div} [a(\mathbf{r}) \nabla u] - q(\mathbf{r})u + \lambda u = 0, \quad (3)$$

$$u = 0 \quad \text{for } \mathbf{r} \in S. \quad (4)$$

The integration in (1) is performed with respect to  $\xi, \eta, \zeta$ ;  $\frac{\partial}{\partial N_\rho}$  denotes the derivative along the outward normal to the surface  $S$  with respect to  $\xi, \eta, \zeta$ .

General properties of the Sturm–Liouville problem (3)–(4):

1°. There are countably many eigenvalues. All eigenvalues are real and can be ordered so that  $\lambda_1 \leq \lambda_2 \leq \lambda_3 \leq \dots$ , with  $\lambda_n \rightarrow \infty$  as  $n \rightarrow \infty$ ; therefore the number of negative eigenvalues is finite.

2°. If  $a(\mathbf{r}) > 0$  and  $q(\mathbf{r}) \geq 0$ , all eigenvalues are positive,  $\lambda_n > 0$ .

3°. The eigenfunctions are defined up to a constant multiplier. Any two eigenfunctions,  $u_n(\mathbf{r})$  and  $u_m(\mathbf{r})$ , corresponding to different eigenvalues,  $\lambda_n$  and  $\lambda_m$ , are orthogonal to each other in  $V$ :

$$\int_V u_n(\mathbf{r})u_m(\mathbf{r}) dV = 0 \quad \text{for } n \neq m.$$

4°. An arbitrary function  $F(\mathbf{r})$  that is twice continuously differentiable and satisfies the boundary condition of the Sturm–Liouville problem ( $F = 0$  for  $\mathbf{r} \in S$ ) can be expanded into an absolutely and uniformly convergent series in the eigenfunctions; specifically,

$$F(\mathbf{r}) = \sum_{n=1}^{\infty} F_n u_n(\mathbf{r}), \quad F_n = \frac{1}{\|u_n\|^2} \int_V F(\mathbf{r})u_n(\mathbf{r}) dV,$$

where the norm squared  $\|u_n\|^2$  is defined in (2).

*Remark.* In a three-dimensional problem, to each eigenvalue  $\lambda_n$  finitely many linearly independent eigenfunctions  $u_n^{(1)}, \dots, u_n^{(m)}$  generally correspond. These functions can always be replaced by their linear combinations

$$\bar{u}_n^{(k)} = A_{k,1}u_n^{(1)} + \dots + A_{k,k-1}u_n^{(k-1)} + u_n^{(k)}, \quad k = 1, 2, \dots, m,$$

such that  $\bar{u}_n^{(1)}, \dots, \bar{u}_n^{(m)}$  are now pairwise orthogonal. Therefore, without loss of generality, we can assume that all eigenfunctions are orthogonal.

#### 8.4.2-2. Second boundary value problem.

A boundary condition of the second kind is imposed,

$$\frac{\partial w}{\partial N} = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in S.$$

It is assumed that  $q(\mathbf{r}) > 0$ .

Solution:

$$w(\mathbf{r}) = \int_V \Phi(\rho)G(\mathbf{r}, \rho) dV_\rho + \int_S f(\rho)a(\rho)G(\mathbf{r}, \rho) dS_\rho. \quad (5)$$

Here, the Green's function is defined by relation (2), where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and eigenfunctions of the Sturm–Liouville problem for the second-order elliptic equation (3) with the following homogeneous boundary condition of the second kind:

$$\frac{\partial u}{\partial N} = 0 \quad \text{for } \mathbf{r} \in S. \quad (6)$$

If  $q(\mathbf{r}) > 0$ , the general properties of the eigenvalue problem (3), (6) are the same as those of the first boundary value problem (see Paragraph 8.4.2-1).

#### 8.4.2-3. Third boundary value problem.

The following boundary condition of the third kind is set:

$$\frac{\partial w}{\partial N} + k(\mathbf{r})w = f(\mathbf{r}) \quad \text{for } \mathbf{r} \in S.$$

The solution of the third boundary value problem is given by relations (5) and (2), where the  $\lambda_n$  and  $u_n(\mathbf{r})$  are the eigenvalues and eigenfunctions of the Sturm–Liouville problem for the

second-order elliptic equation (3) with the following homogeneous boundary condition of the third kind:

$$\frac{\partial u}{\partial N} + k(\mathbf{r})u = 0 \quad \text{for } \mathbf{r} \in S. \quad (7)$$

If  $q(\mathbf{r}) \geq 0$  and  $k(\mathbf{r}) > 0$ , the general properties of the eigenvalue problem (3), (7) are the same as those of the first boundary value problem (see Paragraph 8.4.2-1).

Let  $k(\mathbf{r}) = k = \text{const.}$  Denote the Green's functions of the second and third boundary value problems by  $G_2(\mathbf{r}, \rho)$  and  $G_3(\mathbf{r}, \rho, k)$ , respectively. For  $q(\mathbf{r}) > 0$ , the following limit relation holds:

$$G_2(\mathbf{r}, \rho) = \lim_{k \rightarrow 0} G_3(\mathbf{r}, \rho, k).$$

## 8.5. Equations with $n$ Space Variables

### 8.5.1. Laplace Equation $\Delta_n w = 0$

The  $n$ -dimensional Laplace equation in the rectangular Cartesian system of coordinates  $x_1, \dots, x_n$  has the form

$$\frac{\partial^2 w}{\partial x_1^2} + \frac{\partial^2 w}{\partial x_2^2} + \dots + \frac{\partial^2 w}{\partial x_n^2} = 0.$$

For  $n = 2$  and  $n = 3$ , see Subsections 7.1.1 and 8.1.1.

A regular solution of the Laplace equation is called a harmonic function.

In what follows we use the notation:  $\mathbf{x} = \{x_1, \dots, x_n\}$  and  $|\mathbf{x}| = \sqrt{x_1^2 + \dots + x_n^2}$ .

#### 8.5.1-1. Particular solutions.

1°. Fundamental solution:

$$\mathcal{E}(\mathbf{x}) = -\frac{1}{(n-2)\sigma_n|\mathbf{x}|^{n-2}}, \quad \sigma_n = \frac{2\pi^{n/2}}{\Gamma(n/2)} \quad (n \geq 3).$$

2°. Solution containing arbitrary functions of  $n-1$  variables:

$$w(x_1, \dots, x_n) = \sum_{k=0}^{\infty} (-1)^k \left[ \frac{x_n^{2k}}{(2k)!} \Delta^k f(x_1, \dots, x_{n-1}) + \frac{x_n^{2k+1}}{(2k+1)!} \Delta^k g(x_1, \dots, x_{n-1}) \right],$$

where  $f(x_1, \dots, x_{n-1})$  and  $g(x_1, \dots, x_{n-1})$  are arbitrary infinitely differentiable functions.

3°. Let  $w(x_1, \dots, x_n)$  be a harmonic function. Then the functions

$$w_1 = Aw(\pm\lambda x_1 + C_1, \dots, \pm\lambda x_n + C_n),$$

$$w_2 = \frac{A}{|\mathbf{x}|^{n-2}} w\left(\frac{x_1}{|\mathbf{x}|^2}, \dots, \frac{x_n}{|\mathbf{x}|^2}\right),$$

are also harmonic functions everywhere they are defined;  $A$ ,  $C_1, \dots, C_n$ , and  $\lambda$  are arbitrary constants. The signs at  $\lambda$  in the expression of  $w_1$  can be taken independently of one another.

⊙ *References:* A. V. Bitsadze and D. F. Kalinichenko (1985), R. Courant and D. Hilbert (1989).

#### 8.5.1-2. Domain: $-\infty < x_1 < \infty, \dots, -\infty < x_{n-1} < \infty, 0 \leq x_n < \infty$ .

The first boundary value problem for an  $n$ -dimensional half-space is considered. A boundary condition is prescribed:

$$w = f(x_1, \dots, x_{n-1}) \quad \text{at } x_n = 0.$$

Solution:

$$w(x_1, \dots, x_n) = \frac{\Gamma(n/2)}{\pi^{n/2}} \int_{-\infty}^{\infty} \dots \int_{-\infty}^{\infty} \left[ \sum_{k=1}^{n-1} (y_k - x_k)^2 + x_n^2 \right]^{-n/2} x_n f(y_1, \dots, y_{n-1}) dy_1 \dots dy_{n-1},$$

where  $\Gamma(z)$  is the gamma function.

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

### 8.5.1-3. Domain: $|\mathbf{x}| \leq 1$ . First boundary value problem.

A sphere of unit radius in the  $n$ -dimensional space is considered. A boundary condition is prescribed:

$$w = f(\mathbf{x}) \quad \text{for} \quad |\mathbf{x}| = 1.$$

Solution (Poisson integral):

$$w(\mathbf{x}) = \frac{\Gamma(n/2)}{2\pi^{n/2}} \int_{|\mathbf{y}|=1} \frac{1 - |\mathbf{x}|^2}{|\mathbf{y} - \mathbf{x}|^n} f(\mathbf{y}) dS_{\mathbf{y}}.$$

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

## 8.5.2. Other Equations

### 1. $\Delta_n w = -\Phi(x_1, \dots, x_n)$ .

This is the *Poisson equation* in  $n$  independent variables. For  $n = 2$  and  $n = 3$ , see Sections 7.2 and 8.2.

1°. Solution:

$$w(x_1, \dots, x_n) = \frac{\Gamma(n/2)}{2(n-2)\pi^{n/2}} \int_{\mathbb{R}^n} \frac{\Phi(y_1, \dots, y_n) dy_1 \dots dy_n}{[(x_1 - y_1)^2 + \dots + (x_n - y_n)^2]^{\frac{n-2}{2}}}.$$

⊙ *Reference:* S. G. Krein (1972).

### 2°. Domain: $0 \leq x_k \leq a_k$ ; $k = 1, \dots, n$ . First boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at} \quad x_k = 0, \\ w &= g_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at} \quad x_k = a_k. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x_1, \dots, x_n, y_1, \dots, y_n) &= \frac{2^n}{a_1 \dots a_n} \sum_{k_1=1}^{\infty} \dots \sum_{k_n=1}^{\infty} \frac{\sin(p_{k_1} x_1) \sin(p_{k_1} y_1) \dots \sin(p_{k_n} x_n) \sin(p_{k_n} y_n)}{p_{k_1}^2 + \dots + p_{k_n}^2}, \\ p_{k_1} &= \frac{\pi k_1}{a_1}, \quad p_{k_2} = \frac{\pi k_2}{a_2}, \quad \dots, \quad p_{k_n} = \frac{\pi k_n}{a_n}. \end{aligned}$$

### 3°. Domain: $0 \leq x_k \leq a_k$ ; $k = 1, \dots, n$ . Mixed boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at} \quad x_k = 0, \\ \partial_{x_k} w &= g_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at} \quad x_k = a_k. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x_1, \dots, x_n, y_1, \dots, y_n) &= \frac{2^n}{a_1 \dots a_n} \sum_{k_1=0}^{\infty} \dots \sum_{k_n=0}^{\infty} \frac{\sin(p_{k_1} x_1) \sin(p_{k_1} y_1) \dots \sin(p_{k_n} x_n) \sin(p_{k_n} y_n)}{p_{k_1}^2 + \dots + p_{k_n}^2}, \\ p_{k_1} &= \frac{\pi(2k_1 + 1)}{2a_1}, \quad p_{k_2} = \frac{\pi(2k_2 + 1)}{2a_2}, \quad \dots, \quad p_{k_n} = \frac{\pi(2k_n + 1)}{2a_n}. \end{aligned}$$

### 2. $\Delta_n w + \lambda w = 0$ .

This is the *Helmholtz equation* in  $n$  independent variables. For  $n = 2$  and  $n = 3$ , see Sections 7.3 and 8.3.

1°. Fundamental solution for  $\lambda = k^2 > 0$ :

$$\mathcal{E}(\mathbf{x}, \mathbf{y}) = \frac{k^{\frac{n-2}{2}}}{4(2\pi)^{\frac{n-2}{2}}} r^{-\frac{n-2}{2}} Y_{\frac{n-2}{2}}(kr), \quad r = |\mathbf{x} - \mathbf{y}| \quad \text{for even } n,$$

$$\mathcal{E}(\mathbf{x}, \mathbf{y}) = \frac{k^{\frac{n-2}{2}}}{4(2\pi)^{\frac{n-2}{2}} \sin(\frac{1}{2}\pi n)} r^{-\frac{n-2}{2}} J_{-\frac{n-2}{2}}(kr) \quad \text{for odd } n,$$

where  $J_\nu(z)$  and  $Y_\nu(z)$  are the Bessel functions.

2°. Domain:  $0 \leq x_k \leq a_k$ ;  $k = 1, \dots, n$ . First boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at } x_k = 0, \\ w &= g_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at } x_k = a_k. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x_1, \dots, x_n, y_1, \dots, y_n) &= \frac{2^n}{a_1 a_2 \dots a_n} \sum_{k_1=1}^{\infty} \dots \sum_{k_n=1}^{\infty} \frac{\sin(p_{k_1} x_1) \sin(p_{k_1} y_1) \dots \sin(p_{k_n} x_n) \sin(p_{k_n} y_n)}{p_{k_1}^2 + \dots + p_{k_n}^2 - \lambda}, \\ p_{k_1} &= \frac{\pi k_1}{a_1}, \quad p_{k_2} = \frac{\pi k_2}{a_2}, \quad \dots, \quad p_{k_n} = \frac{\pi k_n}{a_n}. \end{aligned}$$

3°. Domain:  $0 \leq x_k \leq a_k$ ;  $k = 1, \dots, n$ . Second boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} \partial_{x_k} w &= f_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at } x_k = 0, \\ \partial_{x_k} w &= g_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at } x_k = a_k. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x_1, \dots, x_n, y_1, \dots, y_n) &= \sum_{k_1=0}^{\infty} \dots \sum_{k_n=0}^{\infty} A_{k_1 k_2 \dots k_n} \frac{\cos(p_{k_1} x_1) \cos(p_{k_1} y_1) \dots \cos(p_{k_n} x_n) \cos(p_{k_n} y_n)}{p_{k_1}^2 + \dots + p_{k_n}^2 - \lambda}, \\ A_{k_1 k_2 \dots k_n} &= \frac{\varepsilon_{k_1} \varepsilon_{k_2} \dots \varepsilon_{k_n}}{a_1 a_2 \dots a_n}, \quad p_{k_1} = \frac{\pi k_1}{a_1}, \quad p_{k_2} = \frac{\pi k_2}{a_2}, \quad \dots, \quad p_{k_n} = \frac{\pi k_n}{a_n}, \quad \varepsilon_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases} \end{aligned}$$

4°. Domain:  $0 \leq x_k \leq a_k$ ;  $k = 1, \dots, n$ . Mixed boundary value problem.

A rectangular parallelepiped is considered. Boundary conditions are prescribed:

$$\begin{aligned} w &= f_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at } x_k = 0, \\ \partial_{x_k} w &= g_k(x_1, \dots, x_{k-1}, x_{k+1}, \dots, x_n) \quad \text{at } x_k = a_k. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x_1, \dots, x_n, y_1, \dots, y_n) &= \frac{2^n}{a_1 a_2 \dots a_n} \sum_{k_1=0}^{\infty} \dots \sum_{k_n=0}^{\infty} \frac{\sin(p_{k_1} x_1) \sin(p_{k_1} y_1) \dots \sin(p_{k_n} x_n) \sin(p_{k_n} y_n)}{p_{k_1}^2 + \dots + p_{k_n}^2 - \lambda}, \\ p_{k_1} &= \frac{\pi(2k_1 + 1)}{2a_1}, \quad p_{k_2} = \frac{\pi(2k_2 + 1)}{2a_2}, \quad \dots, \quad p_{k_n} = \frac{\pi(2k_n + 1)}{2a_n}. \end{aligned}$$

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

$$3. \quad \sum_{i,j=1}^n a_{ij} \frac{\partial^2 w}{\partial x_i \partial x_j} = 0.$$

It is assumed that for any real numbers  $y_1, \dots, y_n$  the relation  $\left| \sum_{i,j=1}^n a_{ij} y_i y_j \right| \geq k \sum_{i=1}^n y_i^2$  holds, where  $k$  is some positive constant.

Fundamental solution:

$$\mathcal{E}(x_1, \dots, x_n, y_1, \dots, y_n) = \begin{cases} \frac{\Gamma(n/2)}{2(n-2)\pi^{n/2}\sqrt{A}} \left[ \sum_{i,j=1}^n b_{ij}(x_i - y_i)(x_j - y_j) \right]^{-\frac{n-2}{2}} & \text{for } n \geq 3, \\ \frac{1}{2\pi\sqrt{A}} \ln \left[ \sum_{i,j=1}^2 b_{ij}(x_i - y_i)(x_j - y_j) \right]^{-1/2} & \text{for } n = 2, \end{cases}$$

where  $A$  is the determinant of the matrix  $\mathbf{A} = \{a_{ij}\}$  and the  $b_{ij}$  are the entries of the inverse of  $\mathbf{A}$ .

⊙ *Reference:* V. M. Babich, M. B. Kapilevich, S. G. Mikhlin, et al. (1964).

$$4. \quad \sum_{i=1}^{n-1} \frac{\partial^2 w}{\partial x_i^2} + \frac{\partial}{\partial x_n} \left( x_n^\beta \frac{\partial w}{\partial x_n} \right) + \lambda w = 0.$$

Domain:  $a_i \leq x_i \leq b_i$  ( $i = 1, \dots, n-1$ ),  $0 \leq x_n \leq c$ .

1°. Case  $0 < \beta < 1$ . First boundary value problem. The condition  $w = 0$  is set on the entire boundary of the domain.

Eigenvalues and eigenfunctions:

$$\lambda_{k_1, \dots, k_{n-1}, m} = \sum_{i=1}^{n-1} \frac{k_i^2 \pi^2}{(b_i - a_i)^2} + \frac{(2 - \beta)^2 \gamma_{\nu m}}{4c^{2-\beta}},$$

$$w_{k_1, \dots, k_{n-1}, m} = x_n^{\frac{1-\beta}{2}} J_\nu \left( \gamma_{\nu m} \left( \frac{x_n}{a} \right)^{\frac{2-\beta}{2}} \right) \prod_{i=1}^{n-1} \sin \frac{k_i \pi (x_i - a_i)}{b_i - a_i},$$

where  $\gamma_{\nu m}$  is the  $m$ th positive root of the equation  $J_\nu(\gamma) = 0$ ,

$$k_1, \dots, k_{n-1} = 1, 2, \dots; \quad m = 1, 2, \dots; \quad \nu = \frac{1-\beta}{2-\beta}.$$

2°. Case  $1 \leq \beta < 2$ . Boundary conditions: the solution must be bounded at  $x_n = 0$ , and the condition  $w = 0$  must hold on the rest of the boundary of the domain.

The eigenvalues and eigenfunctions of this problem are given by the relations of Item 1° with  $\nu = (\beta - 1)/(2 - \beta)$ .

⊙ *Reference:* M. M. Smirnov (1975).



## Chapter 9

# Higher-Order Partial Differential Equations

---

### 9.1. Third-Order Partial Differential Equations

1. 
$$\frac{\partial w}{\partial t} + \frac{\partial^3 w}{\partial x^3} = 0.$$

*Linearized Cortevég–de Vries equation.*

1°. Particular solutions:

$$w(x, t) = a(x^3 - 6t) + bx^2 + cx + k,$$

$$w(x, t) = a(x^5 - 60x^2t) + b(x^4 - 24xt),$$

$$w(x, t) = a \sin(\lambda x + \lambda^3 t) + b \cos(\lambda x + \lambda^3 t) + c,$$

$$w(x, t) = a \sinh(\lambda x - \lambda^3 t) + b \cosh(\lambda x - \lambda^3 t) + c,$$

$$w(x, t) = \exp(-\lambda^3 t) \left[ a \exp(\lambda x) + b \exp(-\frac{1}{2}\lambda x) \sin\left(\frac{\sqrt{3}}{2}\lambda x + c\right) \right],$$

where  $a, b, c, k$ , and  $\lambda$  are arbitrary constants.

2°. Domain:  $-\infty < x \leq 0$ . Boundary value problem.

Initial and boundary conditions are prescribed:

$$w = 0 \quad \text{at} \quad t = 0, \quad w = f(t) \quad \text{at} \quad x = 0, \quad w \rightarrow 0 \quad \text{as} \quad x \rightarrow -\infty.$$

Solution:

$$w(x, t) = -\frac{3}{2} \int_0^t \text{Ai}''\left(\frac{x}{(t-\tau)^{1/3}}\right) \frac{f(\tau)}{t-\tau} d\tau,$$

where  $\text{Ai}''(z)$  is the second derivative of the Airy function.

3°. Domain:  $0 \leq x < \infty$ . The function

$$w(x, t) = 3 \int_0^t \text{Ai}''\left(\frac{x}{(t-\tau)^{1/3}}\right) \frac{f(\tau)}{t-\tau} d\tau,$$

satisfies the equation and the first two conditions specified in Item 2°.

⊙ *Reference:* A. V. Faminskii (1999).

2. 
$$\frac{\partial w}{\partial t} = ax^6 \frac{\partial^3 w}{\partial x^3}.$$

The transformation

$$u(z, \tau) = wx^{-2}, \quad z = 1/x, \quad \tau = at$$

leads to a constant coefficient equation of the form 9.1.1:

$$\frac{\partial u}{\partial \tau} = -\frac{\partial^3 u}{\partial z^3}.$$

$$3. \quad \frac{\partial w}{\partial t} = k(t) \frac{\partial^3 w}{\partial x^3} + [x f(t) + g(t)] \frac{\partial w}{\partial x} + h(t)w.$$

The transformation

$$w(x, t) = u(z, \tau) \exp \left[ \int h(t) dt \right], \quad z = xF(t) + \int g(t)F(t) dt, \quad \tau = \int k(t)F^3(t) dt,$$

where  $F(t) = \exp \left[ \int f(t) dt \right]$ , leads to a constant coefficient equation of the form 9.1.1:

$$\frac{\partial u}{\partial \tau} = \frac{\partial^3 u}{\partial z^3}.$$

$$4. \quad \frac{\partial w}{\partial t} = (ax^2 + bx + c)^3 \frac{\partial^3 w}{\partial x^3}.$$

This is a special case of equation 9.6.4.4 with  $k = 1$  and  $n = 3$ . The transformation

$$w(x, t) = u(z, t)(ax^2 + bx + c), \quad z = \int \frac{dx}{ax^2 + bx + c}$$

leads to the constant coefficient equation

$$\frac{\partial u}{\partial t} = \frac{\partial^3 u}{\partial z^3} + (4ac - b^2) \frac{\partial u}{\partial z}.$$

$$5. \quad \frac{\partial^2 w}{\partial t^2} = ax^6 \frac{\partial^3 w}{\partial x^3}.$$

The transformation  $z = 1/x$ ,  $u = wx^{-2}$  leads to the constant coefficient equation

$$\frac{\partial^2 u}{\partial t^2} = -a \frac{\partial^3 u}{\partial z^3}.$$

$$6. \quad \frac{\partial^2 w}{\partial t^2} = (ax^2 + bx + c)^3 \frac{\partial^3 w}{\partial x^3}.$$

This is a special case of equation 9.6.4.4 with  $k = 2$  and  $n = 3$ . The transformation

$$w(x, t) = u(z, t)(ax^2 + bx + c), \quad z = \int \frac{dx}{ax^2 + bx + c}$$

leads to the constant coefficient equation

$$\frac{\partial^2 u}{\partial t^2} = \frac{\partial^3 u}{\partial z^3} + (4ac - b^2) \frac{\partial u}{\partial z}.$$

## 9.2. Fourth-Order One-Dimensional Nonstationary Equations

### 9.2.1. Equations of the Form $\frac{\partial w}{\partial t} + a^2 \frac{\partial^4 w}{\partial x^4} = \Phi(x, t)$

9.2.1-1. Particular solutions of the homogeneous equation ( $\Phi \equiv 0$ ):

$$w(x) = Ax^3 + Bx^2 + Cx + D,$$

$$w(x, t) = A(x^5 - 120a^2xt) + B(x^4 - 24a^2t),$$

$$w(x, t) = [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp(-\lambda^4 a^2 t),$$

where  $A$ ,  $B$ ,  $C$ ,  $D$ , and  $\lambda$  are arbitrary constants.

**9.2.1-2. Domain:  $0 \leq x \leq l$ . Solution in terms of the Green's function.**

1°. We consider problems on an interval  $0 \leq x \leq l$  with the general initial condition

$$w = f(x) \quad \text{at} \quad t = 0$$

and various homogeneous boundary conditions. The solution can be represented in terms of the Green's function as

$$w(x, t) = \int_0^l f(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau.$$

2°. Paragraphs 9.2.1-3 through 9.2.1-10 present the Green's functions for various types of boundary conditions. The Green's functions can be evaluated from the formula

$$G(x, \xi, t) = \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\|\varphi_n\|^2} \exp(-\lambda_n^4 a^2 t), \quad (1)$$

where the  $\lambda_n$  and  $\varphi_n(x)$  are determined by solving the self-adjoint eigenvalue problem for the fourth-order ordinary differential equation

$$\varphi'''' - \lambda^4 \varphi = 0$$

subject to appropriate boundary conditions; the prime denotes differentiation with respect to  $x$ . The norms of eigenfunctions can be calculated by the formula

$$\|\varphi_n\|^2 = \int_0^l \varphi_n^2(x) dx = \frac{l}{4} \varphi_n^2(l) + \frac{l}{4\lambda_n^4} [\varphi_n''(l)]^2 - \frac{l}{2\lambda_n^4} \varphi_n'(l) \varphi_n'''(l). \quad (2)$$

Relations (1) and (2) are written under the assumption that  $\lambda = 0$  is not an eigenvalue.

**9.2.1-3. The function and its first derivative are prescribed at the boundaries:**

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_x w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{l} \sum_{n=1}^{\infty} \frac{\lambda_n^4}{[\varphi_n''(l)]^2} \varphi_n(x) \varphi_n(\xi) \exp(-\lambda_n^4 a^2 t),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = 1$ . The numerical values of the roots can be calculated from the formulas given in Paragraph 9.2.3-2.

**9.2.1-4. The function and its second derivative are prescribed at the boundaries:**

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=1}^{\infty} \sin(\lambda_n x) \sin(\lambda_n \xi) \exp(-\lambda_n^4 a^2 t), \quad \lambda_n = \frac{\pi n}{l}.$$

**9.2.1-5. The first and third derivatives are prescribed at the boundaries:**

$$\partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{1}{l} + \frac{2}{l} \sum_{n=1}^{\infty} \cos(\lambda_n x) \cos(\lambda_n \xi) \exp(-\lambda_n^4 a^2 t), \quad \lambda_n = \frac{\pi n}{l}.$$

**9.2.1-6. The second and third derivatives are prescribed at the boundaries:**

$$w_{xx} = \partial_{xxx} w = 0 \quad \text{at} \quad x = 0, \quad w_{xx} = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{1}{l} + \frac{3}{l^3} (2x - l)(2\xi - l) + \frac{4}{l} \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\varphi_n^2(l)} \exp(-\lambda_n^4 a^2 t),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) + \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) + \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = 1$ . The numerical values of the roots can be calculated from the formulas given in Paragraph 9.2.3-2.

**9.2.1-7. Mixed conditions are prescribed at the boundaries (case 1):**

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=1}^{\infty} \lambda_n^4 \frac{\varphi_n(x) \varphi_n(\xi)}{|\varphi_n'(l) \varphi_n'''(l)|} \exp(-\lambda_n^4 a^2 t),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\tan(\lambda l) - \tanh(\lambda l) = 0$ .

**9.2.1-8. Mixed conditions are prescribed at the boundaries (case 2):**

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad \partial_{xx} w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{l} \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\varphi_n^2(l)} \exp(-\lambda_n^4 a^2 t),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) + \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) + \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = -1$ .

### 9.2.1-9. Mixed conditions are prescribed at the boundaries (case 3):

$$w = \partial_{xx}w = 0 \quad \text{at} \quad x = 0, \quad \partial_x w = \partial_{xxx}w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=0}^{\infty} \sin(\lambda_n x) \sin(\lambda_n \xi) \exp(-\lambda_n^4 a^2 t), \quad \lambda_n = \frac{\pi(2n+1)}{2l}.$$

### 9.2.1-10. Mixed conditions are prescribed at the boundaries (case 4):

$$w = \partial_{xx}w = 0 \quad \text{at} \quad x = 0, \quad \partial_{xx}w = \partial_{xxx}w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{l} \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\varphi_n^2(l)} \exp(-\lambda_n^4 a^2 t),$$

where

$$\varphi_n(x) = \sin(\lambda_n l) \sinh(\lambda_n x) + \sinh(\lambda_n l) \sin(\lambda_n x);$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\tan(\lambda l) - \tanh(\lambda l) = 0$ .

## 9.2.2. Equations of the Form $\frac{\partial^2 w}{\partial t^2} + a^2 \frac{\partial^4 w}{\partial x^4} = 0$

This equation is encountered in studying transverse vibration of elastic rods.

### 9.2.2-1. Particular solutions:

$$w(x, t) = (Ax^3 + Bx^2 + Cx + D)t + A_1x^3 + B_1x^2 + C_1x + D_1,$$

$$w(x, t) = [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \sin(\lambda^2 at),$$

$$w(x, t) = [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \cos(\lambda^2 at),$$

where  $A, B, C, D, A_1, B_1, C_1, D_1$ , and  $\lambda$  are arbitrary constants.

### 9.2.2-2. Domain: $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$w = f(x) \quad \text{at} \quad t = 0, \quad \partial_t w = ag''(x) \quad \text{at} \quad t = 0.$$

Boussinesq solution:

$$w(x, t) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x - 2\xi\sqrt{at}) (\cos \xi^2 + \sin \xi^2) d\xi \\ + \frac{1}{a\sqrt{2\pi}} \int_{-\infty}^{\infty} g(x - 2\xi\sqrt{at}) (\cos \xi^2 - \sin \xi^2) d\xi.$$

⊙ Reference: I. Sneddon (1951).

### 9.2.2-3. Domain: $0 \leq x < \infty$ . Free vibration of a semiinfinite rod.

The following conditions are prescribed:

$$w = 0 \quad \text{at} \quad t = 0, \quad \partial_t w = 0 \quad \text{at} \quad t = 0 \quad (\text{initial conditions}),$$

$$w = f(t) \quad \text{at} \quad x = 0, \quad \partial_{xx}w = 0 \quad \text{at} \quad x = 0 \quad (\text{boundary conditions}).$$

Boussinesq solution:

$$w(x, t) = \frac{1}{\sqrt{\pi}} \int_{x/\sqrt{2at}}^{\infty} f\left(t - \frac{x^2}{2a\xi^2}\right) \left(\sin \frac{\xi^2}{2} + \cos \frac{\xi^2}{2}\right) d\xi.$$

⊙ Reference: I. Sneddon (1951).

### 9.2.2-4. Domain: $0 \leq x \leq l$ . Boundary value problems.

For solutions of various boundary value problems, see Subsection 9.2.3 for  $\Phi \equiv 0$ .

## 9.2.3. Equations of the Form $\frac{\partial^2 w}{\partial t^2} + a^2 \frac{\partial^4 w}{\partial x^4} = \Phi(x, t)$

This equation is encountered in studying forced (transverse) vibration of elastic rods.

### 9.2.3-1. Domain: $0 \leq x \leq l$ . Solution in terms of the Green's function.

1°. We consider boundary value problems on an interval  $0 \leq x \leq l$  with the general initial condition

$$w = f(x) \quad \text{at} \quad t = 0, \quad \partial_t w = g(x) \quad \text{at} \quad t = 0$$

and various homogeneous boundary conditions. The solution can be represented in terms of the Green's function as

$$w(x, t) = \frac{\partial}{\partial t} \int_0^l f(\xi) G(x, \xi, t) d\xi + \int_0^l g(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau.$$

2°. Paragraphs 9.2.3-2 through 9.2.3-9 present the Green's functions for various types of boundary conditions. The Green's functions can be evaluated from the formula

$$G(x, \xi, t) = \frac{1}{a} \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\lambda_n^2 \|\varphi_n\|^2} \sin(\lambda_n^2 a t), \quad (1)$$

where the  $\lambda_n$  and  $\varphi_n(x)$  are determined by solving the self-adjoint eigenvalue problem for the fourth-order ordinary differential equation

$$\varphi'''' - \lambda^4 \varphi = 0$$

subject to appropriate boundary conditions; the prime denotes differentiation with respect to  $x$ . The norms of eigenfunctions can be calculated by Krylov's formula [see Krylov (1949)]:

$$\|\varphi_n\|^2 = \int_0^l \varphi_n^2(x) dx = \frac{l}{4} \varphi_n^2(l) + \frac{l}{4\lambda_n^4} [\varphi_n''(l)]^2 - \frac{l}{2\lambda_n^4} \varphi_n'(l) \varphi_n'''(l). \quad (2)$$

Relations (1) and (2) are written under the assumption that  $\lambda = 0$  is not an eigenvalue.

### 9.2.3-2. Both ends of the rod are clamped.

Boundary conditions are prescribed:

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_x w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{al} \sum_{n=1}^{\infty} \frac{\lambda_n^2}{[\varphi_n''(l)]^2} \varphi_n(x) \varphi_n(\xi) \sin(\lambda_n^2 a t),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = 1$ . The numerical values of the roots can be calculated from the formulas

$$\lambda_n = \frac{\mu_n}{l}, \quad \text{where} \quad \mu_1 = 1.875, \quad \mu_2 = 4.694, \quad \mu_n = \frac{\pi}{2}(2n-1) \quad \text{for} \quad n \geq 3.$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 9.2.3-3. Both ends of the rod are hinged.

Boundary conditions are prescribed:

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2l}{a\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \sin(\lambda_n x) \sin(\lambda_n \xi) \sin(\lambda_n^2 at), \quad \lambda_n = \frac{\pi n}{l}.$$

⊙ *References:* A. N. Krylov (1949), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 9.2.3-4. Both ends of the rod are free.

Boundary conditions are prescribed:

$$w_{xx} = \partial_{xxx} w = 0 \quad \text{at} \quad x = 0, \quad w_{xx} = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{t}{l} + \frac{3t}{l^3} (2x - l)(2\xi - l) + \frac{4}{al} \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\lambda_n^2 \varphi_n^2(l)} \sin(\lambda_n^2 at),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) + \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) + \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = 1$ . For the numerical values of the roots, see Paragraph 9.2.3-2.

The first two terms in the expression of the Green's function correspond to the zero eigenvalue  $\lambda_0 = 0$ , to which two orthogonal eigenfunctions  $w_0^{(1)} = 1$  and  $w_0^{(2)} = 2x - l$  correspond with  $\|w_0^{(1)}\|^2 = l$  and  $\|w_0^{(2)}\|^2 = \frac{1}{3}l^3$ .

⊙ *Reference:* A. N. Krylov (1949).

### 9.2.3-5. One end of the rod is clamped and the other is hinged.

Boundary conditions are prescribed:

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{al} \sum_{n=1}^{\infty} \lambda_n^2 \frac{\varphi_n(x) \varphi_n(\xi)}{|\varphi_n'(l) \varphi_n'''(l)|} \sin(\lambda_n^2 at),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\tan(\lambda l) - \tanh(\lambda l) = 0$ .

### 9.2.3-6. One end of the rod is clamped and the other is free.

Boundary conditions are prescribed:

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_{xx} w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{al} \sum_{n=1}^{\infty} \frac{\varphi_n(x)\varphi_n(\xi)}{\lambda_n^2 \varphi_n^2(l)} \sin(\lambda_n^2 at),$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) + \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) + \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = -1$ .

### 9.2.3-7. One end of the rod is hinged and the other is free.

Boundary conditions are prescribed:

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_{xx} w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{al} \sum_{n=1}^{\infty} \frac{\varphi_n(x)\varphi_n(\xi)}{\lambda_n^2 \varphi_n^2(l)} \sin(\lambda_n^2 at),$$

where

$$\varphi_n(x) = \sin(\lambda_n l) \sinh(\lambda_n x) + \sinh(\lambda_n l) \sin(\lambda_n x);$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\tan(\lambda l) - \tanh(\lambda l) = 0$ .

### 9.2.3-8. The first and third derivatives are prescribed at the ends:

$$\partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{t}{l} + \frac{2}{al} \sum_{n=1}^{\infty} \frac{1}{\lambda_n^2} \cos(\lambda_n x) \cos(\lambda_n \xi) \sin(\lambda_n^2 at), \quad \lambda_n = \frac{\pi n}{l}.$$

### 9.2.3-9. Mixed boundary conditions are prescribed at the ends:

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{al} \sum_{n=0}^{\infty} \frac{1}{\lambda_n^2} \sin(\lambda_n x) \sin(\lambda_n \xi) \sin(\lambda_n^2 at), \quad \lambda_n = \frac{\pi(2n+1)}{2l}.$$

## 9.2.4. Equations of the Form $\frac{\partial^2 w}{\partial t^2} + a^2 \frac{\partial^4 w}{\partial x^4} + kw = \Phi(x, t)$

### 9.2.4-1. Particular solutions of the homogeneous equation ( $\Phi \equiv 0$ ):

$$w(x, t) = (Ax^3 + Bx^2 + Cx + D) \sin(t\sqrt{k}),$$

$$w(x, t) = (Ax^3 + Bx^2 + Cx + D) \cos(t\sqrt{k}),$$

$$w(x, t) = [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \sin(t\sqrt{a^2 \lambda^4 + k}),$$

$$w(x, t) = [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \cos(t\sqrt{a^2 \lambda^4 + k}),$$

where  $A, B, C, D$ , and  $\lambda$  are arbitrary constants.



**9.2.4-2. Domain:  $0 \leq x \leq l$ . Solution in terms of the Green's function.**

1°. We consider boundary value problems on an interval  $0 \leq x \leq l$  with the general initial condition

$$w = f(x) \quad \text{at} \quad t = 0, \quad \partial_t w = g(x) \quad \text{at} \quad t = 0$$

and various homogeneous boundary conditions. The solution can be represented in terms of the Green's function as

$$w(x, t) = \frac{\partial}{\partial t} \int_0^l f(\xi) G(x, \xi, t) d\xi + \int_0^l g(\xi) G(x, \xi, t) d\xi + \int_0^t \int_0^l \Phi(\xi, \tau) G(x, \xi, t - \tau) d\xi d\tau.$$

2°. Paragraphs 9.2.4-3 through 9.2.4-10 present the Green's functions for various types of boundary conditions. The Green's functions can be evaluated from the formula

$$G(x, \xi, t) = \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\|\varphi_n\|^2} \frac{\sin(t \sqrt{a^2 \lambda_n^4 + k})}{\sqrt{a^2 \lambda_n^4 + k}},$$

where the  $\lambda_n$  and  $\varphi_n(x)$  are determined by solving the self-adjoint eigenvalue problem for the fourth-order ordinary differential equation  $\varphi'''' - \lambda^4 \varphi = 0$  subject to appropriate boundary conditions. The norms of eigenfunctions can be calculated by formula (2) from Paragraph 9.2.3-1.

**9.2.4-3. The function and its first derivative are prescribed at the ends:**

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_x w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{l} \sum_{n=1}^{\infty} \lambda_n^4 \frac{\varphi_n(x) \varphi_n(\xi)}{[\varphi_n''(l)]^2} \frac{\sin(t \sqrt{a^2 \lambda_n^4 + k})}{\sqrt{a^2 \lambda_n^4 + k}}, \quad \varphi_n''(x) = \frac{d^2 \varphi_n}{dx^2},$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = 1$ .

**9.2.4-4. The function and its second derivative are prescribed at the ends:**

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=1}^{\infty} \sin(\lambda_n x) \sin(\lambda_n \xi) \frac{\sin(t \sqrt{a^2 \lambda_n^4 + k})}{\sqrt{a^2 \lambda_n^4 + k}}, \quad \lambda_n = \frac{\pi n}{l}.$$

**9.2.4-5. The first and third derivatives are prescribed at the ends:**

$$\partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{\sin(t \sqrt{k})}{l \sqrt{k}} + \frac{2}{l} \sum_{n=1}^{\infty} \cos(\lambda_n x) \cos(\lambda_n \xi) \frac{\sin(t \sqrt{a^2 \lambda_n^4 + k})}{\sqrt{a^2 \lambda_n^4 + k}}, \quad \lambda_n = \frac{\pi n}{l}.$$

**9.2.4-6. The second and third derivatives are prescribed at the ends:**

$$w_{xx} = \partial_{xxx}w = 0 \quad \text{at} \quad x = 0, \quad w_{xx} = \partial_{xxx}w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \left[ 1 + \frac{3}{l^2}(2x-l)(2\xi-l) \right] \frac{\sin(t\sqrt{k})}{l\sqrt{k}} + \frac{4}{l} \sum_{n=1}^{\infty} \frac{\varphi_n(x)\varphi_n(\xi)}{\varphi_n^2(l)} \frac{\sin(t\sqrt{a^2\lambda_n^4+k})}{\sqrt{a^2\lambda_n^4+k}},$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) + \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) + \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = 1$ . For the numerical values of the roots, see Paragraph 9.2.3-2.

**9.2.4-7. Mixed boundary conditions are prescribed at the ends (case 1):**

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=1}^{\infty} \lambda_n^4 \frac{\varphi_n(x)\varphi_n(\xi)}{|\varphi_n'(l)\varphi_n'''(l)|} \frac{\sin(t\sqrt{a^2\lambda_n^4+k})}{\sqrt{a^2\lambda_n^4+k}},$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) - \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) - \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\tan(\lambda l) - \tanh(\lambda l) = 0$ .

**9.2.4-8. Mixed boundary conditions are prescribed at the ends (case 2):**

$$w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad \partial_{xx} w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{l} \sum_{n=1}^{\infty} \frac{\varphi_n(x)\varphi_n(\xi)}{\varphi_n^2(l)} \frac{\sin(t\sqrt{a^2\lambda_n^4+k})}{\sqrt{a^2\lambda_n^4+k}},$$

where

$$\varphi_n(x) = [\sinh(\lambda_n l) + \sin(\lambda_n l)] [\cosh(\lambda_n x) - \cos(\lambda_n x)] - [\cosh(\lambda_n l) + \cos(\lambda_n l)] [\sinh(\lambda_n x) - \sin(\lambda_n x)];$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\cosh(\lambda l) \cos(\lambda l) = -1$ .

**9.2.4-9. Mixed boundary conditions are prescribed at the ends (case 3):**

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{2}{l} \sum_{n=0}^{\infty} \sin(\lambda_n x) \sin(\lambda_n \xi) \frac{\sin(t\sqrt{a^2\lambda_n^4+k})}{\sqrt{a^2\lambda_n^4+k}}, \quad \lambda_n = \frac{\pi(2n+1)}{2l}.$$

**9.2.4-10. Mixed boundary conditions are prescribed at the ends (case 4):**

$$w = \partial_{xx}w = 0 \quad \text{at} \quad x = 0, \quad \partial_{xx}w = \partial_{xxx}w = 0 \quad \text{at} \quad x = l.$$

Green's function:

$$G(x, \xi, t) = \frac{4}{l} \sum_{n=1}^{\infty} \frac{\varphi_n(x)\varphi_n(\xi)}{\varphi_n^2(l)} \frac{\sin(t\sqrt{a^2\lambda_n^4 + k})}{\sqrt{a^2\lambda_n^4 + k}},$$

where

$$\varphi_n(x) = \sin(\lambda_n l) \sinh(\lambda_n x) + \sinh(\lambda_n l) \sin(\lambda_n x);$$

the  $\lambda_n$  are positive roots of the transcendental equation  $\tan(\lambda l) - \tanh(\lambda l) = 0$ .

## 9.2.5. Other Equations

**9.2.5-1. Equations containing the first derivative with respect to  $t$ .**

$$1. \quad \frac{\partial w}{\partial t} + a^2 \frac{\partial^4 w}{\partial x^4} + kw = \Phi(x, t).$$

The change of variable  $w(x, t) = e^{-kt}u(x, t)$  leads to the equation

$$\frac{\partial u}{\partial t} + a^2 \frac{\partial^4 u}{\partial x^4} = e^{kt}\Phi(x, t),$$

which is discussed in Subsection 9.2.1.

$$2. \quad \frac{\partial w}{\partial t} = ax^8 \frac{\partial^4 w}{\partial x^4}.$$

This is a special case of equation 9.6.4.2 with  $k = 1$  and  $n = 4$ .

$$3. \quad \frac{\partial w}{\partial t} = k(t) \frac{\partial^4 w}{\partial x^4} + [xf(t) + g(t)] \frac{\partial w}{\partial x} + h(t)w.$$

This is a special case of equation 9.6.4.1 with  $n = 4$ . The transformation

$$w(x, t) = u(z, \tau) \exp \left[ \int h(t) dt \right], \quad z = xF(t) + \int g(t)F(t) dt, \quad \tau = \int k(t)F^4(t) dt,$$

where  $F(t) = \exp \left[ \int f(t) dt \right]$ , leads to the constant coefficient equation

$$\frac{\partial u}{\partial \tau} = \frac{\partial^4 u}{\partial z^4},$$

which is discussed in Subsection 9.2.1.

$$4. \quad \frac{\partial w}{\partial t} = (ax^2 + bx + c)^4 \frac{\partial^4 w}{\partial x^4}.$$

This is a special case of equation 9.6.4.4 with  $k = 1$  and  $n = 4$ .

$$5. \quad \frac{\partial^2 w}{\partial t^2} + k \frac{\partial w}{\partial t} + a^2 \frac{\partial^4 w}{\partial x^4} = \Phi(x, t).$$

With  $\Phi(x, t) \equiv 0$  this equation governs transverse vibration of an elastic rod in a resisting medium with velocity-proportional resistance coefficient.

The change of variable  $w(x, t) = \exp(-\frac{1}{2}kt)u(x, t)$  leads to the equation

$$\frac{\partial^2 u}{\partial t^2} + a^2 \frac{\partial^4 u}{\partial x^4} - \frac{1}{4}k^2 u = \exp(\frac{1}{2}kt)\Phi(x, t),$$

which is discussed in Subsection 9.2.4.

$$6. \quad \frac{\partial^2 w}{\partial t^2} = ax^8 \frac{\partial^4 w}{\partial x^4}.$$

This is a special case of equation 9.6.4.2 with  $k = 2$  and  $n = 4$ .

$$7. \quad \frac{\partial^2 w}{\partial t^2} = (ax^2 + bx + c)^4 \frac{\partial^4 w}{\partial x^4}.$$

This is a special case of equation 9.6.4.4 with  $k = 2$  and  $n = 4$ .

$$8. \quad \left( \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right)^2 w = 0.$$

1°. General solution (two representations):

$$w(x, t) = tu_1(x, t) + u_0(x, t),$$

$$w(x, t) = xu_1(x, t) + u_0(x, t),$$

where  $u_k = u_k(x, t)$  is an arbitrary function satisfying the heat equation  $\partial_t u_k - \partial_{xx} u_k = 0$ ;  $k = 1, 2$ .

2°. Fundamental solution:

$$\mathcal{E}(x, t) = \frac{\sqrt{t}}{2\sqrt{\pi}} \exp\left(-\frac{x^2}{4t}\right).$$

3°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$w = 0 \quad \text{at} \quad t = 0, \quad \partial_t w = f(x) \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, t) = \frac{\sqrt{t}}{2\sqrt{\pi}} \int_{-\infty}^{\infty} \exp\left[-\frac{(x-\xi)^2}{4t}\right] f(\xi) d\xi.$$

⊙ Reference: G. E. Shilov (1965).

$$9. \quad \frac{\partial^4 w}{\partial t^4} - \frac{\partial^4 w}{\partial x^4} = 0.$$

1°. Fundamental solution:

$$\mathcal{E}(x, t) = \frac{1}{2\pi} \left\{ t \ln \sqrt{x^2 + t^2} - x \arctan \frac{x}{t} - \frac{1}{2}(t+x) \ln |t+x| \right. \\ \left. - \frac{1}{2}(t-x) \ln |t-x| + \frac{1}{8}|t+x| + \frac{1}{8}|t-x| \right\}.$$

2°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$w = 0 \quad \text{at} \quad t = 0, \quad \partial_t w = 0 \quad \text{at} \quad t = 0, \quad \partial_{tt} w = f(x) \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, t) = \int_{-\infty}^{\infty} \mathcal{E}(x-\xi, t) f(\xi) d\xi.$$

⊙ Reference: G. E. Shilov (1965).

$$10. \quad \frac{\partial^4 w}{\partial t^4} - 2 \frac{\partial^4 w}{\partial t^2 \partial x^2} + \frac{\partial^4 w}{\partial x^4} = 0.$$

General solution (three representations):

$$\begin{aligned} w(x, t) &= f_1(t-x) + f_2(t+x) + t[g_1(t-x) + g_2(t+x)], \\ w(x, t) &= f_1(t-x) + f_2(t+x) + x[g_1(t-x) + g_2(t+x)], \\ w(x, t) &= f_1(t-x) + f_2(t+x) + (t+x)g_1(t-x) + (t-x)g_2(t+x), \end{aligned}$$

where  $f_1(y)$ ,  $f_2(z)$ ,  $g_1(y)$ , and  $g_2(z)$  are arbitrary functions.

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

## 9.3. Two-Dimensional Nonstationary Fourth-Order Equations

### 9.3.1. Equations of the Form $\frac{\partial w}{\partial t} + a^2 \left( \frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} \right) = \Phi(x, y, t)$

9.3.1-1. Domain:  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$ . Solution in terms of the Green's function.

We consider boundary value problems in a rectangular domain  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$  with the general initial condition

$$w = f(x, y) \quad \text{at} \quad t = 0$$

and various homogeneous boundary conditions. The solution can be represented in terms of the Green's function as

$$w(x, y, t) = \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, t - \tau) d\eta d\xi d\tau.$$

Below are the Green's functions for various types of boundary conditions.

9.3.1-2. The function and its first derivatives are prescribed at the sides of a rectangle:

$$\begin{aligned} w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_x w = 0 \quad \text{at} \quad x = l_1, \\ w = \partial_y w = 0 \quad \text{at} \quad y = 0, \quad w = \partial_y w = 0 \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{16}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{p_n^4 q_m^4}{[\varphi_n''(l_1) \psi_m''(l_2)]^2} \varphi_n(x) \psi_m(y) \varphi_n(\xi) \psi_m(\eta) \exp[-(p_n^4 + q_m^4) a^2 t], \\ \varphi_n''(x) &= \frac{d^2 \varphi_n}{dx^2}, \quad \psi_m''(y) = \frac{d^2 \psi_m}{dy^2}. \end{aligned}$$

Here,

$$\begin{aligned} \varphi_n(x) &= [\sinh(p_n l_1) - \sin(p_n l_1)] [\cosh(p_n x) - \cos(p_n x)] \\ &\quad - [\cosh(p_n l_1) - \cos(p_n l_1)] [\sinh(p_n x) - \sin(p_n x)], \\ \psi_m(y) &= [\sinh(q_m l_2) - \sin(q_m l_2)] [\cosh(q_m y) - \cos(q_m y)] \\ &\quad - [\cosh(q_m l_2) - \cos(q_m l_2)] [\sinh(q_m y) - \sin(q_m y)], \end{aligned}$$

where the  $p_n$  and  $q_m$  are positive roots of the transcendental equations

$$\cosh(pl_1) \cos(pl_1) = 1, \quad \cosh ql_2 \cos ql_2 = 1 \quad (q_m = p_m l_1 / l_2).$$

9.3.1-3. The function and its second derivatives are prescribed at the sides of a rectangle:

$$\begin{aligned} w = \partial_{xx}w = 0 & \quad \text{at} \quad x = 0, & w = \partial_{xx}w = 0 & \quad \text{at} \quad x = l_1, \\ w = \partial_{yy}w = 0 & \quad \text{at} \quad y = 0, & w = \partial_{yy}w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \exp[-(p_n^4 + q_m^4) a^2 t],$$

$$p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}.$$

9.3.1-4. The first and third derivatives are prescribed at the sides of a rectangle:

$$\begin{aligned} w_x = \partial_{xxx}w = 0 & \quad \text{at} \quad x = 0, & w_x = \partial_{xxx}w = 0 & \quad \text{at} \quad x = l_1, \\ w_y = \partial_{yyy}w = 0 & \quad \text{at} \quad y = 0, & w_y = \partial_{yyy}w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{1}{l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \varepsilon_n \varepsilon_m \cos(p_n x) \sin(q_m y) \cos(p_n \xi) \cos(q_m \eta) \exp[-(p_n^4 + q_m^4) a^2 t],$$

$$p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \varepsilon_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}$$

9.3.1-5. The second and third derivatives are prescribed at the sides of a rectangle:

$$\begin{aligned} w_{xx} = \partial_{xxx}w = 0 & \quad \text{at} \quad x = 0, & w_{xx} = \partial_{xxx}w = 0 & \quad \text{at} \quad x = l_1, \\ w_{yy} = \partial_{yyy}w = 0 & \quad \text{at} \quad y = 0, & w_{yy} = \partial_{yyy}w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta, t) = G_1(x, \xi, t) G_2(y, \eta, t),$$

$$G_1(x, \xi, t) = \frac{1}{l_1} + \frac{3}{l_1^3} (2x - l_1)(2\xi - l_1) + \frac{4}{l_1} \sum_{n=1}^{\infty} \frac{\varphi_n(x) \varphi_n(\xi)}{\varphi_n^2(l_1)} \exp(-p_n^4 a^2 t),$$

$$G_2(y, \eta, t) = \frac{1}{l_2} + \frac{3}{l_2^3} (2y - l_2)(2\eta - l_2) + \frac{4}{l_2} \sum_{m=1}^{\infty} \frac{\psi_m(y) \psi_m(\eta)}{\psi_m^2(l_2)} \exp(-q_m^4 a^2 t).$$

Here,

$$\begin{aligned} \varphi_n(x) &= [\sinh(p_n l_1) - \sin(p_n l_1)] [\cosh(p_n x) + \cos(p_n x)] \\ &\quad - [\cosh(p_n l_1) - \cos(p_n l_1)] [\sinh(p_n x) + \sin(p_n x)], \\ \psi_m(y) &= [\sinh(q_m l_2) - \sin(q_m l_2)] [\cosh(q_m y) + \cos(q_m y)] \\ &\quad - [\cosh(q_m l_2) - \cos(q_m l_2)] [\sinh(q_m y) + \sin(q_m y)], \end{aligned}$$

where the  $p_n$  and  $q_m$  are positive roots of the transcendental equations

$$\cosh(pl_1) \cos(pl_1) = 1, \quad \cosh(ql_2) \cos(ql_2) = 1.$$

**9.3.1-6. Mixed boundary conditions are prescribed at the sides of a rectangle:**

$$\begin{aligned} w = \partial_{xx}w = 0 & \quad \text{at} \quad x = 0, & \partial_x w = \partial_{xxx}w = 0 & \quad \text{at} \quad x = l_1, \\ w = \partial_{yy}w = 0 & \quad \text{at} \quad y = 0, & \partial_y w = \partial_{yyy}w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{4}{l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \exp[-(p_n^4 + q_m^4) a^2 t],$$

$$p_n = \frac{\pi(2n+1)}{2l_1}, \quad q_m = \frac{\pi(2m+1)}{2l_2}.$$

**9.3.2. Two-Dimensional Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \Delta \Delta w = 0$**

This equation governs two-dimensional free transverse vibration of a thin elastic plate; the unknown  $w$  is the deflection (transverse displacement) of the plate's midplane points relative to the original plane position. Here,  $\Delta \Delta = \Delta^2$  and  $\Delta$  is the Laplace operator that is defined as

$$\Delta = \begin{cases} \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} & \text{in the Cartesian coordinate system,} \\ \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \varphi^2} & \text{in the polar coordinate system.} \end{cases}$$

**9.3.2-1. Particular solutions:**

$$\begin{aligned} w(x, y, t) &= [A_1 \sin(k_1 x) + B_1 \cos(k_1 x)] [A_2 \sin(k_2 y) + B_2 \cos(k_2 y)] \sin[(k_1^2 + k_2^2)at], \\ w(x, y, t) &= [A_1 \sin(k_1 x) + B_1 \cos(k_1 x)] [A_2 \sin(k_2 y) + B_2 \cos(k_2 y)] \cos[(k_1^2 + k_2^2)at], \\ w(x, y, t) &= [A_1 \sinh(k_1 x) + B_1 \cosh(k_1 x)] [A_2 \sinh(k_2 y) + B_2 \cosh(k_2 y)] \sin[(k_1^2 + k_2^2)at], \\ w(x, y, t) &= [A_1 \sinh(k_1 x) + B_1 \cosh(k_1 x)] [A_2 \sinh(k_2 y) + B_2 \cosh(k_2 y)] \cos[(k_1^2 + k_2^2)at], \\ w(r, \varphi, t) &= [A_1 J_n(kr) + A_2 Y_n(kr) + A_3 I_n(kr) + A_4 K_n(kr)] \cos(n\varphi) \sin(k^2 at), \\ w(r, \varphi, t) &= [A_1 J_n(kr) + A_2 Y_n(kr) + A_3 I_n(kr) + A_4 K_n(kr)] \sin(n\varphi) \cos(k^2 at), \end{aligned}$$

where  $A_1, A_2, A_3, A_4, B_1, B_2, k, k_1, k_2$  are arbitrary constants, the  $J_n(\xi)$  and  $Y_n(\xi)$  are the Bessel functions of the first and second kind, the  $I_n(\xi)$  and  $K_n(\xi)$  are the modified Bessel functions of the first and second kind,  $r = \sqrt{x^2 + y^2}$ , and  $n = 0, 1, 2, \dots$

**9.3.2-2. Domain:  $-\infty < x < \infty, -\infty < y < \infty$ . Cauchy problem.**

Initial conditions are prescribed:

$$w = f(x, y) \quad \text{at} \quad t = 0, \quad \partial_t w = g(x, y) \quad \text{at} \quad t = 0.$$

Poisson solution:

$$\begin{aligned} w(x, y, t) &= \frac{1}{\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x + 2\xi\sqrt{a\tau}, y + 2\eta\sqrt{a\tau}) \sin(\xi^2 + \eta^2) d\xi d\eta \\ &+ \frac{1}{\pi} \int_0^t d\tau \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x + 2\xi\sqrt{a\tau}, y + 2\eta\sqrt{a\tau}) \sin(\xi^2 + \eta^2) d\xi d\eta. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{1}{4\pi a} \int_0^t \sin\left[\frac{(x-\xi)^2 + (y-\eta)^2}{4a\tau}\right] \frac{d\tau}{\tau}.$$

⊙ *References:* A. N. Krylov (1949), I. Sneddon (1951), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**9.3.2-3. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Solution in terms of the Green's function.**

We consider boundary value problems in a rectangular domain  $0 \leq x \leq l_1, 0 \leq y \leq l_2$  with the general initial conditions

$$w = f(x, y) \quad \text{at} \quad t = 0, \quad \partial_t w = g(x, y) \quad \text{at} \quad t = 0$$

and various homogeneous boundary conditions. The solution can be represented in terms of the Green's function as

$$w(x, y, t) = \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} g(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi.$$

Paragraphs 9.3.2-4 through 9.3.2-6 present the Green's functions for three types of boundary conditions.

**9.3.2-4. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . All sides of the plate are hinged.**

Boundary conditions are prescribed:

$$\begin{aligned} w = \partial_{xx} w = 0 & \quad \text{at} \quad x = 0, & w = \partial_{xx} w = 0 & \quad \text{at} \quad x = l_1, \\ w = \partial_{yy} w = 0 & \quad \text{at} \quad y = 0, & w = \partial_{yy} w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{4}{al_1l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \frac{\sin(\lambda_{nm} at)}{\lambda_{nm}}, \\ p_n &= \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = p_n^2 + q_m^2. \end{aligned}$$

**9.3.2-5. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . The 1st and 3rd derivatives are prescribed at the sides:**

$$\begin{aligned} \partial_x w = \partial_{xxx} w = 0 & \quad \text{at} \quad x = 0, & \partial_x w = \partial_{xxx} w = 0 & \quad \text{at} \quad x = l_1, \\ \partial_y w = \partial_{yyy} w = 0 & \quad \text{at} \quad y = 0, & \partial_y w = \partial_{yyy} w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{1}{al_1l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \varepsilon_n \varepsilon_m \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) \frac{\sin(\lambda_{nm} at)}{\lambda_{nm}}, \\ p_n &= \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = p_n^2 + q_m^2, \quad \varepsilon_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases} \end{aligned}$$

If  $n = m = 0$ , the ratio  $\sin(\lambda_{nm} at)/\lambda_{nm}$  must be replaced by  $at$ .

**9.3.2-6. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Mixed boundary conditions are set at the sides:**

$$\begin{aligned} w = \partial_{xx} w = 0 & \quad \text{at} \quad x = 0, & w = \partial_{xx} w = 0 & \quad \text{at} \quad x = l_1, \\ \partial_y w = \partial_{yyy} w = 0 & \quad \text{at} \quad y = 0, & \partial_y w = \partial_{yyy} w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{2}{al_1l_2} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \varepsilon_m \sin(p_n x) \cos(q_m y) \sin(p_n \xi) \cos(q_m \eta) \frac{\sin(\lambda_{nm} at)}{\lambda_{nm}}, \\ p_n &= \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = p_n^2 + q_m^2, \quad \varepsilon_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases} \end{aligned}$$



**9.3.2-7. Domain:  $0 \leq r < \infty$ ,  $0 \leq \varphi \leq 2\pi$ . Cauchy problem.**

Initial conditions for the symmetric case in the polar coordinate system:

$$w = f(r) \quad \text{at} \quad t = 0, \quad \partial_t w = 0 \quad \text{at} \quad t = 0.$$

Solution:

$$w(r, t) = \frac{1}{2at} \int_0^\infty \xi f(\xi) J_0\left(\frac{\xi r}{2at}\right) \sin\left(\frac{\xi^2 + r^2}{4at}\right) d\xi,$$

where  $J_0(z)$  is the zeroth Bessel function.

⊙ *References:* I. Sneddon (1951), B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

**9.3.2-8. Domain:  $0 \leq r \leq R$ ,  $0 \leq \varphi \leq 2\pi$ . Transverse vibration of a circular plate.**

Initial and boundary conditions for symmetric transverse vibrations of a circular plate of radius  $R$  with clamped contour in the polar coordinate system:

$$\begin{aligned} w &= f(r) \quad \text{at} \quad t = 0, & \partial_t w &= g(t) \quad \text{at} \quad t = 0; \\ w &= 0 \quad \text{at} \quad r = R, & \partial_r w &= 0 \quad \text{at} \quad r = R. \end{aligned}$$

Solution:

$$\begin{aligned} w(r, t) &= \sum_{n=1}^{\infty} [A_n \cos(ak_n^2 t) + B_n \sin(ak_n^2 t)] \Psi_n(r), \\ \Psi_n(r) &= I_0(k_n R) J_0(k_n r) - J_0(k_n R) I_0(k_n r), \end{aligned}$$

where the  $k_n$  are positive roots of the transcendental equation (the prime denotes the derivative)

$$J_0(kR) I_0'(kR) - I_0(kR) J_0'(kR) = 0,$$

and the coefficients  $A_n$  and  $B_n$  are given by

$$\begin{aligned} A_n &= \frac{1}{\|\Psi_n\|^2} \int_0^R f(r) \Psi_n(r) r dr, \quad B_n = \frac{1}{ak_n^2 \|\Psi_n\|^2} \int_0^R g(r) \Psi_n(r) r dr, \\ \|\Psi_n\|^2 &= \frac{1}{4} R^6 [\Psi_n''(R)]^2 = R^2 J_0^2(k_n R) I_0^2(k_n R). \end{aligned}$$

⊙ *Reference:* B. M. Budak, A. A. Samarskii, and A. N. Tikhonov (1980).

### 9.3.3. Three- and $n$ -Dimensional Equations of the Form

$$\frac{\partial^2 w}{\partial t^2} + a^2 \Delta \Delta w = 0$$

**9.3.3-1. Three-dimensional case. Cauchy problem.**

Domain:  $-\infty < x < \infty$ ,  $-\infty < y < \infty$ ,  $-\infty < z < \infty$ . Initial conditions are prescribed:

$$w = f(x, y, z) \quad \text{at} \quad t = 0, \quad \partial_t w = 0 \quad \text{at} \quad t = 0.$$

Solution:

$$w(x, y, z, t) = \frac{1}{(2\sqrt{\pi at})^3} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x + \xi, y + \eta, z + \zeta) \cos\left(\frac{\xi^2 + \eta^2 + \zeta^2}{4at} - \frac{3\pi}{4}\right) d\xi d\eta d\zeta.$$

⊙ *Reference:* V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

### 9.3.3-2. Three-dimensional case. Boundary value problem.

Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2, 0 \leq z \leq l_3$  (rectangular parallelepiped).

Initial conditions:

$$w = f(x, y, z) \quad \text{at} \quad t = 0, \quad \partial_t w = g(x, y, z) \quad \text{at} \quad t = 0.$$

Boundary conditions:

$$\begin{aligned} w = \partial_{xx} w = 0 & \quad \text{at} \quad x = 0, & w = \partial_{xx} w = 0 & \quad \text{at} \quad x = l_1, \\ w = \partial_{yy} w = 0 & \quad \text{at} \quad y = 0, & w = \partial_{yy} w = 0 & \quad \text{at} \quad y = l_2, \\ w = \partial_{zz} w = 0 & \quad \text{at} \quad z = 0, & w = \partial_{zz} w = 0 & \quad \text{at} \quad z = l_3. \end{aligned}$$

Solution:

$$\begin{aligned} w(x, y, z, t) = & \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} \int_0^{l_3} f(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\zeta d\eta d\xi \\ & + \int_0^{l_1} \int_0^{l_2} \int_0^{l_3} g(\xi, \eta, \zeta) G(x, y, z, \xi, \eta, \zeta, t) d\zeta d\eta d\xi, \end{aligned}$$

where

$$\begin{aligned} G(x, y, z, \xi, \eta, \zeta, t) = & \frac{8}{al_1 l_2 l_3} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1}{\lambda_{nmk}} \sin(p_n x) \sin(q_m y) \sin(s_k z) \\ & \times \sin(p_n \xi) \sin(q_m \eta) \sin(s_k \zeta) \sin(\lambda_{nmk} at), \\ p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad s_k = \frac{\pi k}{l_3}, \quad \lambda_{nmk} = p_n^2 + q_m^2 + s_k^2. \end{aligned}$$

### 9.3.3-3. $n$ -dimensional case. Cauchy problem.

Domain:  $\mathbb{R}^n = \{-\infty < x_k < \infty; k = 1, \dots, n\}$ . Initial conditions are prescribed:

$$w = f(\mathbf{x}) \quad \text{at} \quad t = 0, \quad \partial_t w = 0 \quad \text{at} \quad t = 0,$$

where  $\mathbf{x} = \{x_1, \dots, x_n\}$ .

Solution:

$$w(\mathbf{x}, t) = \frac{1}{(2\sqrt{\pi at})^n} \int_{\mathbb{R}^n} f(\mathbf{y}) \cos\left(\frac{|\mathbf{x} - \mathbf{y}|}{4at} - \frac{\pi n}{4}\right) d\mathbf{y},$$

where  $\mathbf{y} = \{y_1, \dots, y_n\}$  and  $d\mathbf{y} = dy_1 \dots dy_n$ .

⊙ Reference: V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).

### 9.3.3-4. $n$ -dimensional case. Boundary value problem.

Domain:  $V = \{0 \leq x_k \leq l_k; k = 1, 2, \dots, n\}$  ( $n$ -dimensional rectangular parallelepiped).

Initial conditions:

$$w = f(\mathbf{x}) \quad \text{at} \quad t = 0, \quad \partial_t w = g(\mathbf{x}) \quad \text{at} \quad t = 0.$$

Boundary conditions:

$$w = \partial_{x_k x_k} w = 0 \quad \text{at} \quad x_k = 0, \quad w = \partial_{x_k x_k} w = 0 \quad \text{at} \quad x_k = l_k.$$

Solution:

$$w(\mathbf{x}, t) = \frac{\partial}{\partial t} \int_V f(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y} + \int_V g(\mathbf{y}) G(\mathbf{x}, \mathbf{y}, t) d\mathbf{y},$$

where

$$G(\mathbf{x}, \mathbf{y}, t) = \frac{2^n}{a l_1 l_2 \dots l_n} \sum_{k_1=1}^{\infty} \sum_{k_2=1}^{\infty} \dots \sum_{k_n=1}^{\infty} \frac{1}{\lambda_{k_1, k_2, \dots, k_n}} \sin(p_{k_1} x_1) \sin(p_{k_2} x_2) \dots \sin(p_{k_n} x_n) \\ \times \sin(p_{k_1} y_1) \sin(p_{k_2} y_2) \dots \sin(p_{k_n} y_n) \sin(\lambda_{k_1, k_2, \dots, k_n} a t), \\ p_{k_1} = \frac{\pi k_1}{l_1}, \quad p_{k_2} = \frac{\pi k_2}{l_2}, \quad \dots, \quad p_{k_n} = \frac{\pi k_n}{l_n}, \quad \lambda_{k_1, k_2, \dots, k_n} = p_{k_1}^2 + p_{k_2}^2 + \dots + p_{k_n}^2.$$

### 9.3.4. Equations of the Form $\frac{\partial^2 w}{\partial t^2} + a^2 \Delta \Delta w + k w = \Phi(x, y, t)$

9.3.4-1. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Solution in terms of the Green's function.

We consider boundary value problems in a rectangular domain  $0 \leq x \leq l_1, 0 \leq y \leq l_2$  with the general initial conditions

$$w = f(x, y) \quad \text{at} \quad t = 0, \quad \partial_t w = g(x, y) \quad \text{at} \quad t = 0$$

and various homogeneous boundary conditions. The solution can be represented in terms of the Green's function as

$$w(x, y, t) = \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} g(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, \zeta, t - \tau) d\eta d\xi d\tau.$$

Paragraphs 9.3.4-2 through 9.3.4-4 present the Green's functions for three types of boundary conditions.

9.3.4-2. The function and its second derivatives are prescribed at the sides of a rectangle:

$$w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l_1, \\ w = \partial_{yy} w = 0 \quad \text{at} \quad y = 0, \quad w = \partial_{yy} w = 0 \quad \text{at} \quad y = l_2.$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}}, \\ p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = \sqrt{a^2(p_n^2 + q_m^2)^2 + k}.$$

9.3.4-3. The first and third derivatives are prescribed at the sides of a rectangle:

$$\partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = 0, \quad \partial_x w = \partial_{xxx} w = 0 \quad \text{at} \quad x = l_1, \\ \partial_y w = \partial_{yyy} w = 0 \quad \text{at} \quad y = 0, \quad \partial_y w = \partial_{yyy} w = 0 \quad \text{at} \quad y = l_2.$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{1}{l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \varepsilon_n \varepsilon_m \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}}, \\ p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = \sqrt{a^2(p_n^2 + q_m^2)^2 + k}, \quad \varepsilon_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}$$

**9.3.4-4. Mixed boundary conditions are prescribed at the sides of a rectangle:**

$$\begin{aligned} w = \partial_{xx}w = 0 & \quad \text{at } x = 0, & w = \partial_{xx}w = 0 & \quad \text{at } x = l_1, \\ \partial_y w = \partial_{yyy}w = 0 & \quad \text{at } y = 0, & \partial_y w = \partial_{yyy}w = 0 & \quad \text{at } y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{2}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} \varepsilon_m \sin(p_n x) \cos(q_m y) \sin(p_n \xi) \cos(q_m \eta) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}}, \\ p_n &= \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = \sqrt{a^2(p_n^2 + q_m^2)^2 + k}, \quad \varepsilon_m = \begin{cases} 1 & \text{for } m = 0, \\ 2 & \text{for } m \neq 0. \end{cases} \end{aligned}$$

**9.3.5. Equations of the Form  $\frac{\partial^2 w}{\partial t^2} + a^2 \left( \frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} \right) + kw = \Phi(x, y, t)$**

**9.3.5-1. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Solution in terms of the Green's function.**

We consider boundary value problems in a rectangular domain  $0 \leq x \leq l_1, 0 \leq y \leq l_2$  with the general initial conditions

$$w = f(x, y) \quad \text{at } t = 0, \quad \partial_t w = g(x, y) \quad \text{at } t = 0$$

and various homogeneous boundary conditions. The solution can be represented in terms of the Green's function as

$$\begin{aligned} w(x, y, t) &= \frac{\partial}{\partial t} \int_0^{l_1} \int_0^{l_2} f(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi + \int_0^{l_1} \int_0^{l_2} g(\xi, \eta) G(x, y, \xi, \eta, t) d\eta d\xi \\ &\quad + \int_0^t \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta, \tau) G(x, y, \xi, \eta, \zeta, t - \tau) d\eta d\xi d\tau. \end{aligned}$$

Paragraphs 9.3.5-2 through 9.3.5-4 present the Green's functions for three types of boundary conditions.

**9.3.5-2. The function and its first derivatives are prescribed at the sides of a rectangle:**

$$\begin{aligned} w = \partial_x w = 0 & \quad \text{at } x = 0, & w = \partial_x w = 0 & \quad \text{at } x = l_1, \\ w = \partial_y w = 0 & \quad \text{at } y = 0, & w = \partial_y w = 0 & \quad \text{at } y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta, t) &= \frac{16}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{p_n^4 q_m^4}{[\varphi_n''(l_1) \psi_m''(l_2)]^2} \varphi_n(x) \psi_m(y) \varphi_n(\xi) \psi_m(\eta) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}}, \\ \lambda_{nm} &= \sqrt{a^2(p_n^4 + q_m^4) + k}, \quad \varphi_n''(x) = \frac{d^2 \varphi_n}{dx^2}, \quad \psi_m''(y) = \frac{d^2 \psi_m}{dy^2}. \end{aligned}$$

Here,

$$\begin{aligned} \varphi_n(x) &= [\sinh(p_n l_1) - \sin(p_n l_1)] [\cosh(p_n x) - \cos(p_n x)] \\ &\quad - [\cosh(p_n l_1) - \cos(p_n l_1)] [\sinh(p_n x) - \sin(p_n x)], \\ \psi_m(y) &= [\sinh(q_m l_2) - \sin(q_m l_2)] [\cosh(q_m y) - \cos(q_m y)] \\ &\quad - [\cosh(q_m l_2) - \cos(q_m l_2)] [\sinh(q_m y) - \sin(q_m y)], \end{aligned}$$

where the  $p_n$  and  $q_m$  are positive roots of the transcendental equations

$$\cosh(pl_1) \cos(pl_1) = 1, \quad \cosh ql_2 \cos ql_2 = 1.$$

9.3.5-3. The function and its second derivatives are prescribed at the sides of a rectangle:

$$\begin{aligned} w = \partial_{xx}w = 0 & \quad \text{at } x = 0, & w = \partial_{xx}w = 0 & \quad \text{at } x = l_1, \\ w = \partial_{yy}w = 0 & \quad \text{at } y = 0, & w = \partial_{yy}w = 0 & \quad \text{at } y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}},$$

$$p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = \sqrt{a^2(p_n^4 + q_m^4) + k}.$$

9.3.5-4. The first and third derivatives are prescribed at the sides of a rectangle:

$$\begin{aligned} \partial_x w = \partial_{xxx} w = 0 & \quad \text{at } x = 0, & \partial_x w = \partial_{xxx} w = 0 & \quad \text{at } x = l_1, \\ \partial_y w = \partial_{yyy} w = 0 & \quad \text{at } y = 0, & \partial_y w = \partial_{yyy} w = 0 & \quad \text{at } y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta, t) = \frac{1}{l_1 l_2} \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \varepsilon_n \varepsilon_m \cos(p_n x) \cos(q_m y) \cos(p_n \xi) \cos(q_m \eta) \frac{\sin(\lambda_{nm} t)}{\lambda_{nm}},$$

$$p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}, \quad \lambda_{nm} = \sqrt{a^2(p_n^4 + q_m^4) + k}, \quad \varepsilon_n = \begin{cases} 1 & \text{for } n = 0, \\ 2 & \text{for } n \neq 0. \end{cases}$$

## 9.4. Fourth-Order Stationary Equations

### 9.4.1. Biharmonic Equation $\Delta\Delta w = 0$

The biharmonic equation is encountered in plane problems of elasticity ( $w$  is the Airy stress function). It is also used to describe slow flows of viscous incompressible fluids ( $w$  is the stream function).

All solutions of the Laplace equation  $\Delta w = 0$  (see Sections 7.1 and 8.1) are also solutions of the biharmonic equation.

9.4.1-1. Two-dimensional equation. Particular solutions.

In the rectangular Cartesian system of coordinates, the biharmonic operator has the form

$$\Delta\Delta \equiv \Delta^2 = \frac{\partial^4}{\partial x^4} + 2 \frac{\partial^4}{\partial x^2 \partial y^2} + \frac{\partial^4}{\partial y^4}.$$

1°. Particular solutions:

$$\begin{aligned} w(x, y) &= Ax^3 + Bx^2y + Cxy^2 + Dy^3 + ax^2 + bxy + cy^2 + \alpha x + \beta y + \gamma, \\ w(x, y) &= (A \cosh \beta x + B \sinh \beta x + Cx \cosh \beta x + Dx \sinh \beta x)(a \cos \beta y + b \sin \beta y), \\ w(x, y) &= (A \cos \beta x + B \sin \beta x + Cx \cos \beta x + Dx \sin \beta x)(a \cosh \beta y + b \sinh \beta y), \\ w(x, y) &= Ar^2 \ln r + Br^2 + C \ln r + D, \quad r = \sqrt{(x-a)^2 + (y-b)^2}, \\ w(x, y) &= (Ax + By + C)(D \cosh \beta x + E \sinh \beta x)(a \cos \beta y + b \sin \beta y), \\ w(x, y) &= (Ax + By + C)(D \cosh \beta y + E \sinh \beta y)(a \cos \beta x + b \sin \beta x), \\ w(x, y) &= (x^2 + y^2)(D \cosh \beta x + E \sinh \beta x)(a \cos \beta y + b \sin \beta y), \\ w(x, y) &= (x^2 + y^2)(D \cosh \beta y + E \sinh \beta y)(a \cos \beta x + b \sin \beta x), \end{aligned}$$

where  $A, B, C, D, E, a, b, c, \alpha, \beta$ , and  $\gamma$  are arbitrary constants.

TABLE 30

Particular solutions of the biharmonic equation in some orthogonal curvilinear coordinate systems;  $A, B, C, D, a, b$ , and  $\lambda$  are arbitrary constants

| Transformation                                                                                                                      | Particular solutions                                                                                                                                                                           |
|-------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Polar coordinates $r, \varphi$ :<br>$x = r \cos \varphi, y = r \sin \varphi$                                                        | $w = (Ar^{2+\lambda} + Br^{2-\lambda} + Cr^\lambda + Dr^{-\lambda})(a \cos \lambda \varphi + b \sin \lambda \varphi),$<br>$w = Ar^2 \ln r + Br^2 + C \ln r + D \quad (\text{at } \lambda = 0)$ |
| Bipolar coordinates $\xi, \eta$ :<br>$x = \frac{c \sinh \xi}{\cosh \xi - \cos \eta}, y = \frac{c \sin \eta}{\cosh \xi - \cos \eta}$ | $w = \frac{1}{\cosh \xi - \cos \eta} [A \cosh(\lambda + 1)\xi + B \sinh(\lambda + 1)\xi$<br>$+ C \cosh(\lambda - 1)\xi + D \sinh(\lambda - 1)\xi] (a \cos \lambda \eta + b \sin \lambda \eta)$ |
| Degenerate bipolar coordinates $u, v$ :<br>$x = \frac{u}{u^2 + v^2}, y = -\frac{v}{u^2 + v^2}$                                      | $w = \frac{1}{(u^2 + v^2)^2} [A \cosh(\lambda u) + B \sinh(\lambda u)$<br>$+ Cu \cosh(\lambda u) + Du \sinh(\lambda u)] [a \cos(\lambda v) + b \sin(\lambda v)]$                               |

2°. Fundamental solution:

$$\mathcal{E}(x, y) = \frac{1}{8\pi} r^2 \ln r, \quad r = \sqrt{x^2 + y^2}.$$

3°. Particular solutions of the biharmonic equation in some orthogonal curvilinear coordinate systems are listed in [Table 30](#).

⊙ *Reference:* N. N. Lebedev, I. P. Skal'skaya, and Ya. S. Uflyand (1972).

#### 9.4.1-2. Two-dimensional equation. Various representations of the general solution.

1°. Various representations of the general solution in terms of harmonic functions:

$$\begin{aligned} w(x, y) &= xu_1(x, y) + u_2(x, y), \\ w(x, y) &= yu_1(x, y) + u_2(x, y), \\ w(x, y) &= (x^2 + y^2)u_1(x, y) + u_2(x, y), \end{aligned}$$

where  $u_1$  and  $u_2$  are arbitrary functions satisfying the Laplace equation  $\Delta u_k = 0$  ( $k = 1, 2$ ).

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

2°. Complex form of representation of the general solution:

$$w(x, y) = \operatorname{Re} [\bar{z}f(z) + g(z)],$$

where  $f(z)$  and  $g(z)$  are arbitrary analytic functions of the complex variable  $z = x + iy$ ;  $\bar{z} = x - iy$ ,  $i^2 = -1$ . The symbol  $\operatorname{Re}[A]$  stands for the real part of the complex quantity  $A$ .

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 9.4.1-3. Two-dimensional boundary value problems for the upper half-plane.

1°. Domain:  $-\infty < x < \infty, 0 \leq y < \infty$ . The desired function and its derivative along the normal are prescribed at the boundary:

$$w = 0 \quad \text{at} \quad y = 0, \quad \partial_y w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} f(\xi) G(x - \xi, y) d\xi, \quad G(x, y) = \frac{1}{\pi} \frac{y^2}{x^2 + y^2}.$$

⊙ *Reference:* G. E. Shilov (1965).

2°. Domain:  $-\infty < x < \infty$ ,  $0 \leq y < \infty$ . The derivatives of the desired function are prescribed at the boundary:

$$\partial_x w = f(x) \quad \text{at} \quad y = 0, \quad \partial_y w = g(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(\xi) \left[ \arctan\left(\frac{x-\xi}{y}\right) + \frac{y(x-\xi)}{(x-\xi)^2 + y^2} \right] d\xi + \frac{y^2}{\pi} \int_{-\infty}^{\infty} \frac{g(\xi) d\xi}{(x-\xi)^2 + y^2} + C,$$

where  $C$  is an arbitrary constant.

**Example.** Let us consider the problem of a slow (Stokes) inflow of a viscous fluid into the half-plane through a slit of width  $2a$  with a constant velocity  $U$  that makes an angle  $\beta$  with the normal to the boundary (the angle is reckoned from the normal counterclockwise).

With the stream function  $w$  introduced by the relations  $v_x = -\frac{\partial w}{\partial y}$  and  $v_y = \frac{\partial w}{\partial x}$  ( $v_x$  and  $v_y$  are the fluid velocity components), the problem is reduced to the special case of the previous problem with

$$f(x) = \begin{cases} U \cos \beta & \text{for } |x| < a, \\ 0 & \text{for } |x| > a, \end{cases} \quad g(x) = \begin{cases} U \sin \beta & \text{for } |x| < a, \\ 0 & \text{for } |x| > a. \end{cases}$$

Dean's solution:

$$w(x, y) = \frac{U}{\pi} [(x-a) \cos \beta + y \sin \beta] \arctan\left(\frac{y}{x-a}\right) - \frac{U}{\pi} [(x+a) \cos \beta + y \sin \beta] \arctan\left(\frac{y}{x+a}\right) + C.$$

⊙ *Reference:* I. Sneddon (1951).

#### 9.4.1-4. Two-dimensional boundary value problem for a circle.

Domain:  $0 \leq r \leq a$ ,  $0 \leq \varphi \leq 2\pi$ . Boundary conditions in the polar coordinate system:

$$w = f(\varphi) \quad \text{at} \quad r = a, \quad \partial_r w = g(\varphi) \quad \text{at} \quad r = a.$$

Solution:

$$w(r, \varphi) = \frac{1}{2\pi a} (r^2 - a^2)^2 \left[ \int_0^{2\pi} \frac{[a - r \cos(\eta - \varphi)] f(\eta) d\eta}{[r^2 + a^2 - 2ar \cos(\eta - \varphi)]^2} - \frac{1}{2} \int_0^{2\pi} \frac{g(\eta) d\eta}{r^2 + a^2 - 2ar \cos(\eta - \varphi)} \right].$$

⊙ *Reference:* A. N. Tikhonov and A. A. Samarskii (1990).

#### 9.4.1-5. Three-dimensional equation.

In the rectangular Cartesian coordinate system, the three-dimensional biharmonic operator is expressed as

$$\Delta \Delta \equiv \Delta^2 = \frac{\partial^4}{\partial x^4} + \frac{\partial^4}{\partial y^4} + \frac{\partial^4}{\partial z^4} + 2 \frac{\partial^4}{\partial x^2 \partial y^2} + 2 \frac{\partial^4}{\partial x^2 \partial z^2} + 2 \frac{\partial^4}{\partial y^2 \partial z^2}.$$

1°. Particular solutions in the Cartesian coordinate system:

$$w(x, y, z) = Ar^2 + Br + C + \frac{D}{r}, \quad r = \sqrt{(x-a)^2 + (y-b)^2 + (z-c)^2},$$

$$w(x, y, z) = [Ax \sin(\beta x) + B \sin(\beta x) + Cx \cos(\beta x) + D \cos(\beta x)] \sin(\mu y) \exp(\pm z \sqrt{\beta^2 + \mu^2}),$$

$$w(x, y, z) = [Ax \sin(\beta x) + B \sin(\beta x) + Cx \cos(\beta x) + D \cos(\beta x)] \cos(\mu y) \exp(\pm z \sqrt{\beta^2 + \mu^2}),$$

$$w(x, y, z) = [Ax \sin(\beta x) + B \sin(\beta x) + Cx \cos(\beta x) + D \cos(\beta x)] \sinh(\mu y) \exp(\pm z \sqrt{\beta^2 - \mu^2}),$$

$$w(x, y, z) = [Ax \sin(\beta x) + B \sin(\beta x) + Cx \cos(\beta x) + D \cos(\beta x)] \cosh(\mu y) \exp(\pm z \sqrt{\beta^2 - \mu^2}),$$

$$w(x, y, z) = [Ax \sinh(\beta x) + B \sinh(\beta x) + Cx \cosh(\beta x) + D \cosh(\beta x)] \sinh(\mu y) \sin(z \sqrt{\beta^2 + \mu^2}),$$

$$w(x, y, z) = [Ax \sinh(\beta x) + B \sinh(\beta x) + Cx \cosh(\beta x) + D \cosh(\beta x)] \cosh(\mu y) \cos(z \sqrt{\beta^2 + \mu^2}),$$

where  $A, B, C, D, \beta$ , and  $\mu$  are arbitrary constants.

2°. Particular solutions in the cylindrical coordinate system ( $r = \sqrt{x^2 + y^2}$ ):

$$w(r, \varphi, z) = J_n(\mu r)(Ar \cos \varphi + Br \sin \varphi + C)(a_1 \cos n\varphi + b_1 \sin n\varphi)(a_2 \cosh \mu z + b_2 \sinh \mu z),$$

$$w(r, \varphi, z) = Y_n(\mu r)(Ar \cos \varphi + Br \sin \varphi + C)(a_1 \cos n\varphi + b_1 \sin n\varphi)(a_2 \cosh \mu z + b_2 \sinh \mu z),$$

$$w(r, \varphi, z) = I_n(\mu r)(Ar \cos \varphi + Br \sin \varphi + C)(a_1 \cos n\varphi + b_1 \sin n\varphi)(a_2 \cosh \mu z + b_2 \sinh \mu z),$$

$$w(r, \varphi, z) = K_n(\mu r)(Ar \cos \varphi + Br \sin \varphi + C)(a_1 \cos n\varphi + b_1 \sin n\varphi)(a_2 \cosh \mu z + b_2 \sinh \mu z),$$

$$w(r, \varphi, z) = J_n(\mu r)(A \cos n\varphi + B \sin n\varphi)(a_1 \cosh \mu z + b_1 \sinh \mu z + a_2 z \cosh \mu z + b_2 z \sinh \mu z),$$

$$w(r, \varphi, z) = Y_n(\mu r)(A \cos n\varphi + B \sin n\varphi)(a_1 \cosh \mu z + b_1 \sinh \mu z + a_2 z \cosh \mu z + b_2 z \sinh \mu z),$$

$$w(r, \varphi, z) = I_n(\mu r)(A \cos n\varphi + B \sin n\varphi)(a_1 \cosh \mu z + b_1 \sinh \mu z + a_2 z \cosh \mu z + b_2 z \sinh \mu z),$$

$$w(r, \varphi, z) = K_n(\mu r)(A \cos n\varphi + B \sin n\varphi)(a_1 \cosh \mu z + b_1 \sinh \mu z + a_2 z \cosh \mu z + b_2 z \sinh \mu z),$$

where  $n = 0, 1, 2, \dots$ ;  $A, B, C, a_1, a_2, b_1, b_2$ , and  $\mu$  are arbitrary constants; the  $J_n(\xi)$  and  $Y_n(\xi)$  are the Bessel functions; and the  $I_n(\xi)$  and  $K_n(\xi)$  are the modified Bessel functions.

3°. Particular solutions in the spherical coordinate system ( $r = \sqrt{x^2 + y^2 + z^2}$ ):

$$w(r) = Ar^2 + Br + C + Dr^{-1},$$

$$w(r, \theta) = (Ar^{n+2} + Br^n + Cr^{1-n} + Dr^{-1-n})P_n(\cos \theta),$$

$$w(r, \theta, \varphi) = (Ar^{n+2} + Br^n + Cr^{1-n} + Dr^{-1-n})P_n^m(\cos \theta)(a \cos m\varphi + b \sin m\varphi),$$

where  $n = 0, 1, 2, \dots$ ;  $m = 0, 1, 2, \dots, n$ ;  $A, B, C, D, a$ , and  $b$  are arbitrary constants; the  $P_n(\xi)$  are the Legendre polynomials; and the  $P_n^m(\xi)$  are the associated Legendre functions defined by

$$P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n, \quad P_n^m(x) = (1 - x^2)^{m/2} \frac{d^m}{dx^m} P_n(x).$$

4°. Fundamental solution:

$$\mathcal{G}(x, y, z) = -\frac{1}{8\pi} \sqrt{x^2 + y^2 + z^2}.$$

5°. Representations of solutions to the biharmonic equation in terms of harmonic functions:

$$w(x, y, z) = xu_1(x, y, z) + u_2(x, y, z),$$

$$w(x, y, z) = (x^2 + y^2 + z^2)u_1(x, y, z) + u_2(x, y, z),$$

where  $u_1$  and  $u_2$  are arbitrary functions satisfying the three-dimensional Laplace equation  $\Delta_3 u_k = 0$  ( $k = 1, 2$ ). The coefficient  $x$  of  $u_1$  in the first formula can be replaced by  $y$  or  $z$ .

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

#### 9.4.1-6. $n$ -dimensional equation.

1°. Particular solutions:

$$w(\mathbf{x}) = \sum_{i,j,k=1}^n A_{ijk} x_i x_j x_k + \sum_{i,j=1}^n B_{ij} x_i x_j + \sum_{i=1}^n C_i x_i + D,$$

$$w(\mathbf{x}) = Ar^2 + B + Cr^{4-n} + Dr^{2-n}, \quad r^2 = \sum_{k=1}^n (x_k - \alpha_k)^2,$$

$$w(\mathbf{x}) = (A + Br^{2-n}) \left( \sum_{i=1}^n C_i x_i + D \right), \quad r^2 = \sum_{k=1}^n (x_k - \alpha_k)^2,$$

$$w(\mathbf{x}) = \exp(\pm x_n \sqrt{\lambda_n}) \left( \sum_{i=1}^n A_i x_i + B \right) \prod_{k=1}^{n-1} \sin(\alpha_k x_k + \beta_k), \quad \lambda_n = \sum_{k=1}^{n-1} \alpha_k^2,$$

$$w(\mathbf{x}) = \left( \sum_{i=1}^n A_i x_i + B \right) \left[ \prod_{k=1}^{m-1} \sin(\alpha_k x_k + \beta_k) \right] \left[ \prod_{k=m}^n \sinh(\gamma_k x_k) \right], \quad \sum_{k=1}^{m-1} \alpha_k^2 - \sum_{k=m}^n \gamma_k^2 = 0,$$

where the  $A_{ijk}$ ,  $B_{ij}$ ,  $A_i$ ,  $C_i$ ,  $A, B, C, D, \alpha_k, \beta_k$ , and  $\gamma_k$  are arbitrary constants.



2°. Fundamental solution:

$$\mathcal{E}(\mathbf{x}) = \begin{cases} \frac{\Gamma(n/2)|\mathbf{x}|^{4-n}}{4\pi^{n/2}(n-2)(n-4)} & \text{for } n = 3, 5, 6, 7, \dots; \\ -\frac{1}{8\pi^2} \ln |\mathbf{x}| & \text{for } n = 4. \end{cases}$$

For  $n = 2$ , see Paragraph 9.4.1-1, Item 2°.

⊙ *Reference:* G. E. Shilov (1965).

3°. Various representations of solutions to the biharmonic equation in terms of harmonic functions:

$$\begin{aligned} w(\mathbf{x}) &= x_s u_1(\mathbf{x}) + u_2(\mathbf{x}), & s &= 1, 2, \dots, n; \\ w(\mathbf{x}) &= |\mathbf{x}|^2 u_1(\mathbf{x}) + u_2(\mathbf{x}), & |\mathbf{x}|^2 &= \sum_{k=1}^n x_k^2, \end{aligned}$$

where  $u_1$  and  $u_2$  are arbitrary functions satisfying the  $n$ -dimensional Laplace equation  $\Delta_n u_m = 0$  ( $m = 1, 2$ ).

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

## 9.4.2. Equations of the Form $\Delta\Delta w = \Phi(x, y)$

*Nonhomogeneous biharmonic equation.* It is encountered in plane problems of elasticity and hydrodynamics.

9.4.2-1. Domain:  $-\infty < x < \infty, -\infty < y < \infty$ .

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) \mathcal{E}(x - \xi, y - \eta) d\xi d\eta, \quad \mathcal{E}(x, y) = \frac{1}{8\pi} (x^2 + y^2) \ln \sqrt{x^2 + y^2}.$$

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

9.4.2-2. Domain:  $-\infty < x < \infty, 0 \leq y < \infty$ . Boundary value problem.

The upper half-plane is considered. The derivatives are prescribed at the boundary:

$$\partial_x w = f(x) \quad \text{at } y = 0, \quad \partial_y w = g(x) \quad \text{at } y = 0.$$

Solution:

$$\begin{aligned} w(x, y) &= \frac{1}{\pi} \int_{-\infty}^{\infty} f(\xi) \left[ \arctan\left(\frac{x - \xi}{y}\right) + \frac{y(x - \xi)}{(x - \xi)^2 + y^2} \right] d\xi + \frac{y^2}{\pi} \int_{-\infty}^{\infty} \frac{g(\xi) d\xi}{(x - \xi)^2 + y^2} \\ &+ \frac{1}{8\pi} \int_{-\infty}^{\infty} d\xi \int_0^{\infty} \left[ \frac{1}{2} (R_+^2 - R_-^2) - R_-^2 \ln \frac{R_+}{R_-} \right] \Phi(\xi, \eta) d\eta + C, \end{aligned}$$

where  $C$  is an arbitrary constant,

$$R_+^2 = (x - \xi)^2 + (y + \eta)^2, \quad R_-^2 = (x - \xi)^2 + (y - \eta)^2.$$

⊙ *Reference:* I. Sneddon (1951).

**9.4.2-3. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . The sides of the plate are hinged.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = \partial_{xx} w = 0 & \quad \text{at} \quad x = 0, & w = \partial_{xx} w = 0 & \quad \text{at} \quad x = l_1, \\ w = \partial_{yy} w = 0 & \quad \text{at} \quad y = 0, & w = \partial_{yy} w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Solution:

$$w(x, y) = \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi,$$

where

$$G(x, y, \xi, \eta) = \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{(p_n^2 + q_m^2)^2} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta),$$

$$p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}.$$

### 9.4.3. Equations of the Form $\Delta \Delta w - \lambda w = \Phi(x, y)$

**9.4.3-1. Homogeneous equation ( $\Phi \equiv 0$ ).**

This equation describes the shapes of two-dimensional free transverse vibrations of a thin elastic plate; the function  $w$  defines the deflection (transverse displacement) of the plate's midplane points relative to the original plane position and  $k = \lambda^{1/4}$  is the frequency parameter. Here,  $\Delta \Delta = \Delta^2$  is the biharmonic operator and  $\Delta$  is the Laplace operator defined as

$$\Delta = \begin{cases} \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} & \text{in the Cartesian coordinate system,} \\ \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \varphi^2} & \text{in the polar coordinate system.} \end{cases}$$

1°. Particular solutions ( $A_1, A_2, A_3, A_4, B_1$ , and  $B_2$  are arbitrary constants):

$$\begin{aligned} w(x, y) &= [A_1 \sin(k_1 x) + B_1 \cos(k_1 x)] [A_2 \sin(k_2 y) + B_2 \cos(k_2 y)], & \lambda &= (k_1^2 + k_2^2)^2, \\ w(x, y) &= [A_1 \sin(k_1 x) + B_1 \cos(k_1 x)] [A_2 \sinh(k_2 y) + B_2 \cosh(k_2 y)], & \lambda &= (k_1^2 - k_2^2)^2, \\ w(x, y) &= [A_1 \sinh(k_1 x) + B_1 \cosh(k_1 x)] [A_2 \sin(k_2 y) + B_2 \cos(k_2 y)], & \lambda &= (k_1^2 - k_2^2)^2, \\ w(x, y) &= [A_1 \sinh(k_1 x) + B_1 \cosh(k_1 x)] [A_2 \sinh(k_2 y) + B_2 \cosh(k_2 y)], & \lambda &= (k_1^2 + k_2^2)^2, \\ w(r, \varphi) &= [A_1 J_n(kr) + A_2 Y_n(kr) + A_3 I_n(kr) + A_4 K_n(kr)] \cos(n\varphi), & \lambda &= k^4 > 0, \\ w(r, \varphi) &= [A_1 J_n(kr) + A_2 Y_n(kr) + A_3 I_n(kr) + A_4 K_n(kr)] \sin(n\varphi), & \lambda &= k^4 > 0, \end{aligned}$$

where the  $J_n(\xi)$  and  $Y_n(\xi)$  are the Bessel functions of the first and second kind, the  $I_n(\xi)$  and  $K_n(\xi)$  are the modified Bessel functions of the first and second kind,  $r = \sqrt{x^2 + y^2}$ , and  $n = 0, 1, 2, \dots$

2°. General solution:

$$w(x, y) = u_1(x, y) + u_2(x, y),$$

where  $u_1$  and  $u_2$  are arbitrary functions satisfying the Helmholtz equations

$$\Delta u_1 + \sqrt{\lambda} u_1 = 0, \quad \Delta u_2 - \sqrt{\lambda} u_2 = 0.$$

For solutions to these equations, see Section 7.3.

**9.4.3-2. Domain:  $0 \leq x \leq l_1, 0 \leq y \leq l_2$ . Boundary value problem.**

A rectangle is considered. Boundary conditions are prescribed:

$$\begin{aligned} w = \partial_{xx} w = 0 & \quad \text{at} \quad x = 0, & w = \partial_{xx} w = 0 & \quad \text{at} \quad x = l_1, \\ w = \partial_{yy} w = 0 & \quad \text{at} \quad y = 0, & w = \partial_{yy} w = 0 & \quad \text{at} \quad y = l_2. \end{aligned}$$

Solution:

$$w(x, y) = \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi,$$

where

$$G(x, y, \xi, \eta, t) = \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{\sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta)}{(p_n^2 + q_m^2)^2 - \lambda}, \quad p_n = \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}.$$

**9.4.3-3. Domain:  $0 \leq r \leq R, 0 \leq \varphi \leq 2\pi$ . Eigenvalue problem with  $\Phi \equiv 0$ .**

The unknown and its normal derivative are zero on the boundary of a circular domain:

$$w = \frac{\partial w}{\partial r} = 0 \quad \text{at} \quad r = R.$$

Eigenvalues:

$$\lambda_{nm} = \frac{\beta_{nm}^4}{R^4}, \quad n = 0, 1, 2, \dots, m = 1, 2, 3, \dots,$$

where the  $\beta_{nm}$  are positive roots of the transcendental equation

$$J_n(\beta) I'_n(\beta) - I_n(\beta) J'_n(\beta) = 0.$$

Numerical values of some roots:

$$\begin{aligned} \beta_{01} = 3.196, \quad \beta_{02} = 6.306, \quad \beta_{03} = 9.439, \quad \beta_{04} = 12.58; \\ \beta_{11} = 4.611, \quad \beta_{12} = 7.799, \quad \beta_{13} = 10.96, \quad \beta_{14} = 14.11; \\ \beta_{21} = 5.906, \quad \beta_{22} = 9.197, \quad \beta_{23} = 12.40, \quad \beta_{24} = 15.58, \\ \beta_{31} = 7.144, \quad \beta_{32} = 10.54, \quad \beta_{33} = 13.79, \quad \beta_{34} = 17.01. \end{aligned}$$

Eigenvalues:

$$\begin{aligned} w_{nm}^{(c)}(r, \varphi) &= \left[ I_n(\beta_{nm}) J_n\left(\beta_{nm} \frac{r}{R}\right) - J_n(\beta_{nm}) I_n\left(\beta_{nm} \frac{r}{R}\right) \right] \cos(n\varphi), \\ w_{nm}^{(s)}(r, \varphi) &= \left[ I_n(\beta_{nm}) J_n\left(\beta_{nm} \frac{r}{R}\right) - J_n(\beta_{nm}) I_n\left(\beta_{nm} \frac{r}{R}\right) \right] \sin(n\varphi). \end{aligned}$$

⊙ Reference: V. V. Bolotin (1978).

**9.4.3-4. Domain:  $(x/a)^2 + (y/b)^2 \leq 1$ . Eigenvalue problem with  $\Phi \equiv 0$ .**

The unknown and its normal derivative are zero on the boundary of an elliptic domain:

$$w = \frac{\partial w}{\partial N} = 0 \quad \text{on} \quad (x/a)^2 + (y/b)^2 = 1 \quad (a \geq b).$$

Eigenvalues and eigenfunctions (approximate formulas):

$$\begin{aligned} \lambda_{01} &= \frac{\beta_{01}^4}{8} \left( \frac{3}{a^4} + \frac{3}{b^4} + \frac{2}{a^2 b^2} \right), & w_{01}(\mathcal{R}) &= I_0(\beta_{01}) J_0(\beta_{01} \mathcal{R}) - J_0(\beta_{01}) I_0(\beta_{01} \mathcal{R}), \\ \lambda_{11}^{(c)} &= \frac{\beta_{11}^4}{8} \left( \frac{5}{a^4} + \frac{1}{b^4} + \frac{2}{a^2 b^2} \right), & w_{11}^{(c)}(\mathcal{R}, \varphi) &= [I_1(\beta_{11}) J_1(\beta_{11} \mathcal{R}) - J_1(\beta_{11}) I_1(\beta_{11} \mathcal{R})] \cos \varphi, \\ \lambda_{11}^{(s)} &= \frac{\beta_{11}^4}{8} \left( \frac{1}{a^4} + \frac{5}{b^4} + \frac{2}{a^2 b^2} \right), & w_{11}^{(s)}(\mathcal{R}, \varphi) &= [I_1(\beta_{11}) J_1(\beta_{11} \mathcal{R}) - J_1(\beta_{11}) I_1(\beta_{11} \mathcal{R})] \sin \varphi, \end{aligned}$$

where  $\mathcal{R} = \sqrt{(x/a)^2 + (y/b)^2}$ , and  $\beta_{01} = 3.196$  and  $\beta_{11} = 4.611$  are the least roots of the transcendental equations

$$\begin{aligned} J_0(\beta)I_1(\beta) + J_1(\beta)I_0(\beta) &= 0, \\ J_1(\beta)I_1'(\beta) - J_1'(\beta)I_1(\beta) &= 0. \end{aligned}$$

The above formulas were obtained with the aid of generalized (nonorthogonal) polar coordinates  $\mathcal{R}, \varphi$  defined by

$$x = a\mathcal{R} \cos \varphi, \quad y = b\mathcal{R} \sin \varphi \quad (0 \leq \mathcal{R} \leq 1, 0 \leq \varphi \leq 2\pi)$$

and the variational method.

The maximum error in the eigenvalue  $\lambda_1$  for  $\varepsilon = \sqrt{1 - (b/a)^2} \leq 0.86$  is less than 1%. The errors in  $\lambda_{11}^{(c)}$  and  $\lambda_{11}^{(s)}$  for  $\varepsilon \leq 0.6$  do not exceed 2%. In the limit case  $\varepsilon = 0$  that corresponds to a circular domain, the formulas provide exact results.

⊙ *Reference:* L. D. Akulenko, S. V. Nesterov, and A. L. Popov (2001).

#### 9.4.4. Equations of the Form $\frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} = \Phi(x, y)$

##### 9.4.4-1. Homogeneous equation ( $\Phi \equiv 0$ ).

1°. Particular solutions:

$$\begin{aligned} w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp\left(\frac{1}{\sqrt{2}} \lambda y\right) \sin\left(\frac{1}{\sqrt{2}} \lambda y\right), \\ w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp\left(\frac{1}{\sqrt{2}} \lambda y\right) \cos\left(\frac{1}{\sqrt{2}} \lambda y\right), \\ w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp\left(-\frac{1}{\sqrt{2}} \lambda y\right) \sin\left(\frac{1}{\sqrt{2}} \lambda y\right), \\ w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp\left(-\frac{1}{\sqrt{2}} \lambda y\right) \cos\left(\frac{1}{\sqrt{2}} \lambda y\right), \end{aligned}$$

where  $A, B, C, D$ , and  $\lambda$  are arbitrary constants.

2°. General solution:

$$w(x, y) = \operatorname{Re} [f(z_1) + g(z_2)].$$

Here,  $f(z_1)$  and  $g(z_2)$  are arbitrary analytic functions of the complex variables  $z_1 = x - \frac{1}{\sqrt{2}}(1+i)y$  and  $z_2 = x + \frac{1}{\sqrt{2}}(1+i)y$ . The symbol  $\operatorname{Re}[A]$  stands for the real part of the complex quantity  $A$ .

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

3°. Domain:  $-\infty < x < \infty, 0 \leq y < \infty$ . Boundary value problem.

The upper half-space is considered. Boundary conditions are prescribed:

$$w = 0 \quad \text{at} \quad y = 0, \quad \partial_y w = f(x) \quad \text{at} \quad y = 0.$$

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} f(\xi) G(x - \xi, y) d\xi,$$

where

$$G(x, y) = \frac{1}{\pi\sqrt{2}} \left[ \arctan\left(1 - \frac{x\sqrt{2}}{y}\right) + \arctan\left(1 + \frac{x\sqrt{2}}{y}\right) \right].$$

⊙ *Reference:* G. E. Shilov (1965).

#### 9.4.4-2. Nonhomogeneous equation. Boundary value problems in a rectangle.

We consider problems in a rectangular domain  $0 \leq x \leq l_1$ ,  $0 \leq y \leq l_2$  with different homogeneous boundary conditions. The solution can be expressed in terms of the Green's function as

$$w(x, y) = \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi.$$

Below are the Green's functions for two types of boundary conditions.

1°. The function and its first derivatives are prescribed at the sides of the rectangle:

$$\begin{aligned} w = \partial_x w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_x w = 0 \quad \text{at} \quad x = l_1, \\ w = \partial_y w = 0 \quad \text{at} \quad y = 0, \quad w = \partial_y w = 0 \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta) = \frac{16}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{p_n^4 q_m^4 \varphi_n(x) \psi_m(y) \varphi_n(\xi) \psi_m(\eta)}{(p_n^4 + q_m^4) [\varphi_n''(l_1) \psi_m''(l_2)]^2}.$$

Here,

$$\begin{aligned} \varphi_n(x) &= [\sinh(p_n l_1) - \sin(p_n l_1)] [\cosh(p_n x) - \cos(p_n x)] \\ &\quad - [\cosh(p_n l_1) - \cos(p_n l_1)] [\sinh(p_n x) - \sin(p_n x)], \\ \psi_m(y) &= [\sinh(q_m l_2) - \sin(q_m l_2)] [\cosh(q_m y) - \cos(q_m y)] \\ &\quad - [\cosh(q_m l_2) - \cos(q_m l_2)] [\sinh(q_m y) - \sin(q_m y)], \end{aligned}$$

where the  $p_n$  and  $q_m$  are positive roots of the transcendental equations

$$\cosh(pl_1) \cos(pl_1) = 1, \quad \cosh ql_2 \cos ql_2 = 1.$$

2°. The function and its second derivatives are prescribed at the sides of the rectangle:

$$\begin{aligned} w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l_1, \\ w = \partial_{yy} w = 0 \quad \text{at} \quad y = 0, \quad w = \partial_{yy} w = 0 \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta) &= \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{p_n^4 + q_m^4} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta), \\ p_n &= \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}. \end{aligned}$$

### 9.4.5. Equations of the Form $\frac{\partial^4 w}{\partial x^4} + \frac{\partial^4 w}{\partial y^4} + kw = \Phi(x, y)$

#### 9.4.5-1. Particular solutions of the homogeneous equation ( $\Phi \equiv 0$ ):

$$\begin{aligned} w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp(\beta y) \sin(\beta y), \\ w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp(\beta y) \cos(\beta y), \\ w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp(-\beta y) \sin(\beta y), \\ w(x, y) &= [A \sin(\lambda x) + B \cos(\lambda x) + C \sinh(\lambda x) + D \cosh(\lambda x)] \exp(-\beta y) \cos(\beta y), \end{aligned}$$

where  $\beta = \frac{1}{\sqrt{2}}(\lambda^4 + k)^{1/4}$ ;  $A, B, C, D$ , and  $\lambda$  are arbitrary constants.

#### 9.4.5-2. Domain: $0 \leq x \leq l_1$ , $0 \leq y \leq l_2$ . Boundary value problems.

1°. We consider problems in a rectangular domain with different homogeneous boundary conditions. The solution can be expressed in terms of the Green's function as

$$w(x, y) = \int_0^{l_1} \int_0^{l_2} \Phi(\xi, \eta) G(x, y, \xi, \eta) d\eta d\xi.$$

Below are the Green's functions for two types of boundary conditions.

2°. The function and its first derivatives are prescribed at the sides of the rectangle:

$$\begin{aligned} w = \partial_x w = 0 \quad \text{at} \quad x = 0, & \quad w = \partial_x w = 0 \quad \text{at} \quad x = l_1, \\ w = \partial_y w = 0 \quad \text{at} \quad y = 0, & \quad w = \partial_y w = 0 \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$G(x, y, \xi, \eta) = \frac{16}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{p_n^4 q_m^4 \varphi_n(x) \psi_m(y) \varphi_n(\xi) \psi_m(\eta)}{(p_n^4 + q_m^4 + k) [\varphi_n''(l_1) \psi_m''(l_2)]^2}.$$

Here,

$$\begin{aligned} \varphi_n(x) &= [\sinh(p_n l_1) - \sin(p_n l_1)] [\cosh(p_n x) - \cos(p_n x)] \\ &\quad - [\cosh(p_n l_1) - \cos(p_n l_1)] [\sinh(p_n x) - \sin(p_n x)], \\ \psi_m(y) &= [\sinh(q_m l_2) - \sin(q_m l_2)] [\cosh(q_m y) - \cos(q_m y)] \\ &\quad - [\cosh(q_m l_2) - \cos(q_m l_2)] [\sinh(q_m y) - \sin(q_m y)], \end{aligned}$$

where the  $p_n$  and  $q_m$  are positive roots of the transcendental equations

$$\cosh(pl_1) \cos(pl_1) = 1, \quad \cosh ql_2 \cos ql_2 = 1.$$

3°. The function and its second derivatives are prescribed at the sides of the rectangle:

$$\begin{aligned} w = \partial_{xx} w = 0 \quad \text{at} \quad x = 0, & \quad w = \partial_{xx} w = 0 \quad \text{at} \quad x = l_1, \\ w = \partial_{yy} w = 0 \quad \text{at} \quad y = 0, & \quad w = \partial_{yy} w = 0 \quad \text{at} \quad y = l_2. \end{aligned}$$

Green's function:

$$\begin{aligned} G(x, y, \xi, \eta) &= \frac{4}{l_1 l_2} \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{1}{p_n^4 + q_m^4 + k} \sin(p_n x) \sin(q_m y) \sin(p_n \xi) \sin(q_m \eta), \\ p_n &= \frac{\pi n}{l_1}, \quad q_m = \frac{\pi m}{l_2}. \end{aligned}$$

### 9.4.6. Stokes Equation (Axisymmetric Flows of Viscous Fluids)

#### 9.4.6-1. Stokes equation for the stream function in the spherical coordinate system.

The Stokes equation for the stream function in the axisymmetric case is written as

$$E^2(E^2 w) = 0, \quad E^2 \equiv \frac{\partial^2}{\partial r^2} + \frac{\sin \theta}{r^2} \frac{\partial}{\partial \theta} \left( \frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \right).$$

It governs slow axisymmetric flows of viscous incompressible fluids, with  $w$  being the stream function,  $r$  and  $\theta$  the spherical coordinates. The components of the fluid velocity are related to the stream function by  $v_r = \frac{1}{r^2 \sin \theta} \frac{\partial w}{\partial \theta}$  and  $v_\theta = -\frac{1}{r \sin \theta} \frac{\partial w}{\partial r}$ .

General solution ( $A_n, B_n, C_n, D_n, \tilde{A}_n, \tilde{B}_n, \tilde{C}_n$ , and  $\tilde{D}_n$  are arbitrary constants):

$$w(r, \theta) = \sum_{n=0}^{\infty} (A_n r^n + B_n r^{1-n} + C_n r^{n+2} + D_n r^{3-n}) \mathcal{J}_n(\cos \theta) + \sum_{n=2}^{\infty} (\tilde{A}_n r^n + \tilde{B}_n r^{1-n} + \tilde{C}_n r^{n+2} + \tilde{D}_n r^{3-n}) \mathcal{H}_n(\cos \theta), \quad (1)$$

where the  $\mathcal{J}_n(\zeta)$  and  $\mathcal{H}_n(\zeta)$  are the Gegenbauer functions of the first and second kind, respectively. These are linearly related to the Legendre functions  $P_n(\zeta)$  and  $Q_n(\zeta)$  by

$$\mathcal{J}_n(\zeta) = \frac{P_{n-2}(\zeta) - P_n(\zeta)}{2n-1}, \quad \mathcal{H}_n(\zeta) = \frac{Q_{n-2}(\zeta) - Q_n(\zeta)}{2n-1} \quad (n \geq 2).$$

The Gegenbauer functions of the first kind are represented in the form of a finite power series as

$$\begin{aligned} \mathcal{J}_n(\zeta) &= -\frac{1}{(n-1)!} \left( \frac{d}{d\zeta} \right)^{n-2} \left( \frac{\zeta^2 - 1}{2} \right)^{n-1} \\ &= \frac{1 \cdot 3 \cdots (2n-3)}{1 \cdot 2 \cdots n} \left[ \zeta^n - \frac{n(n-1)}{2(2n-3)} \zeta^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2 \cdot 4(2n-3)(2n-5)} \zeta^{n-4} - \cdots \right]. \end{aligned}$$

In particular,

$$\begin{aligned} \mathcal{J}_0(\zeta) &= 1, \quad \mathcal{J}_1(\zeta) = -\zeta, \quad \mathcal{J}_2(\zeta) = \frac{1}{2}(1 - \zeta^2), \quad \mathcal{J}_3(\zeta) = \frac{1}{2}\zeta(1 - \zeta^2), \\ \mathcal{J}_4(\zeta) &= \frac{1}{8}(1 - \zeta^2)(5\zeta^2 - 1), \quad \mathcal{J}_5(\zeta) = \frac{1}{8}\zeta(1 - \zeta^2)(7\zeta^2 - 3). \end{aligned}$$

The Gegenbauer functions of the second kind are defined as

$$\mathcal{H}_0(\zeta) = -\zeta, \quad \mathcal{H}_1(\zeta) = -1, \quad \mathcal{H}_n(\zeta) = \frac{1}{2} \mathcal{J}_n(\zeta) \ln \frac{1+\zeta}{1-\zeta} + \mathcal{K}_n(\zeta) \quad \text{at } n \geq 2,$$

where the functions  $\mathcal{K}_n(\zeta)$  are expressed in terms of the Gegenbauer functions of the first kind as

$$\mathcal{K}_n(\zeta) = - \sum_k^{\frac{1}{2}n \leq k \leq \frac{1}{2}n + \frac{1}{2}} \frac{(2n-4k+1)}{(2k-1)(n-k)} \left[ 1 - \frac{(2k-1)(n-k)}{n(n-1)} \right] \mathcal{J}_{n-2k+1}(\zeta);$$

the series start with  $\mathcal{J}_0$  or  $\mathcal{J}_1$ , depending on whether  $n$  is odd or even. In particular,

$$\mathcal{K}_2(\zeta) = \frac{1}{2}\zeta, \quad \mathcal{K}_3(\zeta) = \frac{1}{6}(3\zeta^2 - 2), \quad \mathcal{K}_4(\zeta) = \frac{1}{24}\zeta(15\zeta^2 - 13), \quad \mathcal{K}_5(\zeta) = \frac{1}{120}(105\zeta^4 - 115\zeta^2 + 16).$$

For  $n \geq 2$ , the Gegenbauer functions of the second kind assume infinite values at the points  $\zeta = \pm 1$ , which correspond to  $\theta = 0$  and  $\theta = \pi$ . Therefore, if physically there are no singularities in the problem, then the quantities in (1) labeled with a tilde must be set equal to zero. In the overwhelming majority of problems on the flow about particles, drops, or bubbles, the stream function in the spherical coordinates is given by formula (1) with

$$A_1 = A_0 = B_1 = B_0 = C_1 = C_0 = D_1 = D_0 = 0; \quad \tilde{A}_n = \tilde{B}_n = \tilde{C}_n = \tilde{D}_n = 0 \quad \text{for } n = 2, 3, \dots$$

**Example 1.** In the problem on the translational Stokes flow about a solid spherical particle, the following boundary conditions are imposed on the stream function  $w$ :

$$w = 0 \quad \text{at } r = R, \quad \partial_r w = 0 \quad \text{at } r = R, \quad w \rightarrow \frac{1}{2} U r^2 \sin^2 \theta \quad \text{as } r \rightarrow \infty,$$

where  $R$  is the radius of the particle and  $U$  is the unperturbed fluid velocity at infinity.

Stokes solution:

$$w(r, \theta) = \frac{1}{4} U (r - R)^2 \left( 2 + \frac{R}{r} \right) \sin^2 \theta.$$

Here, only the terms for  $n = 2$  in the first sum of (1) remain.

**Example 2.** In the problem on the axisymmetric straining Stokes flow about a solid spherical particle, the following boundary conditions are imposed on the stream function  $w$ :

$$w = 0 \quad \text{at} \quad r = R, \quad \partial_r w = 0 \quad \text{at} \quad r = R, \quad w \rightarrow \frac{1}{2} E r^3 \sin^2 \theta \cos \theta \quad \text{as} \quad r \rightarrow \infty,$$

where  $R$  is the radius of the particle and  $E$  is the shear coefficient.

Solution:

$$w(r, \theta) = \frac{1}{2} E R^3 \left( \frac{r^3}{R^3} - \frac{5}{2} + \frac{3}{2} \frac{R^2}{r^2} \right) \sin^2 \theta \cos \theta.$$

Here, only the terms for  $n = 3$  in the first sum of (1) remain.

**Example 3.** Solving the problem of the translational Stokes flow about a spherical drop (or bubble) is reduced to solving the Stokes equation outside and inside the drop. The boundary condition at infinity is specified in example 1. Conjugate boundary conditions are set at the drop surface; these conditions can be found in the references cited below and are not written out here.

Hadamard–Rybczynski solution:

$$w(r, \theta) = \frac{1}{4} U r^2 \left( 2 - \frac{3\beta + 2}{\beta + 1} \frac{R}{r} + \frac{\beta}{\beta + 1} \frac{R^3}{r^3} \right) \sin^2 \theta \quad \text{for} \quad r > R,$$

$$w(r, \theta) = \frac{U}{4(\beta + 1)} r^2 \left( \frac{r^2}{R^2} - 1 \right) \sin^2 \theta \quad \text{for} \quad r < R,$$

where  $R$  is the radius of the drop,  $U$  the unperturbed fluid velocity at infinity,  $\beta$  the ratio of the dynamic viscosities of the fluids inside and outside the drop (the value  $\beta = 0$  corresponds to a gas bubble and  $\beta = \infty$  to a solid particle).

**Example 4.** Solving the problem of the axisymmetric straining Stokes flow about a spherical drop (or bubble) is reduced to solving the Stokes equation outside and inside the drop. The boundary condition at infinity is specified in example 2. Conjugate boundary conditions are set at the drop surface; these conditions can be found in the references cited below and are not written out here.

Taylor solution:

$$w(r, \theta) = \frac{1}{2} E R^3 \left( \frac{r^3}{R^3} - \frac{1}{2} \frac{5\beta + 2}{\beta + 1} + \frac{3}{2} \frac{\beta}{\beta + 1} \frac{R^2}{r^2} \right) \sin^2 \theta \cos \theta \quad \text{for} \quad r > R,$$

$$w(r, \theta) = \frac{3}{4} \frac{E R^3}{\beta + 1} \frac{r^3}{R^3} \left( \frac{r^2}{R^2} - 1 \right) \sin^2 \theta \cos \theta \quad \text{for} \quad r < R,$$

where  $R$  is the drop radius,  $E$  the shear coefficient,  $\beta$  the ratio of the dynamic viscosities of the fluids inside and outside the drop (the value  $\beta = 0$  corresponds to a gas bubble and  $\beta = \infty$  to a solid particle).

⊙ *References:* G. I. Taylor (1932), V. G. Levich (1962), J. Happel and H. Brenner (1965), A. D. Polyanin, A. M. Kutepov, A. V. Vyazmin, and D. A. Kazenin (2001).

#### 9.4.6-2. Stokes equation in the bipolar coordinate system.

When studying axisymmetric problems of a flow about two spherical particles (drops, bubbles), one uses the bipolar coordinates  $\xi, \eta$ ; these are related to the cylindrical coordinates  $\rho = r \cos \theta$ ,  $z = r \sin \theta$  by

$$\rho = \frac{a \sin \xi}{\cosh \eta - \cos \xi}, \quad z = \frac{a \sinh \eta}{\cosh \eta - \cos \xi}.$$

The general solution of the equation  $E^2(E^2 w) = 0$  in the bipolar coordinate system has the form

$$w(\xi, \eta) = \frac{1}{(\cosh \eta - \cos \xi)^{3/2}} \left[ \sum_{n=0}^{\infty} \mathcal{J}_{n+1}(\cos \xi) f_n(\eta) + \sum_{n=0}^{\infty} \mathcal{H}_{n+1}(\cos \xi) g_n(\eta) \right],$$

$$f_n(\eta) = A_n \cosh \left[ \left( n - \frac{1}{2} \right) \eta \right] + B_n \sinh \left[ \left( n - \frac{1}{2} \right) \eta \right] + C_n \cosh \left[ \left( n + \frac{3}{2} \right) \eta \right] + D_n \sinh \left[ \left( n + \frac{3}{2} \right) \eta \right],$$

$$g_n(\eta) = \tilde{A}_n \cosh \left[ \left( n - \frac{1}{2} \right) \eta \right] + \tilde{B}_n \sinh \left[ \left( n - \frac{1}{2} \right) \eta \right] + \tilde{C}_n \cosh \left[ \left( n + \frac{3}{2} \right) \eta \right] + \tilde{D}_n \sinh \left[ \left( n + \frac{3}{2} \right) \eta \right],$$

where the  $A_n, B_n, C_n, D_n, \tilde{A}_n, \tilde{B}_n, \tilde{C}_n$ , and  $\tilde{D}_n$  are arbitrary constants and the  $\mathcal{J}_n(\zeta)$  and  $\mathcal{H}_n(\zeta)$  are the Gegenbauer functions.

⊙ *Reference:* J. Happel and H. Brenner (1965).



### 9.4.6-3. Stokes equation in the oblate spheroidal coordinate system.

When studying axisymmetric problems of flows about spheroidal particles, one uses the oblate spheroidal coordinates  $\xi, \eta$ ; these are related to the cylindrical coordinates  $\rho = r \cos \theta, z = r \sin \theta$  by

$$\rho = c \cosh \xi \sin \eta, \quad z = c \sinh \xi \cos \eta.$$

The solution of the equation  $E^2(E^2 w) = 0$  that describes the flow of a fluid about a prolate spheroid in the direction parallel to the spheroid axis is expressed as

$$w = \frac{1}{2} U c^2 \cosh^2 \xi \sin^2 \eta \left\{ 1 - \frac{[\lambda/(\lambda^2+1)] - [(\lambda_0^2-1)/(\lambda_0^2+1)] \operatorname{arccot} \lambda}{[\lambda_0/(\lambda_0^2+1)] - [(\lambda_0^2-1)/(\lambda_0^2+1)] \operatorname{arccot} \lambda_0} \right\}, \quad \lambda = \sinh \xi, \quad \lambda_0 = \sinh \xi_0.$$

Here,  $w$  is the stream function,  $U$  is the fluid velocity at infinity,  $c$  and  $\lambda_0$  are the constants related to the spheroid semiaxes  $a$  and  $b$  ( $a > b$ ) by  $c = \sqrt{a^2 - b^2}$  and  $\lambda_0 = b/c$ .

⊙ *Reference:* J. Happel and H. Brenner (1965).

## 9.5. Higher-Order Linear Equations with Constant Coefficients

► Throughout Section 9.5 the following notation is used:

$$\mathbf{x} = \{x_1, \dots, x_n\}, \quad \mathbf{y} = \{y_1, \dots, y_n\}, \quad \boldsymbol{\omega} = \{\omega_1, \dots, \omega_n\}, \quad \boldsymbol{\xi} = \{\xi_1, \dots, \xi_n\},$$

$$|\mathbf{x}| = \sqrt{x_1^2 + \dots + x_n^2}, \quad |\boldsymbol{\omega}| = \sqrt{\omega_1^2 + \dots + \omega_n^2}, \quad \boldsymbol{\omega} \cdot \mathbf{x} = \omega_1 x_1 + \dots + \omega_n x_n.$$

### 9.5.1. Fundamental Solutions. Cauchy Problem

9.5.1-1. Domain:  $\mathbb{R}^n = \{-\infty < x_k < \infty; k = 1, \dots, n\}$ .

Let  $P$  be a constant coefficient linear differential operator such that

$$P\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) \equiv \sum_{s=0}^M a_{s_1, \dots, s_n} \frac{\partial^s}{\partial x_1^{s_1} \dots \partial x_n^{s_n}}, \quad s = s_1 + \dots + s_n,$$

where  $s_1, \dots, s_n$  are nonnegative integers,  $a_{s_1, \dots, s_n}$  are some constants, and  $M$  is the order of the operator. A generalized function (distribution)  $\mathcal{E}(\mathbf{x}) = \mathcal{E}(x_1, \dots, x_n)$  that satisfies the equation

$$P\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) \mathcal{E}(\mathbf{x}) = \delta(\mathbf{x}),$$

where  $\delta(\mathbf{x}) = \delta(x_1) \dots \delta(x_n)$  is the Dirac delta function in the  $n$ -dimensional Euclidian space, is called the fundamental solution corresponding to the operator  $P$ .

Any constant coefficient linear differential operator has a fundamental solution  $\mathcal{E}(\mathbf{x})$ . The fundamental solution is not unique — it is defined up to an additive term  $w_0(\mathbf{x})$  that is an arbitrary solution of the homogeneous equation  $P\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) w_0(\mathbf{x}) = 0$ .

The solution of the nonhomogeneous equation

$$P\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) w = \Phi(\mathbf{x})$$

with an arbitrary right-hand side has the form

$$w(\mathbf{x}) = \mathcal{E}(\mathbf{x}) * \Phi(\mathbf{x}), \quad \mathcal{E}(\mathbf{x}) * \Phi(\mathbf{x}) = \int_{\mathbb{R}^n} \mathcal{E}(\mathbf{x} - \mathbf{y}) \Phi(\mathbf{y}) d\mathbf{y}.$$

Here,  $d\mathbf{y} = dy_1 \dots dy_n$  and the convolution  $\mathcal{E} * \Phi$  is assumed to be meaningful.

⊙ *References:* G. E. Shilov (1965), S. G. Krein (1972), L. Hörmander (1983), V. S. Vladimirov (1988).

9.5.1-2. Domain:  $0 \leq t < \infty$ ,  $-\infty < x_k < \infty$ ;  $k = 1, \dots, n$ . Cauchy problem.

Now let  $P(\frac{\partial}{\partial t}, \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n})$  be a constant coefficient linear differential operator of order  $m$  with respect to  $t$ . Then a distribution  $\mathcal{E}(t, \mathbf{x}) = \mathcal{E}(t, x_1, \dots, x_n)$ , which is a solution of the equation

$$P\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) \mathcal{E}(t, \mathbf{x}) = 0$$

and satisfies the initial conditions\*

$$\mathcal{E}|_{t=0} = 0, \quad \left. \frac{\partial \mathcal{E}}{\partial t} \right|_{t=0} = 0, \quad \dots, \quad \left. \frac{\partial^{m-2} \mathcal{E}}{\partial t^{m-2}} \right|_{t=0} = 0, \quad \left. \frac{\partial^{m-1} \mathcal{E}}{\partial t^{m-1}} \right|_{t=0} = \delta(\mathbf{x}), \quad (1)$$

is called a fundamental solution of the Cauchy problem corresponding to the operators  $P$ .

## The solution of the Cauchy problem for the linear differential equation

$$P\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right)w = 0 \quad (2)$$

with the special initial conditions

$$w|_{t=0} = 0, \quad \left. \frac{\partial w}{\partial t} \right|_{t=0} = 0, \quad \dots, \quad \left. \frac{\partial^{m-2} w}{\partial t^{m-2}} \right|_{t=0} = 0, \quad \left. \frac{\partial^{m-1} w}{\partial t^{m-1}} \right|_{t=0} = f(\mathbf{x})$$

is given by

$$w(t, \mathbf{x}) = \mathcal{E}(t, \mathbf{x}) * f(\mathbf{x}), \quad \mathcal{E}(t, \mathbf{x}) * f(\mathbf{x}) \equiv \int_{\mathbb{R}^n} \mathcal{E}(t, \mathbf{x} - \mathbf{y}) f(\mathbf{y}) d\mathbf{y}.$$

⊙ *Reference:* S. G. Krein (1972).

### 9.5.1-3. Solution of the Cauchy problem for general initial conditions.

If the general initial conditions

$$w|_{t=0} = f_0(\mathbf{x}), \quad \left. \frac{\partial w}{\partial t} \right|_{t=0} = f_1(\mathbf{x}), \quad \dots, \quad \left. \frac{\partial^{m-2} w}{\partial t^{m-2}} \right|_{t=0} = f_{m-2}(\mathbf{x}), \quad \left. \frac{\partial^{m-1} w}{\partial t^{m-1}} \right|_{t=0} = f_{m-1}(\mathbf{x}) \quad (3)$$

are set, the solution of equation (2) is sought in the form

$$w(t, \mathbf{x}) = \mathcal{E}(t, \mathbf{x}) * \varphi_0(\mathbf{x}) + \frac{\partial \mathcal{E}(t, \mathbf{x})}{\partial t} * \varphi_1(\mathbf{x}) + \cdots + \frac{\partial^{m-1} \mathcal{E}(t, \mathbf{x})}{\partial t^{m-1}} * \varphi_{m-1}(\mathbf{x}). \quad (4)$$

Each term in (4) satisfies equation (2), and the functions  $\varphi_{m-1}, \varphi_{m-2}, \dots, \varphi_0$  are determined successively from the linear system

$$f_0(\mathbf{x}) = \varphi_{m-1}(\mathbf{x}),$$

$$f_1(\mathbf{x}) = \varphi_{m-2}(\mathbf{x}) + \frac{\partial^m \mathcal{E}(0, \mathbf{x})}{\partial t^m} * \varphi_{m-1}(\mathbf{x}),$$

$$f_k(\mathbf{x}) = \varphi_{m-k-1}(\mathbf{x}) + \frac{\partial^m \mathcal{E}(0, \mathbf{x})}{\partial t^m} * \varphi_{m-k}(\mathbf{x}) + \cdots + \frac{\partial^{m+k-1} \mathcal{E}(0, \mathbf{x})}{\partial t^{m+k-1}} * \varphi_{m-1}(\mathbf{x}), \quad k = 2, \dots, m-1.$$

This system of equations is obtained by successively differentiating relation (4) followed by substituting  $t = 0$  and taking into account the initial conditions (1) and (3).

⊙ *Reference:* G. E. Shilov (1965).

\* The number of initial conditions can be less than  $m$  (see Paragraph 9.5.4-1).

## 9.5.2. Elliptic Equations

### 9.5.2-1. Homogeneous elliptic differential operator.

A constant coefficient linear homogeneous differential operator of order  $k$  has the form

$$P_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) \equiv \sum a_{s_1, \dots, s_n} \left(\frac{\partial}{\partial x_1}\right)^{s_1} \dots \left(\frac{\partial}{\partial x_n}\right)^{s_n}, \quad \sum_{i=1}^n s_i = k,$$

where  $s_1, \dots, s_n$  are nonnegative integers. From now on, we adopt the notation

$$\left(\frac{\partial}{\partial x_1}\right)^{s_1} \dots \left(\frac{\partial}{\partial x_n}\right)^{s_n} \equiv \frac{\partial^{s_1 + \dots + s_n}}{\partial x_1^{s_1} \dots \partial x_n^{s_n}}.$$

A linear homogeneous differential operator of order  $k$  possesses the property

$$P_k\left(b\frac{\partial}{\partial x_1}, \dots, b\frac{\partial}{\partial x_n}\right) = b^k P_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right), \quad b \neq 0 \text{ is an arbitrary constant.}$$

A linear homogeneous differential operator  $P_k$  is called elliptic if, on replacing in  $P_k$  the symbols  $\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}$  by variables  $\omega_1, \dots, \omega_n$ , one obtains a polynomial  $P_k(\omega_1, \dots, \omega_n)$  that does not vanish if  $\omega \neq 0$ , i.e.,

$$P_k(\omega_1, \dots, \omega_n) \equiv \sum a_{s_1, \dots, s_n} \omega_1^{s_1} \dots \omega_n^{s_n} \neq 0 \quad \text{if } |\omega| \neq 0.$$

A linear differential equation

$$P_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right)w \equiv \sum a_{s_1, \dots, s_n} \frac{\partial^{s_1 + \dots + s_n} w}{\partial x_1^{s_1} \dots \partial x_n^{s_n}} = 0, \quad \sum_{i=1}^n s_i = k \quad (1)$$

is called elliptic if the linear homogeneous differential operator  $P_k$  is elliptic.

### 9.5.2-2. Elliptic differential operator of general form.

In general, a constant coefficient linear differential operator of order  $k$  has the form

$$\mathcal{P}_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) = P_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right) + \sum_{i=0}^{k-1} P_i\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right),$$

where  $P_k$  is the leading part of the operator and  $P_j$  ( $j = 0, 1, \dots, k$ ) is a linear homogeneous differential operator of order  $j$ . The operator  $\mathcal{P}_k$  is said to be elliptic if its leading part  $P_k$  is elliptic.

A linear differential equation

$$\mathcal{P}_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right)w = 0 \quad (2)$$

is called elliptic if the linear differential operator  $\mathcal{P}_k$  is elliptic.

*Remark.* A linear elliptic operator and a linear elliptic differential equation can only be of even order  $k = 2m$ , where  $m$  is a positive integer.

### 9.5.2-3. Fundamental solution of a homogeneous elliptic equation.

The fundamental solution of the homogeneous elliptic equation (1) with  $k = 2m$  is given by

$$\begin{aligned}\mathcal{E}(\mathbf{x}) &= \frac{(-1)^{\frac{n-1}{2}}}{4(2\pi)^{n-1}(2m-n)!} \int_{\Omega_n} |\boldsymbol{\omega} \cdot \mathbf{x}|^{2m-n} \frac{d\Omega_n}{P_{2m}(\boldsymbol{\omega})} && \text{if } n \text{ is odd and } 2m \geq n; \\ \mathcal{E}(\mathbf{x}) &= \frac{(-1)^{\frac{n-2}{2}}}{(2\pi)^n(2m-n)!} \int_{\Omega_n} |\boldsymbol{\omega} \cdot \mathbf{x}|^{2m-n} \ln |\boldsymbol{\omega} \cdot \mathbf{x}| \frac{d\Omega_n}{P_{2m}(\boldsymbol{\omega})} && \text{if } n \text{ is even and } 2m \geq n; \\ \mathcal{E}(\mathbf{x}) &= \frac{(-1)^{\frac{n-1}{2}}}{2(2\pi)^{\frac{n-1}{2}}} \int_{\Omega_n} \delta^{(n-2m-1)}(\boldsymbol{\omega} \cdot \mathbf{x}) \frac{d\Omega_n}{P_{2m}(\boldsymbol{\omega})} && \text{if } n \text{ is odd and } 2m < n; \\ \mathcal{E}(\mathbf{x}) &= \frac{(-1)^{\frac{n}{2}}(n-2m-1)!}{(2\pi)^n} \int_{\Omega_n} |\boldsymbol{\omega} \cdot \mathbf{x}|^{2m-n} \frac{d\Omega_n}{P_{2m}(\boldsymbol{\omega})} && \text{if } n \text{ is even and } 2m < n.\end{aligned}$$

Here, the integration is performed over the surface of the  $n$ -dimensional sphere  $\Omega_n$  of unit radius defined by the equation  $|\boldsymbol{\omega}| = 1$ ;  $\boldsymbol{\omega} \cdot \mathbf{x} = \omega_1 x_1 + \dots + \omega_n x_n$  and  $P_{2m}(\boldsymbol{\omega}) = P_{2m}(\omega_1, \dots, \omega_n)$ .

A fundamental solution is an ordinary function, analytic at any point  $\mathbf{x} \neq 0$ ; this function is described, in a neighborhood of the origin of coordinates (as  $|\mathbf{x}| \rightarrow 0$ ), by the relations

$$\mathcal{E}(\mathbf{x}) = \begin{cases} b_{n,m} |\mathbf{x}|^{2m-n} & \text{if } n \text{ is odd or } n \text{ is even and } n > 2m; \\ c_{n,m} |\mathbf{x}|^{2m-n} \ln |\mathbf{x}| & \text{if } n \text{ is even and } n \leq 2m. \end{cases}$$

Here,  $b_{n,m}$  and  $c_{n,m}$  are some nonzero constants. If  $2m > n$ , the fundamental solution has continuous derivatives up to order  $2m - n - 1$  inclusive at the origin.

### 9.5.2-4. Fundamental solution of a general elliptic equation.

The fundamental solution of the general elliptic equation (2) with  $k = 2m$  is determined from the relation

$$\mathcal{E}(\mathbf{x}) = \int_{\Omega_n} Z_{\omega}(\boldsymbol{\omega} \cdot \mathbf{x}, -n) d\Omega_n, \quad (3)$$

where

$$Z_{\omega}(\xi, \lambda) = \frac{1}{\sigma_n \pi^{\frac{n-1}{2}} \Gamma\left(\frac{\lambda+1}{2}\right)} \int_{-\infty}^{\infty} G(\xi - \eta, \boldsymbol{\omega}) |\eta|^{\lambda} d\eta, \quad \sigma_n = \frac{2\pi^{n/2}}{\Gamma(n/2)}.$$

Here, the function  $G(\xi, \boldsymbol{\omega})$  is a fundamental solution of the constant coefficient linear ordinary differential equation

$$\mathcal{P}_{2m}\left(\omega_1 \frac{d}{d\xi}, \dots, \omega_n \frac{d}{d\xi}\right) G(\xi, \boldsymbol{\omega}) = \delta(\xi).$$

If  $n$  is odd, the fundamental solution (3) can be represented as

$$\mathcal{E}(\mathbf{x}) = A_n \int_{\Omega_n} \left[ \frac{\partial^{n-1}}{\partial \xi^{n-1}} G(\xi, \boldsymbol{\omega}) \right] d\Omega_n, \quad A_n = \frac{(-1)^{\frac{n-1}{2}}}{1 \times 3 \dots (n-2) \sigma_n (2\pi)^{\frac{n-1}{2}}}.$$

⊙ *References:* I. M. Gel'fand, G. E. Shilov (1959), S. G. Krein (1972).

### 9.5.3. Hyperbolic Equations

Let  $P\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right)$  be a constant coefficient linear homogeneous differential operator of order  $m$  with respect to  $t$ . The operator  $P$  is called hyperbolic if for any numbers  $\omega_1, \dots, \omega_n$  such that  $\sum_{s=1}^n \omega_s^2 = 1$ , the  $m$ th-order algebraic equation

$$P(\lambda, \omega_1, \dots, \omega_n) = 0$$

with respect to  $\lambda$  has  $m$  different real roots.

Fundamental solution of the Cauchy problem for  $m \geq n - 1$ :

$$\begin{aligned} \mathcal{E}(t, \mathbf{x}) &= \frac{(-1)^{\frac{n+1}{2}}}{2(2\pi)^{n-1}(m-n-1)!} \int_{H=0} (\boldsymbol{\xi} \cdot \mathbf{x} + t)^{m-n-1} \frac{[\text{sign}(\boldsymbol{\xi} \cdot \mathbf{x} + t)]^{m-1}}{|\nabla H| \text{sign}(\boldsymbol{\xi} \cdot \nabla H)} d\sigma_H \quad \text{if } n \text{ is odd;} \\ \mathcal{E}(t, \mathbf{x}) &= \frac{2(-1)^{\frac{n}{2}}}{(2\pi)^n(m-n-1)!} \int_{H=0} \frac{(\boldsymbol{\xi} \cdot \mathbf{x} + t)^{m-n-1}}{|\nabla H| \text{sign}(\boldsymbol{\xi} \cdot \nabla H)} \ln \left| \frac{\boldsymbol{\xi} \cdot \mathbf{x} + t}{\boldsymbol{\xi} \cdot \mathbf{x}} \right| d\sigma_H \quad \text{if } n \text{ is even,} \end{aligned}$$

where

$$H = P(1, \xi_1, \dots, \xi_n), \quad |\nabla H| = \sqrt{\left(\frac{\partial H}{\partial \xi_1}\right)^2 + \dots + \left(\frac{\partial H}{\partial \xi_n}\right)^2}, \quad \boldsymbol{\xi} \cdot \nabla H = \xi_1 \frac{\partial H}{\partial \xi_1} + \dots + \xi_n \frac{\partial H}{\partial \xi_n},$$

and  $d\sigma_H$  is the element of the surface  $H = 0$ .

Fundamental solution of the Cauchy problem for  $m < n - 1$ :

$$\begin{aligned} \mathcal{E}(t, \mathbf{x}) &= \frac{(-1)^{\frac{n+1}{2}}}{(2\pi)^{n-1}} \int_{H=0} \frac{\delta^{(n-m)}(\boldsymbol{\xi} \cdot \mathbf{x} + t)}{|\nabla H| \text{sign}(\boldsymbol{\xi} \cdot \nabla H)} d\sigma_H \quad \text{if } n \text{ is odd;} \\ \mathcal{E}(t, \mathbf{x}) &= \frac{(-1)^{\frac{n}{2}}(n-m)!}{(2\pi)^n} \int_{H=0} \frac{(\boldsymbol{\xi} \cdot \mathbf{x} + t)^{m-n-1}}{|\nabla H| \text{sign}(\boldsymbol{\xi} \cdot \nabla H)} d\sigma_H \quad \text{if } n \text{ is even.} \end{aligned}$$

⊙ *References:* I. M. Gel'fand, G. E. Shilov (1959), S. G. Krein (1972).

### 9.5.4. Regular Equations. Number of Initial Conditions in the Cauchy Problem

9.5.4-1. Equations with two independent variables ( $0 \leq t < \infty$ ,  $-\infty < x < \infty$ ).

1°. Consider the constant coefficient linear differential equation

$$\frac{\partial^m w}{\partial t^m} = \sum_{k=0}^{m-1} p_k \left( i \frac{\partial}{\partial x} \right) \frac{\partial^k w}{\partial t^k}, \quad (1)$$

where  $p_k(z)$  is a polynomial of degree  $k$ ,  $i^2 = -1$ . Let  $r = r(\sigma)$  be the number of roots (taking into account their multiplicities) of the characteristic equation

$$\lambda^m - \sum_{k=0}^{m-1} p_k(\sigma) \lambda^k = 0 \quad (2)$$

whose real parts are nonpositive (or bounded above) for given a  $\sigma$ . If  $r$  is the same (up to a set of measure zero) for all  $\sigma \in (-\infty, \infty)$ , the equation (1) will be called regular with regularity index  $r$ .

Classical equations such as the heat, wave, and Laplace equations are regular.

2°. In the Cauchy problem for the regular equation (1), one should set  $r$  initial conditions of the form

$$w|_{t=0} = f_0(x), \quad \frac{\partial w}{\partial t} \Big|_{t=0} = f_1(x), \quad \dots, \quad \frac{\partial^{r-2} w}{\partial t^{r-2}} \Big|_{t=0} = f_{r-2}(x), \quad \frac{\partial^{r-1} w}{\partial t^{r-1}} \Big|_{t=0} = f_{r-1}(x). \quad (3)$$

It should be emphasized that the regularity index  $r$  can, in general, differ from the equation order  $m$  with respect to  $t$ . In particular, for the two-dimensional Laplace equation  $\partial_{tt} w = -\partial_{xx} w$ , we have  $r = 1$  and  $m = 2$ ; here,  $y$  is replaced by  $t$  and the first boundary value problem in the upper half-plane  $t \geq 0$  is considered. For the heat equation  $\partial_t w = \partial_{xx} w$  and the wave equation  $\partial_{tt} w = \partial_{xx} w$ , we have  $r = m = 1$  and  $r = m = 2$ , respectively.

**Example 1.** Below are the regularity indices for some fourth-order equations:

$$\begin{aligned} \frac{\partial^2 w}{\partial t^2} - a^2 \frac{\partial^4 w}{\partial x^4} &= 0 & (r = 1, m = 2); & \quad \frac{\partial^2 w}{\partial t^2} + a^2 \frac{\partial^4 w}{\partial x^4} = 0 & (r = 2, m = 2); \\ \left( \frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} \right)^2 w &= 0 & (r = 2, m = 4); & \quad \frac{\partial^4 w}{\partial t^4} - a^2 \frac{\partial^4 w}{\partial x^4} = 0 & (r = 3, m = 4). \end{aligned}$$

3°. The special solution  $\mathcal{E} = \mathcal{E}(t, x)$  that satisfies the initial conditions

$$\mathcal{E}|_{t=0} = 0, \quad \frac{\partial \mathcal{E}}{\partial t} \Big|_{t=0} = 0, \quad \dots, \quad \frac{\partial^{r-2} \mathcal{E}}{\partial t^{r-2}} \Big|_{t=0} = 0, \quad \frac{\partial^{r-1} \mathcal{E}}{\partial t^{r-1}} \Big|_{t=0} = \delta(x) \quad (4)$$

is called fundamental.

The fundamental solution can be found by applying the Fourier transform in the space variable to equation (1) (with  $w = \mathcal{E}$ ) and the initial conditions (4).

**Example 2.** Consider the polyharmonic equation:

$$\left( \frac{\partial^2}{\partial t^2} + \frac{\partial^2}{\partial x^2} \right)^n w = 0. \quad (5)$$

Taking into account the representation  $\frac{\partial^2}{\partial x^2} = -(i \frac{\partial}{\partial x})^2$ , we rewrite the characteristic equation (2) in the form

$$(\lambda^2 - \sigma^2)^n = 0.$$

It has only one solution whose real part is nonpositive, specifically,  $\lambda = -|\sigma|$ . Considering the multiplicity of the root, we find that the regularity index  $r$  is equal to  $n$ .

In equation (5) with  $w = \mathcal{E}$  and the initial conditions (4) with  $r = n$ , we perform the Fourier transform with respect to the space variable,

$$U(t, \sigma) = \int_{-\infty}^{\infty} e^{i\sigma x} \mathcal{E}(t, x) dx.$$

As a result, we arrive at the ordinary differential equation

$$\left( \frac{d^2}{dt^2} - \sigma^2 \right)^n U = 0 \quad (6)$$

and the initial conditions

$$U|_{t=0} = 0, \quad U'|_{t=0} = 0, \quad \dots, \quad U_t^{(n-2)}|_{t=0} = 0, \quad U_t^{(n-1)}|_{t=0} = 1. \quad (7)$$

The bounded solution of problem (6), (7) is given by

$$U(t, \sigma) = \frac{t^{n-1}}{(n-1)!} e^{-|\sigma|t}.$$

By applying the inverse Fourier transform, we obtain the fundamental solution of the polyharmonic equation in the form

$$\begin{aligned} \mathcal{E}(t, x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{-i\sigma x} U(t, \sigma) d\sigma = \frac{1}{2\pi} \frac{t^{n-1}}{(n-1)!} \int_{-\infty}^{\infty} e^{-i\sigma x - |\sigma|t} d\sigma \\ &= \frac{1}{2\pi} \frac{t^{n-1}}{(n-1)!} \left( \int_0^{\infty} e^{-i\sigma x - \sigma t} d\sigma + \int_{-\infty}^0 e^{-i\sigma x + \sigma t} d\sigma \right) \\ &= \frac{1}{2\pi} \frac{t^{n-1}}{(n-1)!} \left( \frac{1}{t+ix} + \frac{1}{t-ix} \right) = \frac{1}{\pi} \frac{t^{n-1}}{(n-1)!} \frac{t}{t^2 + x^2}. \end{aligned}$$

4°. For general initial conditions of the form (3), the solution of equation (1) is determined on the basis of the fundamental solution from the relation

$$w(t, x) = \mathcal{E}(t, x) * \varphi_0(x) + \frac{\partial \mathcal{E}(t, x)}{\partial t} * \varphi_1(x) + \cdots + \frac{\partial^{r-1} \mathcal{E}(t, x)}{\partial t^{r-1}} * \varphi_{r-1}(x). \quad (8)$$

Each term in (8) satisfies equation (1), and the functions  $\varphi_{r-1}, \varphi_{r-2}, \dots, \varphi_0$  are calculated successively by solving the linear system

$$\begin{aligned} f_0(x) &= \varphi_{r-1}(x), \\ f_1(x) &= \varphi_{r-2}(x) + \frac{\partial^r \mathcal{E}(0, x)}{\partial t^r} * \varphi_{r-1}(x), \\ &\dots\dots\dots \\ f_k(x) &= \varphi_{r-k-1}(x) + \frac{\partial^r \mathcal{E}(0, x)}{\partial t^r} * \varphi_{r-k}(x) + \cdots + \frac{\partial^{r+k-1} \mathcal{E}(0, x)}{\partial t^{r+k-1}} * \varphi_{r-1}(x), \quad k = 2, \dots, r-1. \end{aligned}$$

This system of equations is obtained by successively differentiating relation (8) followed by substituting  $t = 0$  and taking into account the initial conditions (3) and (4).

In the special case  $f_0(x) = f_1(x) = \cdots = f_{r-2}(x) = 0$ , one should set  $\varphi_0(x) = f_{r-1}(x)$  and  $\varphi_1(x) = \cdots = \varphi_{r-1}(x) = 0$  in (8).

⊙ *Reference:* G. E. Shilov (1965).

#### 9.5.4-2. Equations with many independent variables ( $0 \leq t < \infty, \mathbf{x} \in \mathbb{R}^n$ ).

Solving the Cauchy problem for the constant coefficient linear differential equation

$$P\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right)w = 0 \quad (9)$$

with arbitrarily many space variables  $x_1, \dots, x_n$  can be reduced to solving the Cauchy problem for an equation with one space variable  $\xi$ . We take an auxiliary linear differential operator

$$P_\omega\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial \xi}\right) \equiv P\left(\frac{\partial}{\partial t}, \omega_1 \frac{\partial}{\partial \xi}, \dots, \omega_n \frac{\partial}{\partial \xi}\right)$$

that depends on two independent variables  $t$  and  $\xi$  so that the Cauchy problem for the equation

$$P_\omega\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial \xi}\right)v = 0 \quad (10)$$

is well posed. Then the fundamental solution of the Cauchy problem for the original equation (9) is given by

$$\mathcal{E}(t, \mathbf{x}) = \int_{\Omega_n} v_\omega(t, \boldsymbol{\omega} \cdot \mathbf{x}, -n) d\Omega_n.$$

Here,

$$v_\omega(t, \xi, \lambda) = \frac{1}{\sigma_n \pi^{\frac{n-1}{2}} \Gamma\left(\frac{\lambda+1}{2}\right)} \int_{-\infty}^{\infty} G_\omega(t, \xi - \eta) |\eta|^\lambda d\eta, \quad \sigma_n = \frac{2\pi^{n/2}}{\Gamma(n/2)},$$

where  $G_\omega(t, \xi)$  is the fundamental solution of the Cauchy problem for the auxiliary equation (10).

If the number of space variables is odd, one can use the simpler formula

$$\mathcal{E}(t, \mathbf{x}) = \frac{(-1)^{\frac{n-1}{2}} \left(\frac{n-1}{2}\right)!}{\sigma_n \pi^{\frac{n-1}{2}} (n-1)!} \int_{\Omega_n} \left[ \frac{d^{n-1}}{d\xi^{n-1}} G_\omega(t, \xi) \right] d\Omega_n, \quad \xi = \boldsymbol{\omega} \cdot \mathbf{x}.$$

*Remark.* The above relations hold for all equations for which the Cauchy problem is well posed.

⊙ *References:* I. M. Gel'fand, G. E. Shilov (1959), S. G. Krein (1972).

### 9.5.4-3. Stationary homogeneous regular equations ( $\mathbf{x} \in \mathbb{R}^n$ ).

A linear differential operator  $P_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right)$  is called regular if it is homogeneous and if the gradient of the function  $P_k(\omega_1, \dots, \omega_n)$  on the set defined by the equation  $P_k(\omega_1, \dots, \omega_n) = 0$  is everywhere nonzero whenever  $|\omega| \neq 0$ .

The fundamental solution of the linear regular differential equation  $P_k\left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n}\right)w = 0$  generated by the linear regular differential operator  $P_k$  is expressed as

$$\mathcal{E}(\mathbf{x}) = \int_{\Omega_n} \frac{\varphi_{nk}(\omega \cdot \mathbf{x})}{P_k(\omega)} d\Omega_n, \quad (11)$$

where the function  $\varphi_{nk}(z)$  is defined by

$$\begin{aligned} \varphi_{nk}(z) &= \frac{(-1)^{\frac{n-2}{2}}}{(2\pi)^n(k-n)!} z^{k-n} \ln|z| && \text{if } n \text{ is even and } k \geq n; \\ \varphi_{nk}(z) &= \frac{(-1)^{\frac{n+2k}{2}}(n-k-1)!}{(2\pi)^n} z^{k-n} && \text{if } n \text{ is even and } k < n; \\ \varphi_{nk}(z) &= \frac{(-1)^{\frac{n-1}{2}}}{4(2\pi)^{n-1}(k-n)!} z^{k-n} \operatorname{sign} z && \text{if } n \text{ is odd and } k \geq n; \\ \varphi_{nk}(z) &= \frac{(-1)^{\frac{n-1}{2}}}{2(2\pi)^{n-1}} \delta^{(n-k-1)}(z) && \text{if } n \text{ is odd and } k < n. \end{aligned}$$

The integral in (11) is understood in the sense of its regularized value, i.e.,

$$\mathcal{E}(\mathbf{x}) = \lim_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(\mathbf{x}), \quad \mathcal{E}_\varepsilon(\mathbf{x}) = \int_{\Omega_n^{(\varepsilon)}} \frac{\varphi_{nk}(\omega \cdot \mathbf{x})}{P_k(\omega)} d\Omega_n^{(\varepsilon)},$$

where  $\Omega_n^{(\varepsilon)}$  is the set of points on a sphere of unit radius for which  $|P_k(\omega)| > \varepsilon$ .

⊙ *References:* I. M. Gel'fand, G. E. Shilov (1959), S. G. Krein (1972).

### 9.5.5. Some Special-Type Equations

$$1. \quad \frac{\partial w}{\partial t} = P_n\left(i \frac{\partial}{\partial x}\right)w, \quad P_n(z) = a_n z^n + \dots + a_1 z + a_0, \quad i^2 = -1.$$

The condition  $\operatorname{Re} P_n(z) \leq C < \infty$  is assumed to be met for all real  $z$ .

1°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

An initial condition is prescribed:

$$w = f(x) \quad \text{at} \quad t = 0. \quad (1)$$

Solution:

$$w(x, t) = \int_{-\infty}^{\infty} G(x - \xi, t) f(\xi) d\xi, \quad G(x, t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \exp[tP_n(\lambda) - i x \lambda] d\lambda.$$

2°. The solution of the Cauchy problem with the initial condition (1) for the nonhomogeneous equation

$$\frac{\partial w}{\partial t} = P_n\left(i \frac{\partial}{\partial x}\right)w + \Phi(x, t)$$

is given by

$$w(x, t) = \int_{-\infty}^{\infty} G(x - \xi, t) f(\xi) d\xi + \int_0^t \int_{-\infty}^{\infty} G(x - \xi, t - \tau) \Phi(\xi, \tau) d\xi d\tau,$$

where the function  $G(x, t)$  is defined in Item 1°.

⊙ *References:* S. G. Krein (1972), V. S. Vladimirov, V. P. Mikhailov, A. A. Vasharin, et al. (1974).



$$2. \quad \left( \frac{\partial}{\partial t} - a \frac{\partial}{\partial x} \right)^n w = 0, \quad n = 1, 2, \dots$$

1°. General solution (two representations):

$$w(x, t) = \sum_{k=0}^{n-1} t^k f_k(x + at),$$

$$w(x, t) = \sum_{k=0}^{n-1} x^k f_k(x + at),$$

where the  $f_k = f_k(z)$  are arbitrary functions.

2°. Fundamental solution:

$$\mathcal{E}(x, t) = \frac{t^{n-1}}{(n-1)!} \delta(x + at).$$

$$3. \quad \left( \frac{\partial}{\partial t} - \frac{\partial^2}{\partial x^2} \right)^n w = 0, \quad n = 1, 2, \dots$$

1°. General solution (two representations):

$$w(x, t) = \sum_{k=0}^{n-1} t^k u_k(x, t),$$

$$w(x, t) = \sum_{k=0}^{n-1} x^k u_k(x, t),$$

where the  $u_k = u_k(x, t)$  are arbitrary functions that satisfy the heat equations  $\partial_t u_k - \partial_{xx} u_k = 0$ .

2°. Fundamental solution:

$$\mathcal{E}(x, t) = \frac{1}{2\sqrt{\pi}(n-1)!} t^{n-3/2} \exp\left(-\frac{x^2}{4t}\right).$$

3°. Domain:  $-\infty < x < \infty$ . Cauchy problem.

Initial conditions are prescribed:

$$w|_{t=0} = 0, \quad \frac{\partial w}{\partial t} \Big|_{t=0} = 0, \quad \dots, \quad \frac{\partial^{n-2} w}{\partial t^{n-2}} \Big|_{t=0} = 0, \quad \frac{\partial^{n-1} w}{\partial t^{n-1}} \Big|_{t=0} = f(x).$$

Solution:

$$w(x, t) = \int_{-\infty}^{\infty} f(\xi) \mathcal{E}(x - \xi, t) d\xi.$$

⊙ *Reference:* G. E. Shilov (1965).

$$4. \quad \left( \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial x^2} \right)^n w = 0, \quad n = 1, 2, \dots$$

1°. General solution (two representations):

$$w(x, t) = \sum_{k=0}^{n-1} t^k [f_k(x+t) + g_k(x-t)],$$

$$w(x, t) = \sum_{k=0}^{n-1} x^k [f_k(x+t) + g_k(x-t)],$$

where the  $f_k = f_k(y)$  and  $g_k = g_k(z)$  are arbitrary functions.

2°. Fundamental solution:

$$\mathcal{E}(x, t) = \frac{(-1)^{n-1}}{4^n(n-1)!} \left[ \text{sign}(x-t) \sum_{k=0}^{n-1} \frac{(2t)^k (x-t)^{2n-k-2}}{k!(n-k-1)!} + (-1)^n \text{sign}(x+t) \sum_{k=0}^{n-1} \frac{(-2t)^k (x+t)^{2n-k-2}}{k!(n-k-1)!} \right].$$

⊙ *Reference:* G. E. Shilov (1965).

$$5. \quad \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)^n w = 0, \quad n = 1, 2, \dots$$

This is the *polyharmonic equation* of order  $n$  with two independent variables.

1°. General solution (two representations):

$$w(x, y) = \sum_{k=0}^{n-1} r^{2k} u_k(x, y), \quad r = \sqrt{x^2 + y^2},$$

$$w(x, y) = \sum_{k=0}^{n-1} x^k u_k(x, y),$$

where the  $u_k(x, y)$  are arbitrary harmonic functions ( $\Delta u_k = 0$ ). In the second relation,  $x^k$  can be replaced by  $y^k$ .

2°. Domain:  $-\infty < x < \infty, -\infty < y < \infty$ . Fundamental solution:

$$\mathcal{E}(x, y) = \frac{1}{\pi 2^{2n-1} [(n-1)!]^2} r^{2n-2} \ln r, \quad r = \sqrt{x^2 + y^2}.$$

3°. Domain:  $-\infty < x < \infty, 0 \leq y < \infty$ . Boundary value problem.

Boundary conditions are prescribed:

$$w|_{y=0} = 0, \quad \frac{\partial w}{\partial y} \Big|_{y=0} = 0, \quad \dots, \quad \frac{\partial^{n-2} w}{\partial y^{n-2}} \Big|_{y=0} = 0, \quad \frac{\partial^{n-1} w}{\partial y^{n-1}} \Big|_{y=0} = f(x).$$

Solution:

$$w(x, y) = \int_{-\infty}^{\infty} f(\xi) G(x - \xi, y) d\xi, \quad G(x, y) = \frac{1}{\pi(n-1)!} \frac{y^n}{x^2 + y^2}.$$

See also Example 2 in Paragraph 9.5.4-1.

⊙ *References:* G. E. Shilov (1965), L. D. Faddeev (1998).

$$6. \quad \left( \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)^n w = \Phi(x, y), \quad n = 1, 2, \dots$$

This is a *nonhomogeneous polyharmonic equation* of order  $n$  with two independent variables.

Particular solution:

$$w(x, y) = \frac{1}{\pi 2^{2n} [(n-1)!]^2} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \Phi(\xi, \eta) [(x-\xi)^2 + (y-\eta)^2]^{n-1} \ln[(x-\xi)^2 + (y-\eta)^2] d\xi d\eta.$$

The general solution is given by the sum of any particular solution of the nonhomogeneous equation and the general solution of the homogeneous equation (see equation 9.5.5.5, Item 1°).

$$7. \quad \Delta_n^m w = 0, \quad \Delta_n = \sum_{k=1}^n \frac{\partial^2}{\partial x_k^2}.$$

This is the *polyharmonic equation* of order  $m$  with  $n$  independent variables. For  $m = 1$ , see Sections 7.1 and 8.1. For  $m = 2$ , see Subsection 9.4.1. For  $n = 2$ , see equations 9.5.5.5 and 9.5.5.6.

1°. Particular solutions:

$$w(\mathbf{x}) = \sum_{s=0}^{m-1} x_j^s u_s(\mathbf{x}), \quad j = 1, 2, \dots, n,$$

where the  $u_s(\mathbf{x})$  are arbitrary harmonic functions ( $\Delta_n u_s = 0$ ).

2°. Fundamental solution for  $m \geq 1$  and  $n \geq 3$ :

$$\mathcal{E}(\mathbf{x}) = \begin{cases} b_{n,m} |\mathbf{x}|^{2m-n} & \text{if } n \text{ is odd or } n \text{ is even and } n > 2m; \\ c_{n,m} |\mathbf{x}|^{2m-n} \ln |\mathbf{x}| & \text{if } n \text{ is even and } n \leq 2m. \end{cases}$$

Here,

$$b_{n,m} = \frac{\Gamma(n/2)}{2^m(m-1)! \pi^{n/2} (2-n)(4-n) \dots (2m-n)},$$

$$c_{n,m} = \frac{\Gamma(n/2)}{2^m(m-1)! \pi^{n/2} (2-n)(4-n) \dots (2m_0-2-n)(2m_0+2-n)(2m_0+4-n) \dots (2m-n)},$$

where  $m_0 = n/2$ . The expression of the coefficient  $c_{n,m}$  can be obtained formally from the expression of  $b_{n,m}$  by removing the multiplier  $(2m_0 - n)$  equal to zero from the denominator.

⊙ *Reference:* G. E. Shilov (1965).

$$8. \quad \sum_{k=0}^m a_k \Delta^k w = 0, \quad \Delta \equiv \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}.$$

Particular solutions:

$$w(x, y) = \sum_{n=1}^m u_n(x, y),$$

where the  $u_n$  are solutions of the Helmholtz equations  $\Delta u_n - \lambda_n u_n = 0$  and the  $\lambda_n$  are roots of the characteristic equation  $\sum_{k=0}^m a_k \lambda^k = 0$ .

⊙ *Reference:* A. V. Bitsadze and D. F. Kalinichenko (1985).

$$9. \quad \sum_{k=0}^m a_k L^k[w] = 0.$$

Here,  $L$  is any constant coefficient linear differential operator with arbitrarily many independent variables  $x_1, \dots, x_n$ .

Particular solutions:

$$w(x_1, \dots, x_n) = \sum_{s=1}^m C_s u_s(x_1, \dots, x_n),$$

where the  $u_s$  are solution of the equations  $L[u_s] - \lambda_s u_s = 0$  the  $\lambda_s$  are roots of the characteristic equation  $\sum_{k=0}^m a_k \lambda^k = 0$ , and the  $C_s$  are arbitrary constants.

## 9.6. Higher-Order Linear Equations with Variable Coefficients

### 9.6.1. Equations Containing the First Time Derivative

9.6.1-1. Statement of the problem for an equation with two independent variables.

Consider the linear nonhomogeneous partial differential equation

$$\frac{\partial w}{\partial t} - L_{x,t}[w] = \Phi(x, t), \quad (1)$$

where  $L_{x,t}$  is a general linear differential operator of order  $n$  with respect to the space variable  $x$ ,

$$L_{x,t}[w] \equiv \sum_{k=0}^n a_k(x, t) \frac{\partial^k w}{\partial x^k}, \quad (2)$$

whose coefficients  $a_k = a_k(x, t)$  are sufficiently smooth functions of both arguments for  $t \geq 0$  and  $x_1 \leq x \leq x_2$ . The subscripts  $x$  and  $t$  indicate that the operator  $L_{x,t}$  is dependent on the variables  $x$  and  $t$ .

We set the initial condition

$$w = f(x) \quad \text{at} \quad t = 0 \quad (3)$$

and the general nonhomogeneous boundary conditions

$$\begin{aligned} \Gamma_m^{(1)}[w] &\equiv \sum_{k=0}^{n-1} b_{mk}^{(1)}(t) \frac{\partial^k w}{\partial x^k} = g_m^{(1)}(t) \quad \text{at} \quad x = x_1 \quad (m = 1, \dots, s), \\ \Gamma_m^{(2)}[w] &\equiv \sum_{k=0}^{n-1} b_{mk}^{(2)}(t) \frac{\partial^k w}{\partial x^k} = g_m^{(2)}(t) \quad \text{at} \quad x = x_2 \quad (m = s+1, \dots, n), \end{aligned} \quad (4)$$

where  $s \geq 1$  and  $n \geq s+1$ . We assume that both sets of the boundary forms  $\Gamma_m^{(1)}[w]$  ( $m = 1, \dots, s$ ) and  $\Gamma_m^{(2)}[w]$  ( $m = s+1, \dots, n$ ) are linearly independent, which means that for any nonzero  $\psi_m = \psi_m(t)$  the following relations hold:

$$\sum_{m=1}^s \psi_m(t) \Gamma_m^{(1)}[w] \neq 0, \quad \sum_{m=s+1}^n \psi_m(t) \Gamma_m^{(2)}[w] \neq 0.$$

In what follows, we deal with the nonstationary boundary value problem (1), (3), (4).

9.6.1-2. The case of general homogeneous boundary conditions. The Green's function.

The solution of equation (1) with the initial condition (3) and the homogeneous boundary conditions

$$\begin{aligned} \Gamma_m^{(1)}[w] &= 0 \quad \text{at} \quad x = x_1 \quad (m = 1, \dots, s), \\ \Gamma_m^{(2)}[w] &= 0 \quad \text{at} \quad x = x_2 \quad (m = s+1, \dots, n) \end{aligned} \quad (5)$$

can be written as

$$w(x, t) = \int_{x_1}^{x_2} f(y) G(x, y, t, 0) dy + \int_0^t \int_{x_1}^{x_2} \Phi(y, \tau) G(x, y, t, \tau) dy d\tau. \quad (6)$$

Here,  $G(x, y, t, \tau)$  is the Green's function that satisfies, for  $t > \tau \geq 0$ , the homogeneous equation

$$\frac{\partial G}{\partial t} - L_{x,t}[G] = 0 \quad (7)$$

with the special nonhomogeneous initial condition

$$G = \delta(x - y) \quad \text{at} \quad t = \tau \quad (8)$$

and the homogeneous boundary conditions

$$\begin{aligned} \Gamma_m^{(1)}[G] &= 0 \quad \text{at} \quad x = x_1 \quad (m = 1, \dots, s), \\ \Gamma_m^{(2)}[G] &= 0 \quad \text{at} \quad x = x_2 \quad (m = s + 1, \dots, n). \end{aligned} \quad (9)$$

The quantities  $y$  and  $\tau$  appear in problem (7)–(9) as free parameters ( $x_1 \leq y \leq x_2$ ), and  $\delta(x)$  is the Dirac delta function.

It should be emphasized that the Green's function  $G$  is independent of the functions  $\Phi(x, t)$ ,  $f(x)$ ,  $g_m^{(1)}(t)$ , and  $g_m^{(2)}(t)$  that characterize various nonhomogeneities of the boundary value problem. If the coefficients  $a_k$ ,  $b_{mk}^{(1)}$ , and  $b_{mk}^{(2)}$  determining the differential operator (2) and boundary conditions (4) are independent of time  $t$ , then the Green's function depends only on three arguments,  $G(x, y, t, \tau) = G(x, y, t - \tau)$ .

⊙ *Reference:* Mathematical Encyclopedia (1977, Vol. 1).

### 9.6.1-3. The case of nonhomogeneous boundary conditions. Preliminary transformations.

To solve the problem with nonhomogeneous boundary conditions (1), (3), (4), we choose a sufficiently smooth “test function”  $\varphi = \varphi(x, t)$  that satisfies the same boundary conditions as the unknown function; thus,

$$\begin{aligned} \Gamma_m^{(1)}[\varphi] &= g_m^{(1)}(t) \quad \text{at} \quad x = x_1 \quad (m = 1, \dots, s), \\ \Gamma_m^{(2)}[\varphi] &= g_m^{(2)}(t) \quad \text{at} \quad x = x_2 \quad (m = s + 1, \dots, n). \end{aligned} \quad (10)$$

Otherwise the choice of the “test function”  $\varphi$  is arbitrary and is not linked to the solution of the equation in question; there are infinitely many such functions.

Let us pass from  $w = w(x, t)$  to the new unknown  $u = u(x, t)$  by the relation

$$w(x, t) = u(x, t) + \varphi(x, t). \quad (11)$$

Substituting (11) into (1), (3), and (4), we arrive at the problem for an equation with a modified right-hand side,

$$\frac{\partial u}{\partial t} - L_{x,t}[u] = \overline{\Phi}(x, t), \quad \overline{\Phi}(x, t) = \Phi(x, t) - \frac{\partial \varphi}{\partial t} + L_{x,t}[\varphi], \quad (12)$$

subject to the nonhomogeneous initial condition

$$u = f(x) - \varphi(x, 0) \quad \text{at} \quad t = 0 \quad (13)$$

and the homogeneous boundary conditions

$$\begin{aligned} \Gamma_m^{(1)}[u] &= 0 \quad \text{at} \quad x = x_1 \quad (m = 1, \dots, s), \\ \Gamma_m^{(2)}[u] &= 0 \quad \text{at} \quad x = x_2 \quad (m = s + 1, \dots, n). \end{aligned} \quad (14)$$

The solution of problem (12)–(14) can be found using the Green's function by formula (6) in which one should replace  $w$  by  $u$ ,  $\Phi(x, t)$  by  $\overline{\Phi}(x, t)$ , and  $f(x)$  by  $\overline{f}(x) = f(x) - \varphi(x, 0)$ . Taking into account relation (11), for  $w$  we obtain the representation

$$\begin{aligned} w(x, t) &= \int_{x_1}^{x_2} f(y)G(x, y, t, 0) dy + \int_0^t \int_{x_1}^{x_2} \Phi(y, \tau)G(x, y, t, \tau) dy d\tau + \varphi(x, t) \\ &\quad - \int_{x_1}^{x_2} \varphi(y, 0)G(x, y, t, 0) dy - \int_0^t \int_{x_1}^{x_2} \frac{\partial \varphi}{\partial \tau}(y, \tau)G(x, y, t, \tau) dy d\tau \\ &\quad + \int_0^t \int_{x_1}^{x_2} G(x, y, t, \tau)L_{y,\tau}[\varphi(y, \tau)] dy d\tau. \end{aligned} \quad (15)$$

Changing the order of integration and integrating by parts with respect to  $\tau$ , we find, with reference to the initial condition (8) for the Green's function,

$$\int_0^t \frac{\partial \varphi}{\partial \tau} G d\tau = \varphi(y, t)\delta(x - y) - \varphi(y, 0)G(x, y, t, 0) - \int_0^t \varphi(y, \tau) \frac{\partial G}{\partial \tau}(x, y, t, \tau) d\tau. \quad (16)$$

We transform the inner integral of the last term in (15) using the Lagrange–Green formula [see Kamke (1977)] to obtain

$$\begin{aligned} \int_{x_1}^{x_2} G L_{y,\tau}[\varphi] dy &= \int_{x_1}^{x_2} \varphi L_{y,\tau}^*[G] dy + \mathcal{L}[\varphi, G] \Big|_{y=x_1}^{y=x_2}, \\ L_{y,\tau}^*[G] &\equiv \sum_{k=0}^n (-1)^k \frac{\partial^k}{\partial y^k} [a_k(y, \tau)G], \quad \mathcal{L}[\varphi, G] \equiv \sum_{r=0}^{n-1} \sum_{p+q=r} (-1)^p \frac{\partial^q \varphi}{\partial y^q} \frac{\partial^p}{\partial y^p} [a_{r+1}(y, \tau)G], \end{aligned} \quad (17)$$

where  $L_{x,t}^*[w]$  is the differential form adjoint with  $L_{x,t}[w]$  of (2);  $\varphi = \varphi(y, \tau)$ ; and  $p$  and  $q$  are nonnegative integers.

Using relations (16) and (17), we rewrite solution (15) in the form

$$w(x, t) = \int_{x_1}^{x_2} f(y)G(x, y, t, 0) dy + \int_0^t \int_{x_1}^{x_2} \Phi(y, \tau)G(x, y, t, \tau) dy d\tau + \int_0^t \mathcal{L}[\varphi, G] \Big|_{y=x_1}^{y=x_2} d\tau. \quad (18)$$

This formula was derived taking into account the fact that the Green's function with respect to  $y$  and  $\tau$  satisfies the adjoint equation\*

$$\frac{\partial G}{\partial \tau} + L_{y,\tau}^*[G] = 0.$$

For subsequent analysis, it is convenient to represent the bilinear differential form  $\mathcal{L}[\varphi, G]$  as

$$\mathcal{L}[\varphi, G] = \sum_{k=0}^{n-1} \frac{\partial^k \varphi}{\partial y^k} \Psi_k[G], \quad \Psi_k[G] = \sum_{s=0}^{n-k-1} (-1)^s \frac{\partial^s}{\partial y^s} [a_{s+k+1}(y, \tau)G]. \quad (19)$$

Note that in the special case where operator (2) is binomial,

$$L_{x,t}[w] = a_n \frac{\partial^n w}{\partial x^n} + a_0(x, t)w, \quad a_n = \text{const},$$

the differential forms in (19) are written as

$$\mathcal{L}[\varphi, G] = a_n \sum_{k=0}^{n-1} (-1)^{n-k-1} \frac{\partial^k \varphi}{\partial y^k} \frac{\partial^{n-k-1} G}{\partial y^{n-k-1}}, \quad \Psi_k[G] = a_n (-1)^{n-k-1} \frac{\partial^{n-k-1} G}{\partial y^{n-k-1}}.$$

#### 9.6.1-4. The case of special nonhomogeneous boundary conditions.

Consider the following nonhomogeneous boundary conditions of special form that are often encountered in applications:

$$\begin{aligned} \frac{\partial^{k_m} w}{\partial x^{k_m}} &= g_{k_m}^{(1)}(t) \quad \text{at} \quad x = x_1 \quad (m = 1, \dots, s), \\ \frac{\partial^{k_m} w}{\partial x^{k_m}} &= g_{k_m}^{(2)}(t) \quad \text{at} \quad x = x_2 \quad (m = s+1, \dots, n). \end{aligned} \quad (20)$$

Without loss of generality, we assume that the following inequalities hold:

$$n-1 \geq k_1 > k_2 > \dots > k_s, \quad n-1 \geq k_{s+1} > k_{s+2} > \dots > k_n.$$

The Green's function satisfies the corresponding homogeneous boundary conditions that can be obtained from (20) by replacing  $w$  by  $G$  and setting  $g_{k_m}^{(1)}(t) = g_{k_m}^{(2)}(t) = 0$ .

---

\* This equation can be derived by considering the case of homogeneous initial and boundary conditions and using arbitrariness in the choice of the test function  $\varphi = \varphi(x, t)$ ; it should be taken into account that the solution itself must be independent of the specific form of  $\varphi$ , because  $\varphi$  does not occur in the original statement of the problem. By appropriately selecting the test function, one can also derive the boundary conditions (21).

The adjoint homogeneous boundary conditions, with respect to (20), which must be met by the Green's function with respect to  $y$  and  $\tau$  have the form

$$\begin{aligned}\Psi_{k_\beta}[G] &= 0 \quad \text{at } x = x_1 \quad (k_\beta \neq k_m, \beta = s+1, \dots, n; m = 1, \dots, s), \\ \Psi_{k_\beta}[G] &= 0 \quad \text{at } x = x_2 \quad (k_\beta \neq k_m, \beta = 1, \dots, s; m = s+1, \dots, n).\end{aligned}\quad (21)$$

These conditions involve the linear differential forms  $\Psi_k[G]$  defined in (19). For each endpoint of the interval in question, the set  $\{k_\beta\}$  of the indices in the boundary operators (21) together with the set  $\{k_m\}$  of the orders of derivatives in the boundary conditions (20) make up a complete set of nonnegative integers from 0 to  $n-1$ .

Taking into account the fact that the test function  $\varphi$  must satisfy the boundary conditions (20) and the Green's function  $G$  to conditions (21), we rewrite solution (18) to obtain

$$\begin{aligned}w(x, t) &= \int_{x_1}^{x_2} f(y)G(x, y, t, 0) dy + \int_0^t \int_{x_1}^{x_2} \Phi(y, \tau)G(x, y, t, \tau) dy d\tau \\ &\quad - \sum_{m=1}^s \int_0^t g_{k_m}^{(1)}(\tau) \Psi_{k_m}[G]|_{y=x_1} d\tau + \sum_{m=s+1}^n \int_0^t g_{k_m}^{(2)}(\tau) \Psi_{k_m}[G]|_{y=x_2} d\tau,\end{aligned}\quad (22)$$

where the  $\Psi_{k_m}[G]$  are differential operators with respect to  $y$ , which are defined in (19).

If the Green's function is known, formula (22) can be used to immediately obtain the solution of the nonhomogeneous boundary value problem (1), (3), (20) for arbitrary  $\Phi(x, t)$ ,  $f(x)$ ,  $g_{k_m}^{(1)}(t)$  ( $m = 1, \dots, s$ ), and  $g_{k_m}^{(2)}(t)$  ( $m = s+1, \dots, n$ ).

#### 9.6.1-5. The case of general nonhomogeneous boundary conditions.

On solving (4) for the highest derivatives, we reduce the boundary conditions (4) to the canonical form

$$\begin{aligned}\frac{\partial^{k_m} w}{\partial x^{k_m}} + \sum_{i=0}^{k_m-1} c_{mi}^{(1)}(t) \frac{\partial^i w}{\partial x^i} &= h_{k_m}^{(1)}(t) \quad \text{at } x = x_1 \quad (m = 1, \dots, s), \\ \frac{\partial^{k_m} w}{\partial x^{k_m}} + \sum_{i=0}^{k_m-1} c_{mi}^{(2)}(t) \frac{\partial^i w}{\partial x^i} &= h_{k_m}^{(2)}(t) \quad \text{at } x = x_2 \quad (m = s+1, \dots, n),\end{aligned}\quad (23)$$

where the leading terms in different boundary conditions are different,

$$n-1 \geq k_1 > k_2 > \dots > k_s, \quad n-1 \geq k_{s+1} > k_{s+2} > \dots > k_n.$$

The sums in (23) do not contain the derivatives of orders  $k_1, \dots, k_s$  (for  $x = x_1$ ) and  $k_{s+1}, \dots, k_n$  (for  $x = x_2$ ); thus,

$$\begin{aligned}c_{mi}^{(1)}(t) &= 0 \quad \text{at } i = k_j \quad (j = 1, \dots, s), \\ c_{mi}^{(2)}(t) &= 0 \quad \text{at } i = k_j \quad (j = s+1, \dots, n).\end{aligned}$$

It can be shown that the solution of problem (1), (3), (23) is given by

$$\begin{aligned}w(x, t) &= \int_{x_1}^{x_2} f(y)G(x, y, t, 0) dy + \int_0^t \int_{x_1}^{x_2} \Phi(y, \tau)G(x, y, t, \tau) dy d\tau \\ &\quad - \sum_{m=1}^s \int_0^t h_{k_m}^{(1)}(\tau) \Psi_{k_m}[G]|_{y=x_1} d\tau + \sum_{m=s+1}^n \int_0^t h_{k_m}^{(2)}(\tau) \Psi_{k_m}[G]|_{y=x_2} d\tau,\end{aligned}\quad (24)$$

where the  $\Psi_{k_m}[G]$  are differential operators with respect to  $y$ , which are defined in (19). Relation (24) is similar to (22) but contains the Green's function satisfying the more complicated boundary conditions that can be obtained from (23) by substituting  $G$  for  $w$  and setting  $h_{k_m}^{(1)}(t) = h_{k_m}^{(2)}(t) = 0$ .

## 9.6.2. Equations Containing the Second Time Derivative

### 9.6.2-1. The case of homogeneous initial and boundary conditions.

Consider the linear nonhomogeneous differential equation

$$\frac{\partial^2 w}{\partial t^2} + \psi(x, t) \frac{\partial w}{\partial t} - \sum_{k=0}^n a_k(x, t) \frac{\partial^k w}{\partial x^k} = \Phi(x, t). \quad (1)$$

We set the homogeneous initial conditions

$$\begin{aligned} w &= 0 & \text{at } t &= 0, \\ \partial_t w &= 0 & \text{at } t &= 0 \end{aligned} \quad (2)$$

and the homogeneous boundary conditions

$$\begin{aligned} \Gamma_m^{(1)}[w] &= 0 & \text{at } x &= x_1 & (m = 1, \dots, s), \\ \Gamma_m^{(2)}[w] &= 0 & \text{at } x &= x_2 & (m = s + 1, \dots, n), \end{aligned} \quad (3)$$

where the boundary operators  $\Gamma_m^{(1)}[w]$  and  $\Gamma_m^{(2)}[w]$  are defined in Paragraph 9.6.1-1.

The solution of problem (1)–(3) can be represented in the form\*

$$w(x, t) = \int_0^t \int_{x_1}^{x_2} \Phi(y, \tau) G(x, y, t, \tau) dy d\tau. \quad (4)$$

Here,  $G = G(x, y, t, \tau)$  is the Green's function; for  $t > \tau \geq 0$ , it satisfies the homogeneous equation

$$\frac{\partial^2 G}{\partial t^2} + \psi(x, t) \frac{\partial G}{\partial t} - \sum_{k=0}^n a_k(x, t) \frac{\partial^k G}{\partial x^k} = 0 \quad (5)$$

with the special semihomogeneous initial conditions

$$\begin{aligned} G &= 0 & \text{at } t &= \tau, \\ \partial_t G &= \delta(x - y) & \text{at } t &= \tau \end{aligned} \quad (6)$$

and the corresponding homogeneous boundary conditions

$$\begin{aligned} \Gamma_m^{(1)}[G] &= 0 & \text{at } x &= x_1 & (m = 1, \dots, s), \\ \Gamma_m^{(2)}[G] &= 0 & \text{at } x &= x_2 & (m = s + 1, \dots, n). \end{aligned} \quad (7)$$

The quantities  $y$  and  $\tau$  appear in problem (5)–(7) as free parameters ( $x_1 \leq y \leq x_2$ ), and  $\delta(x)$  is the Dirac delta function.

One can verify by direct substitution into the equation and the initial and boundary conditions (1)–(3) that formula (4) is correct, taking into account the properties (5)–(7) of the Green's function.

### 9.6.2-2. The case of nonhomogeneous initial and boundary conditions.

Consider the linear nonhomogeneous differential equation (1) with the general nonhomogeneous initial conditions

$$\begin{aligned} w &= f_0(x) & \text{at } t &= 0, \\ \partial_t w &= f_1(x) & \text{at } t &= 0 \end{aligned} \quad (8)$$

---

\* Problem (1)–(3) is assumed to be well posed.



and the nonhomogeneous boundary conditions, reduced to the canonical form (see Paragraph 9.6.1-5):

$$\begin{aligned} \frac{\partial^{k_m} w}{\partial x^{k_m}} + \sum_{i=0}^{k_m-1} c_{mi}^{(1)}(t) \frac{\partial^i w}{\partial x^i} &= h_{k_m}^{(1)}(t) \quad \text{at } x = x_1 \quad (m = 1, \dots, s), \\ \frac{\partial^{k_m} w}{\partial x^{k_m}} + \sum_{i=0}^{k_m-1} c_{mi}^{(2)}(t) \frac{\partial^i w}{\partial x^i} &= h_{k_m}^{(2)}(t) \quad \text{at } x = x_2 \quad (m = s+1, \dots, n). \end{aligned} \quad (9)$$

Introducing a test function  $\varphi = \varphi(x, t)$  that satisfies the nonhomogeneous initial and boundary conditions (8), (9) and using the same line of reasoning as in Paragraph 9.6.1-3 for a simpler equation, we arrive at the solution of problem (1), (8), (9) in the form

$$\begin{aligned} w(x, t) &= \int_0^t \int_{x_1}^{x_2} \Phi(y, \tau) G(x, y, t, \tau) dy d\tau \\ &\quad - \int_{x_1}^{x_2} f_0(y) \frac{\partial}{\partial \tau} \left[ G(x, y, t, \tau) \right]_{\tau=0} dy + \int_{x_1}^{x_2} [f_1(y) + f_0(y)\psi(y, 0)] G(x, y, t, 0) dy \\ &\quad - \sum_{m=1}^s \int_0^t h_{k_m}^{(1)}(\tau) \Psi_{k_m}[G]|_{y=x_1} d\tau + \sum_{m=s+1}^n \int_0^t h_{k_m}^{(2)}(\tau) \Psi_{k_m}[G]|_{y=x_2} d\tau, \end{aligned} \quad (10)$$

where the  $\Psi_{k_m}[G]$  are differential operators with respect to  $y$ , which are defined in relations (19), Paragraph 9.6.1-3.

*Remark.* If the coefficients of equation (1) and those of the boundary conditions (9) are time independent, i.e.,

$$\psi = \psi(x), \quad a_k = a_k(x), \quad c_{mi}^{(1)} = \text{const}, \quad c_{mi}^{(2)} = \text{const},$$

then in solution (10) one should set

$$G(x, y, t, \tau) = \tilde{G}(x, y, t - \tau), \quad \frac{\partial}{\partial \tau} G(x, y, t, \tau) \Big|_{\tau=0} = -\frac{\partial}{\partial t} \tilde{G}(x, y, t).$$

### 9.6.3. Nonstationary Problems with Many Space Variables

#### 9.6.3-1. Equations with the first-order partial derivative with respect to $t$ .

Consider the following linear differential operator with respect to variables  $x_1, \dots, x_n$ :

$$\mathfrak{L}_{\mathbf{x}, t}[w] \equiv \sum A_{k_1, \dots, k_n}(x_1, \dots, x_n, t) \frac{\partial^{k_1 + \dots + k_n} w}{\partial x_1^{k_1} \dots \partial x_n^{k_n}}. \quad (1)$$

The coefficients  $A_{k_1, \dots, k_n}$  of the operator are assumed to be sufficiently smooth functions of  $x_1, \dots, x_n$  and  $t$  (and also bounded if necessary). The coefficients of the highest derivatives are assumed to be everywhere nonzero.

1°. *Cauchy problem* ( $t \geq 0$ ,  $\mathbf{x} \in \mathbb{R}^n$ ). The solution of the Cauchy problem for the linear nonhomogeneous parabolic differential equation with variable coefficients

$$\frac{\partial w}{\partial t} - \mathfrak{L}_{\mathbf{x}, t}[w] = \Phi(\mathbf{x}, t) \quad (2)$$

under the initial conditions

$$w = f(\mathbf{x}) \quad \text{at } t = 0 \quad (3)$$

is given by

$$w(\mathbf{x}, t) = \int_0^t \int_{\mathbb{R}^n} \Phi(\mathbf{y}, \tau) \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau + \int_{\mathbb{R}^n} f(\mathbf{y}) \mathcal{E}(\mathbf{x}, \mathbf{y}, t, 0) d\mathbf{y}, \quad d\mathbf{y} = dy_1 \dots dy_n. \quad (4)$$

Here,  $\mathcal{E} = \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau)$  is the fundamental solution of the Cauchy problem, which satisfies for  $t > \tau \geq 0$  the equation

$$\frac{\partial \mathcal{E}}{\partial t} - \mathfrak{L}_{\mathbf{x}, t}[\mathcal{E}] = 0 \quad (5)$$

and the special initial condition

$$\mathcal{E}|_{t=\tau} = \delta(\mathbf{x} - \mathbf{y}). \quad (6)$$

The quantities  $\mathbf{y}$  and  $\tau$  appear in problem (5), (6) as free parameters ( $\mathbf{y} \in \mathbb{R}^n$ ), and  $\delta(\mathbf{x})$  is the  $n$ -dimensional Dirac delta function.

If the coefficients  $A_{k_1, \dots, k_n}$  of operator (1) are independent of time  $t$ , then the fundamental solution depends on only three arguments,  $\mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) = \mathcal{E}(\mathbf{x}, \mathbf{y}, t - \tau)$ . If the coefficients of operator (1) are constants, then  $\mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) = \mathcal{E}(\mathbf{x} - \mathbf{y}, t - \tau)$ .

2°. *Boundary value problems* ( $t \geq 0, \mathbf{x} \in \mathcal{D}$ ). The solutions of linear boundary value problems in a spatial domain  $\mathcal{D}$  for equation (2) with initial condition (3) and homogeneous boundary conditions for  $\mathbf{x} \in \partial\mathcal{D}$  (these conditions are not written out here) are given by formula (4) in which the domain of integration  $\mathbb{R}^n$  should be replaced by  $\mathcal{D}$ . Here, by  $\mathcal{E}$  we mean the Green's function that must satisfy, apart from equation (5) and the boundary condition (6), the same homogeneous boundary conditions for  $\mathbf{x} \in \partial\mathcal{D}$  as the original equation (2). For boundary value problems, the parameter  $\mathbf{y}$  belongs to the same domain as  $\mathbf{x}$ , i.e.,  $\mathbf{y} \in \mathcal{D}$ .

⊙ *Reference*: Mathematical Encyclopedia (1977, Vol. 1).

### 9.6.3-2. Equations with the second-order partial derivative with respect to $t$ .

1°. *Cauchy problem* ( $t \geq 0, \mathbf{x} \in \mathbb{R}^n$ ). The solution of the Cauchy problem for the linear nonhomogeneous differential equation with variable coefficients

$$\frac{\partial^2 w}{\partial t^2} - \mathfrak{L}_{\mathbf{x}, t}[w] = \Phi(\mathbf{x}, t) \quad (7)$$

under the initial conditions

$$\begin{aligned} w &= f(\mathbf{x}) \quad \text{at} \quad t = 0, \\ \partial_t w &= g(\mathbf{x}) \quad \text{at} \quad t = 0 \end{aligned} \quad (8)$$

is given by

$$\begin{aligned} w(\mathbf{x}, t) &= \int_0^t \int_{\mathbb{R}^n} \Phi(\mathbf{y}, \tau) \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau \\ &\quad - \int_{\mathbb{R}^n} f(\mathbf{y}) \left[ \frac{\partial}{\partial \tau} \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) \right]_{\tau=0} d\mathbf{y} + \int_{\mathbb{R}^n} g(\mathbf{y}) \mathcal{E}(\mathbf{x}, \mathbf{y}, t, 0) d\mathbf{y}. \end{aligned}$$

Here,  $\mathcal{E} = \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau)$  is the fundamental solution of the Cauchy problem

$$\begin{aligned} \frac{\partial^2 \mathcal{E}}{\partial t^2} - \mathfrak{L}_{\mathbf{x}, t}[\mathcal{E}] &= 0, \\ \mathcal{E}|_{t=\tau} &= 0, \quad \frac{\partial \mathcal{E}}{\partial t} \Big|_{t=\tau} = \delta(\mathbf{x} - \mathbf{y}), \end{aligned}$$

where  $\mathbf{y}$  and  $\tau$  play the role of parameters.

If the coefficients  $A_{k_1, \dots, k_n}$  of operator (1) are independent of time  $t$ , then the fundamental solution depends on only three arguments,  $\mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) = \mathcal{E}(\mathbf{x}, \mathbf{y}, t - \tau)$ , and the relation  $\frac{\partial}{\partial \tau} \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) \Big|_{\tau=0} = -\frac{\partial}{\partial t} \mathcal{E}(\mathbf{x}, \mathbf{y}, t)$  holds. If the coefficients of operator (1) are constants, then  $\mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) = \mathcal{E}(\mathbf{x} - \mathbf{y}, t - \tau)$ .

2°. The solution of the Cauchy problem for the more complicated linear nonhomogeneous differential equation with variable coefficients

$$\frac{\partial^2 w}{\partial t^2} + \psi(\mathbf{x}, t) \frac{\partial w}{\partial t} - \mathfrak{L}_{\mathbf{x}, t}[w] = \Phi(\mathbf{x}, t)$$

with initial conditions (8) is expressed as

$$w(\mathbf{x}, t) = \int_0^t \int_{\mathbb{R}^n} \Phi(\mathbf{y}, \tau) \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) d\mathbf{y} d\tau - \int_{\mathbb{R}^n} f(\mathbf{y}) \left[ \frac{\partial}{\partial \tau} \mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau) \right]_{\tau=0} d\mathbf{y} + \int_{\mathbb{R}^n} [g(\mathbf{y}) + \psi(\mathbf{y}, 0)f(\mathbf{y})] \mathcal{E}(\mathbf{x}, \mathbf{y}, t, 0) d\mathbf{y}. \quad (9)$$

Here,  $\mathcal{E}(\mathbf{x}, \mathbf{y}, t, \tau)$  is the corresponding fundamental solution of the Cauchy problem,

$$\begin{aligned} \frac{\partial^2 \mathcal{E}}{\partial t^2} + \psi(\mathbf{x}, t) \frac{\partial \mathcal{E}}{\partial t} - \mathfrak{L}_{\mathbf{x}, t}[\mathcal{E}] &= 0, \\ \mathcal{E}|_{t=\tau} &= 0, \quad \frac{\partial \mathcal{E}}{\partial t} \Big|_{t=\tau} = \delta(\mathbf{x} - \mathbf{y}). \end{aligned}$$

3°. *Boundary value problems* ( $t \geq 0$ ,  $\mathbf{x} \in \mathcal{D}$ ). The solutions of linear boundary value problems in a spatial domain  $\mathcal{D}$  for equation (7) with initial condition (8) and homogeneous boundary conditions for  $\mathbf{x} \in \partial\mathcal{D}$  (these conditions are not written out here) are given by formula (9) in which the domain of integration  $\mathbb{R}^n$  should be replaced by  $\mathcal{D}$ . Here, by  $\mathcal{E}$  we mean the Green's function that must satisfy, apart from equation (7) and the initial conditions (8), the same homogeneous boundary conditions as the original equation (7).

## 9.6.4. Some Special-Type Equations

$$1. \quad \frac{\partial w}{\partial t} = k(t) \frac{\partial^n w}{\partial x^n} + [x f(t) + g(t)] \frac{\partial w}{\partial x} + h(t)w.$$

The transformation

$$w(x, t) = u(z, \tau) \exp \left[ \int h(t) dt \right], \quad z = xF(t) + \int g(t)F(t) dt, \quad \tau = \int k(t)F^n(t) dt,$$

where  $F(t) = \exp \left[ \int f(t) dt \right]$ , leads to the simpler constant coefficient equation

$$\frac{\partial u}{\partial \tau} = \frac{\partial^n u}{\partial z^n}.$$

$$2. \quad \frac{\partial^k w}{\partial t^k} = ax^{2n} \frac{\partial^n w}{\partial x^n}.$$

The transformation  $z = 1/x$ ,  $u = wx^{1-n}$  leads to the constant coefficient equation

$$\frac{\partial^k u}{\partial t^k} = a(-1)^n \frac{\partial^n u}{\partial z^n}.$$

$$3. \quad \frac{\partial^k w}{\partial t^k} = \sum_{m=0}^n a_m x^m \frac{\partial^m w}{\partial x^m}.$$

The change of variable  $z = \ln|x|$  leads to a constant coefficient equation.

$$4. \quad \frac{\partial^k w}{\partial t^k} = (ax^2 + bx + c)^n \frac{\partial^n w}{\partial x^n}.$$

The transformation

$$w(x, t) = u(z, t) |ax^2 + bx + c|^{\frac{n-1}{2}}, \quad z = \int \frac{dx}{ax^2 + bx + c}$$

leads to a constant coefficient equation.

$$5. \quad \left( \frac{\partial}{\partial t} - L_x \right)^n w = 0, \quad n = 1, 2, \dots$$

Here,  $L_x$  is a linear differential operator of any order with respect to the space variable  $x$  whose coefficients can depend on  $x$ .

1°. General solution:

$$w(x, t) = \sum_{k=0}^{n-1} t^k u_k(x, t),$$

where the  $u_k = u_k(x, t)$  are arbitrary functions that satisfy the original equation with  $n = 1$ :  $(\partial_t - L_x)u_k = 0$ .

2°. Fundamental solution:

$$\mathcal{E}_n(x, t) = \frac{t^{n-1}}{(n-1)!} \mathcal{E}_1(x, t),$$

where  $\mathcal{E}_1(x, t)$  is the fundamental solution of the equation with  $n = 1$ .

*Remark.* The linear differential operator  $L_x$  can involve arbitrarily many space variables.

$$6. \quad \left( \frac{\partial^2}{\partial t^2} - L_x \right)^n w = 0, \quad n = 1, 2, \dots$$

Here,  $L_x$  is a linear differential operator of any order with respect to the space variable  $x$  whose coefficients can depend on  $x$ .

1°. General solution:

$$w(x, t) = \sum_{k=0}^{n-1} t^k u_k(x, t),$$

where the  $u_k = u_k(x, t)$  are arbitrary functions that satisfy the original equation with  $n = 1$ :  $(\partial_{tt} - L_x)u_k = 0$ .

2°. Suppose that the Cauchy problem for the special case of the equation with  $n = 1$  is well posed if only one initial condition is set at  $t = 0$ ; this means that the constant coefficient differential operator  $L_x$  is such that the equation with  $n = 1$  is regular with regularity index  $r = 1$ . Then the fundamental solution of the original equation can be found by the formula

$$\mathcal{E}_n(x, t) = \frac{t^{n-1}}{(n-1)!} \mathcal{E}_1(x, t),$$

where  $\mathcal{E}_1(x, t)$  is the fundamental solution for  $n = 1$ .

*Remark.* The linear differential operator  $L_x$  can involve arbitrarily many space variables.

$$7. \quad \sum_{k=0}^m a_k L^k[w] = 0.$$

Here,  $L$  is any linear differential operator with arbitrarily many independent variables  $x_1, \dots, x_n$ .

Particular solutions:

$$w(x_1, \dots, x_n) = \sum_{s=1}^m C_s u_s(x_1, \dots, x_n),$$

where the  $u_s$  are solutions of the equations  $L[u_s] - \lambda_s u_s = 0$ , the  $\lambda_s$  are roots of the characteristic equation  $\sum_{k=0}^m a_k \lambda^k = 0$ , and the  $C_s$  are arbitrary constants.

# REFERENCES

- Abramowitz, M. and Stegun, I. A. (Editors)**, *Handbook of Mathematical Functions with Formulas, Graphs and Mathematical Tables*, National Bureau of Standards Applied Mathematics, Washington, 1964.
- Acrivos, A.**, *A note of the rate of heat or mass transfer from a small sphere freely suspended in linear shear field*, J. Fluid Mech., Vol. 98, No. 2, pp. 299–304, 1980.
- Aksenov, A. V.**, *Linear differential relations between solutions of the equations of Euler–Poisson–Darboux class*, Mechanics of Solids, Vol. 36, No. 1, pp. 11–15, 2001.
- Akulenko, L. D. and Nesterov, S. V.**, *Determination of the frequencies and forms of oscillations of non-uniform distributed systems with boundary conditions of the third kind*, Appl. Math. Mech. (PMM), Vol. 61, No. 4, p. 531–538, 1997.
- Akulenko, L. D. and Nesterov, S. V.**, *Vibration of an nonhomogeneous membrane*, Mechanics of Solids, Vol. 34, No. 6, pp. 112–121, 1999.
- Akulenko, L. D. and Nesterov, S. V.**, *Free vibrations of a homogeneous elliptic membrane*, Mechanics of Solids, Vol. 35, No. 1, pp. 153–162, 2000.
- Akulenko, L. D., Nesterov, S. V., and Popov, A. L.**, *Natural frequencies of an elliptic plate with clamped edge*, Mechanics of Solids, Vol. 36, No. 1, pp. 143–148, 2001.
- Andreev, V. K., Kaptsov, O. V., Pukhnachov, V. V., and Rodionov, A. A.**, *Applications of Group-Theoretical Methods in Hydrodynamics*, Nauka, Moscow, 1994. (English translation: Kluwer, Dordrecht, 1999.)
- Appell, P.**, *Traité de Mécanique Rationnelle, T. 1: Statique. Dynamique du Point (Ed. 6)*, Gauthier-Villars, Paris, 1953.
- Arscott, F.**, *Periodic Differential Equations*, Macmillan (Pergamon), New York, 1964.
- Arscott, F.**, *The Whittaker–Hill equation and the wave equation in paraboloidal coordinates*, Proc. Roy. Soc. Edinburg, Vol. A67, pp. 265–276, 1967.
- Babich, V. M., Kapilevich, M. B., Mikhlin, S. G., et al.**, *Linear Equations of Mathematical Physics* [in Russian], Nauka, Moscow, 1964.
- Batchelor, G. K.**, *Mass transfer from a particle suspended in fluid with a steady linear ambient velocity distribution*, J. Fluid Mech., Vol. 95, No. 2, pp. 369–400, 1979.
- Bateman, H. and Erdélyi, A.**, *Higher Transcendental Functions, Vol. 1 and Vol. 2*, McGraw-Hill, New York, 1953.
- Bateman, H. and Erdélyi, A.**, *Higher Transcendental Functions, Vol. 3*, McGraw-Hill, New York, 1955.
- Bateman, H. and Erdélyi, A.**, *Tables of Integral Transforms, Vol. 1 and Vol. 2*, McGraw-Hill, New York, 1954.
- Belotserkovskii, O. M., and Oparin, A. A.**, *Numerical Experiment in Turbulence*, Nauka, Moscow, 2000.
- Beyer, W. H.**, *CRC Standard Mathematical Tables and Formulae*, CRC Press, Boca Raton, 1991.
- Bitsadze, A. V. and Kalinichenko, D. F.**, *Collection of Problems on Mathematical Physics Equations* [in Russian], Nauka, Moscow, 1985.
- Bolotin V. V. (Editor)**, *Vibration in Engineering: a Handbook. Vol. 1. Vibration of Linear Systems* [in Russian], Mashinostroenie, Moscow, 1978.

- Borzykh, A. A. and Cherepanov, G. P.**, *A plane problem of the theory of convective heat transfer and mass exchange*, PMM [Applied Mathematics and Mechanics], Vol. 42, No. 5, pp. 848–855, 1978.
- Boyer, C.**, *The maximal kinematical invariance group for an arbitrary potential*, Helv. Phys. Acta, Vol. 47, pp. 589–605, 1974.
- Boyer, C.**, *Lie theory and separation of variables for equation  $iU_t + \Delta_2 U - (\alpha/x_1^2 + \beta/x_2^2)U = 0$* , SIAM J. Math. Anal., Vol. 7, pp. 230–263, 1976.
- Bôcher, M.**, *Die Reihenentwicklungen der Potentialtheory*, Leipzig, 1894.
- Brenner, H.**, *Forced convection-heat and mass transfer at small Peclet numbers from particle of arbitrary shape*, Chem. Eng. Sci., Vol. 18, No. 2, pp. 109–122, 1963.
- Brychkov, Yu. A. and Prudnikov, A. P.**, *Integral Transforms of Generalized Functions*, Gordon & Breach Sci. Publ., New York, 1989.
- Budak, B. M., Samarskii, A. A., and Tikhonov, A. N.**, *Collection of Problems on Mathematical Physics* [in Russian], Nauka, Moscow, 1980.
- Burde, G. I.**, *The construction of special explicit solutions of the boundary-layer equations. Steady flows*, Q. J. Mech. Appl. Math., Vol. 47, No. 2, pp. 247–260, 1994.
- Butkov, E.**, *Mathematical Physics*, Addison-Wesley, Reading, Mass., 1968.
- Butkovskiy, A. G.**, *Characteristics of Systems with Distributed Parameters* [in Russian], Nauka, Moscow, 1979.
- Butkovskiy, A. G.**, *Green's Functions and Transfer Functions Handbook*, Halstead Press–John Wiley & Sons, New York, 1982.
- Carslaw, H. S. and Jaeger, J. C.**, *Conduction of Heat in Solids*, Clarendon Press, Oxford, 1984.
- Clarkson, P. A and Kruskal, M. D.**, *New similarity reductions of the Boussinesq equation*, J. Math. Phys., Vol. 30, No. 10, pp. 2201–2213, 1989.
- Colton, D.**, *Partial Differential Equations. An Introduction*, Random House, New York, 1988.
- Courant, R. and Hilbert, D.**, *Methods of Mathematical Physics, Vol. 2*, Wiley–Interscience Publ., New York, 1989.
- Crank, J.**, *The Mathematics of Diffusion*, Clarendon Press, Oxford, 1975.
- Davis, B.**, *Integral Transforms and Their Applications*, Springer-Verlag, New York, 1978.
- Davis, E. J.**, *Exact solutions for a class of heat and mass transfer problems*, Can. J. Chem. Eng., Vol. 51, No. 5, pp. 562–572, 1973.
- Deavours, C. A.**, *An exact solution for the temperature distribution in parallel plate Poiseuille flow*, Trans. ASME, J. Heat Transfer, Vol. 96, No. 4, 1974.
- Dezin, A. A.**, *Partial Differential Equations. An Introduction to a General Theory of Linear Boundary Value Problems*, Springer-Verlag, Berlin-New York, 1987.
- Ditkin, V. A. and Prudnikov, A. P.**, *Integral Transforms and Operational Calculus*, Pergamon Press, New York, 1965.
- Doyle, Ph. W.**, *Separation of variables for scalar evolution equations in one space dimension*, J. Phys. A: Math. Gen., Vol. 29, pp. 7581–7595, 1996.
- Doyle, Ph. W. and Vassiliou, P. J.**, *Separation of variables for the 1-dimensional non-linear diffusion equation*, Int. J. Non-Linear Mech., Vol. 33, No. 2, pp. 315–326, 1998.
- Elrick, D. E.**, *Source functions for diffusion in uniform shear flows*, Australian J. Phys., Vol. 15, No. 3, p. 283–288, 1962.
- Faddeev, L. D. (Editor)**, *Mathematical Physics. Encyclopedia* [in Russian], Bol'shaya Rossiiskaya Entsiklopediya, Moscow, 1998.
- Faminskii, A. V.**, *On mixed problems for the Corteveg–de Vries equation with irregular boundary data*, Doklady Mathematics, Vol. 59, No. 3, pp. 366–367, 1999.

- Farlow, S. J.**, *Partial Differential Equations for Scientists and Engineers*, John Wiley & Sons, New York, 1982.
- Galaktionov, V. A.**, *Invariant subspace and new explicit solutions to evolution equations with quadratic nonlinearities*, Proc. Roy. Soc. Edinburgh, Vol. 125A, No. 2, pp. 225–448, 1995.
- Galaktionov, V. A. and Posashkov, S. A.**, *On new exact solutions of parabolic equations with quadratic nonlinearities*, Zh. Vych. Matem. i Mat. Fiziki, Vol. 29, No. 4, pp. 497–506, 1989.
- Galaktionov, V. A. and Posashkov, S. A.**, *Exact solutions and invariant subspace for nonlinear gradient-diffusion equations*, Zh. Vych. Matem. i Mat. Fiziki, Vol. 34, No. 3, pp. 374–383, 1994.
- Galaktionov, V. A., Posashkov, S. A., and Svirshchevskii, S. R.**, *Generalized separation of variables for differential equations with polynomial right-hand sides*, Dif. Uravneniya, Vol. 31, No. 2, pp. 253–261, 1995.
- Gel'fand, I. M. and Shilov, G. E.**, *Distributions and Operations on Them* [in Russian], Fizmatlit, Moscow, 1959.
- Gradshteyn, I. S. and Ryzhik, I. M.**, *Tables of Integrals, Series, and Products*, Academic Press, Orlando, 2000.
- Graetz, L.**, *Über die Wärmeleitungsfähigkeit von Flüssigkeiten*, Annln. Phys., Bd. 18, S. 79–84, 1883.
- Grundland, A. M. and Infeld, E.**, *A family of nonlinear Klein–Gordon equations and their solutions*, J. Math. Phys., Vol. 33, No. 7, pp. 2498–2503, 1992.
- Guenther, R. B. and Lee, J. W.**, *Partial Differential Equations of Mathematical Physics and Integral Equations*, Dover Publ., Mineola, 1996.
- Gupalo, Yu. P., Polyanin, A. D., and Ryazantsev, Yu. S.**, *Mass Exchange of Reacting Particles with Flow* [in Russian], Nauka, Moscow, 1985.
- Gupalo, Yu. P. and Ryazantsev, Yu. S.**, *Mass and heat transfer from a sphere in a laminar flow*, Chem. Eng. Sci., Vol. 27, pp. 61–68, 1972.
- Haberman, R.**, *Elementary Applied Partial Differential Equations with Fourier Series and Boundary Value Problems*, Prentice-Hall, Englewood Cliffs, 1987.
- Hanna, J. R. and Rowland, J. H.**, *Fourier Series, Transforms, and Boundary Value Problems*, Wiley-Interscience Publ., New York, 1990.
- Happel, J. and Brenner, H.**, *Low Reynolds Number Hydrodynamics*, Prentice-Hall, Englewood Cliffs, 1965.
- Hörmander, L.**, *The Analysis of Linear Partial Differential Operators. II. Differential Operators with Constant Coefficients*, Springer-Verlag, Berlin-New York, 1983.
- Hörmander, L.**, *The Analysis of Linear Partial Differential Operators. I. Distribution Theory and Fourier Analysis*, Springer-Verlag, Berlin, 1990.
- Ibragimov N. H. (Editor)**, *CRC Handbook of Lie Group to Differential Equations, Vol. 1*, CRC Press, Boca Raton, 1994.
- Ignatovich, N. V.**, *Invariant irreducible, partially invariant solutions of stationary boundary layer equations*, Mat. Zametki, Vol. 53, No. 1, pp. 140–143, 1993.
- Ivanov, V. I., and Trubetskov, M. K.**, *Handbook of Conformal Mapping with Computer-Aided Visualization*, Boca Raton, CRC Press, 1994.
- John, F.**, *Partial Differential Equations*, Springer-Verlag, New York, 1982.
- Kalnins, E.**, *On the separation of variables for the Laplace equation in two- and three-dimensional Minkowski space*, SIAM J. Math. Anal., Hung., Vol. 6, pp. 340–373, 1975.
- Kalnins, E. and Miller, W. (Jr.)**, *Lie theory and separation of variables, 5: The equations  $iU_t + U_{xx} = 0$  and  $iU_t + U_{xx} - \frac{c}{x^2}U = 0$* , J. Math. Phys., Vol. 15, pp. 1728–1737, 1974.



- Kalnins, E. and Miller, W. (Jr.),** *Lie theory and separation of variables, 8: Semisubgroup coordinates for  $\Psi_{tt} - \Delta_2 \Psi = 0$* , J. Math. Phys., Vol. 16, pp. 2507–2516, 1975.
- Kalnins, E. and Miller, W. (Jr.),** *Lie theory and separation of variables, 9: Orthogonal R-separable coordinate systems for the wave equation  $\Psi_{tt} - \Delta_2 \Psi = 0$* , J. Math. Phys., Vol. 17, pp. 331–335, 1976.
- Kalnins, E. and Miller, W. (Jr.),** *Lie theory and separation of variables, 10: Nonorthogonal R-separable solutions of the wave equation  $\Psi_{tt} - \Delta_2 \Psi = 0$* , J. Math. Phys., Vol. 17, pp. 356–368, 1976.
- Kamke, E.,** *Differentialgleichungen: Lösungsmethoden und Lösungen, I, Gewöhnliche Differentialgleichungen*, B. G. Teubner, Leipzig, 1977.
- Kamke, E.,** *Differentialgleichungen: Lösungsmethoden und Lösungen, II, Partielle Differentialgleichungen Erster Ordnung für eine gesuchte Funktion*, Akad. Verlagsgesellschaft Geest & Portig, Leipzig, 1965.
- Kanwal, R. P.,** *Generalized Functions. Theory and Technique*, Academic Press, Orlando, 1983.
- Korn, G. A. and Korn, T. M.,** *Mathematical Handbook for Scientists and Engineers*, McGraw-Hill, New York, 1968.
- Koshlyakov, N. S., Gliner, E. B., and Smirnov, M. M.,** *Partial Differential Equations of Mathematical Physics* [in Russian], Vysshaya Shkola, Moscow, 1970.
- Krein, S. G. (Editor),** *Functional Analysis* [in Russian], Nauka, Moscow, 1972.
- Krylov, A. N.,** *Collected Works: III Mathematics, Pt. 2* [in Russian], Izd-vo AN SSSR, Moscow, 1949.
- Lamb, H.,** *Hydrodynamics*, Dover Publ., New York, 1945.
- Lavrent'ev, M. A. and Shabat B. V.,** *Methods of Complex Variable Theory* [in Russian], Nauka, Moscow, 1973.
- Lavrik, V. I. and Savenkov, V. N.,** *Handbook of Conformal Mappings* [in Russian], Naukova Dumka, Kiev, 1970.
- Landau, L. D. and Lifshits, E. M.,** *Quantum Mechanics. Nonrelativistic Theory* [in Russian], Nauka, Moscow, 1974.
- Lebedev, N. N., Skal'skaya, I. P., and Uflyand, Ya. S.,** *Collection of Problems on Mathematical Physics* [in Russian], Gostekhizdat, Moscow, 1955.
- Leis, R.,** *Initial-Boundary Value Problems in Mathematical Physics*, John Wiley & Sons, Chichester, 1986.
- Levich, V. G.,** *Physicochemical Hydrodynamics*, Prentice-Hall, Englewood Cliffs, New Jersey, 1962.
- Levitan, B. M. and Sargsyan, I. S.,** *Sturm–Liouville and Dirac Operators* [in Russian], Nauka, Moscow, 1988.
- Loitsyanskiy, L. G.,** *Mechanics of Liquids and Gases*, Begell House, New York, 1996.
- Lykov, A. V.,** *Theory of Heat Conduction* [in Russian], Vysshaya Shkola, Moscow, 1967.
- Mackie, A. G.,** *Boundary Value Problems*, Scottish Academic Press, Edinburgh, 1989.
- Makarov, A., Smorodinsky, J., Valiev, K., and Winternitz, P.,** *A systematic search for nonrelativistic systems with dynamical symmetries. Part I: The integrals of motion*, Nuovo Cimento, Vol. 52A, pp. 1061–1084, 1967.
- Marchenko, V. A.,** *Sturm–Liouville Operators and Applications*, Birkhauser Verlag, Basel-Boston, 1986.
- Markeev, A. P.,** *Theoretical Mechanics* [in Russian], Nauka, Moscow, 1990.
- Mathematical Encyclopedia* [in Russian], Sovetskaya Entsiklopediya, Moscow, 1977.
- McLachlan, N. W.,** *Theory and Application of Mathieu Functions*, Clarendon Press, Oxford, 1947.

- Meixner, J. and Schäfer, F.**, *Mathieusche Funktionen und Sphäroidfunktionen*, Springer-Verlag, Berlin, 1965.
- Mikhlin, S. G.**, *Variational Methods in Mathematical Physics* [in Russian], Nauka, Moscow, 1970.
- Miles, J. W.**, *Integral Transforms in Applied Mathematics*, Cambridge Univ. Press, Cambridge, 1971.
- Miller, W. (Jr.)**, *Symmetry and Separation of Variables*, Addison-Wesley, London, 1977.
- Miller, J. (Jr.) and Rubel, L. A.**, *Functional separation of variables for Laplace equations in two dimensions*, J. Phys. A, Vol. 26, No. 8, pp. 1901–1913, 1993.
- Moon, P. and Spencer, D.**, *Field Theory Handbook*, Springer-Verlag, Berlin, 1961.
- Morse, P. M. and Feshbach, H.**, *Methods of Theoretical Physics, Vols. 1–2*, McGraw-Hill, New York, 1953.
- Murphy, G.M.**, *Ordinary Differential Equations and Their Solutions*, D. Van Nostrand, New York, 1960.
- Myint-U, T. and Debnath, L.**, *Partial differential equations for scientists and engineers*, North-Holland Publ., New York, 1987.
- Naimark, M. A.**, *Linear Differential Operators* [in Russian], Nauka, Moscow, 1969.
- Niederer, U.**, *The maximal kinematical invariance group of the harmonic oscillator*, Helv. Phys. Acta, Vol. 46, pp. 191–200, 1973.
- Nikiforov, A. F. and Uvarov, V. B.**, *Special Functions of Mathematical Physics. A Unified Introduction with Applications*, Birkhauser Verlag, Basel-Boston, 1988.
- Novikov, E. A.**, *Concerning turbulent diffusion in a stream with a transverse gradient of velocity*, Appl. Math. Mech. (PMM), Vol. 22, No. 3, p. 412–414, 1958.
- Nusselt, W.**, *Abhängigkeit der Wärmeübergangszahl von der Rohrlänge*, VDI Zeitschrift, Bd. 54, No. 28, S. 1154–1158, 1910.
- Olver, P. J.**, *Application of Lie Groups to Differential Equations*, Springer-Verlag, New York, 1986.
- Ovsiannikov, L. V.**, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
- Pavlovskii, Yu. N.**, *Analysis of some invariant solutions of boundary layer equations*, Zh. Bykh. Matem. i Mat. Fiziki, Vol. 1, No. 2, pp. 280–294, 1961.
- Petrovsky, I. G.**, *Lectures on Partial Differential Equations*, Dover Publ., New York, 1991.
- Pinsky, M. A.**, *Introduction to Partial Differential Equations with Applications*, McGraw-Hill, New York, 1984.
- Polozhii, G. N.**, *Mathematical Physics Equations* [in Russian], Vysshaya Shkola, Moscow, 1964.
- Polyanin, A. D.**, *The structure of solutions of linear nonstationary boundary-value problems of mechanics and mathematical physics*, Doklady Physics, Vol. 45, No. 8, pp. 415–418, 2000a.
- Polyanin, A. D.**, *Partial separation of variables in unsteady problems of mechanics and mathematical physics*, Doklady Physics, Vol. 45, No. 12, pp. 680–684, 2000b.
- Polyanin, A. D.**, *Linear problems of heat and mass transfer: general relations and results*, Theor. Found. Chem. Eng., Vol. 34, No. 6, pp. 509–520, 2000c.
- Polyanin, A. D.**, *Handbook of Linear Mathematical Physics Equations* [in Russian], Fizmatlit, Moscow, 2001a.
- Polyanin, A. D.**, *Transformations and exact solutions of boundary layer equations with arbitrary functions*, Doklady AN, Vol. 379, No. 3, 2001b.
- Polyanin, A. D.**, *Exact solutions and transformations of the equations of a stationary laminar boundary layer*, Theor. Found. Chem. Eng., Vol. 35, No. 4, pp. 319–328, 2001c.
- Polyanin, A. D.**, *Generalized separable solutions of Navier–Stokes equations*, Doklady AN, Vol. 380, No. 4, 2001d.

- Polyanin, A. D. and Dilman, V. V.**, *Methods of Modeling Equations and Analogies in Chemical Engineering*, CRC Press, Boca Raton, 1994.
- Polyanin, A. D., Kutepov, A. M., Vyazmin, A. V., and Kazenin, D. A.**, *Hydrodynamics, Mass and Heat Transfer in Chemical Engineering*, Gordon & Breach Sci. Publ., London, 2001.
- Polyanin, A. D. and Manzhairov, A. V.**, *Handbook of Integral Equations*, CRC Press, Boca Raton, 1998.
- Polyanin, A. D., Vyazmin, A. V., Zhurov, A. I., and Kazenin, D. A.**, *Handbook of Exact Solutions of Heat and Mass Transfer Equations* [in Russian], Faktorial, Moscow, 1998.
- Polyanin, A. D. and Zaitsev, V. F.**, *Handbook of Exact Solutions for Ordinary Differential Equations*, CRC Press, Boca Raton, 1995.
- Polyanin, A. D., Zaitsev, V. F., and Moussiaux, A.**, *Handbook of First Order Partial Differential Equations*, Gordon & Breach, London, 2001.
- Polyanin, A. D. and Zhurov, A. I.**, *Exact solutions to nonlinear equations of mechanics and mathematical physics*, Doklady Physics, Vol. 43, No. 6, pp. 381–385, 1998.
- Polyanin, A. D., Zhurov, A. I., and Vyazmin, A. V.**, *Generalized separation of variables in nonlinear heat and mass transfer equations*, J. Non-Equilibrium Thermodynamics, Vol. 25, No. 3/4, pp. 251–267, 2000.
- Prudnikov, A. P., Brychkov, Yu. A., and Marichev, O. I.**, *Integrals and Series, Vol. 1, Elementary Functions*, Gordon & Breach Sci. Publ., New York, 1986.
- Prudnikov, A. P., Brychkov, Yu. A., and Marichev, O. I.**, *Integrals and Series, Vol. 2, Special Functions*, Gordon & Breach Sci. Publ., New York, 1986.
- Pukhnachev, V. V.**, *Group properties of the Navier–Stokes equations in the plane case*, Zh. Prikl. Mekh. i Tekhn. Fiziki, No. 1, pp. 83–90, 1960.
- Rimmer, P. L.**, *Heat transfer from a sphere in a stream of small Reynolds number*, J. Fluid Mech., Vol. 32, No. 1, pp. 1–7, 1968.
- Rotem, Z., and Neilson, J. E.**, *Exact solution for diffusion to flow down an incline*, Can. J. Chem. Engng., Vol. 47, pp. 341–346, 1966.
- Schlichting, H.**, *Boundary Layer Theory*, McGraw-Hill, New York, 1981.
- Sedov, L. I.**, *Plane Problems of Hydrodynamics and Airdynamics* [in Russian], Nauka, Moscow, 1980.
- Shilov, G. E.**, *Mathematical Analysis: A Second Special Course* [in Russian], Nauka, Moscow, 1965.
- Smirnov, V. I.**, *A Course of Higher Mathematics. Vols. 2–3* [in Russian], Nauka, Moscow, 1974.
- Smirnov, M. M.**, *Second Order Partial Differential Equations* [in Russian], Nauka, Moscow, 1964.
- Smirnov, M. M.**, *Problems on Mathematical Physics Equations* [in Russian], Nauka, Moscow, 1975.
- Sneddon, I.**, *Fourier Transformations*, McGraw-Hill, New York, 1951.
- Stakgold, I.**, *Boundary Value Problems of Mathematical Physics. Vols. I, II*, SIAM, Philadelphia, 2000.
- Strauss, W. A.**, *Partial Differential Equations. An Introduction*, John Wiley & Sons, New York, 1992.
- Sutton, W. G. L.**, *On the equation of diffusion in a turbulent medium*, Proc. Poy. Soc., Ser. A, Vol. 138, No. 988, pp. 48–75, 1943.
- Svirshchetskii, S. R.**, *Lie–Bäcklund symmetries of linear ODEs and generalized separation of variables in nonlinear equations*, Phys. Letters A, Vol. 199, pp. 344–348, 1995.
- Taylor, M.**, *Partial Differential Equations, Vol. 3*, Springer-Verlag, New York, 1996.
- Temme, N. M.**, *Special Functions. An Introduction to the Classical Functions of Mathematical Physics*, Wiley-Interscience Publ., New York, 1996.

- Tikhonov, A. N. and Samarskii, A. A.,** *Equations of Mathematical Physics*, Dover Publ., New York, 1990.
- Thomas, H. C.,** *Heterogeneous ion exchange in a flowing system*, J. Amer. Chem. Soc., Vol. 66, pp. 1664–1666, 1944.
- Tomotika, S. and Tamada, K.,** *Studies on two-dimensional transonic flows of compressible fluid, Part I*, Quart. Appl. Math., Vol. 7, p. 381, 1950.
- Urvin, K. and Arscott, F.,** *Theory of the Whittaker–Hill equation*, Proc. Roy. Soc., Vol. A69, pp. 28–44, 1970.
- Vereshchagina, L. I.,** *Group fibering of the spatial nonstationary boundary layer equations*, Vestnik LGU, Vol. 13, No. 3, pp. 82–86, 1973.
- Vladimirov, V. S., Mikhailov, V. P., Vasharin A. A., et al.,** *Collection of Problems on Mathematical Physics Equations* [in Russian], Nauka, Moscow, 1974.
- Vladimirov, V. S.,** *Mathematical Physics Equations* [in Russian], Nauka, Moscow, 1988.
- Vvedensky, D.,** *Partial Differential Equations*, Addison-Wesley, Wokingham, 1993.
- Whittaker, E. T. and Watson, G. N.,** *A Course of Modern Analysis, Vols. 1–2*, Cambridge Univ. Press, Cambridge, 1952.
- Zachmanoglou, E. C. and Thoe, D. W.,** *Introduction to Partial Differential Equations with Applications*, Dover Publ., New York, 1986.
- Zaitsev, V. F. and Polyanin, A. D.,** *Handbook of Partial Differential Equations: Exact Solutions* [in Russian], MP Obrazovaniya, Moscow, 1996.
- Zauderer, E.,** *Partial Differential Equations of Applied Mathematics*, Wiley–Interscience Publ., New York, 1989.
- Zhdanov, R. Z.,** *Separation of variables in the non-linear wave equation*, J. Phys. A, Vol. 27, pp. L291–L297, 1994.
- Zwillinger, D.,** *Handbook of Differential Equations*, Academic Press, San Diego, 1998.

# Supplement A

## Special Functions and Their Properties

---

Throughout Supplement A it is assumed that  $n$  is a positive integer, unless otherwise specified.

### A.1. Some Symbols and Coefficients

#### A.1.1. Factorials

Definitions and some properties:

$$\begin{aligned} 0! &= 1! = 1, & n! &= 1 \cdot 2 \cdot 3 \cdots (n-1)n, & n &= 2, 3, \dots, \\ (2n)!! &= 2 \cdot 4 \cdot 6 \cdots (2n-2)(2n) = 2^n n!, \\ (2n+1)!! &= 1 \cdot 3 \cdot 5 \cdots (2n-1)(2n+1) = \frac{2^{n+1}}{\sqrt{\pi}} \Gamma\left(n + \frac{3}{2}\right), \\ n!! &= \begin{cases} (2k)!! & \text{if } n = 2k, \\ (2k+1)!! & \text{if } n = 2k+1, \end{cases} & 0!! &= 1. \end{aligned}$$

#### A.1.2. Binomial Coefficients

Definition:

$$\begin{aligned} C_n^k &= \frac{n!}{k!(n-k)!}, & \text{where } k &= 1, \dots, n, \\ C_a^k &= (-1)^k \frac{(-a)_k}{k!} = \frac{a(a-1) \cdots (a-k+1)}{k!}, & \text{where } k &= 1, 2, \dots \end{aligned}$$

General case:

$$C_a^b = \frac{\Gamma(a+1)}{\Gamma(b+1)\Gamma(a-b+1)}, \quad \text{where } \Gamma(x) \text{ is the gamma function.}$$

Properties:

$$\begin{aligned} C_a^0 &= 1, & C_n^k &= 0 \quad \text{for } k = -1, -2, \dots \text{ or } k > n, \\ C_a^{b+1} &= \frac{a}{b+1} C_{a-1}^b = \frac{a-b}{b+1} C_a^b, & C_a^b + C_a^{b+1} &= C_{a+1}^{b+1}, \\ C_{-1/2}^n &= \frac{(-1)^n}{2^{2n}} C_{2n}^n = (-1)^n \frac{(2n-1)!!}{(2n)!!}, \\ C_{1/2}^n &= \frac{(-1)^{n-1}}{n 2^{2n-1}} C_{2n-2}^{n-1} = \frac{(-1)^{n-1}}{n} \frac{(2n-3)!!}{(2n-2)!!}, \\ C_{n+1/2}^{2n+1} &= (-1)^n 2^{-4n-1} C_{2n}^n, & C_{2n+1/2}^n &= 2^{-2n} C_{4n+1}^{2n}, \\ C_n^{1/2} &= \frac{2^{2n+1}}{\pi C_{2n}^n}, & C_n^{n/2} &= \frac{2^{2n}}{\pi} C_n^{(n-1)/2}. \end{aligned}$$

### A.1.3. Pochhammer Symbol

Definition and some properties ( $k = 1, 2, \dots$ ):

$$\begin{aligned}(a)_n &= a(a+1) \dots (a+n-1) = \frac{\Gamma(a+n)}{\Gamma(a)} = (-1)^n \frac{\Gamma(1-a)}{\Gamma(1-a-n)}, \\(a)_0 &= 1, \quad (a)_{n+k} = (a)_n (a+n)_k, \quad (n)_k = \frac{(n+k-1)!}{(n-1)!}, \\(a)_{-n} &= \frac{\Gamma(a-n)}{\Gamma(a)} = \frac{(-1)^n}{(1-a)_n}, \quad \text{where } a \neq 1, \dots, n; \\(1)_n &= n!, \quad (1/2)_n = 2^{-2n} \frac{(2n)!}{n!}, \quad (3/2)_n = 2^{-2n} \frac{(2n+1)!}{n!}, \\(a+mk)_{nk} &= \frac{(a)_{mk+nk}}{(a)_{mk}}, \quad (a+n)_n = \frac{(a)_{2n}}{(a)_n}, \quad (a+n)_k = \frac{(a)_k (a+k)_n}{(a)_n}.\end{aligned}$$

### A.1.4. Bernoulli Numbers

Definition:

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}.$$

The numbers:

$$\begin{aligned}B_0 &= 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_4 = -\frac{1}{30}, \quad B_6 = \frac{1}{42}, \quad B_8 = -\frac{1}{30}, \quad B_{10} = \frac{5}{66}, \quad \dots, \\B_{2m+1} &= 0 \quad \text{for } m = 1, 2, \dots\end{aligned}$$

## A.2. Error Functions and Exponential Integral

### A.2.1. Error Function and Complementary Error Function

Definitions:

$$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_0^x \exp(-t^2) dt, \quad \operatorname{erfc} x = 1 - \operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_x^{\infty} \exp(-t^2) dt.$$

Expansion of  $\operatorname{erf} x$  into series in powers of  $x$  as  $x \rightarrow 0$ :

$$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \sum_{k=0}^{\infty} (-1)^k \frac{x^{2k+1}}{(k)!(2k+1)} = \frac{2}{\sqrt{\pi}} \exp(-x^2) \sum_{k=0}^{\infty} \frac{2^k x^{2k+1}}{2k+1}!!.$$

Asymptotic expansion of  $\operatorname{erfc} x$  as  $x \rightarrow \infty$ :

$$\operatorname{erfc} x = \frac{1}{\sqrt{\pi}} \exp(-x^2) \left[ \sum_{m=0}^{M-1} (-1)^m \frac{\left(\frac{1}{2}\right)_m}{x^{2m+1}} + O(|x|^{-2M-1}) \right], \quad M = 1, 2, \dots$$

### A.2.2. Exponential Integral

Definition:

$$\begin{aligned}\operatorname{Ei}(x) &= \int_{-\infty}^x \frac{e^t}{t} dt \quad \text{for } x < 0, \\ \operatorname{Ei}(x) &= \lim_{\varepsilon \rightarrow +0} \left( \int_{-\infty}^{-\varepsilon} \frac{e^t}{t} dt + \int_{\varepsilon}^x \frac{e^t}{t} dt \right) \quad \text{for } x > 0.\end{aligned}$$

Other integral representations:

$$\begin{aligned}\operatorname{Ei}(-x) &= -e^{-x} \int_0^{\infty} \frac{x \sin t + t \cos t}{x^2 + t^2} dt & \text{for } x > 0, \\ \operatorname{Ei}(-x) &= e^{-x} \int_0^{\infty} \frac{x \sin t - t \cos t}{x^2 + t^2} dt & \text{for } x < 0, \\ \operatorname{Ei}(-x) &= -x \int_1^{\infty} e^{-xt} \ln t dt & \text{for } x > 0.\end{aligned}$$

Expansion into series in powers of  $x$  as  $x \rightarrow 0$ :

$$\operatorname{Ei}(x) = \begin{cases} \mathcal{C} + \ln(-x) + \sum_{k=1}^{\infty} \frac{x^k}{k \cdot k!} & \text{if } x < 0, \\ \mathcal{C} + \ln x + \sum_{k=1}^{\infty} \frac{x^k}{k \cdot k!} & \text{if } x > 0, \end{cases}$$

where  $\mathcal{C} = 0.5772 \dots$  is the Euler constant.

Asymptotic expansion as  $x \rightarrow \infty$ :

$$\operatorname{Ei}(-x) = e^{-x} \sum_{k=1}^n (-1)^k \frac{(k-1)!}{x^k} + R_n, \quad R_n < \frac{n!}{x^n}.$$

### A.2.3. Logarithmic Integral

Definition:

$$\operatorname{li}(x) = \begin{cases} \int_0^x \frac{dt}{\ln t} = \operatorname{Ei}(\ln x) & \text{if } 0 < x < 1, \\ \lim_{\varepsilon \rightarrow +0} \left( \int_0^{1-\varepsilon} \frac{dt}{\ln t} + \int_{1+\varepsilon}^x \frac{dt}{\ln t} \right) & \text{if } x > 1. \end{cases}$$

For small  $x$ ,

$$\operatorname{li}(x) \approx \frac{x}{\ln(1/x)}.$$

Asymptotic expansion as  $x \rightarrow 1$ :

$$\operatorname{li}(x) = \mathcal{C} + \ln |\ln x| + \sum_{k=1}^{\infty} \frac{\ln^k x}{k \cdot k!}.$$

## A.3. Sine Integral and Cosine Integral. Fresnel Integrals

### A.3.1. Sine Integral

Definition:

$$\operatorname{Si}(x) = \int_0^x \frac{\sin t}{t} dt, \quad \operatorname{si}(x) = - \int_x^{\infty} \frac{\sin t}{t} dt = \operatorname{Si}(x) - \frac{\pi}{2}.$$

Specific values:

$$\operatorname{Si}(0) = 0, \quad \operatorname{Si}(\infty) = \frac{\pi}{2}, \quad \operatorname{si}(\infty) = 0.$$

Properties:

$$\operatorname{Si}(-x) = -\operatorname{Si}(x), \quad \operatorname{si}(x) + \operatorname{si}(-x) = -\pi, \quad \lim_{x \rightarrow -\infty} \operatorname{si}(x) = -\pi.$$

Expansion into series in powers of  $x$  as  $x \rightarrow 0$ :

$$\text{Si}(x) = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} x^{2k-1}}{(2k-1)(2k-1)!}.$$

Asymptotic expansion as  $x \rightarrow \infty$ :

$$\text{si}(x) = -\cos x \left[ \sum_{m=0}^{M-1} \frac{(-1)^m (2m)!}{x^{2m+1}} + O(|x|^{-2M-1}) \right] + \sin x \left[ \sum_{m=1}^{N-1} \frac{(-1)^m (2m-1)!}{x^{2m}} + O(|x|^{-2N}) \right],$$

where  $M, N = 1, 2, \dots$

### A.3.2. Cosine Integral

Definition:

$$\text{Ci}(x) = -\int_x^{\infty} \frac{\cos t}{t} dt = C + \ln x + \int_0^x \frac{\cos t - 1}{t} dt, \quad C = 0.5772 \dots$$

Expansion into series in powers of  $x$  as  $x \rightarrow 0$ :

$$\text{Ci}(x) = C + \ln x + \sum_{k=1}^{\infty} \frac{(-1)^k x^{2k}}{2k(2k)!}.$$

Asymptotic expansion as  $x \rightarrow \infty$ :

$$\text{Ci}(x) = \cos x \left[ \sum_{m=1}^{M-1} \frac{(-1)^m (2m-1)!}{x^{2m}} + O(|x|^{-2M}) \right] + \sin x \left[ \sum_{m=0}^{N-1} \frac{(-1)^m (2m)!}{x^{2m+1}} + O(|x|^{-2N-1}) \right],$$

where  $M, N = 1, 2, \dots$

### A.3.3. Fresnel Integrals

Definitions:

$$S(x) = \frac{1}{\sqrt{2\pi}} \int_0^x \frac{\sin t}{\sqrt{t}} dt = \sqrt{\frac{2}{\pi}} \int_0^{\sqrt{x}} \sin t^2 dt,$$

$$C(x) = \frac{1}{\sqrt{2\pi}} \int_0^x \frac{\cos t}{\sqrt{t}} dt = \sqrt{\frac{2}{\pi}} \int_0^{\sqrt{x}} \cos t^2 dt.$$

Expansion into series in powers of  $x$  as  $x \rightarrow 0$ :

$$S(x) = \sqrt{\frac{2}{\pi}} x \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{(4k+3)(2k+1)!},$$

$$C(x) = \sqrt{\frac{2}{\pi}} x \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{(4k+1)(2k)!}.$$

Asymptotic expansion as  $x \rightarrow \infty$ :

$$S(x) = \frac{1}{2} - \frac{\cos x}{\sqrt{2\pi x}} P(x) - \frac{\sin x}{\sqrt{2\pi x}} Q(x),$$

$$C(x) = \frac{1}{2} + \frac{\sin x}{\sqrt{2\pi x}} P(x) - \frac{\cos x}{\sqrt{2\pi x}} Q(x),$$

$$P(x) = 1 - \frac{1 \cdot 3}{(2x)^2} + \frac{1 \cdot 3 \cdot 5 \cdot 7}{(2x)^4} - \dots, \quad Q(x) = \frac{1}{2x} - \frac{1 \cdot 3 \cdot 5}{(2x)^3} + \dots$$



## A.4. Gamma and Beta Functions

### A.4.1. Gamma Function

#### A.4.1-1. Definition. Integral representations.

The gamma function,  $\Gamma(z)$ , is an analytic function of the complex argument  $z$  everywhere, except for the points  $z = 0, -1, -2, \dots$

For  $\operatorname{Re} z > 0$ ,

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt.$$

For  $-(n+1) < \operatorname{Re} z < -n$ , where  $n = 0, 1, 2, \dots$ ,

$$\Gamma(z) = \int_0^{\infty} \left[ e^{-t} - \sum_{m=0}^n \frac{(-1)^m}{m!} t^m \right] t^{z-1} dt.$$

#### A.4.1-2. Some formulas.

Euler formula

$$\Gamma(z) = \lim_{n \rightarrow \infty} \frac{n! n^z}{z(z+1) \dots (z+n)} \quad (z \neq 0, -1, -2, \dots).$$

Simplest properties:

$$\Gamma(z+1) = z\Gamma(z), \quad \Gamma(n+1) = n!, \quad \Gamma(1) = \Gamma(2) = 1.$$

Symmetry formulas:

$$\begin{aligned} \Gamma(z)\Gamma(-z) &= -\frac{\pi}{z \sin(\pi z)}, & \Gamma(z)\Gamma(1-z) &= \frac{\pi}{\sin(\pi z)}, \\ \Gamma\left(\frac{1}{2} + z\right)\Gamma\left(\frac{1}{2} - z\right) &= \frac{\pi}{\cos(\pi z)}. \end{aligned}$$

Multiple argument formulas:

$$\begin{aligned} \Gamma(2z) &= \frac{2^{2z-1}}{\sqrt{\pi}} \Gamma(z)\Gamma\left(z + \frac{1}{2}\right), \\ \Gamma(3z) &= \frac{3^{3z-1/2}}{2\pi} \Gamma(z)\Gamma\left(z + \frac{1}{3}\right)\Gamma\left(z + \frac{2}{3}\right), \\ \Gamma(nz) &= (2\pi)^{(1-n)/2} n^{nz-1/2} \prod_{k=0}^{n-1} \Gamma\left(z + \frac{k}{n}\right). \end{aligned}$$

Fractional values of the argument:

$$\begin{aligned} \Gamma\left(\frac{1}{2}\right) &= \sqrt{\pi}, & \Gamma\left(n + \frac{1}{2}\right) &= \frac{\sqrt{\pi}}{2^n} (2n-1)!!, \\ \Gamma\left(-\frac{1}{2}\right) &= -2\sqrt{\pi}, & \Gamma\left(\frac{1}{2} - n\right) &= (-1)^n \frac{2^n \sqrt{\pi}}{(2n-1)!!}. \end{aligned}$$

Asymptotic expansion (Stirling formula):

$$\Gamma(z) = \sqrt{2\pi} e^{-z} z^{z-1/2} \left[ 1 + \frac{1}{12} z^{-1} + \frac{1}{288} z^{-2} + O(z^{-3}) \right] \quad (\arg |z| < \pi).$$

### A.4.1-3. Logarithmic derivative of the gamma function.

Definition:

$$\psi(z) = \frac{d \ln \Gamma(z)}{dz} = \frac{\Gamma'_z(z)}{\Gamma(z)}.$$

Functional relations:

$$\psi(z) - \psi(1+z) = -\frac{1}{z},$$

$$\psi(z) - \psi(1-z) = -\pi \cot(\pi z),$$

$$\psi(z) - \psi(-z) = -\pi \cot(\pi z) - \frac{1}{z},$$

$$\psi\left(\frac{1}{2} + z\right) - \psi\left(\frac{1}{2} - z\right) = \pi \tan(\pi z),$$

$$\psi(mz) = \ln m + \frac{1}{m} \sum_{k=0}^{m-1} \psi\left(z + \frac{k}{m}\right).$$

Integral representations ( $\operatorname{Re} z > 0$ ):

$$\psi(z) = \int_0^\infty [e^{-t} - (1+t)^{-z}] t^{-1} dt,$$

$$\psi(z) = \ln z + \int_0^\infty [t^{-1} - (1-e^{-t})^{-1}] e^{-tz} dt,$$

$$\psi(z) = -\mathcal{C} + \int_0^1 \frac{1-t^{z-1}}{1-t} dt,$$

where  $\mathcal{C} = -\psi(1) = 0.5772 \dots$  is the Euler constant.

Values for integer argument:

$$\psi(1) = -\mathcal{C}, \quad \psi(n) = -\mathcal{C} + \sum_{k=1}^{n-1} k^{-1} \quad (n = 2, 3, \dots)$$

## A.4.2. Beta Function

Definition:

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt,$$

where  $\operatorname{Re} x > 0$  and  $\operatorname{Re} y > 0$ .

Relationship with the gamma function:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}.$$

## A.5. Incomplete Gamma and Beta Functions

### A.5.1. Incomplete Gamma Function

Definitions (integral representations):

$$\gamma(\alpha, x) = \int_0^x e^{-t} t^{\alpha-1} dt, \quad \operatorname{Re} \alpha > 0,$$

$$\Gamma(\alpha, x) = \int_x^\infty e^{-t} t^{\alpha-1} dt = \Gamma(\alpha) - \gamma(\alpha, x).$$

Recurrent formulas:

$$\begin{aligned}\gamma(\alpha + 1, x) &= \alpha \gamma(\alpha, x) - x^\alpha e^{-x}, \\ \Gamma(\alpha + 1, x) &= \alpha \Gamma(\alpha, x) + x^\alpha e^{-x}.\end{aligned}$$

Asymptotic expansions as  $x \rightarrow 0$ :

$$\begin{aligned}\gamma(\alpha, x) &= \sum_{n=0}^{\infty} \frac{(-1)^n x^{\alpha+n}}{n! (\alpha + n)}, \\ \Gamma(\alpha, x) &= \Gamma(\alpha) - \sum_{n=0}^{\infty} \frac{(-1)^n x^{\alpha+n}}{n! (\alpha + n)}.\end{aligned}$$

Asymptotic expansions as  $x \rightarrow \infty$ :

$$\begin{aligned}\gamma(\alpha, x) &= \Gamma(\alpha) - x^{\alpha-1} e^{-x} \left[ \sum_{m=0}^{M-1} \frac{(1-\alpha)_m}{(-x)^m} + O(|x|^{-M}) \right], \\ \Gamma(\alpha, x) &= x^{\alpha-1} e^{-x} \left[ \sum_{m=0}^{M-1} \frac{(1-\alpha)_m}{(-x)^m} + O(|x|^{-M}) \right] \quad \left( -\frac{3}{2}\pi < \arg x < \frac{3}{2}\pi \right).\end{aligned}$$

Integral functions related to the gamma function:

$$\operatorname{erf} x = \frac{1}{\sqrt{\pi}} \gamma\left(\frac{1}{2}, x^2\right), \quad \operatorname{erfc} x = \frac{1}{\sqrt{\pi}} \Gamma\left(\frac{1}{2}, x^2\right), \quad \operatorname{Ei}(-x) = -\Gamma(0, x).$$

## A.5.2. Incomplete Beta Function

Definition:

$$B_x(p, q) = \int_0^1 t^{p-1} (1-t)^{q-1} dt,$$

where  $\operatorname{Re} x > 0$  and  $\operatorname{Re} y > 0$ .

## A.6. Bessel Functions

### A.6.1. Definitions and Basic Formulas

A.6.1-1. The Bessel functions of the first and the second kinds.

The Bessel function of the first kind,  $J_\nu(x)$ , and the Bessel function of the second kind,  $Y_\nu(x)$  (also called the Neumann function), are solutions of the Bessel equation

$$x^2 y''_{xx} + x y'_x + (x^2 - \nu^2) y = 0$$

and are defined by the formulas

$$J_\nu(x) = \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{\nu+2k}}{k! \Gamma(\nu + k + 1)}, \quad Y_\nu(x) = \frac{J_\nu(x) \cos \pi \nu - J_{-\nu}(x)}{\sin \pi \nu}. \quad (1)$$

The formula for  $Y_\nu(x)$  is valid for  $\nu \neq 0, \pm 1, \pm 2, \dots$  (the cases  $\nu \neq 0, \pm 1, \pm 2, \dots$  are discussed in what follows).

The general solution of the Bessel equation has the form  $Z_\nu(x) = C_1 J_\nu(x) + C_2 Y_\nu(x)$  and is called the cylinder function.

**A.6.1-2. Some formulas.**

$$\begin{aligned}
 2\nu Z_\nu(x) &= x[Z_{\nu-1}(x) + Z_{\nu+1}(x)], \\
 \frac{d}{dx} Z_\nu(x) &= \frac{1}{2}[Z_{\nu-1}(x) - Z_{\nu+1}(x)] = \pm \left[ \frac{\nu}{x} Z_\nu(x) - Z_{\nu\pm 1}(x) \right], \\
 \frac{d}{dx} [x^\nu Z_\nu(x)] &= x^\nu Z_{\nu-1}(x), \quad \frac{d}{dx} [x^{-\nu} Z_\nu(x)] = -x^{-\nu} Z_{\nu+1}(x), \\
 \left( \frac{1}{x} \frac{d}{dx} \right)^n [x^\nu J_\nu(x)] &= x^{\nu-n} J_{\nu-n}(x), \quad \left( \frac{1}{x} \frac{d}{dx} \right)^n [x^{-\nu} J_\nu(x)] = (-1)^n x^{-\nu-n} J_{\nu+n}(x), \\
 J_{-n}(x) &= (-1)^n J_n(x), \quad Y_{-n}(x) = (-1)^n Y_n(x), \quad n = 0, 1, 2, \dots
 \end{aligned}$$

**A.6.1-3. The Bessel functions for  $\nu = \pm n \pm \frac{1}{2}$ , where  $n = 0, 1, 2, \dots$  :**

$$\begin{aligned}
 J_{1/2}(x) &= \sqrt{\frac{2}{\pi x}} \sin x, & J_{-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \cos x, \\
 J_{3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left( \frac{1}{x} \sin x - \cos x \right), & J_{-3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left( -\frac{1}{x} \cos x - \sin x \right), \\
 J_{n+1/2}(x) &= \sqrt{\frac{2}{\pi x}} \left[ \sin \left( x - \frac{n\pi}{2} \right) \sum_{k=0}^{[n/2]} \frac{(-1)^k (n+2k)!}{(2k)! (n-2k)! (2x)^{2k}} \right. \\
 &\quad \left. + \cos \left( x - \frac{n\pi}{2} \right) \sum_{k=0}^{[(n-1)/2]} \frac{(-1)^k (n+2k+1)!}{(2k+1)! (n-2k-1)! (2x)^{2k+1}} \right], \\
 J_{-n-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \left[ \cos \left( x + \frac{n\pi}{2} \right) \sum_{k=0}^{[n/2]} \frac{(-1)^k (n+2k)!}{(2k)! (n-2k)! (2x)^{2k}} \right. \\
 &\quad \left. - \sin \left( x + \frac{n\pi}{2} \right) \sum_{k=0}^{[(n-1)/2]} \frac{(-1)^k (n+2k+1)!}{(2k+1)! (n-2k-1)! (2x)^{2k+1}} \right], \\
 Y_{1/2}(x) &= -\sqrt{\frac{2}{\pi x}} \cos x, & Y_{-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \sin x, \\
 Y_{n+1/2}(x) &= (-1)^{n+1} J_{-n-1/2}(x), & Y_{-n-1/2}(x) &= (-1)^n J_{n+1/2}(x).
 \end{aligned}$$

**A.6.1-4. The Bessel functions for  $\nu = \pm n$ , where  $n = 0, 1, 2, \dots$**

Let  $\nu = n$  be an arbitrary integer. The relations

$$J_{-n}(x) = (-1)^n J_n(x), \quad Y_{-n}(x) = (-1)^n Y_n(x)$$

are valid. The function  $J_n(x)$  is given by the first formula in (1) with  $\nu = n$ , and  $Y_n(x)$  can be obtained from the second formula in (1) by proceeding to the limit  $\nu \rightarrow n$ . For nonnegative  $n$ ,  $Y_n(x)$  can be represented in the form

$$Y_n(x) = \frac{2}{\pi} J_n(x) \ln \frac{x}{2} - \frac{1}{\pi} \sum_{k=0}^{n-1} \frac{(n-k-1)!}{k!} \left( \frac{2}{x} \right)^{n-2k} - \frac{1}{\pi} \sum_{k=0}^{\infty} (-1)^k \left( \frac{x}{2} \right)^{n+2k} \frac{\psi(k+1) + \psi(n+k+1)}{k! (n+k)!},$$

where  $\psi(1) = -\mathcal{C}$ ,  $\psi(n) = -\mathcal{C} + \sum_{k=1}^{n-1} k^{-1}$ ,  $\mathcal{C} = 0.5772 \dots$  is the Euler constant,  $\psi(x) = [\ln \Gamma(x)]'_x$  is the logarithmic derivative of the gamma function.

### A.6.1-5. Wronskians and similar formulas:

$$W(J_\nu, J_{-\nu}) = -\frac{2}{\pi x} \sin(\pi\nu), \quad W(J_\nu, Y_\nu) = \frac{2}{\pi x},$$

$$J_\nu(x)J_{-\nu+1}(x) + J_{-\nu}(x)J_{\nu-1}(x) = \frac{2 \sin(\pi\nu)}{\pi x}, \quad J_\nu(x)Y_{\nu+1}(x) - J_{\nu+1}(x)Y_\nu(x) = -\frac{2}{\pi x}.$$

Here, the notation  $W(f, g) = fg' - f'_xg$  is used.

## A.6.2. Integral Representations and Asymptotic Expansions

### A.6.2-1. Integral representations.

The functions  $J_\nu$  and  $Y_\nu$  can be represented in the form of definite integrals (for  $x > 0$ ):

$$\pi J_\nu(x) = \int_0^\pi \cos(x \sin \theta - \nu\theta) d\theta - \sin \pi\nu \int_0^\infty \exp(-x \sinh t - \nu t) dt,$$

$$\pi Y_\nu(x) = \int_0^\pi \sin(x \sin \theta - \nu\theta) d\theta - \int_0^\infty (e^{\nu t} + e^{-\nu t} \cos \pi\nu) e^{-x \sinh t} dt.$$

For  $|\nu| < \frac{1}{2}$ ,  $x > 0$ ,

$$J_\nu(x) = \frac{2^{1+\nu} x^{-\nu}}{\pi^{1/2} \Gamma(\frac{1}{2} - \nu)} \int_1^\infty \frac{\sin(xt) dt}{(t^2 - 1)^{\nu+1/2}},$$

$$Y_\nu(x) = -\frac{2^{1+\nu} x^{-\nu}}{\pi^{1/2} \Gamma(\frac{1}{2} - \nu)} \int_1^\infty \frac{\cos(xt) dt}{(t^2 - 1)^{\nu+1/2}}.$$

For  $\nu > -\frac{1}{2}$ ,

$$J_\nu(x) = \frac{2(x/2)^\nu}{\pi^{1/2} \Gamma(\frac{1}{2} + \nu)} \int_0^{\pi/2} \cos(x \cos t) \sin^{2\nu} t dt \quad (\text{Poisson's formula}).$$

For  $\nu = 0$ ,  $x > 0$ ,

$$J_0(x) = \frac{2}{\pi} \int_0^\infty \sin(x \cosh t) dt, \quad Y_0(x) = -\frac{2}{\pi} \int_0^\infty \cos(x \cosh t) dt.$$

For integer  $\nu = n = 0, 1, 2, \dots$ ,

$$J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(nt - x \sin t) dt \quad (\text{Bessel's formula}),$$

$$J_{2n}(x) = \frac{2}{\pi} \int_0^{\pi/2} \cos(x \sin t) \cos(2nt) dt,$$

$$J_{2n+1}(x) = \frac{2}{\pi} \int_0^{\pi/2} \sin(x \sin t) \sin[(2n+1)t] dt.$$

### A.6.2-2. Integrals with Bessel functions:

$$\int_0^x x^\lambda J_\nu(x) dx = \frac{x^{\lambda+\nu+1}}{2^\nu (\lambda + \nu + 1) \Gamma(\nu + 1)} F\left(\frac{\lambda + \nu + 1}{2}, \frac{\lambda + \nu + 3}{2}, \nu + 1; -\frac{x^2}{4}\right), \quad \operatorname{Re}(\lambda + \nu) > -1,$$

where  $F(a, b, c; x)$  is the hypergeometric series (see Section 10.9 of this supplement),

$$\int_0^x x^\lambda Y_\nu(x) dx = -\frac{\cos(\nu\pi) \Gamma(-\nu)}{2^\nu \pi (\lambda + \nu + 1)} x^{\lambda+\nu+1} F\left(\frac{\lambda + \nu + 1}{2}, \nu + 1, \frac{\lambda + \nu + 3}{2}, -\frac{x^2}{4}\right)$$

$$- \frac{2^\nu \Gamma(\nu)}{\lambda - \nu + 1} x^{\lambda-\nu+1} F\left(\frac{\lambda - \nu + 1}{2}, 1 - \nu, \frac{\lambda - \nu + 3}{2}, -\frac{x^2}{4}\right), \quad \operatorname{Re} \lambda > |\operatorname{Re} \nu| - 1.$$

### A.6.2-3. Asymptotic expansions as $|x| \rightarrow \infty$ :

$$J_\nu(x) = \sqrt{\frac{2}{\pi x}} \left\{ \cos\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[ \sum_{m=0}^{M-1} (-1)^m (\nu, 2m) (2x)^{-2m} + O(|x|^{-2M}) \right] \right. \\ \left. - \sin\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[ \sum_{m=0}^{M-1} (-1)^m (\nu, 2m+1) (2x)^{-2m-1} + O(|x|^{-2M-1}) \right] \right\},$$

$$Y_\nu(x) = \sqrt{\frac{2}{\pi x}} \left\{ \sin\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[ \sum_{m=0}^{M-1} (-1)^m (\nu, 2m) (2x)^{-2m} + O(|x|^{-2M}) \right] \right. \\ \left. + \cos\left(\frac{4x - 2\nu\pi - \pi}{4}\right) \left[ \sum_{m=0}^{M-1} (-1)^m (\nu, 2m+1) (2x)^{-2m-1} + O(|x|^{-2M-1}) \right] \right\},$$

where  $(\nu, m) = \frac{1}{2^{2m} m!} (4\nu^2 - 1)(4\nu^2 - 3^2) \dots [4\nu^2 - (2m-1)^2] = \frac{\Gamma(\frac{1}{2} + \nu + m)}{m! \Gamma(\frac{1}{2} + \nu - m)}.$

For nonnegative integer  $n$  and large  $x$ ,

$$\sqrt{\pi x} J_{2n}(x) = (-1)^n (\cos x + \sin x) + O(x^{-2}),$$

$$\sqrt{\pi x} J_{2n+1}(x) = (-1)^{n+1} (\cos x - \sin x) + O(x^{-2}).$$

### A.6.2-4. Asymptotic for large $\nu$ ( $\nu \rightarrow \infty$ ):

$$J_\nu(x) \simeq \frac{1}{\sqrt{2\pi\nu}} \left(\frac{ex}{2\nu}\right)^\nu, \quad Y_\nu(x) \simeq -\sqrt{\frac{2}{\pi\nu}} \left(\frac{ex}{2\nu}\right)^{-\nu},$$

where  $x$  is fixed,

$$J_\nu(\nu) \simeq \frac{2^{1/3}}{3^{2/3} \Gamma(2/3)} \frac{1}{\nu^{1/3}}, \quad Y_\nu(\nu) \simeq -\frac{2^{1/3}}{3^{1/6} \Gamma(2/3)} \frac{1}{\nu^{1/3}}.$$

## A.6.3. Zeros and Orthogonality Properties of Bessel Functions

### A.6.3-1. Zeros of Bessel functions.

Each of the functions  $J_\nu(x)$  and  $Y_\nu(x)$  has infinitely many real zeros (for real  $\nu$ ). All zeros are simple, except possibly for the point  $x = 0$ .

The zeros  $\gamma_m$  of  $J_0(x)$ , i.e., the roots of the equation  $J_0(\gamma_m) = 0$ , are approximately given by

$$\gamma_m = 2.4 + 3.13(m-1) \quad (m = 1, 2, \dots),$$

with maximum error 0.2%.

### A.6.3-2. Orthogonality properties of Bessel functions.

1°. Let  $\mu = \mu_m$  be positive roots of the Bessel function  $J_\nu(\mu)$ , where  $\nu > -1$  and  $m = 1, 2, 3, \dots$ . Then the set of functions  $J_\nu(\mu_m r/a)$  is orthogonal on the interval  $0 \leq r \leq a$  with weight  $r$ :

$$\int_0^a J_\nu\left(\frac{\mu_m r}{a}\right) J_\nu\left(\frac{\mu_k r}{a}\right) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{1}{2} a^2 [J'_\nu(\mu_m)]^2 = \frac{1}{2} a^2 J_{\nu+1}^2(\mu_m) & \text{if } m = k. \end{cases}$$

2°. Let  $\mu = \mu_m$  be positive zeros of the Bessel function derivative  $J'_\nu(\mu)$ , where  $\nu > -1$  and  $m = 1, 2, 3, \dots$ . Then the set of functions  $J_\nu(\mu_m r/a)$  is orthogonal on the interval  $0 \leq r \leq a$  with weight  $r$ :

$$\int_0^a J_\nu\left(\frac{\mu_m r}{a}\right) J_\nu\left(\frac{\mu_k r}{a}\right) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{1}{2}a^2\left(1 - \frac{\nu^2}{\mu_m^2}\right) J_\nu^2(\mu_m) & \text{if } m = k. \end{cases}$$

3°. Let  $\mu = \mu_m$  be positive roots of the transcendental equation  $\mu J'_\nu(\mu) + s J_\nu(\mu) = 0$ , where  $\nu > -1$  and  $m = 1, 2, 3, \dots$ . Then the set of functions  $J_\nu(\mu_m r/a)$  is orthogonal on the interval  $0 \leq r \leq a$  with weight  $r$ :

$$\int_0^a J_\nu\left(\frac{\mu_m r}{a}\right) J_\nu\left(\frac{\mu_k r}{a}\right) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{1}{2}a^2\left(1 + \frac{s^2 - \nu^2}{\mu_m^2}\right) J_\nu^2(\mu_m) & \text{if } m = k. \end{cases}$$

4°. Let  $\mu = \mu_m$  be positive roots of the transcendental equation

$$J_\nu(\lambda_m b) Y_\nu(\lambda_m a) - J_\nu(\lambda_m a) Y_\nu(\lambda_m b) = 0 \quad (\nu > -1, m = 1, 2, 3, \dots).$$

Then the set of functions

$$Z_\nu(\lambda_m r) = J_\nu(\lambda_m r) Y_\nu(\lambda_m a) - J_\nu(\lambda_m a) Y_\nu(\lambda_m r), \quad m = 1, 2, 3, \dots,$$

satisfying the conditions  $Z_\nu(\lambda_m a) = Z_\nu(\lambda_m b) = 0$  is orthogonal on the interval  $a \leq r \leq b$  with weight  $r$ :

$$\int_a^b Z_\nu(\lambda_m r) Z_\nu(\lambda_k r) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{2}{\pi^2 \lambda_m^2} \frac{J_\nu^2(\lambda_m a) - J_\nu^2(\lambda_m b)}{J_\nu^2(\lambda_m b)} & \text{if } m = k. \end{cases}$$

5°. Let  $\mu = \mu_m$  be positive roots of the transcendental equation

$$J'_\nu(\lambda_m b) Y'_\nu(\lambda_m a) - J'_\nu(\lambda_m a) Y'_\nu(\lambda_m b) = 0 \quad (\nu > -1, m = 1, 2, 3, \dots)$$

Then the set of functions

$$Z_\nu(\lambda_m r) = J_\nu(\lambda_m r) Y'_\nu(\lambda_m a) - J'_\nu(\lambda_m a) Y_\nu(\lambda_m r), \quad m = 1, 2, 3, \dots,$$

satisfying the conditions  $Z'_\nu(\lambda_m a) = Z'_\nu(\lambda_m b) = 0$  is orthogonal on the interval  $a \leq r \leq b$  with weight  $r$ :

$$\int_a^b Z_\nu(\lambda_m r) Z_\nu(\lambda_k r) r dr = \begin{cases} 0 & \text{if } m \neq k, \\ \frac{2}{\pi^2 \lambda_m^2} \left[ \left(1 - \frac{\nu^2}{b^2 \lambda_m^2}\right) \frac{[J'_\nu(\lambda_m a)]^2}{[J'_\nu(\lambda_m b)]^2} - \left(1 - \frac{\nu^2}{a^2 \lambda_m^2}\right) \right] & \text{if } m = k. \end{cases}$$

## A.6.4. Hankel Functions (Bessel Functions of the Third Kind)

The Hankel functions of the first kind and the second kind are related to Bessel functions by

$$H_\nu^{(1)}(z) = J_\nu(z) + i Y_\nu(z), \quad H_\nu^{(2)}(z) = J_\nu(z) - i Y_\nu(z), \quad i^2 = -1.$$

Asymptotics for  $z \rightarrow 0$ :

$$H_0^{(1)}(z) \simeq \frac{2i}{\pi} \ln z, \quad H_\nu^{(1)}(z) \simeq -\frac{i}{\pi} \frac{\Gamma(\nu)}{(z/2)^\nu} \quad (\operatorname{Re} \nu > 0),$$

$$H_0^{(2)}(z) \simeq -\frac{2i}{\pi} \ln z, \quad H_\nu^{(2)}(z) \simeq \frac{i}{\pi} \frac{\Gamma(\nu)}{(z/2)^\nu} \quad (\operatorname{Re} \nu > 0).$$

Asymptotics for  $|z| \rightarrow \infty$ :

$$H_\nu^{(1)}(z) \simeq \sqrt{\frac{2}{\pi z}} \exp\left[i\left(z - \frac{1}{2}\pi\nu - \frac{1}{4}\pi\right)\right] \quad (-\pi < \arg z < 2\pi),$$

$$H_\nu^{(2)}(z) \simeq \sqrt{\frac{2}{\pi z}} \exp\left[-i\left(z - \frac{1}{2}\pi\nu - \frac{1}{4}\pi\right)\right] \quad (-2\pi < \arg z < \pi).$$

## A.7. Modified Bessel Functions

### A.7.1. Definitions. Basic Formulas

A.7.1-1. The modified Bessel functions of the first and the second kinds.

The modified Bessel functions of the first kind,  $I_\nu(x)$ , and the second kind,  $K_\nu(x)$  (also called the Macdonald function), of order  $\nu$  are solutions of the modified Bessel equation

$$x^2 y''_{xx} + x y'_x - (x^2 + \nu^2) y = 0$$

and are defined by the formulas

$$I_\nu(x) = \sum_{k=0}^{\infty} \frac{(x/2)^{2k+\nu}}{k! \Gamma(\nu + k + 1)}, \quad K_\nu(x) = \frac{\pi}{2} \frac{I_{-\nu} - I_\nu}{\sin \pi \nu},$$

(see below for  $K_\nu(x)$  with  $\nu = 0, 1, 2, \dots$ ).

A.7.1-2. Some formulas.

The modified Bessel functions possess the properties

$$\begin{aligned} K_{-\nu}(x) &= K_\nu(x); & I_{-n}(x) &= (-1)^n I_n(x), \quad n = 0, 1, 2, \dots \\ 2\nu I_\nu(x) &= x[I_{\nu-1}(x) - I_{\nu+1}(x)], & 2\nu K_\nu(x) &= -x[K_{\nu-1}(x) - K_{\nu+1}(x)], \\ \frac{d}{dx} I_\nu(x) &= \frac{1}{2}[I_{\nu-1}(x) + I_{\nu+1}(x)], & \frac{d}{dx} K_\nu(x) &= -\frac{1}{2}[K_{\nu-1}(x) + K_{\nu+1}(x)]. \end{aligned}$$

A.7.1-3. Modified Bessel functions for  $\nu = \pm n \pm \frac{1}{2}$ , where  $n = 0, 1, 2, \dots$ :

$$\begin{aligned} I_{1/2}(x) &= \sqrt{\frac{2}{\pi x}} \sinh x, & I_{-1/2}(x) &= \sqrt{\frac{2}{\pi x}} \cosh x, \\ I_{3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left( -\frac{1}{x} \sinh x + \cosh x \right), & I_{-3/2}(x) &= \sqrt{\frac{2}{\pi x}} \left( -\frac{1}{x} \cosh x + \sinh x \right), \\ I_{n+1/2}(x) &= \frac{1}{\sqrt{2\pi x}} \left[ e^x \sum_{k=0}^n \frac{(-1)^k (n+k)!}{k! (n-k)! (2x)^k} - (-1)^n e^{-x} \sum_{k=0}^n \frac{(n+k)!}{k! (n-k)! (2x)^k} \right], \\ I_{-n-1/2}(x) &= \frac{1}{\sqrt{2\pi x}} \left[ e^x \sum_{k=0}^n \frac{(-1)^k (n+k)!}{k! (n-k)! (2x)^k} + (-1)^n e^{-x} \sum_{k=0}^n \frac{(n+k)!}{k! (n-k)! (2x)^k} \right], \\ K_{\pm 1/2}(x) &= \sqrt{\frac{\pi}{2x}} e^{-x}, & K_{\pm 3/2}(x) &= \sqrt{\frac{\pi}{2x}} \left( 1 + \frac{1}{x} \right) e^{-x}, \\ K_{n+1/2}(x) &= K_{-n-1/2}(x) = \sqrt{\frac{\pi}{2x}} e^{-x} \sum_{k=0}^n \frac{(n+k)!}{k! (n-k)! (2x)^k}. \end{aligned}$$

A.7.1-4. Modified Bessel functions for  $\nu = n$ , where  $n = 0, 1, 2, \dots$

If  $\nu = n$  is a nonnegative integer, then

$$\begin{aligned} K_n(x) &= (-1)^{n+1} I_n(x) \ln \frac{x}{2} + \frac{1}{2} \sum_{m=0}^{n-1} (-1)^m \left( \frac{x}{2} \right)^{2m-n} \frac{(n-m-1)!}{m!} \\ &\quad + \frac{1}{2} (-1)^n \sum_{m=0}^{\infty} \left( \frac{x}{2} \right)^{n+2m} \frac{\psi(n+m+1) + \psi(m+1)}{m! (n+m)!}; \quad n = 0, 1, 2, \dots, \end{aligned}$$

where  $\psi(z)$  is the logarithmic derivative of the gamma function; for  $n = 0$ , the first sum is dropped.



### A.7.1-5. Wronskians and similar formulas:

$$W(I_\nu, I_{-\nu}) = -\frac{2}{\pi x} \sin(\pi\nu), \quad W(I_\nu, K_\nu) = -\frac{1}{x},$$

$$I_\nu(x)I_{-\nu+1}(x) - I_{-\nu}(x)I_{\nu-1}(x) = -\frac{2 \sin(\pi\nu)}{\pi x}, \quad I_\nu(x)K_{\nu+1}(x) + I_{\nu+1}(x)K_\nu(x) = \frac{1}{x},$$

where  $W(f, g) = fg'_x - f'_xg$ .

## A.7.2. Integral Representations and Asymptotic Expansions

### A.7.2-1. Integral representations.

The functions  $I_\nu(x)$  and  $K_\nu(x)$  can be represented in terms of definite integrals:

$$I_\nu(x) = \frac{x^\nu}{\pi^{1/2} 2^\nu \Gamma(\nu + \frac{1}{2})} \int_{-1}^1 \exp(-xt) (1-t^2)^{\nu-1/2} dt \quad (x > 0, \nu > -\frac{1}{2}),$$

$$K_\nu(x) = \int_0^\infty \exp(-x \cosh t) \cosh(\nu t) dt \quad (x > 0),$$

$$K_\nu(x) = \frac{1}{\cos(\frac{1}{2}\pi\nu)} \int_0^\infty \cos(x \sinh t) \cosh(\nu t) dt \quad (x > 0, -1 < \nu < 1),$$

$$K_\nu(x) = \frac{1}{\sin(\frac{1}{2}\pi\nu)} \int_0^\infty \sin(x \sinh t) \sinh(\nu t) dt \quad (x > 0, -1 < \nu < 1).$$

For integer  $\nu = n$ ,

$$I_n(x) = \frac{1}{\pi} \int_0^\pi \exp(x \cos t) \cos(nt) dt \quad (n = 0, 1, 2, \dots),$$

$$K_0(x) = \int_0^\infty \cos(x \sinh t) dt = \int_0^\infty \frac{\cos(xt)}{\sqrt{t^2 + 1}} dt \quad (x > 0).$$

### A.7.2-2. Integrals with modified Bessel functions:

$$\int_0^x x^\lambda I_\nu(x) dx = \frac{x^{\lambda+\nu+1}}{2^\nu (\lambda + \nu + 1) \Gamma(\nu + 1)} F\left(\frac{\lambda + \nu + 1}{2}, \frac{\lambda + \nu + 3}{2}, \nu + 1; \frac{x^2}{4}\right), \quad \operatorname{Re}(\lambda + \nu) > -1,$$

where  $F(a, b, c; x)$  is the hypergeometric series (see Section 10.9 of this supplement),

$$\int_0^x x^\lambda K_\nu(x) dx = \frac{2^{\nu-1} \Gamma(\nu)}{\lambda - \nu + 1} x^{\lambda-\nu+1} F\left(\frac{\lambda - \nu + 1}{2}, 1 - \nu, \frac{\lambda - \nu + 3}{2}, \frac{x^2}{4}\right)$$

$$+ \frac{2^{-\nu-1} \Gamma(-\nu)}{\lambda + \nu + 1} x^{\lambda+\nu+1} F\left(\frac{\lambda + \nu + 1}{2}, 1 + \nu, \frac{\lambda + \nu + 3}{2}, \frac{x^2}{4}\right), \quad \operatorname{Re} \lambda > |\operatorname{Re} \nu| - 1.$$

### A.7.2-3. Asymptotic expansions as $x \rightarrow \infty$ :

$$I_\nu(x) = \frac{e^x}{\sqrt{2\pi x}} \left\{ 1 + \sum_{m=1}^M (-1)^m \frac{(4\nu^2 - 1)(4\nu^2 - 3^2) \dots [4\nu^2 - (2m-1)^2]}{m! (8x)^m} \right\},$$

$$K_\nu(x) = \sqrt{\frac{\pi}{2x}} e^{-x} \left\{ 1 + \sum_{m=1}^M \frac{(4\nu^2 - 1)(4\nu^2 - 3^2) \dots [4\nu^2 - (2m-1)^2]}{m! (8x)^m} \right\}.$$

The terms of the order of  $O(x^{-M-1})$  are omitted in the braces.

## A.8. Airy Functions

### A.8.1. Definition and Basic Formulas

A.8.1-1. The Airy functions of the first and the second kinds.

The Airy function of the first kind,  $\text{Ai}(x)$ , and the Airy function of the second kind,  $\text{Bi}(x)$ , are solutions of the Airy equation

$$y''_{xx} - xy = 0$$

and are defined by the formulas

$$\text{Ai}(x) = \frac{1}{\pi} \int_0^{\infty} \cos\left(\frac{1}{3}t^3 + xt\right) dt,$$

$$\text{Bi}(x) = \frac{1}{\pi} \int_0^{\infty} \left[ \exp\left(-\frac{1}{3}t^3 + xt\right) + \sin\left(\frac{1}{3}t^3 + xt\right) \right] dt.$$

Wronskian:  $W\{\text{Ai}(x), \text{Bi}(x)\} = 1/\pi$ .

A.8.1-2. Connection with the Bessel functions and the modified Bessel functions:

$$\text{Ai}(x) = \frac{1}{3}\sqrt{x} [I_{-1/3}(z) - I_{1/3}(z)] = \pi^{-1} \sqrt{\frac{1}{3}x} K_{1/3}(z), \quad z = \frac{2}{3}x^{3/2},$$

$$\text{Ai}(-x) = \frac{1}{3}\sqrt{x} [J_{-1/3}(z) + J_{1/3}(z)],$$

$$\text{Bi}(x) = \sqrt{\frac{1}{3}x} [I_{-1/3}(z) + I_{1/3}(z)],$$

$$\text{Bi}(-x) = \sqrt{\frac{1}{3}x} [J_{-1/3}(z) - J_{1/3}(z)].$$

### A.8.2. Power Series and Asymptotic Expansions

A.8.2-1. Power series expansions as  $x \rightarrow 0$ :

$$\text{Ai}(x) = c_1 f(x) - c_2 g(x),$$

$$\text{Bi}(x) = \sqrt{3} [c_1 f(x) + c_2 g(x)],$$

$$f(x) = 1 + \frac{1}{3!}x^3 + \frac{1 \cdot 4}{6!}x^6 + \frac{1 \cdot 4 \cdot 7}{9!}x^9 + \dots = \sum_{k=0}^{\infty} 3^k \left(\frac{1}{3}\right)_k \frac{x^{3k}}{(3k)!},$$

$$g(x) = x + \frac{2}{4!}x^4 + \frac{2 \cdot 5}{7!}x^7 + \frac{2 \cdot 5 \cdot 8}{10!}x^{10} + \dots = \sum_{k=0}^{\infty} 3^k \left(\frac{2}{3}\right)_k \frac{x^{3k+1}}{(3k+1)!},$$

where  $c_1 = 3^{-2/3}/\Gamma(2/3) \approx 0.3550$  and  $c_2 = 3^{-1/3}/\Gamma(1/3) \approx 0.2588$ .

A.8.2-2. Asymptotic expansions as  $x \rightarrow \infty$ .

For large values of  $x$ , the leading terms of asymptotic expansions of the Airy functions are

$$\text{Ai}(x) \simeq \frac{1}{2}\pi^{-1/2}x^{-1/4}\exp(-z), \quad z = \frac{2}{3}x^{3/2},$$

$$\text{Ai}(-x) \simeq \pi^{-1/2}x^{-1/4}\sin\left(z + \frac{\pi}{4}\right),$$

$$\text{Bi}(x) \simeq \pi^{-1/2}x^{-1/4}\exp(z),$$

$$\text{Bi}(-x) \simeq \pi^{-1/2}x^{-1/4}\cos\left(z + \frac{\pi}{4}\right).$$

⊙ *Reference:* M. Abramowitz and I. Stegun (1964).

TABLE A1  
Special cases of the Kummer function  $\Phi(a, b; z)$

| $a$                 | $b$           | $z$             | $\Phi$                                                                                                | Conventional notation                                                                                                                                                 |
|---------------------|---------------|-----------------|-------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $a$                 | $a$           | $x$             | $e^x$                                                                                                 |                                                                                                                                                                       |
| 1                   | 2             | $2x$            | $\frac{1}{x}e^x \sinh x$                                                                              |                                                                                                                                                                       |
| $a$                 | $a+1$         | $-x$            | $ax^{-a}\gamma(a, x)$                                                                                 | Incomplete gamma function<br>$\gamma(a, x) = \int_0^x e^{-t}t^{a-1} dt$                                                                                               |
| $\frac{1}{2}$       | $\frac{3}{2}$ | $-x^2$          | $\frac{\sqrt{\pi}}{2} \operatorname{erf} x$                                                           | Error function<br>$\operatorname{erf} x = \frac{2}{\sqrt{\pi}} \int_0^x \exp(-t^2) dt$                                                                                |
| $-n$                | $\frac{1}{2}$ | $\frac{x^2}{2}$ | $\frac{n!}{(2n)!} \left(-\frac{1}{2}\right)^{-n} H_{2n}(x)$                                           | Hermite polynomials<br>$H_n = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2}),$<br>$n = 0, 1, 2, \dots$                                                                    |
| $-n$                | $\frac{3}{2}$ | $\frac{x^2}{2}$ | $\frac{n!}{(2n+1)!} \left(-\frac{1}{2}\right)^{-n} H_{2n+1}(x)$                                       |                                                                                                                                                                       |
| $-n$                | $b$           | $x$             | $\frac{n!}{(b)_n} L_n^{(b-1)}(x)$                                                                     | Laguerre polynomials<br>$L_n^{(\alpha)}(x) = \frac{e^x x^{-\alpha}}{n!} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}),$<br>$\alpha = b-1,$<br>$(b)_n = b(b+1) \dots (b+n-1)$ |
| $\nu + \frac{1}{2}$ | $2\nu+1$      | $2x$            | $\Gamma(1+\nu) e^x \left(\frac{x}{2}\right)^{-\nu} I_\nu(x)$                                          | Modified Bessel functions<br>$I_\nu(x)$                                                                                                                               |
| $n+1$               | $2n+2$        | $2x$            | $\Gamma\left(n+\frac{3}{2}\right) e^x \left(\frac{x}{2}\right)^{-n-\frac{1}{2}} I_{n+\frac{1}{2}}(x)$ |                                                                                                                                                                       |

## A.9. Degenerate Hypergeometric Functions

### A.9.1. Definitions and Basic Formulas

A.9.1-1. The degenerate hypergeometric functions  $\Phi(a, b; x)$  and  $\Psi(a, b; x)$ .

The degenerate hypergeometric functions  $\Phi(a, b; x)$  and  $\Psi(a, b; x)$  are solutions of the degenerate hypergeometric equation

$$xy''_{xx} + (b-x)y'_x - ay = 0.$$

In the case  $b \neq 0, -1, -2, -3, \dots$ , the function  $\Phi(a, b; x)$  can be represented as Kummer's series:

$$\Phi(a, b; x) = 1 + \sum_{k=1}^{\infty} \frac{(a)_k}{(b)_k} \frac{x^k}{k!},$$

where  $(a)_k = a(a+1) \dots (a+k-1)$ ,  $(a)_0 = 1$ .

Table A1 presents some special cases where  $\Phi$  can be expressed in terms of simpler functions.

The function  $\Psi(a, b; x)$  is defined as follows:

$$\Psi(a, b; x) = \frac{\Gamma(1-b)}{\Gamma(a-b+1)} \Phi(a, b; x) + \frac{\Gamma(b-1)}{\Gamma(a)} x^{1-b} \Phi(a-b+1, 2-b; x).$$

### A.9.1-2. Kummer transformation and linear relations.

Kummer transformation:

$$\Phi(a, b; x) = e^x \Phi(b - a, b; -x), \quad \Psi(a, b; x) = x^{1-b} \Psi(1 + a - b, 2 - b; x).$$

Linear relations for  $\Phi$ :

$$\begin{aligned} (b-a)\Phi(a-1, b; x) + (2a-b+x)\Phi(a, b; x) - a\Phi(a+1, b; x) &= 0, \\ b(b-1)\Phi(a, b-1; x) - b(b-1+x)\Phi(a, b; x) + (b-a)x\Phi(a, b+1; x) &= 0, \\ (a-b+1)\Phi(a, b; x) - a\Phi(a+1, b; x) + (b-1)\Phi(a, b-1; x) &= 0, \\ b\Phi(a, b; x) - b\Phi(a-1, b; x) - x\Phi(a, b+1; x) &= 0, \\ b(a+x)\Phi(a, b; x) - (b-a)x\Phi(a, b+1; x) - ab\Phi(a+1, b; x) &= 0, \\ (a-1+x)\Phi(a, b; x) + (b-a)\Phi(a-1, b; x) - (b-1)\Phi(a, b-1; x) &= 0. \end{aligned}$$

Linear relations for  $\Psi$ :

$$\begin{aligned} \Psi(a-1, b; x) - (2a-b+x)\Psi(a, b; x) + a(a-b+1)\Psi(a+1, b; x) &= 0, \\ (b-a-1)\Psi(a, b-1; x) - (b-1+x)\Psi(a, b; x) + x\Psi(a, b+1; x) &= 0, \\ \Psi(a, b; x) - a\Psi(a+1, b; x) - \Psi(a, b-1; x) &= 0, \\ (b-a)\Psi(a, b; x) - x\Psi(a, b+1; x) + \Psi(a-1, b; x) &= 0, \\ (a+x)\Psi(a, b; x) + a(b-a-1)\Psi(a+1, b; x) - x\Psi(a, b+1; x) &= 0, \\ (a-1+x)\Psi(a, b; x) - \Psi(a-1, b; x) + (a-c+1)\Psi(a, b-1; x) &= 0. \end{aligned}$$

### A.9.1-3. Differentiation formulas and Wronskian.

Differentiation formulas:

$$\begin{aligned} \frac{d}{dx} \Phi(a, b; x) &= \frac{a}{b} \Phi(a+1, b+1; x), & \frac{d^n}{dx^n} \Phi(a, b; x) &= \frac{(a)_n}{(b)_n} \Phi(a+n, b+n; x), \\ \frac{d}{dx} \Psi(a, b; x) &= -a\Psi(a+1, b+1; x), & \frac{d^n}{dx^n} \Psi(a, b; x) &= (-1)^n (a)_n \Psi(a+n, b+n; x). \end{aligned}$$

Wronskian:

$$W(\Phi, \Psi) = \Phi \Psi'_x - \Phi'_x \Psi = -\frac{\Gamma(b)}{\Gamma(a)} x^{-b} e^x.$$

### A.9.1-4. Degenerate hypergeometric functions for $n = 0, 1, 2, \dots$

$$\begin{aligned} \Psi(a, n+1; x) &= \frac{(-1)^{n-1}}{n! \Gamma(a-n)} \left\{ \Phi(a, n+1; x) \ln x \right. \\ &\quad \left. + \sum_{r=0}^{\infty} \frac{(a)_r}{(n+1)_r} [\psi(a+r) - \psi(1+r) - \psi(1+n+r)] \frac{x^r}{r!} \right\} + \frac{(n-1)!}{\Gamma(a)} \sum_{r=0}^{n-1} \frac{(a-n)_r}{(1-n)_r} \frac{x^{r-n}}{r!}, \end{aligned}$$

where  $n = 0, 1, 2, \dots$  (the last sum is dropped for  $n = 0$ ),  $\psi(z) = [\ln \Gamma(z)]'_z$  is the logarithmic derivative of the gamma function,

$$\psi(1) = -\mathcal{C}, \quad \psi(n) = -\mathcal{C} + \sum_{k=1}^{n-1} k^{-1},$$

where  $\mathcal{C} = 0.5772 \dots$  is the Euler constant.

If  $b < 0$ , then the formula

$$\Psi(a, b; x) = x^{1-b} \Psi(a - b + 1, 2 - b; x)$$

is valid for any  $x$ .

For  $b \neq 0, -1, -2, -3, \dots$ , the general solution of the degenerate hypergeometric equation can be represented in the form

$$y = C_1 \Phi(a, b; x) + C_2 \Psi(a, b; x),$$

and for  $b = 0, -1, -2, -3, \dots$ , in the form

$$y = x^{1-b} [C_1 \Phi(a - b + 1, 2 - b; x) + C_2 \Psi(a - b + 1, 2 - b; x)].$$

## A.9.2. Integral Representations and Asymptotic Expansions

### A.9.2-1. Integral representations:

$$\begin{aligned} \Phi(a, b; x) &= \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 e^{xt} t^{a-1} (1-t)^{b-a-1} dt & (\text{for } b > a > 0), \\ \Psi(a, b; x) &= \frac{1}{\Gamma(a)} \int_0^\infty e^{-xt} t^{a-1} (1+t)^{b-a-1} dt & (\text{for } a > 0, x > 0), \end{aligned}$$

where  $\Gamma(a)$  is the gamma function.

### A.9.2-2. Integrals with degenerate hypergeometric functions:

$$\begin{aligned} \int \Phi(a, b; x) dx &= \frac{b-1}{a-1} \Psi(a-1, b-1; x) + C, \\ \int \Psi(a, b; x) dx &= \frac{1}{1-a} \Psi(a-1, b-1; x) + C, \\ \int x^n \Phi(a, b; x) dx &= n! \sum_{k=1}^{n+1} \frac{(-1)^{k+1} (1-b)_k x^{n-k+1}}{(1-a)_k (n-k+1)!} \Phi(a-k, b-k; x) + C, \\ \int x^n \Psi(a, b; x) dx &= n! \sum_{k=1}^{n+1} \frac{(-1)^{k+1} x^{n-k+1}}{(1-a)_k (n-k+1)!} \Psi(a-k, b-k; x) + C. \end{aligned}$$

### A.9.2-3. Asymptotic expansion as $|x| \rightarrow \infty$ :

$$\begin{aligned} \Phi(a, b; x) &= \frac{\Gamma(b)}{\Gamma(a)} e^x x^{a-b} \left[ \sum_{n=0}^N \frac{(b-a)_n (1-a)_n}{n!} x^{-n} + \varepsilon \right], & x > 0, \\ \Phi(a, b; x) &= \frac{\Gamma(b)}{\Gamma(b-a)} (-x)^{-a} \left[ \sum_{n=0}^N \frac{(a)_n (a-b+1)_n}{n!} (-x)^{-n} + \varepsilon \right], & x < 0, \\ \Psi(a, b; x) &= x^{-a} \left[ \sum_{n=0}^N (-1)^n \frac{(a)_n (a-b+1)_n}{n!} x^{-n} + \varepsilon \right], & -\infty < x < \infty, \end{aligned}$$

where  $\varepsilon = O(x^{-N-1})$ .

## A.10. Hypergeometric Functions

### A.10.1. Definition and Some Formulas

The hypergeometric function  $F(\alpha, \beta, \gamma; x)$  is a solution of the Gaussian hypergeometric equation

$$x(x-1)y''_{xx} + [(\alpha + \beta + 1)x - \gamma]y'_x + \alpha\beta y = 0.$$

For  $\gamma \neq 0, -1, -2, -3, \dots$ , the function  $F(\alpha, \beta, \gamma; x)$  can be expressed in terms of the hypergeometric series:

$$F(\alpha, \beta, \gamma; x) = 1 + \sum_{k=1}^{\infty} \frac{(\alpha)_k(\beta)_k}{(\gamma)_k} \frac{x^k}{k!}, \quad (\alpha)_k = \alpha(\alpha+1)\dots(\alpha+k-1),$$

which certainly converges for  $|x| < 1$ .

Table A2 shows some special cases where  $F$  can be expressed in term of elementary functions.

### A.10.2. Basic Properties and Integral Representations

#### A.10.2-1. Some properties.

The function  $F$  possesses the following properties:

$$F(\alpha, \beta, \gamma; x) = F(\beta, \alpha, \gamma; x),$$

$$F(\alpha, \beta, \gamma; x) = (1-x)^{\gamma-\alpha-\beta} F(\gamma-\alpha, \gamma-\beta, \gamma; x),$$

$$F(\alpha, \beta, \gamma; x) = (1-x)^{-\alpha} F\left(\alpha, \gamma-\beta, \gamma; \frac{x}{x-1}\right),$$

$$\frac{d^n}{dx^n} F(\alpha, \beta, \gamma; x) = \frac{(\alpha)_n(\beta)_n}{(\gamma)_n} F(\alpha+n, \beta+n, \gamma+n; x).$$

If  $\gamma$  is not an integer, then the general solution of the hypergeometric equation can be written in the form

$$y = C_1 F(\alpha, \beta, \gamma; x) + C_2 x^{1-\gamma} F(\alpha-\gamma+1, \beta-\gamma+1, 2-\gamma; x).$$

#### A.10.2-2. Integral representations.

For  $\gamma > \beta > 0$ , the hypergeometric function can be expressed in terms of a definite integral:

$$F(\alpha, \beta, \gamma; x) = \frac{\Gamma(\gamma)}{\Gamma(\beta)\Gamma(\gamma-\beta)} \int_0^1 t^{\beta-1} (1-t)^{\gamma-\beta-1} (1-tx)^{-\alpha} dt,$$

where  $\Gamma(\beta)$  is the gamma function.

See M. Abramowitz and I. Stegun (1964) and H. Bateman and A. Erdélyi (1953, Vol. 1) for more detailed information about hypergeometric functions.

## A.11. Whittaker Functions

The Whittaker functions  $M_{k,\mu}(x)$  and  $W_{k,\mu}(x)$  are linearly independent solutions of the Whittaker equation:

$$y''_{xx} + \left[-\frac{1}{4} + \frac{1}{2}k + \left(\frac{1}{4} - \mu^2\right)x^{-2}\right]y = 0.$$

The Whittaker functions are expressed in terms of degenerate hypergeometric functions as

$$M_{k,\mu}(x) = x^{\mu+1/2} e^{-x/2} \Phi\left(\frac{1}{2} + \mu - k, 1 + 2\mu, x\right),$$

$$W_{k,\mu}(x) = x^{\mu+1/2} e^{-x/2} \Psi\left(\frac{1}{2} + \mu - k, 1 + 2\mu, x\right).$$

TABLE A2

Some special cases where the hypergeometric function  $F(\alpha, \beta, \gamma; z)$   
can be expressed in terms of elementary functions

| $\alpha$      | $\beta$                | $\gamma$            | $z$        | $F$                                                                                                                                                                                    |
|---------------|------------------------|---------------------|------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $-n$          | $\beta$                | $\gamma$            | $x$        | $\sum_{k=0}^n \frac{(-n)_k (\beta)_k}{(\gamma)_k} \frac{x^k}{k!}, \quad \text{where } n = 1, 2, \dots$                                                                                 |
| $-n$          | $\beta$                | $-n-m$              | $x$        | $\sum_{k=0}^n \frac{(-n)_k (\beta)_k}{(-n-m)_k} \frac{x^k}{k!}, \quad \text{where } n = 1, 2, \dots$                                                                                   |
| $\alpha$      | $\beta$                | $\beta$             | $x$        | $(1-x)^{-\alpha}$                                                                                                                                                                      |
| $\alpha$      | $\alpha + \frac{1}{2}$ | $\frac{1}{2}$       | $x^2$      | $\frac{1}{2} [(1+x)^{-2\alpha} + (1-x)^{-2\alpha}]$                                                                                                                                    |
| $\alpha$      | $\alpha + \frac{1}{2}$ | $\frac{3}{2}$       | $x^2$      | $\frac{(1+x)^{-2\alpha} - (1-x)^{-2\alpha}}{2x(1-2\alpha)}$                                                                                                                            |
| $\alpha$      | $-\alpha$              | $\frac{1}{2}$       | $-x^2$     | $\frac{1}{2} [(\sqrt{1+x^2}+x)^{2\alpha} + (\sqrt{1+x^2}-x)^{2\alpha}]$                                                                                                                |
| $\alpha$      | $1-\alpha$             | $\frac{1}{2}$       | $-x^2$     | $\frac{(\sqrt{1+x^2}+x)^{2\alpha-1} + (\sqrt{1+x^2}-x)^{2\alpha-1}}{2\sqrt{1+x^2}}$                                                                                                    |
| $\alpha$      | $\alpha - \frac{1}{2}$ | $2\alpha - 1$       | $x$        | $2^{2\alpha-2} (1+\sqrt{1-x})^{-2\alpha}$                                                                                                                                              |
| $\alpha$      | $1-\alpha$             | $\frac{3}{2}$       | $\sin^2 x$ | $\frac{\sin[(2\alpha-1)x]}{(\alpha-1)\sin(2x)}$                                                                                                                                        |
| $\alpha$      | $2-\alpha$             | $\frac{3}{2}$       | $\sin^2 x$ | $\frac{\sin[(2\alpha-2)x]}{(\alpha-1)\sin(2x)}$                                                                                                                                        |
| $\alpha$      | $1-\alpha$             | $\frac{1}{2}$       | $\sin^2 x$ | $\frac{\cos[(2\alpha-1)x]}{\cos x}$                                                                                                                                                    |
| $\alpha$      | $\alpha + 1$           | $\frac{1}{2}\alpha$ | $x$        | $(1+x)(1-x)^{-\alpha-1}$                                                                                                                                                               |
| $\alpha$      | $\alpha + \frac{1}{2}$ | $2\alpha + 1$       | $x$        | $\left(\frac{1+\sqrt{1-x}}{2}\right)^{-2\alpha}$                                                                                                                                       |
| $\alpha$      | $\alpha + \frac{1}{2}$ | $2\alpha$           | $x$        | $\frac{1}{\sqrt{1-x}} \left(\frac{1+\sqrt{1-x}}{2}\right)^{1-2\alpha}$                                                                                                                 |
| $\frac{1}{2}$ | $\frac{1}{2}$          | $\frac{3}{2}$       | $x^2$      | $\frac{1}{x} \arcsin x$                                                                                                                                                                |
| $\frac{1}{2}$ | $1$                    | $\frac{3}{2}$       | $-x^2$     | $\frac{1}{x} \arctan x$                                                                                                                                                                |
| $1$           | $1$                    | $2$                 | $-x$       | $\frac{1}{x} \ln(x+1)$                                                                                                                                                                 |
| $\frac{1}{2}$ | $1$                    | $\frac{3}{2}$       | $x^2$      | $\frac{1}{2x} \ln \frac{1+x}{1-x}$                                                                                                                                                     |
| $n+1$         | $n+m+1$                | $n+m+l+2$           | $x$        | $\frac{(-1)^m (n+m+l+1)!}{n! l! (n+m)! (m+l)!} \frac{d^{n+m}}{dx^{n+m}} \left\{ (1-x)^{m+l} \frac{d^l F}{dx^l} \right\},$<br>$F = -\frac{\ln(1-x)}{x}, \quad n, m, l = 0, 1, 2, \dots$ |

## A.12. Legendre Polynomials and Legendre Functions

### A.12.1. Definitions. Basic Formulas

The Legendre polynomials  $P_n = P_n(x)$  and the Legendre functions  $Q_n(x)$  are solutions of the equation

$$(1 - x^2)y''_{xx} - 2xy'_x + n(n+1)y = 0.$$

The Legendre polynomials  $P_n(x)$  and the Legendre functions  $Q_n(x)$  are defined by the formulas

$$P_n(x) = \frac{1}{n! 2^n} \frac{d^n}{dx^n} (x^2 - 1)^n, \quad Q_n(x) = \frac{1}{2} P_n(x) \ln \frac{1+x}{1-x} - \sum_{m=1}^n \frac{1}{m} P_{m-1}(x) P_{n-m}(x).$$

The polynomials  $P_n = P_n(x)$  can be calculated recursively using the relations

$$P_0(x) = 1, \quad P_1(x) = x, \quad P_2(x) = \frac{1}{2}(3x^2 - 1), \quad \dots, \quad P_{n+1}(x) = \frac{2n+1}{n+1} x P_n(x) - \frac{n}{n+1} P_{n-1}(x).$$

The first three functions  $Q_n = Q_n(x)$  have the form

$$Q_0(x) = \frac{1}{2} \ln \frac{1+x}{1-x}, \quad Q_1(x) = \frac{x}{2} \ln \frac{1+x}{1-x} - 1, \quad Q_2(x) = \frac{3x^2-1}{4} \ln \frac{1+x}{1-x} - \frac{3}{2}x.$$

The polynomials  $P_n(x)$  have the implicit representation

$$P_n(x) = 2^{-n} \sum_{m=0}^{[n/2]} (-1)^m C_n^m C_{2n-2m}^n x^{n-2m},$$

where  $[A]$  is the integer part of a number  $A$ .

### A.12.2. Zeros of Legendre Polynomials and the Generating Function

All zeros of  $P_n(x)$  are real and lie on the interval  $-1 < x < +1$ ; the functions  $P_n(x)$  form an orthogonal system on the interval  $-1 \leq x \leq +1$ , with

$$\int_{-1}^{+1} P_n(x) P_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{2}{2n+1} & \text{if } n = m. \end{cases}$$

The generating function is

$$\frac{1}{\sqrt{1-2sx+s^2}} = \sum_{n=0}^{\infty} P_n(x) s^n \quad (|s| < 1).$$

### A.12.3. Associated Legendre Functions

The associated Legendre functions  $P_n^m(x)$  of order  $m$  are defined by the formulas

$$P_n^m(x) = (1-x^2)^{m/2} \frac{d^m}{dx^m} P_n(x), \quad n = 1, 2, 3, \dots, \quad m = 0, 1, 2, \dots$$

It is assumed by definition that  $P_n^0(x) = P_n(x)$ .

The functions  $P_n^m(x)$  form an orthogonal system on the interval  $-1 \leq x \leq +1$ , with

$$\int_{-1}^{+1} P_n^m(x) P_k^m(x) dx = \begin{cases} 0 & \text{if } n \neq k, \\ \frac{2}{2n+1} \frac{(n+m)!}{(n-m)!} & \text{if } n = k. \end{cases}$$

The functions  $P_n^m(x)$  (with  $m \neq 0$ ) are orthogonal on the interval  $-1 \leq x \leq +1$  with weight  $(1-x^2)^{-1}$ , that is,

$$\int_{-1}^{+1} \frac{P_n^m(x) P_k^m(x)}{(1-x^2)} dx = \begin{cases} 0 & \text{if } n \neq k, \\ \frac{(n+m)!}{m(n-m)!} & \text{if } n = k. \end{cases}$$



## A.13. Parabolic Cylinder Functions

### A.13.1. Definitions. Basic Formulas

The Weber parabolic cylinder function  $D_\nu(z)$  is a solution of the linear differential equation:

$$y''_{zz} + \left(-\frac{1}{4}z^2 + \nu + \frac{1}{2}\right)y = 0,$$

where the parameter  $\nu$  and the variable  $z$  can assume arbitrary real or complex values. Another linearly independent solution of this equation is the function  $D_{-\nu-1}(iz)$ ; if  $\nu$  is noninteger, then  $D_\nu(-z)$  can also be taken as a linearly independent solution.

The parabolic cylinder functions can be expressed in terms of degenerate hypergeometric functions as

$$D_\nu(z) = 2^{1/2} \exp\left(-\frac{1}{4}z^2\right) \left[ \frac{\Gamma\left(\frac{1}{2}\right)}{\Gamma\left(\frac{1}{2} - \frac{\nu}{2}\right)} \Phi\left(-\frac{\nu}{2}, \frac{1}{2}, \frac{1}{2}z^2\right) + 2^{-1/2} \frac{\Gamma\left(-\frac{1}{2}\right)}{\Gamma\left(-\frac{\nu}{2}\right)} z \Phi\left(\frac{1}{2} - \frac{\nu}{2}, \frac{3}{2}, \frac{1}{2}z^2\right) \right].$$

For nonnegative integer  $\nu = n$ , we have

$$D_n(z) = 2^{-n/2} \exp\left(-\frac{1}{4}z^2\right) H_n\left(2^{-1/2}z\right), \quad n = 0, 1, 2, \dots;$$

$$H_n(z) = (-1)^n \exp(z^2) \frac{d^n}{dz^n} \exp(-z^2),$$

where  $H_n(z)$  is the Hermitean polynomial of order  $n$ .

### A.13.2. Integral Representations and Asymptotic Expansions

Integral representations:

$$D_\nu(z) = \sqrt{2/\pi} \exp\left(\frac{1}{4}z^2\right) \int_0^\infty t^\nu \exp\left(-\frac{1}{2}t^2\right) \cos\left(z t - \frac{1}{2}\pi\nu\right) dt \quad \text{for } \operatorname{Re} \nu > -1,$$

$$D_\nu(z) = \frac{1}{\Gamma(-\nu)} \exp\left(-\frac{1}{4}z^2\right) \int_0^\infty t^{-\nu-1} \exp\left(-zt - \frac{1}{2}t^2\right) dt \quad \text{for } \operatorname{Re} \nu < 0.$$

Asymptotic expansion as  $|z| \rightarrow \infty$ :

$$D_\nu(z) = z^\nu \exp\left(-\frac{1}{4}z^2\right) \left[ \sum_{n=0}^N \frac{(-2)^n \left(-\frac{\nu}{2}\right)_n \left(\frac{1}{2} - \frac{\nu}{2}\right)_n}{n!} \frac{1}{z^{2n}} + O(|z|^{-2N-2}) \right] \quad \text{for } |\arg z| < \frac{3\pi}{4},$$

where  $(a)_0 = 1$ ,  $(a)_n = a(a+1) \dots (a+n-1)$  for  $n = 1, 2, 3, \dots$

## A.14. Mathieu Functions

### A.14.1. Definitions and Basic Formulas

#### A.14.1-1. Mathieu equation and Mathieu functions.

The Mathieu functions  $\operatorname{ce}_n(x, q)$  and  $\operatorname{se}_n(x, q)$  are periodical solutions of the Mathieu equation

$$y''_{xx} + (a - 2q \cos 2x)y = 0.$$

Such solutions exist for definite values of parameters  $a$  and  $q$  (those values of  $a$  are referred to as eigenvalues). The Mathieu functions are listed in [Table A3](#).

TABLE A3

The Mathieu functions  $\text{ce}_n = \text{ce}_n(x, q)$  and  $\text{se}_n = \text{se}_n(x, q)$  (for odd  $n$ , functions  $\text{ce}_n$  and  $\text{se}_n$  are  $2\pi$ -periodical, and for even  $n$ , they are  $\pi$ -periodical); definite eigenvalues  $a = a_n(q)$  and  $a = b_n(q)$  correspond to each value of parameter  $q$ .

| Mathieu functions                                                                 | Recurrence relations for coefficients                                                                                                                           | Normalization conditions                                                                                                            |
|-----------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|
| $\text{ce}_{2n} = \sum_{m=0}^{\infty} A_{2m}^{2n} \cos 2mx$                       | $qA_2^{2n} = a_{2n}A_0^{2n};$<br>$qA_4^{2n} = (a_{2n}-4)A_2^{2n} - 2qA_0^{2n};$<br>$qA_{2m+2}^{2n} = (a_{2n}-4m^2)A_{2m}^{2n} - qA_{2m-2}^{2n}, \quad m \geq 2$ | $(A_0^{2n})^2 + \sum_{m=0}^{\infty} (A_{2m}^{2n})^2$<br>$= \begin{cases} 2 & \text{if } n=0 \\ 1 & \text{if } n \geq 1 \end{cases}$ |
| $\text{ce}_{2n+1} = \sum_{m=0}^{\infty} A_{2m+1}^{2n+1} \cos(2m+1)x$              | $qA_3^{2n+1} = (a_{2n+1}-1-q)A_1^{2n+1};$<br>$qA_{2m+3}^{2n+1} = [a_{2n+1}-(2m+1)^2]A_{2m+1}^{2n+1} - qA_{2m-1}^{2n+1}, \quad m \geq 1$                         | $\sum_{m=0}^{\infty} (A_{2m+1}^{2n+1})^2 = 1$                                                                                       |
| $\text{se}_{2n} = \sum_{m=0}^{\infty} B_{2m}^{2n} \sin 2mx,$<br>$\text{se}_0 = 0$ | $qB_4^{2n} = (b_{2n}-4)B_2^{2n};$<br>$qB_{2m+2}^{2n} = (b_{2n}-4m^2)B_{2m}^{2n} - qB_{2m-2}^{2n}, \quad m \geq 2$                                               | $\sum_{m=0}^{\infty} (B_{2m}^{2n})^2 = 1$                                                                                           |
| $\text{se}_{2n+1} = \sum_{m=0}^{\infty} B_{2m+1}^{2n+1} \sin(2m+1)x$              | $qB_3^{2n+1} = (b_{2n+1}-1-q)B_1^{2n+1};$<br>$qB_{2m+3}^{2n+1} = [b_{2n+1}-(2m+1)^2]B_{2m+1}^{2n+1} - qB_{2m-1}^{2n+1}, \quad m \geq 1$                         | $\sum_{m=0}^{\infty} (B_{2m+1}^{2n+1})^2 = 1$                                                                                       |

#### A.14.1-2. Properties of the Mathieu functions.

The Mathieu functions possess the following properties:

$$\begin{aligned} \text{ce}_{2n}(x, -q) &= (-1)^n \text{ce}_{2n}\left(\frac{\pi}{2} - x, q\right), & \text{ce}_{2n+1}(x, -q) &= (-1)^n \text{se}_{2n+1}\left(\frac{\pi}{2} - x, q\right), \\ \text{se}_{2n}(x, -q) &= (-1)^{n-1} \text{se}_{2n}\left(\frac{\pi}{2} - x, q\right), & \text{se}_{2n+1}(x, -q) &= (-1)^n \text{ce}_{2n+1}\left(\frac{\pi}{2} - x, q\right). \end{aligned}$$

Selecting sufficiently large number  $m$  and omitting the term with the maximum number in the recurrence relations (indicated in Table A3), we can obtain approximate relations for eigenvalues  $a_n$  (or  $b_n$ ) with respect to parameter  $q$ . Then, equating the determinant of the corresponding homogeneous linear system of equations for coefficients  $A_m^n$  (or  $B_m^n$ ) to zero, we obtain an algebraic equation for finding  $a_n(q)$  (or  $b_n(q)$ ).

For fixed real  $q \neq 0$ , eigenvalues  $a_n$  and  $b_n$  are all real and different, while

$$\begin{aligned} \text{if } q > 0 \quad \text{then} \quad & a_0 < b_1 < a_1 < b_2 < a_2 < \dots \\ \text{if } q < 0 \quad \text{then} \quad & a_0 < a_1 < b_1 < b_2 < a_2 < a_3 < b_3 < b_4 < \dots \end{aligned}$$

The eigenvalues possess the properties

$$a_{2n}(-q) = a_{2n}(q), \quad b_{2n}(-q) = b_{2n}(q), \quad a_{2n+1}(-q) = b_{2n+1}(q).$$

Tables of the eigenvalues  $a_n = a_n(q)$  and  $b_n = b_n(q)$  can be found in Abramowitz and Stegun (1964, Chapter 20).

The solution of the Mathieu equation corresponding to eigenvalue  $a_n$  (or  $b_n$ ) has  $n$  zeros on the interval  $0 \leq x < \pi$  ( $q$  is a real number).

### A.14.1-3. Asymptotic expansions at $q \rightarrow 0$ and $q \rightarrow \infty$ .

Listed below are two leading terms of asymptotic expansions of the Mathieu functions  $\text{ce}_n(x, q)$  and  $\text{se}_n(x, q)$ , as well as of the corresponding eigenvalues  $a_n(q)$  and  $b_n(q)$ , as  $q \rightarrow 0$ :

$$\text{ce}_0(x, q) = \frac{1}{\sqrt{2}} \left( 1 - \frac{q}{2} \cos 2x \right), \quad a_0(q) = -\frac{q^2}{2} + \frac{7q^4}{128};$$

$$\text{ce}_1(x, q) = \cos x - \frac{q}{8} \cos 3x, \quad a_1(q) = 1 + q;$$

$$\text{ce}_2(x, q) = \cos 2x + \frac{q}{4} \left( 1 - \frac{\cos 4x}{3} \right), \quad a_2(q) = 4 + \frac{5q^2}{12};$$

$$\text{ce}_n(x, q) = \cos nx + \frac{q}{4} \left[ \frac{\cos(n+2)x}{n+1} - \frac{\cos(n-2)x}{n-1} \right], \quad a_n(q) = n^2 + \frac{q^2}{2(n^2-1)} \quad (n \geq 3);$$

$$\text{se}_1(x, q) = \sin x - \frac{q}{8} \sin 3x, \quad b_1(q) = 1 - q;$$

$$\text{se}_2(x, q) = \sin 2x - q \frac{\sin 4x}{12}, \quad b_2(q) = 4 - \frac{q^2}{12};$$

$$\text{se}_n(x, q) = \sin nx - \frac{q}{4} \left[ \frac{\sin(n+2)x}{n+1} - \frac{\sin(n-2)x}{n-1} \right], \quad b_n(q) = n^2 + \frac{q^2}{2(n^2-1)} \quad (n \geq 3).$$

Asymptotic results as  $q \rightarrow \infty$  ( $-\pi/2 < x < \pi/2$ ):

$$a_n(q) \approx -2q + 2(2n+1)\sqrt{q} + \frac{1}{4}(2n^2 + 2n + 1),$$

$$b_{n+1}(q) \approx -2q + 2(2n+1)\sqrt{q} + \frac{1}{4}(2n^2 + 2n + 1),$$

$$\text{ce}_n(x, q) \approx \lambda_n q^{-1/4} \cos^{-n-1} x [\cos^{2n+1} \xi \exp(2\sqrt{q} \sin x) + \sin^{2n+1} \xi \exp(-2\sqrt{q} \sin x)], \quad \xi = \frac{1}{2}x + \frac{\pi}{4},$$

$$\text{se}_{n+1}(x, q) \approx \mu_{n+1} q^{-1/4} \cos^{-n-1} x [\cos^{2n+1} \xi \exp(2\sqrt{q} \sin x) - \sin^{2n+1} \xi \exp(-2\sqrt{q} \sin x)],$$

where the  $\lambda_n$  and  $\mu_n$  some constants independent of the parameter  $q$ .

⊙ *References:* H. Bateman and A. Erdélyi (1955, Vol. 3), M. Abramowitz and I. Stegun (1964).

## A.15. Modified Mathieu Functions

The modified Mathieu functions  $\text{Ce}_n(x, q)$  and  $\text{Se}_n(x, q)$  are solutions of the modified Mathieu equation

$$y''_{xx} - (a - 2q \cosh 2x)y = 0,$$

with  $a = a_n(q)$  and  $a = b_n(q)$  being the eigenvalues of the Mathieu equation (see Section A.12).

The modified Mathieu functions are defined as

$$\text{Ce}_{2n+p}(x, q) = \text{ce}_{2n+p}(ix, q) = \sum_{k=0}^{\infty} A_{2k+p}^{2n+p} \cosh[(2k+p)x],$$

$$\text{Se}_{2n+p}(x, q) = -i \text{se}_{2n+p}(ix, q) = \sum_{k=0}^{\infty} B_{2k+p}^{2n+p} \sinh[(2k+p)x],$$

where  $p$  may be equal to 0 and 1, and coefficients  $A_{2k+p}^{2n+p}$  and  $B_{2k+p}^{2n+p}$  are indicated in Subsection A.12.

⊙ *References:* H. Bateman and A. Erdélyi (1955, Vol. 3), M. Abramowitz and I. Stegun (1964).

## A.16. Orthogonal Polynomials

All zeros of each of the orthogonal polynomials  $\mathcal{P}_n(x)$  considered in this section are real and simple. The zeros of the polynomials  $\mathcal{P}_n(x)$  and  $\mathcal{P}_{n+1}(x)$  are alternating.

For Legendre polynomials see Section A.12.

## A.16.1. Laguerre Polynomials and Generalized Laguerre Polynomials

### A.16.1-1. Laguerre polynomials.

The Laguerre polynomials  $L_n = L_n(x)$  satisfy the equation

$$xy''_{xx} + (1-x)y'_x + ny = 0$$

and are defined by the formulas

$$L_n(x) = \frac{1}{n!} e^x \frac{d^n}{dx^n} (x^n e^{-x}) = \frac{(-1)^n}{n!} \left[ x^n - n^2 x^{n-1} + \frac{n^2(n-1)^2}{2!} x^{n-2} + \dots \right].$$

The first four polynomials have the form

$$L_0 = 1, \quad L_1 = -x + 1, \quad L_2 = \frac{1}{2}(x^2 - 4x + 2), \quad L_3 = \frac{1}{6}(-x^3 + 9x^2 - 18x + 6).$$

To calculate  $L_n(x)$  for  $n \geq 2$ , one can use the recurrent formulas

$$L_{n+1}(x) = \frac{1}{n+1} [(2n+1-x)L_n(x) - nL_{n-1}(x)].$$

The functions  $L_n(x)$  form an orthonormal system on the interval  $0 < x < \infty$  with weight  $e^{-x}$ :

$$\int_0^\infty e^{-x} L_n(x) L_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ 1 & \text{if } n = m. \end{cases}$$

The generating function is

$$\frac{1}{1-s} \exp\left(-\frac{sx}{1-s}\right) = \sum_{n=0}^{\infty} L_n(x) s^n, \quad |s| < 1.$$

### A.16.1-2. Generalized Laguerre polynomials.

The generalized Laguerre polynomials  $L_n^\alpha = L_n^\alpha(x)$  ( $\alpha > -1$ ) satisfy the equation

$$xy''_{xx} + (\alpha + 1 - x)y'_x + ny = 0$$

and are defined by the formulas

$$L_n^\alpha(x) = \frac{1}{n!} x^{-\alpha} e^x \frac{d^n}{dx^n} (x^{n+\alpha} e^{-x}) = \sum_{m=0}^n C_{n+\alpha}^{n-m} \frac{(-x)^m}{m!}.$$

The first two polynomials have the form

$$L_0^\alpha = 1, \quad L_1^\alpha = \alpha + 1 - x.$$

To calculate  $L_n^\alpha(x)$  for  $n \geq 2$ , one can use the recurrent formulas

$$L_{n+1}^\alpha(x) = \frac{1}{n+1} [(2n+\alpha+1-x)L_n^\alpha(x) - (n+\alpha)L_{n-1}^\alpha(x)].$$

The functions  $L_n^\alpha(x)$  form an orthogonal system on the interval  $0 < x < \infty$  with weight  $x^\alpha e^{-x}$ :

$$\int_0^\infty x^\alpha e^{-x} L_n^\alpha(x) L_m^\alpha(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{\Gamma(\alpha+n+1)}{n!} & \text{if } n = m. \end{cases}$$

The generating function is

$$(1-s)^{-\alpha-1} \exp\left(-\frac{sx}{1-s}\right) = \sum_{n=0}^{\infty} L_n^\alpha(x) s^n, \quad |s| < 1.$$

## A.16.2. Chebyshev Polynomials and Functions

### A.16.2-1. Chebyshev polynomials.

The Chebyshev polynomials  $T_n = T_n(x)$  satisfy the equation

$$(1 - x^2)y''_{xx} - xy'_x + n^2y = 0 \quad (1)$$

and are defined by the formulas

$$\begin{aligned} T_n(x) &= \cos(n \arccos x) = \frac{(-2)^n n!}{(2n)!} \sqrt{1-x^2} \frac{d^n}{dx^n} [(1-x^2)^{n-\frac{1}{2}}] \\ &= \frac{n}{2} \sum_{m=0}^{[n/2]} (-1)^m \frac{(n-m-1)!}{m! (n-2m)!} (2x)^{n-2m} \quad (n = 0, 1, 2, \dots), \end{aligned}$$

where  $[A]$  stands for the integer part of a number  $A$ .

The first four polynomials are

$$T_0 = 1, \quad T_1 = x, \quad T_2 = 2x^2 - 1, \quad T_3 = 4x^3 - 3x.$$

The recurrent formulas:

$$T_{n+1}(x) = 2xT_n(x) - T_{n-1}(x), \quad n \geq 2.$$

The functions  $T_n(x)$  form an orthogonal system on the interval  $-1 < x < +1$ , with

$$\int_{-1}^{+1} \frac{T_n(x)T_m(x)}{\sqrt{1-x^2}} dx = \begin{cases} 0 & \text{if } n \neq m, \\ \frac{1}{2}\pi & \text{if } n = m \neq 0, \\ \pi & \text{if } n = m = 0. \end{cases}$$

### A.16.2-2. Chebyshev functions of the second kind.

The Chebyshev functions of the second kind,

$$U_0(x) = \arcsin x,$$

$$U_n(x) = \sin(n \arcsin x) = \frac{\sqrt{1-x^2}}{n} \frac{dT_n(x)}{dx} \quad (n = 1, 2, \dots),$$

just as the Chebyshev polynomials, also satisfy the differential equation (1).

The generating function is

$$\frac{1-sx}{1-2sx+s^2} = \sum_{n=0}^{\infty} T_n(x)s^n \quad (|s| < 1).$$

## A.16.3. Hermite Polynomial

The Hermite polynomial  $H_n = H_n(x)$  satisfies the equation

$$y''_{xx} - 2xy'_x + 2ny = 0$$

and is defined by the formulas

$$H_n(x) = (-1)^n \exp(x^2) \frac{d^n}{dx^n} \exp(-x^2).$$

The first four polynomials are

$$H_0 = 1, \quad H_1 = x, \quad H_2 = 4x^2 - 2, \quad H_3 = 8x^3 - 12x.$$

The recurrent formulas:

$$H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x), \quad n \geq 2.$$

The functions  $H_n(x)$  form an orthogonal system on the interval  $-\infty < x < \infty$  with weight  $e^{-x^2}$ :

$$\int_{-\infty}^{\infty} \exp(-x^2) H_n(x) H_m(x) dx = \begin{cases} 0 & \text{if } n \neq m, \\ \sqrt{\pi} 2^n n! & \text{if } n = m. \end{cases}$$

The Hermite functions  $\psi_n(x)$  are introduced by the formula  $\psi_n(x) = \exp(-\frac{1}{2}x^2) H_n(x)$ , where  $n = 0, 1, 2, \dots$

The generating function:

$$\exp(-s^2 + 2sx) = \sum_{n=0}^{\infty} H_n(x) \frac{s^n}{n!}.$$

#### A.16.4. Jacobi Polynomials

The Jacobi polynomials  $P_n^{\alpha, \beta} = P_n^{\alpha, \beta}(x)$  satisfy the equation

$$(1-x^2)y''_{xx} + [\beta - \alpha - (\alpha + \beta + 2)x]y'_x + n(n + \alpha + \beta + 1)y = 0$$

and are defined by the formulas

$$P_n^{\alpha, \beta} = \frac{(-1)^n}{2^n n!} (1-x)^{-\alpha} (1+x)^{-\beta} \frac{d^n}{dx^n} \left[ (1-x)^{\alpha+n} (1+x)^{\beta+n} \right] = 2^{-n} \sum_{m=0}^n C_{n+\alpha}^m C_{n+\beta}^{n-m} (x-1)^{n-m} (x+1)^m,$$

where  $C_b^a$  are binomial coefficients.

☉ References for Supplement: H. Bateman and A. Erdélyi (1953, 1955), M. Abramowitz and I. A. Stegun (1964).

## Supplement B

# Methods of Generalized and Functional Separation of Variables in Nonlinear Equations of Mathematical Physics\*

---

## B.1. Introduction

### B.1.1. Preliminary Remarks

Separation of variables is the most common approach to solve linear equations of mathematical physics. This approach involves searching for exact solutions in the form of the product of functions depending on different arguments (see Section 0.4).

As far as nonlinear equations with two independent variables  $x, y$  and a dependent variable  $w$  are concerned, some of these equations also have solutions with the form

$$w(x, y) = \varphi(x)\psi(y) \quad \text{or} \quad w(x, y) = \varphi(x) + \psi(y)$$

that are called *multiplicatively* and *additively separable*, respectively. We call such solutions *ordinary separable solutions*. In particular, integrating a few classes of first-order nonlinear partial differential equations is based on searching for additively separable solutions [e.g., see Appell (1953), Kamke (1965), Markeev (1990), Zwillinger (1998), Polyanin, Zaitsev, and Moussiaux (2001)].

Over the last decade, more sophisticated, *generalized* and *functional separable solutions* have been obtained for a number of second-order nonlinear equations of mathematical physics. For example, Galaktionov and Posashkov (1989) and Galaktionov, Posashkov, and Svirshchevskii (1995) obtained generalized separable solutions with the forms  $w(x, t) = \varphi(t)\psi(x) + \chi(t)$  and  $w(x, t) = \varphi(t)\psi(x) + \chi(x)$  for some classes of parabolic and hyperbolic equations with quadratic nonlinearities. In Galaktionov and Posashkov (1994), Galaktionov (1995), and Svirshchevskii (1995), more complicated generalized separable solutions are presented. The results of Galaktionov and Posashkov (1994) and Galaktionov (1995) are based on finding finite-dimensional subspaces that are invariant under appropriate nonlinear differential operators (in practice, the authors had to find a system of coordinate functions in one of the variables by the method of undetermined coefficients).

In Grundland and Infeld (1992), Miller and Rubel (1993), Zhdanov (1994), and Andreev, Kaptsov, Pukhnachev, and Rodionov (1994), all nonlinear heat (diffusion) and wave equations of the form  $\partial_{xx}w \pm \partial_{yy}w = f(w)$  which admit functional separable solutions having the form  $w(x, y) = F(z)$ , where  $z = \varphi(x) + \psi(y)$ , are described. Doyle and Vassiliou (1998) indicated all one-dimensional nonstationary heat equations  $\partial_t w = \partial_x[f(w)\partial_x w]$  which admit solutions of the form  $w(x, t) = F(z)$ ,  $z = \varphi(x) + \psi(t)$ . In Zaitsev and Polyanin (1996), Polyanin and Zhurov (1998), Polyanin, Vyazmin, Zhurov, and Kazenin (1998), and Polyanin, Zhurov, and Vyazmin (2000), many nonlinear mathematical physics equations of various types that admit generalized and functional separable solutions are described (special attention was paid to equations of general form which depend on arbitrary functions).

---

\* Sections B.1–B.5 were written with A. I. Zhurov.

Functional differential equations that involve unknown functions (and their derivatives) with different arguments arise when searching for ordinary, generalized, and functional separable solutions. The current supplement presents direct methods for and examples of constructing such solutions and reviews application of these methods to solving various classes of the second-, third-, fourth-, and higher-order partial differential equations (in total, about 150 nonlinear equations with solutions are described). Special attention is paid to equations of heat and mass transfer theory, wave theory, and hydrodynamics, as well as mathematical physics equations of general form that involve arbitrary functions.

It should be noted that often exact generalized and functional separable solutions cannot be obtained by group theoretic methods or other well-known methods.

### B.1.2. Simple Cases of Variable Separation in Nonlinear Equations

In isolated cases, the separation of variables in nonlinear equations is carried out following the same technique as in linear equations. Specifically, an exact solution is sought in the form of the product or sum of functions depending on different arguments. On substituting it into the equation and performing elementary algebraic manipulations, one obtains an equation with the two sides dependent on different variables (for equations with two variables). Then one concludes that the expressions on each side must be equal to the same constant quantity, called a separation constant. Below we consider specific examples.

**Example 1.** The heat equation with a power nonlinearity

$$\frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left( w^k \frac{\partial w}{\partial x} \right) \quad (1)$$

has an exact solution in the product form

$$w = \varphi(x)\psi(t). \quad (2)$$

Substituting (2) into (1) yields

$$\varphi\psi'_t = a\psi^{k+1}(\varphi^k\varphi'_x)'_x.$$

Separating the variables by dividing both sides by  $\varphi\psi^{k+1}$ , we obtain

$$\frac{\psi'_t}{\psi^{k+1}} = \frac{a(\varphi^k\varphi'_x)'_x}{\varphi}.$$

The left-hand side depends of  $t$  alone and the right-hand side on  $x$  alone. This is possible only if

$$\frac{\psi'_t}{\psi^{k+1}} = C, \quad \frac{a(\varphi^k\varphi'_x)'_x}{\varphi} = C, \quad (3)$$

where  $C$  is an arbitrary constant (separation constant). On solving the ordinary differential equations (3), we obtain a solution of equation (1) with the form (2).

The procedure for constructing a separable solution (2) of the nonlinear equation (1) is identical to that used in solving linear equations [in particular, equation (1) with  $k = 0$ ]. We refer to the cases of similar separation of variables as *simple separable* cases.

**Example 2.** The wave equation with an exponential nonlinearity

$$\frac{\partial^2 w}{\partial t^2} = a \frac{\partial}{\partial x} \left( e^{\lambda w} \frac{\partial w}{\partial x} \right) \quad (4)$$

has an additively separable solution

$$w = \varphi(x) + \psi(t). \quad (5)$$

On substituting (5) into (4) and dividing by  $e^{\lambda\psi}$ , we arrive at the equation

$$e^{-\lambda\psi}\psi''_{tt} = a(e^{\lambda\varphi}\varphi'_x)'_x,$$

whose left-hand side depends on  $t$  alone and the right-hand side on  $x$  alone. This is possible only if

$$e^{-\lambda\psi}\psi''_{tt} = C, \quad a(e^{\lambda\varphi}\varphi'_x)'_x = C, \quad (6)$$



where  $C$  is an arbitrary constant. Solving the ordinary differential equations (6) yields a solution of equation (4) with the form (5).

**Example 3.** The heat equation in an anisotropic medium with a logarithmic source

$$\frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(y) \frac{\partial w}{\partial y} \right] = aw \ln w \quad (7)$$

has a multiplicatively separable solution

$$w = \varphi(x)\psi(y). \quad (8)$$

On substituting (8) into (7), dividing by  $\varphi\psi$ , and transposing individual terms of the resulting equation, we obtain

$$\frac{1}{\varphi} [f(x)\varphi'_x]'_x - a \ln \varphi = -\frac{1}{\psi} [g(y)\psi'_y]'_y + a \ln \psi.$$

The left-hand side of this equation depends only on  $x$  and the right-hand only on  $y$ . By equating both sides to a constant quantity, one obtains ordinary differential equations for  $\varphi(x)$  and  $\psi(y)$ .

### B.1.3. Examples of Nontrivial Variable Separation in Nonlinear Equations

Unlike linear equations, the variables in nonlinear equations often separate differently. We exemplify this below.

**Example 4.** Consider the equation with a cubic nonlinearity

$$\frac{\partial w}{\partial t} = f(t) \frac{\partial^2 w}{\partial x^2} + w \left( \frac{\partial w}{\partial x} \right)^2 - aw^3, \quad (9)$$

where  $f(t)$  is an arbitrary function. We look for exact solutions in the product form. We substitute (2) into (9) and divide the resulting equation by  $f(t)\varphi(x)\psi(t)$  to obtain

$$\frac{\psi'_t}{f\psi} = \frac{\varphi''_{xx}}{\varphi} + \frac{\psi^2}{f} [(\varphi'_x)^2 - a\varphi^2]. \quad (10)$$

In the general case, this expression cannot be represented as the sum of two functions depending on different arguments. This however does not mean that equation (9) has no solutions of the form (2).

1°. One can make sure by direct check that, for  $a > 0$ , the functional differential equation (10) has solutions

$$\varphi(x) = C \exp(\pm x\sqrt{a}), \quad \psi(t) = \exp \left[ a \int f(t) dt \right], \quad (11)$$

where  $C$  is an arbitrary constant. Solution (11) for  $\varphi$  makes the expression in square brackets in (10) vanish, which allows separation of variables.

2°. There is a more general solution of the functional differential equation (10) for  $a > 0$ :

$$\begin{aligned} \varphi(x) &= C_1 \exp(x\sqrt{a}) + C_2 \exp(-x\sqrt{a}), \\ \psi(t) &= e^F \left( C_3 + 8aC_1C_2 \int e^{2F} dt \right)^{-1/2}, \quad F = a \int f(t) dt, \end{aligned}$$

where  $C_1$ ,  $C_2$ , and  $C_3$  are arbitrary constants. The function  $\varphi = \varphi(x)$  makes each of the terms in (10) that depend on  $x$  constant, namely,

$$\varphi''_{xx}/\varphi = \text{const}, \quad (\varphi'_x)^2 - a\varphi^2 = \text{const}.$$

It is this circumstance that makes it possible to separate the variables.

Note that the function  $\psi = \psi(t)$  satisfies the Bernoulli equation  $\psi'_t = af(t)\psi - 4aC_1C_2\psi^3$ .

3°. There is another solution of the functional differential equation (10) for  $a < 0$ :

$$\begin{aligned} \varphi(x) &= C_1 \sin(x\sqrt{-a}) + C_2 \cos(x\sqrt{-a}), \\ \psi(t) &= e^F \left[ C_3 + 2a(C_1^2 + C_2^2) \int e^{2F} dt \right]^{-1/2}, \quad F = a \int f(t) dt, \end{aligned}$$

where  $C_1$ ,  $C_2$ , and  $C_3$  are arbitrary constants. The function  $\varphi = \varphi(x)$  makes both terms in (10) that depend on  $x$  constant. Note that the function  $\psi = \psi(t)$  is determined by the Bernoulli equation  $\psi'_t = af(t)\psi - a(C_1^2 + C_2^2)\psi^3$ .

**Example 5.** Consider the third-order equation

$$\frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x^2} + a \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = b \frac{\partial^3 w}{\partial x^3} + c \frac{\partial^3 w}{\partial y^3}. \quad (12)$$

We look for additive separable solutions

$$w = f(x) + g(y). \quad (13)$$

Substituting (13) into (12) yields

$$g_y' f_{xx}'' + a f_x' g_{yy}'' = b f_{xxx}''' + c g_{yyy}'''. \quad (14)$$

This expression cannot be rewritten as the sum of two functions depending on different arguments.

It is not difficult to see that the functional differential equation (14) is satisfied if

$$\begin{aligned} g_y' = C_1 &\implies g(y) = C_1 y + C_2, & f(x) = C_3 \exp(C_1 x / b) + C_4 x &\quad (\text{case 1}), \\ f_x' = C_1 &\implies f(x) = C_1 x + C_2, & g(y) = C_3 \exp(a C_1 y / c) + C_4 y &\quad (\text{case 2}), \end{aligned}$$

where  $C_1, C_2, C_3$ , and  $C_4$  are arbitrary constants. In both cases, two terms of the four in (14) vanish, which makes it possible to separate the variables.

In addition, equation (12) has a more complicated solution of the form (13):

$$w = C_1 e^{-a\lambda x} + \frac{c\lambda}{a} x + C_2 e^{\lambda y} - ab\lambda y + C_3,$$

where  $C_1, C_2, C_3$ , and  $\lambda$  are arbitrary constants. The mechanism of separation of variables is different here: both nonlinear terms on the left-hand side in (14) contain terms which cannot be rewritten in additive form but are equal in magnitude and have unlike signs. In adding, the two terms cancel, thus resulting in separation of variables:

$$\begin{aligned} &g_y' f_{xx}'' = C_1 C_2 a^2 \lambda^3 e^{\lambda y - a\lambda x} - C_1 b(a\lambda)^3 e^{-a\lambda x} \\ &+ a f_x' g_{yy}'' = -C_1 C_2 a^2 \lambda^3 e^{\lambda y - a\lambda x} + C_2 c \lambda^3 e^{\lambda y} \\ \hline &g_y' f_{xx}'' + a f_x' g_{yy}'' = -C_1 b(a\lambda)^3 e^{-a\lambda x} + C_2 c \lambda^3 e^{\lambda y} = b f_{xxx}''' + c g_{yyy}'''. \end{aligned}$$

**Example 6.** Consider the second-order equation with a cubic nonlinearity

$$(1 + w^2) \left( \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} \right) - 2w \left( \frac{\partial w}{\partial x} \right)^2 - 2w \left( \frac{\partial w}{\partial y} \right)^2 = aw(1 - w^2). \quad (15)$$

We seek an exact solution of this equation in the product form

$$w = f(x)g(y). \quad (16)$$

Substituting (16) into (15) yields

$$(1 + f^2 g^2)(g f_{xx}'' + f g_{yy}'') - 2fg[g^2(f_x')^2 + f^2(g_y')^2] = afg(1 - f^2 g^2). \quad (17)$$

This expression cannot be rewritten as the sum of two functions with different arguments. Nevertheless, equation (15) has solutions of the form (16). One can make sure by direct check that the functions  $f = f(x)$  and  $g = g(y)$  satisfying the nonlinear ordinary differential equations

$$\begin{aligned} (f_x')^2 &= Af^4 + Bf^2 + C, \\ (g_y')^2 &= Cg^4 + (a - B)g^2 + A, \end{aligned} \quad (18)$$

where  $A, B$ , and  $C$  are arbitrary constants, reduce equation (17) to an identity; to verify this, one should use the relations  $f_{xx}'' = 2Af^3 + Bf$  and  $g_{yy}'' = 2Cg^3 + (a - B)g$  that follow from (18).

*Remark.* By the variable change  $u = 4 \arctan w$  equation (15) can be reduced to a nonlinear heat equation with a sinusoidal source,  $\Delta u = a \sin u$ .

The examples considered above illustrate some specific features of separable solutions to nonlinear equations. Sections B.2 and B.3 outline fairly general methods for constructing similar and more complicated solutions to nonlinear partial differential equations.

## B.2. Methods of Generalized Separation of Variables

### B.2.1. Structure of Generalized Separable Solutions

B.2.1-1. General form of solutions. The classes of nonlinear equations considered.

To simplify the presentation, we confine ourselves to the case of mathematical physics equations with two independent variables  $x, y$  and a dependent variable  $w$  (one of the independent variables can play the role of time).

Linear separable equations of mathematical physics admit exact solutions in the form

$$w(x, y) = \varphi_1(x)\psi_1(y) + \varphi_2(x)\psi_2(y) + \cdots + \varphi_n(x)\psi_n(y), \quad (1)$$

where the  $w_i = \varphi_i(x)\psi_i(y)$  are particular solutions; the functions  $\varphi_i(x)$ , as well as the functions  $\psi_i(y)$ , with different numbers  $i$  are not related to one another.

Also having exact solutions of the form (1) are many nonlinear partial differential equations with quadratic or power nonlinearities

$$f_1(x)g_1(y)\Pi_1[w] + f_2(x)g_2(y)\Pi_2[w] + \cdots + f_m(x)g_m(y)\Pi_m[w] = 0, \quad (2)$$

where the  $\Pi_i[w]$  are differential forms that are the products of nonnegative integer powers of the function  $w$  and its partial derivatives  $\partial_x w, \partial_y w, \partial_{xx} w, \partial_{xy} w, \partial_{yy} w, \partial_{xxx} w$ , etc. We will refer to solutions (1) of nonlinear equations (2) as *generalized separable solutions*. Unlike linear equations, in nonlinear equations the functions  $\varphi_i(x)$  with different subscripts  $i$  are usually related to one another [and to the functions  $\psi_j(y)$ ]. Subsections B.1.2 and B.1.3 give examples of exact solutions (1) to nonlinear equations (2) for some simple cases with  $n = 1$  or  $n = 2$  (for  $\psi_1 = \varphi_2 = 1$ ).

*Remark.* If the  $f_s(x)$  and  $g_s(y)$  in (2) are all constant, then one can seek solutions in the more general form

$$w(x, y) = \sum_{m=1}^n \varphi_m(\xi)\psi_m(\eta), \quad \xi = a_1x + a_2y, \quad \eta = b_1x + b_2y,$$

where  $a_1, a_2, b_1$ , and  $b_2$  are constants. Some solutions of this sort are discussed in Subsections B.7.1 and B.8.1.

B.2.1-2. General form of functional differential equations.

In general, on substituting expression (1) into the differential equation (2), one arrives at a functional differential equation

$$\Phi_1(X)\Psi_1(Y) + \Phi_2(X)\Psi_2(Y) + \cdots + \Phi_k(X)\Psi_k(Y) = 0 \quad (3)$$

for the  $\varphi_i(x)$  and  $\psi_i(y)$ . The functionals  $\Phi_j(X)$  and  $\Psi_j(Y)$  depend only on  $x$  and  $y$ , respectively,

$$\begin{aligned} \Phi_j(X) &\equiv \Phi_j(x, \varphi_1, \varphi_1', \varphi_1'', \dots, \varphi_n, \varphi_n', \varphi_n''), \\ \Psi_j(Y) &\equiv \Psi_j(y, \psi_1, \psi_1', \psi_1'', \dots, \psi_n, \psi_n', \psi_n''). \end{aligned} \quad (4)$$

Here, for simplicity, the formulas are written out for the case of a second-order equation (2); for higher-order equations, the right-hand sides of relations (4) will contain higher-order derivatives of  $\varphi_i$  and  $\psi_j$ .

## B.2.2. Solution of Functional Differential Equations by Differentiation

### B.2.2-1. Description of the method.

1°. Assume that  $\Psi_k \neq 0$ . We divide equation (3) by  $\Psi_k$  and differentiate with respect to  $y$ . This results in a similar equation but with fewer terms:

$$\begin{aligned}\tilde{\Phi}_1(X)\tilde{\Psi}_1(Y) + \tilde{\Phi}_2(X)\tilde{\Psi}_2(Y) + \cdots + \tilde{\Phi}_{k-1}(X)\tilde{\Psi}_{k-1}(Y) &= 0, \\ \tilde{\Phi}_j(X) &= \Phi_j(X), \quad \tilde{\Psi}_j(Y) = [\Psi_j(Y)/\Psi_k(Y)]'_y.\end{aligned}$$

We continue the above procedure until we obtain a separable two-term equation

$$\hat{\Phi}_1(X)\hat{\Psi}_1(Y) + \hat{\Phi}_2(X)\hat{\Psi}_2(Y) = 0. \quad (5)$$

Three cases must be considered.

*Nondegenerate case:*  $|\hat{\Phi}_1(X)| + |\hat{\Phi}_2(X)| \neq 0$  and  $|\hat{\Psi}_1(Y)| + |\hat{\Psi}_2(Y)| \neq 0$ . Then equation (5) is equivalent to the ordinary differential equations

$$\hat{\Phi}_1(X) + C\hat{\Phi}_2(X) = 0, \quad C\hat{\Psi}_1(Y) - \hat{\Psi}_2(Y) = 0,$$

where  $C$  is an arbitrary constant. The equations  $\hat{\Phi}_2 = 0$  and  $\hat{\Psi}_1 = 0$  correspond to the limit case  $C = \infty$ .

*Two degenerate cases:*

$$\begin{aligned}\hat{\Phi}_1(X) \equiv 0, \quad \hat{\Phi}_2(X) \equiv 0 &\implies \hat{\Psi}_{1,2}(Y) \text{ are any;} \\ \hat{\Psi}_1(Y) \equiv 0, \quad \hat{\Psi}_2(Y) \equiv 0 &\implies \hat{\Phi}_{1,2}(X) \text{ are any.}\end{aligned}$$

2°. The solutions of the two-term equation (5) should be substituted into the original functional differential equation (3) to “remove” redundant constants of integration [these arise because equation (5) is obtained from (3) by differentiation].

3°. The case  $\Psi_k \equiv 0$  should be treated separately (since we divided the equation by  $\Psi_k$  at the first stage). Likewise, we have to study all other cases where the functionals by which the intermediate functional differential equations were divided vanish.

*Remark 1.* The functional differential equation (3) can happen to have no solutions.

*Remark 2.* At each subsequent stage, the number of terms in the functional differential equation can be reduced by differentiation with respect to either  $y$  or  $x$ . For example, we can assume at the first stage that  $\Phi_k \neq 0$ . On dividing equation (3) by  $\Phi_k$  and differentiating with respect to  $x$ , we again obtain a similar equation that has fewer terms.

### B.2.2-2. Examples of constructing exact generalized separable solutions.

Below we consider specific examples illustrating the application of the above method to constructing exact generalized separable solutions of nonlinear equations.

**Example 1.** The two-dimensional stationary equations of motion of a viscous incompressible fluid are reduced to a single fourth-order nonlinear equation for the stream function (see equation 1 in Subsection B.7.1), specifically,

$$\frac{\partial w}{\partial y} \frac{\partial}{\partial x} (\Delta w) - \frac{\partial w}{\partial x} \frac{\partial}{\partial y} (\Delta w) = \nu \Delta \Delta w, \quad \Delta w = \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2}. \quad (6)$$

We seek exact separable solutions of equation (6) in the form

$$w = f(x) + g(y). \quad (7)$$

Substituting (7) into (6) yields

$$g'_y f'''_{xxx} - f'_x g'''_{yyy} = \nu f''_{xxx} + \nu g''_{yyy}. \quad (8)$$

Differentiating (8) with respect to  $x$  and  $y$ , we obtain

$$g''_{yy} f''''_{xxx} - f''_{xx} g''''_{yyy} = 0. \quad (9)$$

*Nondegenerate case.* If  $f''_{xx} \neq 0$  and  $g''_{yy} \neq 0$ , we separate the variables in (9) to obtain the ordinary differential equations

$$f''''_{xxx} = C f''_{xx}, \quad (10)$$

$$g''''_{yyy} = C g''_{yy}, \quad (11)$$

which have different solutions depending on the value of the integration constant  $C$ .

1°. Solutions of equations (10) and (11) for  $C = 0$ :

$$\begin{aligned} f(x) &= A_1 + A_2x + A_3x^2 + A_4x^3, \\ g(y) &= B_1 + B_2y + B_3y^2 + B_4y^3, \end{aligned} \quad (12)$$

where  $A_k$  and  $B_k$  are arbitrary constants ( $k = 1, 2, 3, 4$ ). On substituting (12) into (8), we evaluate the integration constants. Three cases are possible:

$$\begin{aligned} A_4 &= B_4 = 0, & A_n, B_n &\text{ are any numbers } (n = 1, 2, 3); \\ A_k &= 0, & B_k &\text{ are any numbers } (k = 1, 2, 3, 4); \\ B_k &= 0, & A_k &\text{ are any numbers } (k = 1, 2, 3, 4). \end{aligned}$$

The first two sets of constants determine two simple solutions (7) of equation (6):

$$\begin{aligned} w &= C_1x^2 + C_2x + C_3y^2 + C_4y + C_5, \\ w &= C_1y^3 + C_2y^2 + C_3y + C_4, \end{aligned}$$

where  $C_1, \dots, C_5$  are arbitrary constants.

2°. Solutions of equations (10) and (11) for  $C = \lambda^2 > 0$ :

$$\begin{aligned} f(x) &= A_1 + A_2x + A_3e^{\lambda x} + A_4e^{-\lambda x}, \\ g(y) &= B_1 + B_2y + B_3e^{\lambda y} + B_4e^{-\lambda y}. \end{aligned} \quad (13)$$

Substituting (13) into (8), dividing by  $\lambda^3$ , and collecting terms, we obtain

$$A_3(\nu\lambda - B_2)e^{\lambda x} + A_4(\nu\lambda + B_2)e^{-\lambda x} + B_3(\nu\lambda + A_2)e^{\lambda y} + B_4(\nu\lambda - A_2)e^{-\lambda y} = 0.$$

Equating the coefficients of the exponentials to zero, we find

$$\begin{aligned} A_3 &= A_4 = B_3 = 0, & A_2 &= \nu\lambda & (\text{case 1}), \\ A_3 &= B_3 = 0, & A_2 &= \nu\lambda, & B_2 &= -\nu\lambda & (\text{case 2}), \\ A_3 &= B_4 = 0, & A_2 &= -\nu\lambda, & B_2 &= -\nu\lambda & (\text{case 3}). \end{aligned}$$

(The other constants are arbitrary.) These sets of constants determine three solutions (7) of equation (6):

$$\begin{aligned} w &= C_1e^{-\lambda y} + C_2y + C_3 + \nu\lambda x, \\ w &= C_1e^{-\lambda x} + \nu\lambda x + C_2e^{-\lambda y} - \nu\lambda y + C_3, \\ w &= C_1e^{-\lambda x} - \nu\lambda x + C_2e^{\lambda y} - \nu\lambda y + C_3, \end{aligned}$$

where  $C_1, C_2, C_3$ , and  $\lambda$  are arbitrary constants.

3°. Solution of equations (10) and (11) for  $C = -\lambda^2 < 0$ :

$$\begin{aligned} f(x) &= A_1 + A_2x + A_3 \cos(\lambda x) + A_4 \sin(\lambda x), \\ g(y) &= B_1 + B_2y + B_3 \cos(\lambda y) + B_4 \sin(\lambda y). \end{aligned} \quad (14)$$

Substituting (14) into (8) does not yield new real solutions.

*Degenerate cases.* If  $f''_{xx} \equiv 0$  or  $g''_{yy} \equiv 0$ , equation (9) becomes an identity for any  $g = g(y)$  or  $f = f(x)$ , respectively. These cases should be treated separately from the nondegenerate case. For example, if  $f''_{xx} \equiv 0$ , we have  $f(x) = Ax + B$ , where  $A$  and  $B$  are arbitrary numbers. Substituting this  $f$  into (8), we arrive at the equation  $-Ag'''_{yyy} = \nu g'''_{yyy}$ . Its general solution is given by  $g(y) = C_1 \exp(-Ay/\nu) + C_2y^2 + C_3y + C_4$ . Thus, we obtain another solution (7) of equation (6):

$$w = C_1e^{-\lambda y} + C_2y^2 + C_3y + C_4 + \nu\lambda x \quad (A = \nu\lambda, B = 0).$$

**Example 2.** Consider the second-order nonlinear parabolic equation

$$\frac{\partial w}{\partial t} = aw \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + c. \quad (15)$$

We look for exact separable solutions of equation (15) in the form

$$w = \varphi(t) + \psi(t)\theta(x). \quad (16)$$

Substituting (15) into (16) and collecting terms yields

$$\varphi'_t - c + \psi'_t \theta = a\varphi\psi\theta''_{xx} + \psi^2 [a\theta\theta''_{xx} + b(\theta'_x)^2]. \quad (17)$$

On dividing this relation by  $\psi^2$  and differentiating with respect to  $t$  and  $x$ , we obtain

$$(\psi'_t/\psi^2)'_t \theta'_x = a(\varphi/\psi)'_t \theta'''_{xxx}.$$

Separating the variables, we arrive at the ordinary differential equations

$$\theta'''_{xxx} = K\theta'_x, \quad (18)$$

$$(\psi'_t/\psi^2)'_t = aK(\varphi/\psi)'_t, \quad (19)$$

where  $K$  is an arbitrary constant. The general solution of equation (18) is given by

$$\theta = \begin{cases} A_1 x^2 + A_2 x + A_3 & \text{if } K = 0, \\ A_1 e^{\lambda x} + A_2 e^{-\lambda x} + A_3 & \text{if } K = \lambda^2 > 0, \\ A_1 \sin(\lambda x) + A_2 \cos(\lambda x) + A_3 & \text{if } K = -\lambda^2 < 0, \end{cases} \quad (20)$$

where  $A_1$ ,  $A_2$ , and  $A_3$  are arbitrary constants. Integrating (19) yields

$$\begin{aligned} \psi &= \frac{B}{t + C_1}, \quad \varphi(t) \text{ is any} & \text{if } K = 0, \\ \varphi &= B\psi + \frac{1}{aK} \frac{\psi'_t}{\psi}, \quad \psi(t) \text{ is any} & \text{if } K \neq 0, \end{aligned} \quad (21)$$

where  $B$  is an arbitrary constant. On substituting solutions (20) and (21) into (17), one can “remove” the redundant constants and define the functions  $\varphi$  and  $\psi$ . Below we summarize the results.

1°. Solution for  $a \neq -b$  and  $a \neq -2b$ :

$$w = \frac{c(a+2b)}{2(a+b)}(t+C_1) + C_2(t+C_1)^{-\frac{a}{a+2b}} - \frac{(x+C_3)^2}{2(a+2b)(t+C_1)} \quad (\text{corresponds to } K=0),$$

where  $C_1$ ,  $C_2$ , and  $C_3$  are arbitrary constants.

2°. Solution for  $b = -a$ :

$$w = \frac{1}{a\lambda^2} \frac{\psi'_t}{\psi} + \psi(A_1 e^{\lambda x} + A_2 e^{-\lambda x}) \quad (\text{corresponds to } K = \lambda^2 > 0),$$

where the function  $\psi = \psi(t)$  is determined from the autonomous ordinary differential equation

$$Z''_{tt} = ac\lambda^2 + 4a^2\lambda^4 A_1 A_2 e^{2Z}, \quad \psi = e^Z,$$

whose solution can be found in implicit form. In the special case  $A_1 = 0$  or  $A_2 = 0$ , we have  $\psi = C_1 \exp(\frac{1}{2}ac\lambda^2 t^2 + C_2 t)$ .

3°. Solution for  $b = -a$ :

$$w = -\frac{1}{a\lambda^2} \frac{\psi'_t}{\psi} + \psi[A_1 \sin(\lambda x) + A_2 \cos(\lambda x)] \quad (\text{corresponds to } K = -\lambda^2 < 0).$$

where the function  $\psi = \psi(t)$  is determined from the autonomous ordinary differential equation

$$Z''_{tt} = -ac\lambda^2 + a^2\lambda^4(A_1^2 + A_2^2)e^{2Z}, \quad \psi = e^Z,$$

whose solution can be found in implicit form.

## B.2.3. Solution of Functional Differential Equations by Splitting

### B.2.3-1. Preliminary remarks. Description of the method.

As one reduces the number of terms in the functional differential equation (3) by differentiation, redundant constants of integration arise. These constants must be “removed” at the final stage. Furthermore, the resulting equation can be of a higher-order than the original equation. To avoid these difficulties, it is convenient to reduce the solution of the functional differential equation to the solution of a linear functional equation of a standard form and solution of a system of ordinary differential equations. Thus, the original problem splits into two simpler problems. Below we outline the basic stages of the splitting method.

*The case of even number of terms in equation (3),  $k = 2s$ .*

1°. At the first stage, we treat equation (3) as a purely functional equation that depends on two variables  $X$  and  $Y$ , where  $\Phi_1(X), \dots, \Phi_s(X), \Psi_{s+1}(Y), \dots, \Psi_{2s}(Y)$  are unknown quantities and the functions  $\Phi_{s+1}(X), \dots, \Phi_{2s}(X), \Psi_1(Y), \dots, \Psi_s(Y)$  are assumed to be known.

It can be shown (by induction and differentiation) that the functional equation (3) has a solution depending on  $s^2$  arbitrary constants

$$\begin{aligned}\Phi_i(X) &= C_{i1}\Phi_{s+1}(X) + C_{i2}\Phi_{s+2}(X) + \dots + C_{is}\Phi_{2s}(X) & (i = 1, \dots, s), \\ \Psi_{s+i}(Y) &= -C_{1i}\Psi_1(Y) - C_{2i}\Psi_2(Y) - \dots - C_{si}\Psi_s(Y) & (i = 1, \dots, s),\end{aligned}\quad (22)$$

where the  $C_{ij}$  are arbitrary constants. Note that there are also “degenerate” solutions depending on fewer arbitrary constants (see Item 2° in Paragraph B.2.3-2).

2°. At the second stage, we substitute the  $\Phi_i(X)$  and  $\Psi_j(Y)$  of (4) into (22). This results in an overdetermined system of ordinary differential equations for the unknown functions  $\varphi_p(x)$  and  $\psi_q(y)$ .

*The case of odd number of terms in equation (3),  $k = 2s - 1$ .*

1°. If the number of terms is odd ( $k = 2s - 1$ ), the functional equation (3) has two different solutions with  $s(s - 1)$  arbitrary constants. One of them can be obtained from formulas (22) by setting  $\Phi_{2s} \equiv 0$  and discarding the last term with  $\Psi_{2s}$ . The other solution can be obtained from the first one by renaming  $\Phi_i(X) \rightleftharpoons \Psi_i(Y)$ .

2°. Further analysis for each solution should be performed following the same scheme as in the case of even number of terms in (3).

### B.2.3-2. Solutions of simple functional equations and their application.

Below we give solutions of two simple functional equations of the form (3) that will be used subsequently for solving specific nonlinear partial differential equations.

1°. The functional equation

$$\Phi_1\Psi_1 + \Phi_2\Psi_2 + \Phi_3\Psi_3 = 0 \quad (23)$$

where the  $\Phi_i$  are all functions of the same argument and the  $\Psi_i$  are all functions of another argument, has two solutions:

$$\begin{aligned}\Phi_1 &= A_1\Phi_3, & \Phi_2 &= A_2\Phi_3, & \Psi_3 &= -A_1\Psi_1 - A_2\Psi_2, \\ \Psi_1 &= A_1\Psi_3, & \Psi_2 &= A_2\Psi_3, & \Phi_3 &= -A_1\Phi_1 - A_2\Phi_2,\end{aligned}\quad (24)$$

where  $A_1$  and  $A_2$  are arbitrary constants.

2°. The functional equation

$$\Phi_1\Psi_1 + \Phi_2\Psi_2 + \Phi_3\Psi_3 + \Phi_4\Psi_4 = 0, \quad (25)$$

where the  $\Phi_i$  are all functions of the same argument and the  $\Psi_i$  are all functions of another argument, has a solution

$$\begin{aligned}\Phi_1 &= A_1\Phi_3 + A_2\Phi_4, & \Phi_2 &= A_3\Phi_3 + A_4\Phi_4, \\ \Psi_3 &= -A_1\Psi_1 - A_3\Psi_2, & \Psi_4 &= -A_2\Psi_1 - A_4\Psi_2\end{aligned}\quad (26a)$$

depending on four arbitrary constants  $A_m$  [see solution (22) with  $s=2$ ,  $C_{11} = A_1$ ,  $C_{12} = A_2$ ,  $C_{21} = A_3$ , and  $C_{22} = A_4$ ].

Equation (25) has also two “degenerate” solutions

$$\begin{aligned}\Phi_1 &= A_1\Phi_4, & \Phi_2 &= A_2\Phi_4, & \Phi_3 &= A_3\Phi_4, & \Psi_4 &= -A_1\Psi_1 - A_2\Psi_2 - A_3\Psi_3, \\ \Psi_1 &= A_1\Psi_4, & \Psi_2 &= A_2\Psi_4, & \Psi_3 &= A_3\Psi_4, & \Phi_4 &= -A_1\Phi_1 - A_2\Phi_2 - A_3\Phi_3\end{aligned}\quad (26b)$$

involving three arbitrary constants.

**Example 3.** Consider the nonlinear hyperbolic equation

$$\frac{\partial^2 w}{\partial t^2} = a \frac{\partial}{\partial x} \left( w \frac{\partial w}{\partial x} \right) + f(t)w + g(t), \quad (27)$$

where  $f(t)$  and  $g(t)$  are arbitrary functions. We look for generalized separable solutions with the form

$$w(x, t) = \varphi(x)\psi(t) + \chi(t). \quad (28)$$

Substituting (28) into (27) and collecting terms yield

$$a\psi^2(\varphi\varphi'_x)'_x + a\psi\chi\varphi''_{xx} + (f\psi - \psi''_{tt})\varphi + f\chi + g - \chi''_{tt} = 0.$$

This equation can be represented as a functional equation (25) in which

$$\begin{aligned} \Phi_1 &= (\varphi\varphi'_x)'_x, & \Phi_2 &= \varphi''_{xx}, & \Phi_3 &= \varphi, & \Phi_4 &= 1, \\ \Psi_1 &= a\psi^2, & \Psi_2 &= a\psi\chi, & \Psi_3 &= f\psi - \psi''_{tt}, & \Psi_4 &= f\chi + g - \chi''_{tt}. \end{aligned} \quad (29)$$

On substituting (29) into (26a), we obtain the following overdetermined system of ordinary differential equations for the functions  $\varphi = \varphi(x)$ ,  $\psi = \psi(t)$ , and  $\chi = \chi(t)$ :

$$\begin{aligned} (\varphi\varphi'_x)'_x &= A_1\varphi + A_2, & \varphi''_{xx} &= A_3\varphi + A_4, \\ f\psi - \psi''_{tt} &= -A_1a\psi^2 - A_3a\psi\chi, & f\chi + g - \chi''_{tt} &= -A_2a\psi^2 - A_4a\psi\chi. \end{aligned} \quad (30)$$

The first two equations in (30) are consistent only if

$$A_1 = 6B_2, \quad A_2 = B_1^2 - 4B_0B_2, \quad A_3 = 0, \quad A_4 = 2B_2, \quad (31)$$

where  $B_0$ ,  $B_1$ , and  $B_2$  are arbitrary constants, and the solution is given by

$$\varphi(x) = B_2x^2 + B_1x + B_0. \quad (32)$$

On substituting the expressions (31) into the last two equations in (30), we obtain the following system of equations for  $\psi(t)$  and  $\chi(t)$ :

$$\begin{aligned} \psi''_{tt} &= 6aB_2\psi^2 + f(t)\psi, \\ \chi''_{tt} &= [2aB_2\psi + f(t)]\chi + a(B_1^2 - 4B_0B_2)\psi^2 + g(t), \end{aligned} \quad (33)$$

Relations (28), (32) and system (33) determine a generalized separable solution of equation (27). The first equation in (33) can be solved independently; it is linear if  $B_2 = 0$  and is integrable in quadrature for  $f(t) = \text{const}$ . The second equation in (33) is linear in  $\chi$  (for  $\psi$  known).

Equation (27) does not have other solutions with the form (28) if  $f$  and  $g$  are arbitrary function and  $\varphi \not\equiv 0$ ,  $\psi \not\equiv 0$ , and  $\chi \not\equiv 0$ .

*Remark.* It can be shown that equation (27) has a more general solution with the form

$$w(x, y) = \varphi_1(x)\psi_1(t) + \varphi_2(x)\psi_2(t) + \psi_3(t), \quad \varphi_1(x) = x^2, \quad \varphi_2(x) = x, \quad (34)$$

where the functions  $\psi_i = \psi_i(t)$  are determined by the ordinary differential equations

$$\begin{aligned} \psi_1'' &= 6a\psi_1^2 + f(t)\psi_1, \\ \psi_2'' &= [6a\psi_1 + f(t)]\psi_2, \\ \psi_3'' &= [2a\psi_1 + f(t)]\psi_3 + a\psi_2^2 + g(t). \end{aligned} \quad (35)$$

(The prime denotes the derivative with respect to  $t$ .) The second equation in (35) has a particular solution  $\psi_2 = \psi_1$ . Hence, its general solution can be represented as (see Polyanin and Zaitsev, 1995)

$$\psi_2 = C_1\psi_1 + C_2\psi_1 \int \frac{dt}{\psi_1^2}.$$

The solution obtained in Example 3 corresponds to the special case  $C_2 = 0$ .

**Example 4.** Consider the nonlinear equation

$$\frac{\partial^2 w}{\partial x \partial t} + \left( \frac{\partial w}{\partial x} \right)^2 - w \frac{\partial^2 w}{\partial x^2} = \nu \frac{\partial^3 w}{\partial x^3}, \quad (36)$$

which arises in hydrodynamics (see equations B.6.2.1, Item 3° and B.7.2.1, Item 2°).

We look for exact solutions of the form

$$w = \varphi(t)\theta(x) + \psi(t). \quad (37)$$



Substituting (37) into (36) yields

$$\varphi'_t \theta'_x - \varphi \psi \theta''_{xx} + \varphi^2 [(\theta'_x)^2 - \theta \theta''_{xx}] - \nu \varphi \theta'''_{xxx} = 0.$$

This functional differential equation can be reduced to the functional equation (25) by setting

$$\begin{aligned} \Phi_1 &= \varphi'_t, & \Phi_2 &= \varphi \psi, & \Phi_3 &= \varphi^2, & \Phi_4 &= \nu \varphi, \\ \Psi_1 &= \theta'_x, & \Psi_2 &= -\theta''_{xx}, & \Psi_3 &= (\theta'_x)^2 - \theta \theta''_{xx}, & \Psi_4 &= -\theta'''_{xxx}. \end{aligned}$$

On substituting these expressions into (26a), we obtain the system of equations

$$\begin{aligned} \varphi'_t &= A_1 \varphi^2 + A_2 \nu \varphi, & \varphi \psi &= A_3 \varphi^2 + A_4 \nu \varphi, \\ (\theta'_x)^2 - \theta \theta''_{xx} &= -A_1 \theta'_x + A_3 \theta''_{xx}, & \theta'''_{xxx} &= A_2 \theta'_x - A_4 \theta''_{xx}. \end{aligned} \quad (38)$$

It can be shown that the last two equations in (38) are consistent only if the function  $\theta$  and its derivative are linearly dependent,

$$\theta'_x = B_1 \theta + B_2. \quad (39)$$

The six constants  $B_1, B_2, A_1, A_2, A_3$ , and  $A_4$  must satisfy the three conditions

$$\begin{aligned} B_1(A_1 + B_2 - A_3 B_1) &= 0, \\ B_2(A_1 + B_2 - A_3 B_1) &= 0, \\ B_1^2 + A_4 B_1 - A_2 &= 0. \end{aligned} \quad (40)$$

Integrating (39) yields

$$\theta = \begin{cases} B_3 \exp(B_1 x) - \frac{B_2}{B_1} & \text{if } B_1 \neq 0, \\ B_2 x + B_3 & \text{if } B_1 = 0, \end{cases} \quad (41)$$

where  $B_3$  is an arbitrary constant.

The first two equations in (38) lead to the following expressions for  $\varphi$  and  $\psi$ :

$$\varphi = \begin{cases} \frac{A_2 \nu}{C \exp(-A_2 \nu t) - A_1} & \text{if } A_2 \neq 0, \\ -\frac{1}{A_1 t + C} & \text{if } A_2 = 0, \end{cases} \quad \psi = A_3 \varphi + A_4 \nu, \quad (42)$$

where  $C$  is an arbitrary constant.

Formulas (41), (42) and relations (40) allow us to find the following solutions of equation (36) with the form (37):

$$\begin{aligned} w &= \frac{x + C_1}{t + C_2} + C_3 & \text{if } A_2 &= B_1 = 0, B_2 = -A_1; \\ w &= \frac{C_1 e^{-\lambda x} + 1}{\lambda t + C_2} + \nu \lambda & \text{if } A_2 &= 0, B_1 = -A_4, B_2 = -A_1 - A_3 A_4; \\ w &= C_1 e^{-\lambda(x + \beta \nu t)} + \nu(\lambda + \beta) & \text{if } A_1 &= A_3 = B_2 = 0, A_2 = B_1^2 + A_4 B_1; \\ w &= \frac{\nu \beta + C_1 e^{-\lambda x}}{1 + C_2 e^{-\nu \lambda \beta t}} + \nu(\lambda - \beta) & \text{if } A_1 &= A_3 B_1 - B_2, A_2 = B_1^2 + A_4 B_1, \end{aligned}$$

where  $C_1, C_2, C_3, \beta$ , and  $\lambda$  are arbitrary constants (these can be expressed in terms of the  $A_k$  and  $B_k$ ).

The analysis of the second degenerate solution (26b) of the functional equation (25) leads to the following two more general solutions of the differential equation (36):

$$\begin{aligned} w &= \frac{x}{t + C_1} + \psi(t), \\ w &= \varphi(t) e^{-\lambda x} - \frac{\varphi'_t(t)}{\lambda \varphi(t)} + \nu \lambda, \end{aligned}$$

where  $\varphi(t)$  and  $\psi(t)$  are arbitrary functions, and  $C_1$  and  $\lambda$  are arbitrary constants.

## B.2.4. Simplified Scheme for Constructing Exact Solutions of Equations with Quadratic Nonlinearities

### B.2.4-1. Description of the simplified scheme.

To construct exact solutions of equations (2) with quadratic or power nonlinearities that do not depend explicitly on  $x$  (all  $f_i$  constant), it is reasonable to use the following simplified approach. As

before, we seek solutions in the form of finite sums (1). We assume that the system of coordinate functions  $\{\varphi_i(x)\}$  is governed by linear differential equations with constant coefficients. The most common solutions of such equations are of the form

$$\varphi_i(x) = x^i, \quad \varphi_i(x) = e^{\lambda_i x}, \quad \varphi_i(x) = \sin(\alpha_i x), \quad \varphi_i(x) = \cos(\beta_i x). \quad (43)$$

Finite chains of these functions (in various combinations) can be used to search for separable solutions (1), where the quantities  $\lambda_i$ ,  $\alpha_i$ , and  $\beta_i$  are regarded as free parameters. The other system of functions  $\{\psi_i(y)\}$  is determined by solving the nonlinear equations resulting from substituting (1) into the equation under consideration.

This simplified approach lacks the generality of the methods outlined in Subsections B.2.2 and B.2.3. However, specifying one of the systems of coordinate functions,  $\{\varphi_i(x)\}$ , simplifies the procedure of finding exact solutions substantially. The drawback of this approach is that some solutions of the form (1) can be overlooked. It is significant that the overwhelming majority of generalized separable solutions known to date, for partial differential equations with quadratic nonlinearities, are determined by coordinate functions (36) (usually with  $n = 2$ ).

#### B.2.4-2. Examples of constructing exact solutions of higher-order equations.

Below we consider specific examples that illustrate the application of the above simplified scheme to constructing generalized separable solutions of higher-order nonlinear equations.

**Example 5.** The equations of laminar boundary layer on a flat plate are reduced to a single third-order nonlinear equation for the stream function (see Schlichting 1981, Loitsyanskiy 1996):

$$\frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = \nu \frac{\partial^3 w}{\partial y^3}. \quad (44)$$

We look for generalized separable solutions with the form

$$w(x, y) = x\psi(y) + \theta(y), \quad (45)$$

which corresponds to the simplest set of functions  $\varphi_1(x) = x$ ,  $\varphi_2(x) = 1$  with  $n = 2$  in formula (1). On substituting (45) into (44) and collecting terms, we obtain

$$x[(\psi')^2 - \psi\psi'' - \nu\psi'''] + [\psi'\theta' - \psi\theta'' - \nu\theta'''] = 0.$$

(The prime denotes the derivative with respect to  $y$ .) To meet this equation for any  $x$ , one should equate both expressions in square brackets to zero. This results in a system of ordinary differential equations for  $\psi = \psi(y)$  and  $\theta = \theta(y)$ :

$$\begin{aligned} (\psi')^2 - \psi\psi'' - \nu\psi''' &= 0, \\ \psi'\theta' - \psi\theta'' - \nu\theta''' &= 0. \end{aligned}$$

For example, this system has an exact solution

$$\psi = \frac{6\nu}{y + C_1}, \quad \theta = \frac{C_2}{y + C_1} + \frac{C_3}{(y + C_1)^2} + C_4,$$

where  $C_1$ ,  $C_2$ ,  $C_3$ , and  $C_4$  are arbitrary constants.

Other generalized separable solutions of equation (44) can be found in Polyanin (2001b, 2001c) and Subsection B.6.1.

**Example 6.** Consider the  $n$ th-order nonlinear equation

$$\frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = f(x) \frac{\partial^n w}{\partial y^n}, \quad (46)$$

where  $f(x)$  is an arbitrary function. In the special case  $n = 3$  with  $f(x) = \nu = \text{const}$ , this equation coincides with the boundary layer equation (44).

We look for generalized separable solutions of the form

$$w(x, y) = \varphi(x)e^{\lambda y} + \theta(x), \quad (47)$$

which correspond to the set of functions  $\psi_1(y) = e^{\lambda y}$ ,  $\psi_2(y) = 1$  in (1). On substituting (47) into (46) and rearranging terms, we obtain

$$\lambda^2 e^{\lambda y} \varphi[\theta'_x + \lambda^{n-2} f(x)] = 0.$$

This equation is met if

$$\theta(x) = -\lambda^{n-2} \int f(x) dx + C, \quad \varphi(x) \text{ is any}, \quad (48)$$

where  $C$  is an arbitrary constant. (The other case  $\varphi = 0$  and  $\theta$  is any is of little interest.) Formulas (47) and (48) define an exact solution of equation (46),

$$w(x, y) = \varphi(x)e^{\lambda y} - \lambda^{n-2} \int f(x) dx + C, \quad (49)$$

which involves an arbitrary function  $\varphi(x)$  and two arbitrary constants  $C$  and  $\lambda$ .

Note that solution (49) with  $n = 3$  and  $f(x) = \text{const}$  was obtained by Ignatovich (1993) by a more complicated approach.

## B.3. Methods of Functional Separation of Variables

### B.3.1. Structure of Functional Separable Solutions

#### B.3.1-1. Functional separable solutions.

Suppose a nonlinear equation for  $w = w(x, y)$  is obtained from a separable linear mathematical physics equation for  $z = z(x, y)$  by a nonlinear change of variable  $w = F(z)$ . Then, obviously, the former has exact solutions of the form

$$w(x, y) = F(z), \quad \text{where} \quad z = \sum_{m=1}^n \varphi_m(x) \psi_m(y). \quad (1)$$

It is noteworthy that many nonlinear partial differential equations that are not reduced to linear equation have exact solutions of the form (1) as well. We will call such solutions *functional separable solutions*. In general, the functions  $\varphi_m(x)$ ,  $\psi_m(y)$ , and  $F(z)$  in (1) are not known in advance and are to be identified.

*Remark 1.* In functional separation of variables, searching for solutions in the forms  $w = F(\varphi(x) + \psi(y))$  and  $w = F(\varphi(x)\psi(y))$  leads to equivalent results, because the two forms are functionally equivalent. Indeed, we have  $F(\varphi(x)\psi(y)) = F_1(\varphi_1(x) + \psi_1(y))$ , where  $F_1(z) = F(e^z)$ ,  $\varphi_1(x) = \ln \varphi(x)$ , and  $\psi_1(y) = \ln \psi(y)$ .

*Remark 2.* In constructing functional separable solutions with the form  $w = F(\varphi(x) + \psi(y))$ , it is assumed that  $\varphi \neq \text{const}$  and  $\psi \neq \text{const}$ .

#### B.3.1-2. Various modifications.

Below we give three more general modifications of solution structure (1):

$$w(x, y) = F(z), \quad z = \sum_{m=1}^n \varphi_m(\xi) \psi_m(\eta), \quad \xi = a_1 x + a_2 y, \quad \eta = b_1 x + b_2 y; \quad (2)$$

$$w(x, y) = \theta_1(x) F(z) + \theta_2(x), \quad z = \sum_{m=1}^n \varphi_m(x) \psi_m(y); \quad (3)$$

$$w(x, y) = \theta_1(y) F(z) + \theta_2(y), \quad z = \sum_{m=1}^n \varphi_m(x) \psi_m(y). \quad (4)$$

These can also be used to find exact solutions of nonlinear mathematical physics equations.

The solution structures of (1)–(4) cover all most common types of solutions—traveling wave, self-similar, and additively and multiplicatively separable solutions (as well as many invariant solutions). In general, the functions  $\varphi_m(\xi)$ ,  $\psi_m(\eta)$ ,  $\varphi_m(x)$ ,  $\psi_m(y)$ ,  $F(z)$ ,  $\theta_m(x)$ , and  $\theta_m(y)$  are not known in advance and are to be determined in the analysis.

Note that Miller and Rubel (1993) studied functional separable solutions of a different form (for a stationary heat equation with a nonlinear source); see also Clarkson and Kruskal (1989) and Burde (1994).

### B.3.2. Special Functional Separable Solutions

To simplify the analysis, some of the functions in (1) can be specified a priori and the other functions will be defined in the analysis. We call such solutions *special functional separable solutions*.

### B.3.2-1. Solutions of the form (1) with $z$ linear in one of the independent variables.

Consider functional separable solutions of the form (1) in the special case where the composite argument  $z$  is linear in one of the independent variables (e.g., in  $x$ ). We substitute (1) into the equation under study and eliminate  $x$  using the expression of  $z$  to obtain a functional differential equation with two arguments. In many cases, this equation can be solved by the methods outlined in Section B.2.

**Example 1.** Consider the nonstationary heat equation with a nonlinear source

$$\frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \mathcal{F}(w). \quad (5)$$

We look for functional separable solutions of the special form

$$w = w(z), \quad z = \varphi(t)x + \psi(t). \quad (6)$$

The functions  $w(z)$ ,  $\varphi(t)$ ,  $\psi(t)$ , and  $\mathcal{F}(w)$  are to be determined.

On substituting (6) into (5) and on dividing by  $w'_z$ , we have

$$\varphi'_t x + \psi'_t = \varphi^2 \frac{w''_{zz}}{w'_z} + \frac{\mathcal{F}(w)}{w'_z}. \quad (7)$$

We express  $x$  from (6) in terms of  $z$  and substitute into (7) to obtain a functional differential equation with two variables  $t$  and  $z$ ,

$$-\psi'_t + \frac{\psi}{\varphi} \varphi'_t - \frac{\varphi'_t}{\varphi} z + \varphi^2 \frac{w''_{zz}}{w'_z} + \frac{\mathcal{F}(w)}{w'_z} = 0,$$

which can be treated as the functional equation (25) in Section B.2 where

$$\begin{aligned} \Phi_1 &= -\psi'_t + \frac{\psi}{\varphi} \varphi'_t, & \Phi_2 &= -\frac{\varphi'_t}{\varphi}, & \Phi_3 &= \varphi^2, & \Phi_4 &= 1, \\ \Psi_1 &= 1, & \Psi_2 &= z, & \Psi_3 &= \frac{w''_{zz}}{w'_z}, & \Psi_4 &= \frac{\mathcal{F}(w)}{w'_z}. \end{aligned}$$

Substituting these expressions into relations (26a) of Section B.2 yields the system of ordinary differential equations

$$\begin{aligned} -\psi'_t + \frac{\psi}{\varphi} \varphi'_t &= A_1 \varphi^2 + A_2, & -\frac{\varphi'_t}{\varphi} &= A_3 \varphi^2 + A_4, \\ \frac{w''_{zz}}{w'_z} &= -A_1 - A_3 z, & \frac{\mathcal{F}(w)}{w'_z} &= -A_2 - A_4 z, \end{aligned} \quad (8)$$

where  $A_1$ ,  $A_2$ ,  $A_3$ , and  $A_4$  are arbitrary constants.

The solution of system (8) is given by

$$\begin{aligned} \varphi(t) &= \pm \left( C_1 e^{-2A_4 t} - \frac{A_3}{A_4} \right)^{-1/2}, \\ \psi(t) &= -\varphi(t) \left[ A_1 \int \varphi(t) dt + A_2 \int \frac{dt}{\varphi(t)} + C_2 \right], \\ w(z) &= C_3 \int \exp\left(-\frac{1}{2} A_3 z^2 - A_1 z\right) dz + C_4, \\ \mathcal{F}(w) &= -C_3(A_4 z + A_2) \exp\left(-\frac{1}{2} A_3 z^2 - A_1 z\right), \end{aligned} \quad (9)$$

where  $C_1$ ,  $C_2$ ,  $C_3$ , and  $C_4$  are arbitrary constants. The dependence  $\mathcal{F} = \mathcal{F}(w)$  is defined by the last two relations in parametric form ( $z$  is considered the parameter).

In the special case  $A_3 = C_4 = 0$ ,  $A_1 = -1$ , and  $C_3 = 1$ , the source function can be represented in explicit form as

$$\mathcal{F}(w) = -w(A_4 \ln w + A_2). \quad (10)$$

If  $A_3 \neq 0$  in (9), the source function is expressed in terms of elementary functions and the inverse of the error function.

**Example 2.** Consider the more general equation

$$\frac{\partial w}{\partial t} = a(t) \frac{\partial^2 w}{\partial x^2} + b(t) \frac{\partial w}{\partial x} + c(t) \mathcal{F}(w).$$

We look for solutions in the form (6). In this case, only the first two equations in system (8) will change, and the functions  $w(z)$  and  $\mathcal{F}(w)$  will be given by (9).

**Example 3.** The nonlinear heat equation

$$\frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ \mathcal{G}(w) \frac{\partial w}{\partial x} \right] + \mathcal{F}(w)$$

has also solutions of the form (6). The unknown quantities are governed by system (8) in which  $w''_{zz}$  must be replaced by  $[\mathcal{G}(w)w'_z]_z$ . The functions  $\varphi(t)$  and  $\psi(t)$  are determined by the first two formulas in (9). One of the two functions  $\mathcal{G}(w)$  and  $\mathcal{F}(w)$  can be assumed arbitrary and the other is identified in the course of the solution. The special case  $\mathcal{F}(w) = \text{const}$  yields  $\mathcal{G}(w) = C_1 e^{2kw} + (C_2 w + C_3) e^{kw}$ .

**Example 4.** Likewise, we can treat the  $n$ th-order nonlinear equation

$$\frac{\partial w}{\partial t} = \frac{\partial^n w}{\partial x^n} + \mathcal{F}(w).$$

As before, we look for solutions in the form (6). In this case, the quantities  $\varphi^2$  and  $w''_{zz}$  in (8) must be replaced by  $\varphi^n$  and  $w_z^{(n)}$ , respectively. In particular, for  $A_3 = 0$ , apart from equations with logarithmic nonlinearities of the form (10), we obtain other equations.

**Example 5.** For the  $n$ th-order nonlinear equation

$$\frac{\partial w}{\partial t} = \frac{\partial^n w}{\partial x^n} + \mathcal{F}(w) \frac{\partial w}{\partial x},$$

the search for exact solutions of the form (6) leads to the following system of equations for  $\varphi(t)$ ,  $\psi(t)$ ,  $w(z)$ , and  $\mathcal{F}(w)$ :

$$\begin{aligned} -\psi'_t + \frac{\psi}{\varphi} \varphi'_t &= A_1 \varphi^n + A_2 \varphi, & -\frac{\varphi'_t}{\varphi} &= A_3 \varphi^n + A_4 \varphi, \\ \frac{w_z^{(n)}}{w'_z} &= -A_1 - A_3 z, & \mathcal{F}(w) &= -A_2 - A_4 z, \end{aligned}$$

where  $A_1$ ,  $A_2$ ,  $A_3$ , and  $A_4$  are arbitrary constants.

In the case  $n = 3$ , we assume  $A_3 = 0$  and  $A_1 > 0$  to find in particular that  $\mathcal{F}(w) = -A_2 - A_4 \arcsin(kw)$ .

**Example 6.** In addition, searching for solutions of equation (5) with  $z$  quadratically dependent on  $x$ ,

$$w = w(z), \quad z = \varphi(t)x^2 + \psi(t), \tag{11}$$

also makes sense here. Indeed, on substituting (11) into (5), we arrive at an equation that contains terms with  $x^2$  and does not contain terms linear in  $x$ . Eliminating  $x^2$  from the resulting equation with the aid of (11), we obtain

$$-\psi'_t + \frac{\psi}{\varphi} \varphi'_t + 2\varphi - \frac{\varphi'_t}{\varphi} z + 4\varphi z \frac{w''_{zz}}{w'_z} - 4\varphi \psi \frac{w''_{zz}}{w'_z} + \frac{\mathcal{F}(w)}{w'_z} = 0.$$

To solve this functional differential equation with two arguments, we apply the splitting method outlined in Subsection B.2.3. It can be shown that, for equations (5), this equation has a solution with a logarithmic nonlinearity of the form (10).

### B.3.2-2. Solution by reduction to equations with quadratic nonlinearities.

In some cases, solutions of the form (1) can be searched for in two stages. First, one looks for a transformation that would reduce the original equation to an equation with a quadratic (or power) nonlinearity. Then the methods outlined in Section B.2 are used to find solutions of the resulting equation.

Sometimes, quadratically nonlinear equations can be obtained using the substitutions

$$\begin{aligned} w(z) &= z^\lambda && \text{(for equations with power nonlinearities),} \\ w(z) &= \lambda \ln z && \text{(for equations with exponential nonlinearities),} \\ w(z) &= e^{\lambda z} && \text{(for equations with logarithmic nonlinearities),} \end{aligned}$$

where  $\lambda$  is a constant to be determined. This approach is equivalent to specifying the form of the function  $F(z)$  in (1) a priori.

**Example 6.** The nonlinear heat equation with a logarithmic source

$$\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(t)w \ln w + g(t)w$$

can be reduced by the change of variable  $w = e^z$  to the quadratically nonlinear equation

$$\frac{\partial z}{\partial t} = a \frac{\partial^2 z}{\partial x^2} + a \left( \frac{\partial z}{\partial x} \right)^2 + f(t)z + g(t),$$

which admits separable solutions with the form

$$z = \varphi_1(x)\psi_1(t) + \varphi_2(x)\psi_2(t) + \psi_3(t),$$

where  $\varphi_1(x) = x^2$ ,  $\varphi_2(x) = x$ , and the functions  $\psi_k(t)$  are determined by an appropriate system of ordinary differential equations.

### B.3.3. Differentiation Method

#### B.3.3-1. Basic ideas of the method. Reduction to a standard equation.

In general, the substitution of expression (1) into the nonlinear partial differential equation under study leads to a functional differential equation with three arguments—two arguments are usual,  $x$  and  $y$ , and the third is composite,  $z$ . In many cases, the resulting equation can be reduced by differentiation to a standard functional differential equation with two arguments (either  $x$  or  $y$  is eliminated). To solve the two-argument equation, one can use the methods outlined in Section B.2.

#### B.3.3-2. Examples of constructing functional separable solutions.

Below we consider specific examples illustrating the application of the differentiation method for constructing functional separable solutions of nonlinear equations.

**Example 7.** Consider the nonlinear heat equation

$$\frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ f(w) \frac{\partial w}{\partial x} \right]. \quad (12)$$

We look for exact solutions with the form

$$w = w(z), \quad z = \varphi(x) + \psi(t). \quad (13)$$

On substituting (13) into (12) and dividing by  $w'_z$ , we obtain the functional differential equation

$$\psi'_t = \varphi''_{xx} f(w) + (\varphi'_x)^2 Q(z), \quad (14)$$

where

$$Q(z) = f(w) \frac{w''_{zz}}{w'_z} + f'_z(w), \quad w = w(z). \quad (15)$$

Differentiating (14) with respect to  $x$  yields

$$\varphi'''_{xxx} f(w) + \varphi'_x \varphi''_{xx} [f'_z(w) + 2Q(z)] + (\varphi'_x)^3 Q'_z = 0. \quad (16)$$

This functional differential equation with two variables can be treated as the functional equation (23) of Section B.2. This three-term functional equation has two different solutions. Accordingly, we consider two cases.

*Case 1.* The solutions of the functional differential equation (16) are determined from the system of ordinary differential equations

$$\begin{aligned} f'_z + 2Q &= 2A_1 f, & Q'_z &= A_2 f, \\ \varphi'''_{xxx} + 2A_1 \varphi'_x \varphi''_{xx} + A_2 (\varphi'_x)^3 &= 0, \end{aligned} \quad (17)$$

where  $A_1$  and  $A_2$  are arbitrary constants.

The first two equations (17) are linear and independent of the third equation. Their general solution is given by

$$f = \begin{cases} e^{A_1 z} (B_1 e^{kz} + B_2 e^{-kz}) & \text{if } A_1^2 > 2A_2, \\ e^{A_1 z} (B_1 + B_2 z) & \text{if } A_1^2 = 2A_2, \\ e^{A_1 z} [B_1 \sin(kz) + B_2 \cos(kz)] & \text{if } A_1^2 < 2A_2, \end{cases} \quad Q = A_1 f - \frac{1}{2} f'_z, \quad k = \sqrt{|A_1^2 - 2A_2|}. \quad (18)$$

Substituting  $Q$  of (18) into (15) yields a differential equation for  $w = w(z)$ . On integrating this equation, we obtain

$$w = C_1 \int e^{A_1 z} |f(z)|^{-3/2} dz + C_2, \quad (19)$$

where  $C_1$  and  $C_2$  are arbitrary constants. The expression of  $f$  in (18) together with expression (19) define the function  $f = f(w)$  in parametric form.

Without full analysis, we will study the case  $A_2 = 0$  ( $k = A_1$ ) and  $A_1 \neq 0$  in more detail. It follows from (18) and (19) that

$$f(z) = B_1 e^{2A_1 z} + B_2, \quad Q = A_1 B_2, \quad w(z) = C_3 (B_1 + B_2 e^{-2A_1 z})^{-1/2} + C_2 \quad (C_1 = A_1 B_2 C_3). \quad (20)$$

Eliminating  $z$  yields

$$f(w) = \frac{B_2 C_3^2}{C_3^2 - B_1 w^2}. \quad (21)$$

The last equation in (17) with  $A_2 = 0$  has the first integral  $\varphi'_{xx} + A_1 (\varphi'_x)^2 = \text{const}$ . The corresponding general solution is given by

$$\begin{aligned} \varphi(x) &= -\frac{1}{2A_1} \ln \left[ \frac{D_2}{D_1} \frac{1}{\sinh^2(A_1 \sqrt{D_2} x + D_3)} \right] & \text{for } D_1 > 0 \text{ and } D_2 > 0; \\ \varphi(x) &= -\frac{1}{2A_1} \ln \left[ -\frac{D_2}{D_1} \frac{1}{\cos^2(A_1 \sqrt{-D_2} x + D_3)} \right] & \text{for } D_1 > 0 \text{ and } D_2 < 0; \\ \varphi(x) &= -\frac{1}{2A_1} \ln \left[ -\frac{D_2}{D_1} \frac{1}{\cosh^2(A_1 \sqrt{D_2} x + D_3)} \right] & \text{for } D_1 < 0 \text{ and } D_2 > 0; \end{aligned} \quad (22)$$

where  $D_1$ ,  $D_2$ , and  $D_3$  are constants of integration. In all three cases, the following relations hold:

$$(\varphi'_x) = D_1 e^{-2A_1 \varphi} + D_2, \quad \varphi''_{xx} = -A_1 D_1 e^{-2A_1 \varphi}. \quad (23)$$

We substitute (20) and (23) into the original functional differential equation (14). With reference to the expression of  $z$  in (13), we obtain the following equation for  $\psi = \psi(t)$ :

$$\psi'_t = -A_1 B_1 D_1 e^{2A_1 \psi} + A_1 B_2 D_2.$$

Its general solution is given by

$$\psi(t) = \frac{1}{2A_1} \ln \frac{B_2 D_2}{D_4 \exp(-2A_1^2 B_2 D_2 t) + B_1 D_1}, \quad (24)$$

where  $D_4$  is an arbitrary constant.

Formulas (13), (20) for  $w$ , (22), and (24) define three solutions of the nonlinear equation (12) with  $f(w)$  of the form (21) [recall that these solutions correspond to the special case  $A_2 = 0$  in (18) and (19)].

**Case 2.** The solutions of the functional differential equation (16) are determined from the system of ordinary differential equations

$$\begin{aligned} \varphi'''_{xxx} &= A_1 (\varphi'_x)^3, & \varphi'_x \varphi''_{xx} &= A_2 (\varphi'_x)^3, \\ A_1 f + A_2 (f'_z + 2Q) + Q'_z &= 0. \end{aligned} \quad (25)$$

The first two equations in (25) are consistent in the two cases

$$\begin{aligned} A_1 = A_2 = 0 &\implies \varphi(x) = B_1 x + B_2, \\ A_1 = 2A_2^2 &\implies \varphi(x) = -\frac{1}{A_2} \ln |B_1 x + B_2|. \end{aligned} \quad (26)$$

The first solution in (26) eventually leads to the traveling wave solution  $w = w(B_1 x + B_2 t)$  of equation (12) and the second solution to the self-similar solution of the form  $w = \tilde{w}(x^2/t)$ . In both cases, the function  $f(w)$  in (12) is arbitrary.

A more detailed analysis of functional separable solutions (13) of equation (12) can be found in the reference cited below.

⊙ *Reference:* P. W. Doyle and P. J. Vassiliou (1998).

**Example 8.** One can look for more complicated functional separable solutions of equation (12) with the form

$$w = w(z), \quad z = \varphi(\xi) + \psi(t), \quad \xi = x + at \quad (a = \text{const}).$$

We substitute this into (12), divide the resulting functional differential equation by  $w'_z$ , and differentiate with respect to  $x$  to obtain

$$-a\varphi''_{\xi\xi} + \varphi'''_{\xi\xi\xi} f(w) + \varphi'_\xi \varphi''_{\xi\xi} [f'_z(w) + 2Q(z)] + (\varphi'_\xi)^3 Q'_z = 0,$$

where the function  $Q = Q(z)$  is defined by (15). This functional differential equation with two variables  $\xi$  and  $z$  can be treated as the functional equation (25) of Section B.2. The solution of (25) is given by relations (26), thus representing a system of ordinary differential equations for  $f$ ,  $Q$ , and  $\varphi$ .

**Example 9.** Consider the nonlinear Klein–Gordon equation

$$\frac{\partial^2 w}{\partial t^2} - \frac{\partial^2 w}{\partial x^2} = \mathcal{F}(w). \quad (27)$$

We look for functional separable solutions in additive form:

$$w = w(z), \quad z = \varphi(x) + \psi(t). \quad (28)$$

Substituting (28) into (27) yields

$$\psi_{tt}' - \varphi_{xx}'' + [(\psi_t')^2 - (\varphi_x')^2] g(z) = h(z), \quad (29)$$

where

$$g(z) = w_{zz}''/w_z', \quad h(z) = \mathcal{F}(w(z))/w_z'. \quad (30)$$

On differentiating (29) first with respect to  $t$  and then with respect to  $x$  and on dividing by  $\psi_t' \varphi_x'$ , we have

$$2(\psi_{tt}'' - \varphi_{xx}'') g_z' + [(\psi_t')^2 - (\varphi_x')^2] g_{zz}'' = h_{zz}''.$$

Eliminating  $\psi_{tt}'' - \varphi_{xx}''$  from this equation with the aid of (29), we obtain

$$[(\psi_t')^2 - (\varphi_x')^2] (g_{zz}'' - 2g g_z') = h_{zz}'' - 2g_z' h. \quad (31)$$

This relation holds in the following cases:

$$\begin{aligned} g_{zz}'' - 2g g_z' &= 0, \quad h_{zz}'' - 2g_z' h = 0 & (\text{case 1}), \\ (\psi_t')^2 &= A\psi + B, \quad (\varphi_x')^2 = -A\varphi + B - C, \quad h_{zz}'' - 2g_z' h = (Az + C)(g_{zz}'' - 2g g_z') & (\text{case 2}^*), \end{aligned} \quad (32)$$

where  $A$ ,  $B$ , and  $C$  are arbitrary constants. We consider both cases.

*Case 1.* The first two equations in (32) enable one to determine  $g(z)$  and  $h(z)$ . Integrating the first equation once yields  $g_z' = g^2 + \text{const}$ . Further, the following cases are possible:

$$g = k, \quad (33a)$$

$$g = -1/(z + C_1), \quad (33b)$$

$$g = -k \tanh(kz + C_1), \quad (33c)$$

$$g = -k \coth(kz + C_1), \quad (33d)$$

$$g = k \tan(kz + C_1), \quad (33e)$$

where  $C_1$  and  $k$  are arbitrary constants.

The second equation in (32) has the particular solution  $h = g(z)$ . Hence, its general solution is expressed by (e.g., see Polyanin and Zaitsev 1995)

$$h = C_2 g(z) + C_3 g(z) \int \frac{dz}{g^2(z)}, \quad (34)$$

where  $C_2$  and  $C_3$  are arbitrary constants.

The functions  $w(z)$  and  $\mathcal{F}(w)$  are found from (30) as

$$w(z) = B_1 \int G(z) dz + B_2, \quad \mathcal{F}(w) = B_1 h(z) G(z), \quad \text{where} \quad G(z) = \exp \left[ \int g(z) dz \right], \quad (35)$$

and  $B_1$  and  $B_2$  are arbitrary constants ( $\mathcal{F}$  is defined parametrically).

Let us dwell on the case (33b). According to (34),

$$h = A_1(z + C_1)^2 + \frac{A_2}{z + C_1}, \quad (36)$$

where  $A_1 = -C_3/3$  and  $A_2 = -C_2$  are any numbers. Substituting (33b) and (36) into (35) yields

$$w = B_1 \ln |z + C_1| + B_2, \quad \mathcal{F} = A_1 B_1 (z + C_1) + \frac{A_2 B_1}{(z + C_1)^2}.$$

Eliminating  $z$ , we arrive at the explicit form of the right-hand side of equation (27):

$$\mathcal{F}(w) = A_1 B_1 e^u + A_2 B_1 e^{-2u}, \quad \text{where} \quad u = \frac{w - B_2}{B_1}. \quad (37)$$

For simplicity, we set  $C_1 = 0$ ,  $B_1 = 1$ , and  $B_2 = 0$  and denote  $A_1 = a$  and  $A_2 = b$ . Thus, we have

$$w(z) = \ln |z|, \quad \mathcal{F}(w) = ae^w + be^{-2w}, \quad g(z) = -1/z, \quad h(z) = az^2 + b/z. \quad (38)$$

---

\* In case 2, equation (31) can be represented as the functional equation considered in Paragraph B.3.5-1.



TABLE B1

Nonlinear Klein–Gordon equations  $\partial_{tt}w - \partial_{xx}w = \mathcal{F}(w)$  admitting functional separable solutions of the form  $w = w(z)$ ,  $z = \varphi(x) + \psi(t)$ . Notation:  $A$ ,  $C_1$ , and  $C_2$  are arbitrary constants;  $\sigma = 1$  for  $z > 0$  and  $\sigma = -1$  for  $z < 0$

| No | Right-hand side $\mathcal{F}(w)$                                                   | Solution $w(z)$                          | Equations for $\psi(t)$ and $\varphi(x)$                                                                                                                   |
|----|------------------------------------------------------------------------------------|------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | $aw \ln w + bw$                                                                    | $e^z$                                    | $(\psi'_t)^2 = C_1 e^{-2\psi} + a\psi - \frac{1}{2}a + b + A,$<br>$(\varphi'_x)^2 = C_2 e^{-2\varphi} - a\varphi + \frac{1}{2}a + A$                       |
| 2  | $ae^w + be^{-2w}$                                                                  | $\ln  z $                                | $(\psi'_t)^2 = 2a\psi^3 + A\psi^2 + C_1\psi + C_2,$<br>$(\varphi'_x)^2 = -2a\varphi^3 + A\varphi^2 - C_1\varphi + C_2 + b$                                 |
| 3  | $a \sin w + b \left( \sin w \ln \tan \frac{w}{4} + 2 \sin \frac{w}{4} \right)$     | $4 \arctan e^z$                          | $(\psi'_t)^2 = C_1 e^{2\psi} + C_2 e^{-2\psi} + b\psi + a + A,$<br>$(\varphi'_x)^2 = -C_2 e^{2\varphi} - C_1 e^{-2\varphi} - b\varphi + A$                 |
| 4  | $a \sinh w + b \left( \sinh w \ln \tanh \frac{w}{4} + 2 \sinh \frac{w}{2} \right)$ | $2 \ln \left  \coth \frac{z}{2} \right $ | $(\psi'_t)^2 = C_1 e^{2\psi} + C_2 e^{-2\psi} - \sigma b\psi + a + A,$<br>$(\varphi'_x)^2 = C_2 e^{2\varphi} + C_1 e^{-2\varphi} + \sigma b\varphi + A$    |
| 5  | $a \sinh w + 2b \left( \sinh w \arctan e^{w/2} + \cosh \frac{w}{2} \right)$        | $2 \ln \left  \tan \frac{z}{2} \right $  | $(\psi'_t)^2 = C_1 \sin 2\psi + C_2 \cos 2\psi + \sigma b\psi + a + A,$<br>$(\varphi'_x)^2 = -C_1 \sin 2\varphi + C_2 \cos 2\varphi - \sigma b\varphi + A$ |

It remains to determine  $\psi(t)$  and  $\varphi(x)$ . We substitute (38) into the functional differential equation (29). Taking into account (28), we find

$$[\psi'''_{ttt}\psi - (\psi'_t)^2 - a\psi^3 - b] - [\varphi'''_{xxx}\varphi - (\varphi'_x)^2 + a\varphi^3] + (\psi'''_{tt} - 3a\psi^2)\varphi - \psi(\varphi'''_{xxx} + 3a\varphi^2) = 0. \quad (39)$$

Differentiating (39) with respect to  $t$  and  $x$  yields the separable equation\*

$$(\psi'''_{ttt} - 6a\psi\psi'_t)\varphi'_x - (\varphi'''_{xxx} + 6a\varphi\varphi'_x)\psi'_t = 0,$$

whose solution is determined by the ordinary differential equations

$$\psi'''_{ttt} - 6a\psi\psi'_t = A\psi'_t,$$

$$\varphi'''_{xxx} + 6a\varphi\varphi'_x = A\varphi'_x,$$

where  $A$  is the separation constant. Each equation can be integrated twice, thus resulting in

$$(\psi'_t)^2 = 2a\psi^3 + A\psi^2 + C_1\psi + C_2,$$

$$(\varphi'_x)^2 = -2a\varphi^3 + A\varphi^2 + C_3\varphi + C_4, \quad (40)$$

where  $C_1$ ,  $C_2$ ,  $C_3$ , and  $C_4$  are arbitrary constants. Eliminating the derivatives from (39) with the aid of (40), we find that the arbitrary constants are related by  $C_3 = -C_1$  and  $C_4 = C_2 + b$ . So, the functions  $\psi(t)$  and  $\varphi(x)$  are determined by the first-order nonlinear autonomous equations

$$(\psi'_t)^2 = 2a\psi^3 + A\psi^2 + C_1\psi + C_2,$$

$$(\varphi'_x)^2 = -2a\varphi^3 + A\varphi^2 - C_1\varphi + C_2 + b.$$

The solutions of these equations are expressed in terms of elliptic functions.

For the other cases in (33), the analysis is performed in a similar way. Table B1 presents the final results for the cases (33a)–(33e).

Case 2. Integrating the third and fourth equations in (32) yields

$$\begin{aligned} \psi &= \pm\sqrt{B}t + D_1, & \varphi &= \pm\sqrt{B-C}t + D_2 & \text{if } A &= 0; \\ \psi &= \frac{1}{4A}(At + D_1)^2 - \frac{B}{A}, & \varphi &= -\frac{1}{4A}(Ax + D_2)^2 + \frac{B-C}{A} & \text{if } A &\neq 0; \end{aligned} \quad (41)$$

where  $D_1$  and  $D_2$  are arbitrary constants. In both cases, the function  $\mathcal{F}(w)$  in equation (27) is arbitrary. The first row in (41) corresponds to the traveling wave solution  $w = w(kx + \lambda t)$ . The second row leads to a solution of the form  $w = w(x^2 - t^2)$ .

⊙ *References:* A. M. Grundland and E. Infeld (1992), J. Miller and L. A. Rubel (1993), R. Z. Zhdanov (1994), V. K. Andreev, O. V. Kaptsov, V. V. Pukhnachev, and A. A. Rodionov (1994).

**Example 10.** The nonlinear stationary heat (diffusion) equation

$$\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = \mathcal{F}(w)$$

is analyzed just as the nonlinear Klein–Gordon equation considered in Example 9. The final results are listed in Table B2; the traveling wave solutions  $w = w(kx + \lambda t)$  and solutions of the form  $w = w(x^2 + y^2)$ , existing for any  $\mathcal{F}(w)$ , are omitted.

⊙ *References:* A. M. Grundland and E. Infeld (1992), J. Miller and L. A. Rubel (1993), R. Z. Zhdanov (1994), V. K. Andreev, O. V. Kaptsov, V. V. Pukhnachev, and A. A. Rodionov (1994).

\* To solve equation (39), one can use the solution of equation (25) in Section B.2 [see (26a)].

TABLE B2

Nonlinear equations  $\partial_{xx}w + \partial_{yy}w = \mathcal{F}(w)$  admitting functional separable solutions of the form  $w = w(z)$ ,  $z = \varphi(x) + \psi(y)$ . Notation:  $A$ ,  $C_1$ , and  $C_2$  are arbitrary constants;  $\sigma = 1$  for  $z > 0$ ,  $\sigma = -1$  for  $z < 0$

| No | Right-hand side $\Phi(w)$                                                          | Solution $w(z)$                          | Equations for $\varphi(x)$ and $\psi(y)$                                                                                                                  |
|----|------------------------------------------------------------------------------------|------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | $aw \ln w + bw$                                                                    | $e^z$                                    | $(\varphi'_x)^2 = C_1 e^{-2\varphi} + a\varphi - \frac{1}{2}a + b + A,$<br>$(\psi'_y)^2 = C_2 e^{-2\psi} + a\psi - \frac{1}{2}a - A$                      |
| 2  | $ae^w + be^{-2w}$                                                                  | $\ln z $                                 | $(\varphi'_x)^2 = 2a\varphi^3 + A\varphi^2 + C_1\varphi + C_2,$<br>$(\psi'_y)^2 = 2a\psi^3 - A\psi^2 + C_1\psi - C_2 - b$                                 |
| 3  | $a \sin w + b \left( \sin w \ln \tan \frac{w}{4} + 2 \sin \frac{w}{4} \right)$     | $4 \arctan e^z$                          | $(\varphi'_x)^2 = C_1 e^{2\varphi} + C_2 e^{-2\varphi} + b\varphi + a + A,$<br>$(\psi'_y)^2 = C_2 e^{2\psi} + C_1 e^{-2\psi} + b\psi - A$                 |
| 4  | $a \sinh w + b \left( \sinh w \ln \tanh \frac{w}{4} + 2 \sinh \frac{w}{2} \right)$ | $2 \ln \left  \coth \frac{z}{2} \right $ | $(\varphi'_x)^2 = C_1 e^{2\varphi} + C_2 e^{-2\varphi} - \sigma b\varphi + a + A,$<br>$(\psi'_y)^2 = -C_2 e^{2\psi} - C_1 e^{-2\psi} - \sigma b\psi - A$  |
| 5  | $a \sinh w + 2b \left( \sinh w \arctan e^{w/2} + \cosh \frac{w}{2} \right)$        | $2 \ln \left  \tan \frac{z}{2} \right $  | $(\varphi'_x)^2 = C_1 \sin 2\varphi + C_2 \cos 2\varphi + \sigma b\varphi + a + A,$<br>$(\psi'_y)^2 = C_1 \sin 2\psi - C_2 \cos 2\psi + \sigma b\psi - A$ |

### B.3.4. Splitting Method. Reduction to a Functional Equation with Two Variables

#### B.3.4-1. Splitting method. Reduction to a standard functional equation.

The general procedure for constructing functional separable solutions, which is based on the splitting method, involves several stages outlined below.

1°. Substitute expression (1) into the nonlinear partial differential equation under study. This results in a functional differential equation with three arguments—the first two are usual,  $x$  and  $y$ , and the third is composite,  $z$ .

2°. Reduce the functional differential equation to a purely functional equation with three arguments  $x$ ,  $y$ , and  $z$  with the aid of elementary differential substitutions (by selecting and renaming terms with derivatives).

3°. Reduce the three-argument functional differential equation by the differentiation method to the standard functional equation with two arguments (either  $x$  or  $y$  is eliminated) considered in Section B.2.

4°. Construct the solution of the two-argument functional equation using the formulas given in Subsection B.2.3.

5°. Solve the (overdetermined) system formed by the solution of Item 4° and the differential substitutions of Item 2°.

6°. Substitute the solution of Item 5° into the original functional differential equation of Item 1° to establish the relations for the constants of integration and determine all unknown quantities.

7°. Consider all degenerate cases possibly arising due to violation of assumptions adopted in the previous analysis.

The splitting method reduces solving the three-argument functional differential equation to (i) solving a purely functional equation with three arguments (by reducing it to a standard functional equation with two arguments) and (ii) solving a system of ordinary differential equations. Thus, the initial problem splits into several simpler problems. Examples of constructing functional separable solutions by the splitting method are given in Subsection B.3.5.

### B.3.4-2. Three-argument functional equations of special form.

The substitution of expression (1) with  $n = 2$  into nonlinear partial differential equation often leads to functional differential equations of the form

$$\begin{aligned} &\Phi_1(x)\Psi_1(y, z) + \Phi_2(x)\Psi_2(y, z) + \cdots + \Phi_k(x)\Psi_k(y, z) \\ &+ \Psi_{k+1}(y, z) + \Psi_{k+2}(y, z) + \cdots + \Psi_n(y, z) = 0, \end{aligned} \quad (42)$$

where  $\Phi_j(x)$  and  $\Psi_j(y, z)$  are functionals dependent on the variables  $x$  and  $y, z$ , respectively,

$$\Phi_j(x) \equiv \Phi_j(x, \varphi, \varphi'_x, \varphi''_{xx}), \quad \Psi_j(y, z) \equiv \Psi_j(y, \psi, \psi'_y, \psi''_{yy}, F, F'_z, F''_{zz}). \quad (43)$$

(These expressions correspond to a second-order equation.)

It is reasonable to solve equation (42) by the splitting method. To this end, we treat (42) at the first stage as a purely functional equation, thus disregarding (43). Assuming that  $\Psi_1 \neq 0$ , we divide (42) by  $\Psi_1$  and differentiate with respect to  $y$  to obtain a similar equation but with fewer terms:

$$\Phi_2(x)\Psi_2^{(2)}(y, z) + \cdots + \Phi_k(x)\Psi_k^{(2)}(y, z) + \Psi_{k+1}^{(2)}(y, z) + \cdots + \Psi_n^{(2)}(y, z) = 0, \quad (44)$$

where  $\Psi_m^{(2)} = \frac{\partial}{\partial y}(\Psi_m/\Psi_1) + \psi'_y \frac{\partial}{\partial z}(\Psi_m/\Psi_1)$ . We continue this procedure until we arrive at an equation independent of  $x$  explicitly:

$$\Psi_{k+1}^{(k+1)}(y, z) + \cdots + \Psi_n^{(k+1)}(y, z) = 0, \quad (45)$$

where  $\Psi_m^{(k+1)} = \frac{\partial}{\partial y}(\Psi_m^{(k)}/\Psi_k^{(k)}) + \psi'_y \frac{\partial}{\partial z}(\Psi_m^{(k)}/\Psi_k^{(k)})$ .

Relation (45) can be regarded as an equation with two independent variables  $y$  and  $z$ . If  $\Psi_m^{(k+1)}(y, z) = Q_m(y)R_m(z)$  for all  $m = k+1, \dots, n$ , then equation (45) can be solved using the results of Section B.2.

## B.3.5. Some Functional Equations and Their Solutions. Exact Solutions of Heat and Wave Equations

In this subsection, we discuss several types of three-argument functional equations that arise most frequently in functional separation of variables in nonlinear equations of mathematical physics. The results are used to construct exact solutions for some classes of nonlinear heat and wave equations.

### B.3.5-1. The functional equation $f(x) + g(y) = Q(z)$ , where $z = \varphi(x) + \psi(y)$ .

Here, one of the two functions  $f(x)$  and  $\varphi(x)$  is prescribed and the other is assumed unknown, also one of the functions  $g(y)$  and  $\psi(y)$  is prescribed and the other is unknown, and the function  $Q(z)$  is assumed unknown.\*

Differentiating the equation with respect to  $x$  and  $y$  yields  $Q''_{zz} = 0$ . Consequently, the solution is given by

$$f(x) = A\varphi(x) + B, \quad g(y) = A\psi(y) - B + C, \quad Q(z) = Az + C, \quad (46)$$

where  $A, B$ , and  $C$  are arbitrary constants.

### B.3.5-2. The functional equation $f(t) + g(x) + h(x)Q(z) + R(z) = 0$ , where $z = \varphi(x) + \psi(t)$ .

Differentiating the equation with respect to  $x$  yields the two-argument equation

$$g'_x + h'_x Q + h\varphi'_x Q'_z + \varphi'_x R'_z = 0. \quad (47)$$

\* In similar equations with a composite argument, it is assumed that  $\varphi(x) \neq \text{const}$  and  $\psi(y) \neq \text{const}$ .

Such equations were discussed in Section B.2. Hence, the following relations hold [see formulas (25) and (26a) in Section B.2]:

$$\begin{aligned} g'_x &= A_1 h \varphi'_x + A_2 \varphi'_x, \\ h'_x &= A_3 h \varphi'_x + A_4 \varphi'_x, \\ Q'_z &= -A_1 - A_3 Q, \\ R'_z &= -A_2 - A_4 Q, \end{aligned} \quad (48)$$

where  $A_1, A_2, A_3$ , and  $A_4$  are arbitrary constants. By integrating system (48) and substituting the resulting solutions into the original functional equation, one obtains the results given below.

*Case 1.* If  $A_3 = 0$  in (48), the corresponding solution of the functional equation is given by

$$\begin{aligned} f &= -\frac{1}{2} A_1 A_4 \psi^2 + (A_1 B_1 + A_2 + A_4 B_3) \psi - B_2 - B_1 B_3 - B_4, \\ g &= \frac{1}{2} A_1 A_4 \varphi^2 + (A_1 B_1 + A_2) \varphi + B_2, \\ h &= A_4 \varphi + B_1, \\ Q &= -A_1 z + B_3, \\ R &= \frac{1}{2} A_1 A_4 z^2 - (A_2 + A_4 B_3) z + B_4, \end{aligned} \quad (49)$$

where the  $A_k$  and  $B_k$  are arbitrary constants and  $\varphi = \varphi(x)$  and  $\psi = \psi(t)$  are arbitrary functions.

*Case 2.* If  $A_3 \neq 0$  in (48), the corresponding solution of the functional equation is

$$\begin{aligned} f &= -B_1 B_3 e^{-A_3 \psi} + \left( A_2 - \frac{A_1 A_4}{A_3} \right) \psi - B_2 - B_4 - \frac{A_1 A_4}{A_3^2}, \\ g &= \frac{A_1 B_1}{A_3} e^{A_3 \varphi} + \left( A_2 - \frac{A_1 A_4}{A_3} \right) \varphi + B_2, \\ h &= B_1 e^{A_3 \varphi} - \frac{A_4}{A_3}, \\ Q &= B_3 e^{-A_3 z} - \frac{A_1}{A_3}, \\ R &= \frac{A_4 B_3}{A_3} e^{-A_3 z} + \left( \frac{A_1 A_4}{A_3} - A_2 \right) z + B_4, \end{aligned} \quad (50)$$

where the  $A_k$  and  $B_k$  are arbitrary constants and  $\varphi = \varphi(x)$  and  $\psi = \psi(t)$  are arbitrary functions.

*Case 3.* In addition, the functional equation has the two degenerate solutions:

$$f = A_1 \psi + B_1, \quad g = A_1 \varphi + B_2, \quad h = A_2, \quad R = -A_1 z - A_2 Q - B_1 - B_2, \quad (51a)$$

where  $\varphi = \varphi(x)$ ,  $\psi = \psi(t)$ , and  $Q = Q(z)$  are arbitrary functions,  $A_1, A_2, B_1$ , and  $B_2$  are arbitrary constants; and

$$f = A_1 \psi + B_1, \quad g = A_1 \varphi + A_2 h + B_2, \quad Q = -A_2, \quad R = -A_1 z - B_1 - B_2, \quad (51b)$$

where  $\varphi = \varphi(x)$ ,  $\psi = \psi(t)$ , and  $h = h(x)$  are arbitrary functions,  $A_1, A_2, B_1$ , and  $B_2$  are arbitrary constants. The degenerate solutions (51a) and (51b) can be obtained directly from the original equation or its consequence (47) using formulas (26b) in Section B.2.

**Example 11.** Consider the nonstationary heat equation with a nonlinear source

$$\frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial x^2} + \mathcal{F}(w). \quad (52)$$

We look for exact solutions of the form

$$w = w(z), \quad z = \varphi(x) + \psi(t). \quad (53)$$

Substituting (53) into (52) and dividing by  $w'_z$  yields the functional differential equation

$$\psi'_t = \varphi''_{xx} + (\varphi'_x)^2 \frac{w''_{zz}}{w'_z} + \frac{\mathcal{F}(w(z))}{w'_z}.$$

We rewrite it as the functional equation B.3.5-2 in which

$$f(t) = -\psi'_t, \quad g(x) = \varphi''_{xx}, \quad h(x) = (\varphi'_x)^2, \quad Q(z) = w''_{zz}/w'_z, \quad R(z) = f(w(z))/w'_z. \quad (54)$$

We now use the solutions of equation B.3.5-2. On substituting the expressions of  $g$  and  $h$  of (54) into (49)–(51), we arrive at overdetermined systems of equations for  $\varphi = \varphi(x)$ .

*Case 1.* The system

$$\begin{aligned} \varphi''_{xx} &= \frac{1}{2} A_1 A_4 \varphi^2 + (A_1 B_1 + A_2) \varphi + B_2, \\ (\varphi'_x)^2 &= A_4 \varphi + B_1 \end{aligned}$$

following from (49) and corresponding to  $A_3 = 0$  in (48) is consistent in the cases

$$\begin{aligned} \varphi &= C_1 x + C_2 & \text{for } A_2 &= -A_1 C_1^2, \quad A_4 = B_2 = 0, \quad B_1 = C_1^2, \\ \varphi &= \frac{1}{4} A_4 x^2 + C_1 x + C_2 & \text{for } A_1 &= A_2 = 0, \quad B_1 = C_1^2 - A_4 C_2, \quad B_2 = \frac{1}{2} A_4, \end{aligned} \quad (55)$$

where  $C_1$  and  $C_2$  are arbitrary constants.

The first solution in (55) with  $A_1 \neq 0$  leads to a right-hand side of equation (52) containing the inverse of the error function [the form of the right-hand side is identified from the last two relations in (49) and (54)]. The second solution in (55) corresponds to the right-hand side  $\mathcal{F}(w) = k_1 w \ln w + k_2 w$  in (52). In both cases, the first relation in (49) is, taking into account that  $f = -\psi'_t$ , a first-order linear solution with constant coefficients, whose solution is an exponential plus a constant.

*Case 2.* The system

$$\begin{aligned} \varphi''_{xx} &= \frac{A_1 B_1}{A_3} e^{A_3 \varphi} + \left( A_2 - \frac{A_1 A_4}{A_3} \right) \varphi + B_2, \\ (\varphi'_x)^2 &= B_1 e^{A_3 \varphi} - \frac{A_4}{A_3}, \end{aligned}$$

following from (50) and corresponding to  $A_3 \neq 0$  in (48) is consistent in the following cases:

$$\begin{aligned} \varphi &= \pm \sqrt{-A_4/A_3} x + C_1 & \text{for } A_2 &= A_1 A_4/A_3, \quad B_1 = B_2 = 0, \\ \varphi &= -\frac{2}{A_3} \ln|x| + C_1 & \text{for } A_1 &= \frac{1}{2} A_3^2, \quad A_2 = A_4 = B_2 = 0, \quad B_1 = 4 A_3^{-2} e^{-A_3 C_1}, \\ \varphi &= -\frac{2}{A_3} \ln \left| \cos \left( \frac{1}{2} \sqrt{A_3 A_4} x + C_1 \right) \right| + C_2 & \text{for } A_1 &= \frac{1}{2} A_3^2, \quad A_2 = \frac{1}{2} A_3 A_4, \quad B_2 = 0, \quad A_3 A_4 > 0, \\ \varphi &= -\frac{2}{A_3} \ln \left| \sinh \left( \frac{1}{2} \sqrt{-A_3 A_4} x + C_1 \right) \right| + C_2 & \text{for } A_1 &= \frac{1}{2} A_3^2, \quad A_2 = \frac{1}{2} A_3 A_4, \quad B_2 = 0, \quad A_3 A_4 < 0, \\ \varphi &= -\frac{2}{A_3} \ln \left| \cosh \left( \frac{1}{2} \sqrt{-A_3 A_4} x + C_1 \right) \right| + C_2 & \text{for } A_1 &= \frac{1}{2} A_3^2, \quad A_2 = \frac{1}{2} A_3 A_4, \quad B_2 = 0, \quad A_3 A_4 < 0, \end{aligned}$$

where  $C_1$  and  $C_2$  are arbitrary constants. The right-hand sides of equation (52) corresponding to these solutions are represented in parametric form.

*Case 3.* Traveling wave solutions of the nonlinear heat equation (52) and solutions of the linear equation (52) with  $\mathcal{F}'_w = \text{const}$  correspond to the degenerate solutions of the functional equation (51).

**Example 12.** Likewise, one can analyze the more general equation

$$\frac{\partial w}{\partial t} = a(x) \frac{\partial^2 w}{\partial x^2} + b(x) \frac{\partial w}{\partial x} + \mathcal{F}(w). \quad (56)$$

It arises in convective heat/mass exchange problems ( $a = \text{const}$  and  $b = \text{const}$ ), problems of heat transfer in inhomogeneous media ( $b = a'_x \neq \text{const}$ ), and spatial heat transfer problems with axial or central symmetry ( $a = \text{const}$  and  $b = \text{const}/x$ ).

Searching for exact solutions of equation (56) in the form (53) leads to the functional equation B.3.5-2 in which

$$f(t) = -\psi'_t, \quad g(x) = a(x) \varphi''_{xx} + b(x) \varphi'_x, \quad h(x) = a(x) (\varphi'_x)^2, \quad Q(z) = w''_{zz}/w'_z, \quad R(z) = f(w(z))/w'_z.$$

Substituting these expressions into (49)–(51) yields a system of ordinary differential equations for the unknowns.

**Example 13.** Equation (52) also admits more complicated functional separable solutions with the form

$$w = w(z), \quad z = \varphi(\xi) + \psi(t), \quad \xi = x + at.$$

Substituting these expressions into equation (52) yields the functional equation B.3.5-2 again, in which ( $x$  must be replaced by  $\xi$ )

$$f(t) = -\psi'_t, \quad g(\xi) = \varphi''_{\xi\xi} - a \varphi'_\xi, \quad h(\xi) = (\varphi'_\xi)^2, \quad Q(z) = w''_{zz}/w'_z, \quad R(z) = f(w(z))/w'_z.$$

Further, one should follow the same procedure of constructing the solution as in Example 11.

*Remark.* In Examples 11–13, different equations were all reduced to the same functional equation. This demonstrates the utility of isolation and independent analysis of individual types of functional equations, as well as the expedience of developing methods for solving functional equations with a composite argument.

**B.3.5-3.** The functional equation  $f(t) + g(x)Q(z) + h(x)R(z) = 0$ , where  $z = \varphi(x) + \psi(t)$ .

Differentiating with respect to  $x$  yields the two-argument functional differential equation

$$g'_x Q + g\varphi'_x Q'_z + h'_x R + h\varphi'_x R'_z = 0, \quad (57)$$

which coincides with equation (25) in Section B.2, up to notation.

*Nondegenerate case.* Equation (57) can be solved using formulas (49) in Section B.2, just as was the case for equation (25). In this way, we arrive at the system of ordinary differential equations

$$\begin{aligned} g'_x &= (A_1 g + A_2 h)\varphi'_x, \\ h'_x &= (A_3 g + A_4 h)\varphi'_x, \\ Q'_z &= -A_1 Q - A_3 R, \\ R'_z &= -A_2 Q - A_4 R, \end{aligned} \quad (58)$$

where  $A_1, A_2, A_3$ , and  $A_4$  are arbitrary constants.

The solution of equation (58) is given by

$$\begin{aligned} g(x) &= A_2 B_1 e^{k_1 \varphi} + A_2 B_2 e^{k_2 \varphi}, \\ h(x) &= (k_1 - A_1) B_1 e^{k_1 \varphi} + (k_2 - A_1) B_2 e^{k_2 \varphi}, \\ Q(z) &= A_3 B_3 e^{-k_1 z} + A_3 B_4 e^{-k_2 z}, \\ R(z) &= (k_1 - A_1) B_3 e^{-k_1 z} + (k_2 - A_1) B_4 e^{-k_2 z}, \end{aligned} \quad (59)$$

where  $B_1, B_2, B_3$ , and  $B_4$  are arbitrary constants and  $k_1$  and  $k_2$  are roots of the quadratic equation

$$(k - A_1)(k - A_4) - A_2 A_3 = 0. \quad (60)$$

In the degenerate case  $k_1 = k_2$  the terms  $e^{k_2 \varphi}$  and  $e^{-k_2 z}$  in (59) must be replaced by  $\varphi e^{k_1 \varphi}$  and  $z e^{-k_1 z}$ , respectively. In the case of purely imaginary or complex roots, one should extract the real (or imaginary) part of the roots in solution (59).

On substituting (59) into the original functional equation, one obtains conditions that must be met by the free coefficients and identifies the function  $f(t)$ , specifically,

$$\begin{aligned} B_2 = B_4 = 0 &\implies f(t) = [A_2 A_3 + (k_1 - A_1)^2] B_1 B_3 e^{-k_1 \psi}, \\ B_1 = B_3 = 0 &\implies f(t) = [A_2 A_3 + (k_2 - A_1)^2] B_2 B_4 e^{-k_2 \psi}, \\ A_1 = 0 &\implies f(t) = (A_2 A_3 + k_1^2) B_1 B_3 e^{-k_1 \psi} + (A_2 A_3 + k_2^2) B_2 B_4 e^{-k_2 \psi}. \end{aligned} \quad (61)$$

Solution (59), (61) involves arbitrary functions  $\varphi = \varphi(x)$  and  $\psi = \psi(t)$ .

*Degenerate case.* In addition, the functional equation has the two degenerate solutions:

$$f = B_1 B_2 e^{A_1 \psi}, \quad g = A_2 B_1 e^{-A_1 \varphi}, \quad h = B_1 e^{-A_1 \varphi}, \quad R = -B_2 e^{A_1 z} - A_2 Q,$$

where  $\varphi = \varphi(x)$ ,  $\psi = \psi(t)$ , and  $Q = Q(z)$  are arbitrary functions,  $A_1, A_2, B_1$ , and  $B_2$  are arbitrary constants; and

$$f = B_1 B_2 e^{A_1 \psi}, \quad h = -B_1 e^{-A_1 \varphi} - A_2 g, \quad Q = A_2 B_2 e^{A_1 z}, \quad R = B_2 e^{A_1 z},$$

where  $\varphi = \varphi(x)$ ,  $\psi = \psi(t)$ , and  $g = g(x)$  are arbitrary functions,  $A_1, A_2, B_1$ , and  $B_2$  are arbitrary constants. The degenerate solutions can be obtained immediately from the original equation or its consequence (57) using formulas (26b) in Section B.2.

**Example 14.** For the first-order nonlinear equation

$$\frac{\partial w}{\partial t} = \mathcal{F}(w) \left( \frac{\partial w}{\partial x} \right)^2 + \mathcal{G}(x),$$

the search for exact solutions in the form (53) leads to the functional equation B.3.5-3 in which

$$f(t) = -\psi'_t, \quad g(x) = (\varphi'_x)^2, \quad h(x) = \mathcal{G}(x), \quad Q(z) = \mathcal{F}(w)w'_z, \quad R(z) = 1/w'_z, \quad w = w(z).$$

### B.3.5-4. Equation $f_1(x) + f_2(y) + g_1(x)P(z) + g_2(y)Q(z) + R(z) = 0$ , $z = \varphi(x) + \psi(y)$ .

Differentiating with respect to  $y$  and dividing the resulting relation by  $\psi'_y P'_z$  and differentiating with respect to  $y$ , one arrives at the functional equation with two arguments  $y$  and  $z$  that is discussed in Section B.2 [see equation (3) and its solution (22)].

**Example 15.** Consider the following equation of steady-state heat transfer in an anisotropic inhomogeneous medium with a nonlinear source:

$$\frac{\partial}{\partial x} \left[ a(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ b(y) \frac{\partial w}{\partial y} \right] = \mathcal{F}(w). \quad (62)$$

The search for exact solutions in the form  $w = w(z)$ ,  $z = \varphi(x) + \psi(y)$ , leads to the functional equation B.3.5-4 in which

$$f_1(x) = a(x)\varphi''_{xx} + a'_x(x)\varphi'_x, \quad f_2(y) = b(y)\psi''_{yy} + b'_y(y)\psi'_y, \quad g_1(x) = a(x)(\varphi'_x)^2, \quad g_2(y) = b(y)(\psi'_y)^2, \\ P(z) = Q(z) = w''_{zz}/w'_z, \quad R(z) = -\mathcal{F}(w)/w'_z, \quad w = w(z).$$

Here we confine ourselves to studying functional separable solutions existing for arbitrary right-hand side  $\mathcal{F}(w)$ .

With the change of variable  $z = \zeta^2$ , we look for solutions of equation (62) in the form

$$w = w(\zeta), \quad \zeta^2 = \varphi(x) + \psi(y). \quad (63)$$

Taking into account that  $\frac{\partial \zeta}{\partial x} = \frac{\varphi'_x}{2\zeta}$  and  $\frac{\partial \zeta}{\partial y} = \frac{\psi'_y}{2\zeta}$ , we find from (62)

$$[(a\varphi'_x)'_x + (b\psi'_y)'_y] \frac{w'_\zeta}{2\zeta} + [a(\varphi'_x)^2 + b(\psi'_y)^2] \frac{\zeta w''_{\zeta\zeta} - w'_\zeta}{4\zeta^3} = \mathcal{F}(w), \quad \mathcal{F}(w) = \mathcal{F}(w(\zeta)). \quad (64)$$

For this functional differential equation to be solvable we require that the expressions in square brackets be functions of  $\zeta$ :

$$(a\varphi'_x)'_x + (b\psi'_y)'_y = M(\zeta), \quad a(\varphi'_x)^2 + b(\psi'_y)^2 = N(\zeta).$$

Differentiating the first relation with respect to  $x$  and  $y$  yields the equation  $(M'_\zeta/\zeta)'_\zeta = 0$ , whose general solution is  $M(\zeta) = C_1\zeta^2 + C_2$ . Likewise, we find  $N(\zeta) = C_3\zeta^2 + C_4$ . Here,  $C_1$ ,  $C_2$ ,  $C_3$ , and  $C_4$  are arbitrary constants. As a result, we have

$$(a\varphi'_x)'_x + (b\psi'_y)'_y = C_1(\varphi + \psi) + C_2, \quad a(\varphi'_x)^2 + b(\psi'_y)^2 = C_3(\varphi + \psi) + C_4.$$

The separation of variables results in a system of ordinary differential equations for  $\varphi(x)$ ,  $a(x)$ ,  $\psi(y)$ , and  $b(y)$ :

$$(a\varphi'_x)'_x - C_1\varphi - C_2 = k_1, \quad (b\psi'_y)'_y - C_1\psi = -k_1, \\ a(\varphi'_x)^2 - C_3\varphi - C_4 = k_2, \quad b(\psi'_y)^2 - C_3\psi = -k_2.$$

This system is always integrable in quadrature and can be rewritten as

$$(C_3\varphi + C_4 + k_2)\varphi''_{xx} + (C_1\varphi + C_2 + k_1 - C_3)(\varphi'_x)^2 = 0, \quad a = (C_3\varphi + C_4 + k_2)(\varphi'_x)^{-2}; \\ (C_3\psi - k_2)\psi''_{yy} + (C_1\psi - k_1 - C_3)(\psi'_y)^2 = 0, \quad b = (C_3\psi - k_2)(\psi'_y)^{-2}. \quad (65)$$

Here, the equations for  $\varphi$  and  $\psi$  do not involve  $a$  and  $b$  and, hence, can be solved independently. Without full analysis of system (65), we note a special case where the system can be solved in explicit form.

For  $C_1 = C_2 = C_4 = k_1 = k_2 = 0$  and  $C_3 = C \neq 0$ , we find

$$a(x) = \alpha e^{\mu x}, \quad b(y) = \beta e^{\nu y}, \quad \varphi(x) = \frac{C e^{-\mu x}}{\alpha \mu^2}, \quad \psi(y) = \frac{C e^{-\nu y}}{\beta \nu^2},$$

where  $\alpha$ ,  $\beta$ ,  $\mu$ , and  $\nu$  are arbitrary constants. Substituting these expressions into (64) and taking into account (63), we obtain the ordinary differential equation for  $w(\zeta)$

$$w''_{\zeta\zeta} - \frac{1}{\zeta} w'_\zeta = \frac{4}{C} \mathcal{F}(w).$$

System (65) has other solutions as well; these lead to various expressions of  $a(x)$  and  $b(y)$ . Table B3 lists the cases where these functions can be written in explicit form (the traveling wave solution, which corresponds to  $a = \text{const}$  and  $b = \text{const}$ , is omitted). In general, the solution of system (64) enables one to represent  $a(x)$  and  $b(y)$  in parametric form.

⊙ *Reference:* V. F. Zaitsev and A. D. Polyanin (1996), A. D. Polyanin and A. I. Zhurov (1998).

TABLE B3

Functional separable solutions of the form  $w = w(\zeta)$ ,  $\zeta^2 = \varphi(x) + \psi(y)$ , for heat equations in an anisotropic inhomogeneous medium with an arbitrary nonlinear source. Notation:  $C$ ,  $\alpha$ ,  $\beta$ ,  $\mu$ ,  $\nu$ ,  $n$ , and  $k$  are free parameters ( $C \neq 0$ ,  $\mu \neq 0$ ,  $\nu \neq 0$ ,  $n \neq 2$ , and  $k \neq 2$ )

| Heat equation                                                                                                                                                                                           | Functions $\varphi(x)$ and $\psi(y)$                                                       | Equation for $w = w(\zeta)$                                                                          |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------|
| $\frac{\partial}{\partial x} \left( \alpha x^m \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( \beta y^n \frac{\partial w}{\partial y} \right) = \mathcal{F}(w)$             | $\varphi = \frac{Cx^{2-m}}{\alpha(2-m)^2}, \quad \psi = \frac{Cy^{2-n}}{\beta(2-n)^2}$     | $w''_{\zeta\zeta} + \frac{4-mn}{(2-m)(2-n)} \frac{1}{\zeta} w'_{\zeta} = \frac{4}{C} \mathcal{F}(w)$ |
| $\frac{\partial}{\partial x} \left( \alpha e^{\mu x} \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( \beta e^{\nu y} \frac{\partial w}{\partial y} \right) = \mathcal{F}(w)$ | $\varphi = \frac{C}{\alpha\mu^2} e^{-\mu x}, \quad \psi = \frac{C}{\beta\nu^2} e^{-\nu y}$ | $w''_{\zeta\zeta} - \frac{1}{\zeta} w'_{\zeta} = \frac{4}{C} \mathcal{F}(w)$                         |
| $\frac{\partial}{\partial x} \left( \alpha e^{\mu x} \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( \beta y^n \frac{\partial w}{\partial y} \right) = \mathcal{F}(w)$       | $\varphi = \frac{C}{\alpha\mu^2} e^{-\mu x}, \quad \psi = \frac{Cy^{2-n}}{\beta(2-n)^2}$   | $w''_{\zeta\zeta} + \frac{n}{2-n} \frac{1}{\zeta} w'_{\zeta} = \frac{4}{C} \mathcal{F}(w)$           |
| $\frac{\partial}{\partial x} \left( \alpha x^2 \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( \beta y^2 \frac{\partial w}{\partial y} \right) = \mathcal{F}(w)$             | $\varphi = \mu \ln  x , \quad \psi = \nu \ln  y $                                          | Equation (64); both expressions in square brackets are constant                                      |
| $\alpha \frac{\partial^2 w}{\partial x^2} + \frac{\partial}{\partial y} \left( \beta y^2 \frac{\partial w}{\partial y} \right) = \mathcal{F}(w)$                                                        | $\varphi = \mu x, \quad \psi = \nu \ln  y $                                                | Equation (64); both expressions in square brackets are constant                                      |

## B.4. First-Order Nonlinear Equations

### B.4.1. Preliminary Remarks

For first-order partial differential equations with two independent variables, an exact solution

$$w = \Phi(x, y, C_1, C_2) \quad (1)$$

that depends on two arbitrary constants  $C_1$  and  $C_2$  is called a complete integral. The general integral (general solution) can be represented in parametric form by using the complete integral (1) and the two equations

$$\begin{aligned} C_2 &= f(C_1), \\ \frac{\partial \Phi}{\partial C_1} + \frac{\partial \Phi}{\partial C_2} f'(C_1) &= 0, \end{aligned} \quad (2)$$

where  $f$  is an arbitrary function and the prime stands for the derivative. For details, see Kamke (1965), Courant and Hilbert (1989), and Polyanin, Zaitsev, and Moussiaux (2001).

The first-order equations with two independent variables considered below are purely illustrative. The book by Polyanin, Zaitsev, and Moussiaux (2001) presents many more first-order nonlinear equations that admit generalized separable solutions (without specifying the method for obtaining them).

### B.4.2. Individual Equations

$$1. \quad \frac{\partial w}{\partial x} - f(y)w \frac{\partial w}{\partial y} = g(x)w + h(x).$$

Exact solution:

$$w = \varphi(x) \int \frac{dy}{f(y)} + \psi(x),$$

where

$$\begin{aligned} \varphi(x) &= G(x) \left[ C_1 - \int G(x) dx \right]^{-1}, & G(x) &= \exp \left[ \int g(x) dx \right], \\ \psi(x) &= S(x) \left[ C_2 + \int \frac{h(x)}{S(x)} dx \right], & S(x) &= G(x) \exp \left[ \int \varphi(x) dx \right]. \end{aligned}$$



$$2. \quad \frac{\partial w}{\partial x} + f(y)w \frac{\partial w}{\partial y} = aw^2 + g(x)w + h(x).$$

Exact solution:

$$w = \varphi(x) + \psi(x) \exp \left[ a \int \frac{dy}{f(y)} \right], \quad \psi(x) = C_1 \exp \left\{ \int [a\varphi(x) + g(x)] dx \right\},$$

where  $C_1$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the Riccati equation

$$\varphi'_x = a\varphi^2 + g(x)\varphi + h(x).$$

This equation is integrable in quadrature for a lot of specific functions  $g(x)$  and  $h(x)$  [e.g., for  $h(x) \equiv 0$  and any  $g(x)$ ]. For details, see the books by Kamke (1977) and Polyanin and Zaitsev (1995).

$$3. \quad \frac{\partial w}{\partial x} \frac{\partial w}{\partial y} = f(x)y^k + g(x)y^{2k+1}.$$

Exact solutions:

$$w = \varphi(x)y^{k+1} + \frac{1}{k+1} \int \frac{f(x)}{\varphi(x)} dx + C_1, \quad \varphi(x) = \pm \left[ \frac{2}{k+1} \int g(x) dx + C_2 \right]^{1/2}.$$

$$4. \quad \frac{\partial w}{\partial x} \frac{\partial w}{\partial y} = f(x)e^{\lambda y} + g(x)e^{2\lambda y}.$$

Exact solutions:

$$w = \varphi(x)e^{\lambda y} + \frac{1}{\lambda} \int \frac{f(x)}{\varphi(x)} dx + C_1, \quad \varphi(x) = \pm \left[ \frac{2}{\lambda} \int g(x) dx + C_2 \right]^{1/2}.$$

$$5. \quad \frac{\partial w}{\partial x} + a \left( \frac{\partial w}{\partial y} \right)^2 = f(x)y + g(x).$$

Exact solution:

$$w = \varphi(x)y + \int [g(x) - a\varphi^2(x)] dx + C_1, \quad \varphi(x) = \int f(x) dx + C_2.$$

$$6. \quad \frac{\partial w}{\partial x} + a \left( \frac{\partial w}{\partial y} \right)^2 = f(x)y^2 + g(x)y + h(x).$$

Exact solution:

$$w = \varphi(x)y^2 + \psi(x)y + \chi(x),$$

where the functions  $\varphi(x)$ ,  $\psi(x)$ , and  $\chi(x)$  are determined by solving the following system of ordinary differential equations:

$$\varphi'_x = -4a\varphi^2 + f(x), \tag{1}$$

$$\psi'_x = -4a\varphi\psi + g(x), \tag{2}$$

$$\chi'_x = -a\psi^2 + h(x). \tag{3}$$

The Riccati equation (1) can be integrated in quadrature for numerous  $f(x)$ . For details, see Kamke (1977) and Polyanin and Zaitsev (1995). Given a solution of equation (1), equations (2) and (3) are easy to integrate, because they are linear in the unknowns  $\psi$  and  $\chi$ .

$$7. \quad \frac{\partial w}{\partial x} + a \left( \frac{\partial w}{\partial y} \right)^2 = bw^2 + f(x)w + g(x).$$

1°. Exact solution for  $b = 0$ :

$$w = F(x)(C_1 + C_2 y) + F(x) \int [g(x) - aC_2^2 F^2(x)] \frac{dx}{F(x)}, \quad F(x) = \exp \left[ \int f(x) dx \right].$$

2°. Exact solution for  $b \neq 0$ :

$$w = \varphi(x) + \psi(y) \exp(\pm y \sqrt{b/a}), \quad \psi(y) = C_1 \exp \left\{ \int [2b\varphi(x) + f(x)] dx \right\}.$$

The function  $\varphi = \varphi(x)$  is determined by the Riccati equation

$$\varphi' = b\varphi^2 + f(x)\varphi + g(x).$$

This equation can be integrated in quadrature for various  $f$  and  $g$ , in particular, for  $g(x) \equiv 0$  and arbitrary  $f(x)$  and for  $f(x) \equiv \text{const}$  and  $g(x) \equiv \text{const}$ . For details, see the books by Kamke (1977) and Polyanin and Zaitsev (1995).

$$8. \quad f_1(x) \left( \frac{\partial w}{\partial x} \right)^2 + f_2(y) \left( \frac{\partial w}{\partial y} \right)^2 = g_1(x) + g_2(y).$$

This equation is encountered in differential geometry in studying geodesic lines of Liouville surfaces. Exact solutions:

$$w = \pm \int \sqrt{\frac{g_1(x) + C_1}{f_1(x)}} dx \pm \int \sqrt{\frac{g_2(y) - C_1}{f_2(y)}} dy + C_2.$$

The signs before each of the integrals can be chosen independently of each other.

☉ *References:* P. Appell (1953), E. Kamke (1965).

$$9. \quad \frac{\partial w}{\partial x} + f \left( \frac{\partial w}{\partial y} \right) = g(x)y + h(x).$$

Exact solution:

$$w = \varphi(x)y + \int [h(x) - f(\varphi(x))] dx + C_1, \quad \varphi(x) = \int g(x) dx + C_2.$$

$$10. \quad \frac{\partial w}{\partial x} + f \left( \frac{\partial w}{\partial y} \right) = g(x)w + h(x).$$

Exact solution:

$$w = (C_1 y + C_2) \varphi(x) + \varphi(x) \int [h(x) - f(C_1 \varphi(x))] \frac{dx}{\varphi(x)}, \quad \varphi(x) = \exp \left[ \int g(x) dx \right].$$

$$11. \quad F_1 \left( x, \frac{\partial w}{\partial x} \right) + e^{\lambda w} F_2 \left( y, \frac{\partial w}{\partial y} \right) = 0.$$

Exact solution:

$$w = \varphi(x) + \psi(y).$$

The functions  $\varphi = \varphi(x)$  and  $\psi = \psi(y)$  are determined by solving the ordinary differential equations

$$e^{-\lambda \varphi} F_1(x, \varphi'_x) = C, \quad e^{\lambda \psi} F_2(y, \psi'_y) = -C,$$

where  $C$  is an arbitrary constant.

## B.5. Second-Order Nonlinear Equations

### B.5.1. Parabolic Equations

B.5.1-1. Equations of the form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + F(x, t, w)$ .

1. 
$$\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + bw \ln w + [f(x) + g(t)]w.$$

Exact solution with multiplicative form:

$$w(x, t) = \exp \left[ C e^{bt} + e^{bt} \int e^{-bt} g(t) dt \right] \varphi(x),$$

where  $C$  is an arbitrary constant and the function  $\varphi(t)$  is determined by solving the ordinary differential equation

$$a\varphi''_{xx} + b\varphi \ln \varphi + f(x)\varphi = 0.$$

2. 
$$\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(t)w \ln w + g(t)w.$$

1°. Exact solution:

$$w(x, t) = \exp [\Phi(t)x + \Psi(t)],$$

where the functions  $\Phi(t)$  and  $\Psi(t)$  are given by

$$\Phi(t) = Ae^F, \quad \Psi(t) = Be^F + e^F \int e^{-F} (aA^2 e^{2F} + g) dt, \quad F = \int f dt,$$

and  $A$  and  $B$  are arbitrary constants.

2°. Exact solution:

$$w(x, t) = \exp [\varphi(t)x^2 + \psi(t)],$$

where  $\varphi(t)$  and  $\psi(t)$  are given by

$$\varphi(t) = e^F \left( A - 4a \int e^F dt \right)^{-1}, \quad \psi(t) = Be^F + e^F \int e^{-F} (2a\varphi + g) dt, \quad F = \int f dt,$$

and  $A$  and  $B$  are arbitrary constants.

3°. There are also exact solutions of the more general form

$$w(x, t) = \exp [\varphi_2(t)x^2 + \varphi_1(t)x + \varphi_0(t)],$$

where the functions  $\varphi_2(t)$ ,  $\varphi_1(t)$ , and  $\varphi_0(t)$  are determined by a system of ordinary differential equations that can be integrated.

⊙ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

3. 
$$\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(t)w \ln w + [g(t)x^2 + h(t)x + s(t)]w.$$

Exact solution:

$$w(x, t) = \exp [\varphi_2(t)x^2 + \varphi_1(t)x + \varphi_0(t)],$$

where the functions  $\varphi_n(t)$  ( $n = 1, 2, 3$ ) are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\begin{aligned} \varphi'_2 &= 4a\varphi_2^2 + f\varphi_2 + g, \\ \varphi'_1 &= 4a\varphi_2\varphi_1 + f\varphi_1 + h, \\ \varphi'_0 &= f\varphi_0 + a\varphi_1^2 + 2a\varphi_2 + s \end{aligned}$$

(the arguments of  $f$ ,  $g$ ,  $h$ , and  $s$  are not specified and the prime denotes the derivative with respect to  $t$ ).

$$4. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + [k \ln^2 w + f(t) \ln w + g(t)]w.$$

The change of variable  $w = \exp u$  leads to an equation of the form B.5.1.14,

$$\frac{\partial u}{\partial t} = a \frac{\partial^2 u}{\partial x^2} + a \left( \frac{\partial u}{\partial x} \right)^2 + ku^2 + f(t)u + g(t),$$

which has exponential and sinusoidal solutions in  $x$ .

B.5.1-2. Equations of the form  $\frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + F(x, t, w, \frac{\partial w}{\partial x})$ .

$$5. \quad \frac{\partial w}{\partial t} = \frac{a}{x^n} \frac{\partial}{\partial x} \left( x^n \frac{\partial w}{\partial x} \right) + f(t)w \ln w.$$

Exact solution:

$$w(x, t) = \exp[\varphi(t)x^2 + \psi(t)],$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients (the arguments of  $f$  and  $g$  are not specified):

$$\begin{aligned} \varphi'_t &= 4a\varphi^2 + f\varphi, \\ \psi'_t &= 2a(n+1)\varphi + f\psi. \end{aligned}$$

Integrating successively yields

$$\varphi(t) = e^F \left( A - 4a \int e^F dt \right)^{-1}, \quad \psi(t) = Be^F + 2a(n+1)e^F \int \varphi e^{-F} dt, \quad F = \int f dt,$$

where  $A$  and  $B$  are arbitrary constants.

$$6. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + [xf(t) + g(t)] \frac{\partial w}{\partial x} + h(t)w \ln w + [xp(t) + s(t)]w.$$

Exact solution:

$$w(x, t) = \exp[x\varphi(t) + \psi(t)],$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = [f(t) + h(t)]\varphi + p(t), \tag{1}$$

$$\psi'_t = h(t)\psi + a\varphi^2 + g(t)\varphi + s(t). \tag{2}$$

Integrating first (1) and then (2), we obtain ( $C_1$  and  $C_2$  are arbitrary constants)

$$\begin{aligned} \varphi(t) &= C_1 E(t) + E(t) \int \frac{p(t)}{E(t)} dt, & E(t) &= \exp \left[ \int f(t) dt + \int h(t) dt \right], \\ \psi(t) &= C_2 H(t) + H(t) \int \frac{a\varphi^2(t) + g(t)\varphi(t) + s(t)}{H(t)} dt, & H(t) &= \exp \left[ \int h(t) dt \right]. \end{aligned}$$

$$7. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + [xf(t) + g(t)] \frac{\partial w}{\partial x} + h(t)w \ln w + [x^2 r(t) + xp(t) + s(t)]w.$$

Exact solution:

$$w(x, t) = \exp[x^2\varphi(t) + x\psi(t) + \chi(t)],$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\begin{aligned}\varphi'_t &= 4a\varphi^2 + (2f + h)\varphi + r, \\ \psi'_t &= (4a\varphi + f + h)\psi + 2g\varphi + p, \\ \chi'_t &= h\chi + 2a\varphi + a\psi^2 + g\psi + s.\end{aligned}$$

$$8. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + \left[ x f(t) + \frac{g(t)}{x} \right] \frac{\partial w}{\partial x} + h(t)w \ln w + [x^2 p(t) + s(t)]w.$$

Exact solution:

$$w(x, t) = \exp[\varphi(t)x^2 + \psi(t)],$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\begin{aligned}\varphi'_t &= 4a\varphi^2 + (2f + h)\varphi + p, \\ \psi'_t &= h\psi + 2(a + g)\varphi + s.\end{aligned}$$

$$9. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + f(t)x^2 + g(t)x + h(t).$$

Exact solution:

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\begin{aligned}\varphi'_t &= 4b\varphi^2 + f, \\ \psi'_t &= 4b\varphi\psi + g, \\ \chi'_t &= 2a\varphi + b\psi^2 + h.\end{aligned}$$

$$10. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw + f(x) + g(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + Ae^{ct} + e^{ct} \int e^{-ct} g(t) dt.$$

Here,  $A$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the nonlinear ordinary differential equation

$$a\varphi''_{xx} + b(\varphi'_x)^2 + c\varphi + f(x) = 0.$$

By changing variable  $\varphi'_x = \frac{a}{b} \frac{\psi'_x}{\psi}$  this equation is reduced to the second-order linear equation

$$a^2\psi''_{xx} + ac\psi'_x + bf(x)\psi = 0.$$

$$11. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + f(x) \frac{\partial w}{\partial x} + kw + g(x) + h(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + Ce^{kt} + e^{kt} \int e^{-kt} h(t) dt,$$

where  $C$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the second-order ordinary differential equation with variable coefficients

$$a\varphi''_{xx} + b(\varphi'_x)^2 + f(x)\varphi'_x + k\varphi + g(x) = 0.$$

$$12. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw \frac{\partial w}{\partial x} + kw^2 + f(t)w + g(t).$$

The equation has exact solutions of the form

$$w(x, t) = \varphi(t) + \psi(t) \exp(\lambda x),$$

where  $\lambda$  is a root of the quadratic equation  $b\lambda^2 + c\lambda + k = 0$ .

$$13. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + g(t)w + h(t).$$

Exact solution:

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = 4f\varphi^2 + g\varphi, \quad (1)$$

$$\psi'_t = (4f\varphi + g)\psi, \quad (2)$$

$$\chi'_t = g\chi + 2a\varphi + f\psi^2 + h. \quad (3)$$

Equation (1) for  $\varphi$  is a Bernoulli equation, which is easy to integrate. After that, equations (2) and (3), which are linear in  $\psi$  and  $\chi$ , are integrated successively. As a result, we find

$$\varphi = e^G \left( A_1 - 4 \int e^G f dt \right)^{-1}, \quad G = \int g dt,$$

$$\psi = A_2 \exp \left[ \int (4f\varphi + g) dt \right],$$

$$\chi = A_3 e^G + e^G \int e^{-G} (2a\varphi + f\psi^2 + h) dt,$$

where  $A_1$ ,  $A_2$ , and  $A_3$  are arbitrary constants. A degenerate solution with  $\varphi \equiv 0$  corresponds to the limit case  $A_1 \rightarrow \infty$ .

☉ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

$$14. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + bf(t)w^2 + g(t)w + h(t).$$

1°. Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\pm x\sqrt{-b}), \quad b < 0, \quad (1)$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following first-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\varphi'_t = bf\varphi^2 + g\varphi + h, \quad (2)$$

$$\psi'_t = (2bf\varphi + g - ab)\psi. \quad (3)$$

Equation (2) for  $\varphi(t)$  is a Riccati equation; it can be reduced to a second-order linear equation. Many solutions of equation (2) for various  $f$ ,  $g$ , and  $h$  can be found in Kamke (1977) and Polyanin and Zaitsev (1995).

Whenever a solution of equation (2) is known, the solution of equation (3) for  $\psi(t)$  can be evaluated from

$$\psi(t) = C \exp \left[ -abt + \int (2bf\varphi + g) dt \right],$$

where  $C$  is an arbitrary constant.

2°. Exact solution of a more general form:

$$w(x, t) = \varphi(t) + \psi(t) [A \exp(x\sqrt{-b}) + B \exp(-x\sqrt{-b})], \quad b < 0,$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = bf(\varphi^2 + 4AB\psi^2) + g\varphi + h, \quad (4)$$

$$\psi'_t = 2bf\varphi\psi + g\psi - ab\psi. \quad (5)$$

One can express  $\varphi$  in terms of  $\psi$  from (5) and then substitute into (4). As a result, one obtains a second-order nonlinear equation for  $\psi$ ; if  $f, g, h = \text{const}$ , this equation is autonomous and, hence, admits reduction of order.

3°. Exact solution ( $c$  is an arbitrary constant):

$$w(x, t) = \varphi(t) + \psi(t) \cos(x\sqrt{b} + c), \quad b > 0, \quad (6)$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = bf(\varphi^2 + \psi^2) + g\varphi + h, \quad (7)$$

$$\psi'_t = 2bf\varphi\psi + g\psi - ab\psi. \quad (8)$$

One can express  $\varphi$  in terms of  $\psi$  from (8) and then substitute into (7). As a result, one obtains a second-order nonlinear equation for  $\psi$ ; if  $f, g, h = \text{const}$ , this equation is autonomous and, hence, admits reduction of order.

⊙ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

$$15. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + bf(t)w \frac{\partial w}{\partial x} + cf(t)w^2 + g(t)w + h(t).$$

The equation has exact solutions of the form

$$w(x, t) = \varphi(t) + \psi(t) \exp(\lambda x),$$

where  $\lambda$  is a root of the quadratic equation  $\lambda^2 + b\lambda + c = 0$ .

$$16. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(x) \left( \frac{\partial w}{\partial x} \right)^2 + g(x) \frac{\partial w}{\partial x} + bw + h(x) + p(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + Ce^{bt} + e^{bt} \int e^{-bt} p(t) dt,$$

where  $C$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the following second-order ordinary differential equation with variable coefficients:

$$a\varphi''_{xx} + f(x)(\varphi'_x)^2 + g(x)\varphi'_x + b\varphi + h(x) = 0.$$

$$17. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + [g_1(t)x + g_0(t)] \frac{\partial w}{\partial x} + h(t)w + p(t)x^2 + q(t)x + s(t).$$

Exact solution:

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = 4f\varphi^2 + (2g_1 + h)\varphi + p, \quad (1)$$

$$\psi'_t = (4f\varphi + g_1 + h)\psi + 2g_0\varphi + q, \quad (2)$$

$$\chi'_t = h\chi + 2a\varphi + f\psi^2 + g_0\psi + s. \quad (3)$$

Equation (1) for  $\varphi(t)$  is a Riccati equation; it can be reduced to a second-order linear equation. For solutions of Riccati equations, see Kamke (1977) and Polyanin and Zaitsev (1995). Whenever a solution of equation (1) is known, the solutions of equations (2) and (3) can be obtained successively (the equations are linear in  $\psi$  and  $\chi$ ).

$$18. \quad \frac{\partial w}{\partial t} = a \frac{\partial^2 w}{\partial x^2} + f\left(x, \frac{\partial w}{\partial x}\right) + bw + g(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + Ce^{bt} + e^{bt} \int e^{-bt} g(t) dt,$$

where  $C$  is an arbitrary constant and the function  $\varphi(x)$  is described by the second-order ordinary differential equation

$$a\varphi''_{xx} + f(x, \varphi'_x) + b\varphi = 0.$$

|                                                                                                                                                          |
|----------------------------------------------------------------------------------------------------------------------------------------------------------|
| B.5.1-3. Equations of the form $\frac{\partial w}{\partial t} = f(x, t) \frac{\partial^2 w}{\partial x^2} + g(x, t, w, \frac{\partial w}{\partial x})$ . |
|----------------------------------------------------------------------------------------------------------------------------------------------------------|

$$19. \quad \frac{\partial w}{\partial t} = \frac{f(t)}{x^n} \frac{\partial}{\partial x} \left( x^n \frac{\partial w}{\partial x} \right) + g(t)w \ln w.$$

Exact solution:

$$w(x, t) = \exp[\varphi(t)x^2 + \psi(t)],$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients (the arguments of  $f$  and  $g$  are not specified):

$$\begin{aligned} \varphi'_t &= 4f\varphi^2 + g\varphi, \\ \psi'_t &= 2(n+1)f\varphi + g\psi. \end{aligned}$$

Integrating successively yields

$$\varphi(t) = e^G \left( A - 4 \int f e^G dt \right)^{-1}, \quad \psi(t) = B e^G + 2(n+1) e^G \int f \varphi e^{-G} dt, \quad G = \int g dt,$$

where  $A$  and  $B$  are arbitrary constants.

$$20. \quad \frac{\partial w}{\partial t} = f(t) \frac{\partial^2 w}{\partial x^2} + \left[ xg(t) + \frac{h(t)}{x} \right] \frac{\partial w}{\partial x} + s(t)w \ln w + [x^2 p(t) + q(t)]w.$$

Exact solution:

$$w(x, t) = \exp[\varphi(t)x^2 + \psi(t)],$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = 4f\varphi^2 + (2g + s)\varphi + p, \tag{1}$$

$$\psi'_t = s\psi + 2(f + h)\varphi + q. \tag{2}$$

The Riccati equation (1) for the function  $\varphi(t)$  can be reduced to a second-order linear equation. For solutions of the Riccati equation, see Kamke (1977) and Polyanin and Zaitsev (1995). On solving (1), one can determine the solution of the linear equation (2) for  $\psi(t)$ .

$$21. \quad \frac{\partial w}{\partial t} = f(t) \frac{\partial}{\partial x} \left( e^{\lambda x} \frac{\partial w}{\partial x} \right) + g(t)w \ln w + h(t)w.$$

Exact solution:

$$w(x, t) = \exp[\varphi(t)e^{-\lambda x} + \psi(t)],$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by the ordinary differential equations

$$\varphi'_t = \lambda^2 f(t)\varphi^2 + g(t)\varphi,$$

$$\psi'_t = g(t)\psi + h(t).$$



$$\varphi(t) = G(t) \left[ A - \lambda^2 \int f(t)G(t) dt \right]^{-1}, \quad G(t) = \exp \left[ \int g(t) dt \right],$$

$$\psi(t) = BG(t) + G(t) \int \frac{h(t)}{G(t)} dt,$$

where  $A$  and  $B$  are arbitrary constants.

**22.**  $\frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + aw \ln w.$

Exact solution:

$$w(x, t) = \exp [Ae^{at} + \varphi(x)],$$

where  $A$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the ordinary differential equation

$$(f\varphi'_x)'_x + f(\varphi'_x)^2 + a\varphi = 0.$$

**23.**  $\frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + aw \ln w + [g(x) + h(t)]w.$

Exact solution in multiplicative form:

$$w(x, t) = \exp \left[ Ce^{at} + e^{at} \int e^{-at} h(t) dt \right] \varphi(x),$$

where  $C$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the ordinary differential equation

$$(f\varphi'_x)'_x + a\varphi \ln \varphi + g(x)\varphi = 0.$$

**24.**  $\frac{\partial w}{\partial t} = f(x) \frac{\partial^2 w}{\partial x^2} + g(x) \frac{\partial w}{\partial x} + aw \ln w + [h(x) + s(t)]w.$

Exact solution in multiplicative form:

$$w(x, t) = \exp \left[ Ce^{at} + e^{at} \int e^{-at} s(t) dt \right] \varphi(x),$$

where  $C$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the ordinary differential equation

$$f(x)\varphi''_{xx} + g(x)\varphi'_x + a\varphi \ln \varphi + h(x)\varphi = 0.$$

**25.**  $\frac{\partial w}{\partial t} = f(x) \frac{\partial^2 w}{\partial x^2} + g \left( x, \frac{\partial w}{\partial x} \right) + aw + h(t).$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + Ce^{at} + e^{at} \int e^{-at} h(t) dt,$$

where  $C$  is an arbitrary constant and the function  $\varphi(x)$  is determined by the following second-order ordinary differential equation:

$$f(x)\varphi''_{xx} + g(x, \varphi'_x) + a\varphi = 0.$$

$$26. \quad \frac{\partial w}{\partial t} = aw\frac{\partial^2 w}{\partial x^2} + f(x)w + bx + c.$$

Exact solution:

$$w(x, t) = (bx + c)t + Ax + B - \frac{1}{a} \int_{x_0}^x (x - \xi)f(\xi) d\xi,$$

where  $A$ ,  $B$ , and  $x_0$  are arbitrary constants.

$$27. \quad \frac{\partial w}{\partial t} = aw\frac{\partial^2 w}{\partial x^2} + f(t)w + g(t).$$

1°. Exact solution:

$$w(x, t) = F(t)(Ax + B) + F(t) \int \frac{g(t)}{F(t)} dt, \quad F(t) = \exp \left[ \int f(t) dt \right],$$

where  $A$  and  $B$  are arbitrary constants.

2°. Exact solution:

$$w(x, t) = \varphi(t)(x^2 + Ax + B) + \varphi(t) \int \frac{g(t)}{\varphi(t)} dt,$$

$$\varphi(t) = F(t) \left[ C - 2a \int F(t) dt \right]^{-1}, \quad F(t) = \exp \left[ \int f(t) dt \right],$$

where  $A$ ,  $B$ , and  $C$  are arbitrary constants.

$$28. \quad \frac{\partial w}{\partial t} = aw\frac{\partial^2 w}{\partial x^2} + f(x)w\frac{\partial w}{\partial x} + g(t)w + h(t).$$

Exact solution:

$$w(x, t) = \varphi(t)\Theta(x) + \psi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\Theta(x)$  are described by ordinary differential equations

$$\begin{aligned} \varphi'_t &= C\varphi^2 + g(t)\varphi, \\ \psi'_t &= [C\varphi + g(t)]\psi + h(t), \\ a\Theta''_{xx} + f(x)\Theta'_x &= C, \end{aligned}$$

where  $C$  is an arbitrary constant. Integrating successively yields

$$\begin{aligned} \varphi(t) &= G(t) \left[ A_1 - C \int G(t) dt \right]^{-1}, \quad G(t) = \exp \left[ \int g(t) dt \right], \\ \psi(t) &= A_2\varphi(t) + \varphi(t) \int \frac{h(t)}{\varphi(t)} dt, \\ \Theta(x) &= B_1 \int \frac{dx}{F(x)} + B_2 + \frac{C}{a} \int \left[ \int F(x) dx \right] \frac{dx}{F(x)}, \quad F(x) = \exp \left[ \frac{1}{a} \int f(x) dx \right], \end{aligned}$$

where  $A_1$ ,  $A_2$ ,  $B_1$ , and  $B_2$  are arbitrary constants.

$$29. \quad \frac{\partial w}{\partial t} = aw \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + g(t) \frac{\partial w}{\partial x} + h(t)w + s(t).$$

Exact solution:

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ ,  $\chi(t)$  are determined by solving the following system of first-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ ,  $h$ , and  $s$  are not specified):

$$\varphi'_t = 2(2f + a)\varphi^2 + h\varphi, \quad (1)$$

$$\psi'_t = (4f\varphi + 2a\varphi + h)\psi + 2g\varphi, \quad (2)$$

$$\chi'_t = (2a\varphi + h)\chi + f\psi^2 + g\psi + s. \quad (3)$$

Equation (1) for  $\varphi = \varphi(t)$  is a Bernoulli equation; it is easy to integrate. After that, one can successively construct the solutions of equations (2) and (3); each equation is linear in the unknown function.

⊙ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

$$30. \quad \frac{\partial w}{\partial t} = aw \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw^2 + f(t)w + g(t).$$

1°. Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\pm \lambda x), \quad \lambda = \left( \frac{-c}{a+b} \right)^{1/2}, \quad (1)$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following first-order ordinary differential equations with variable coefficients (the arguments of  $f$  and  $g$  are not specified):

$$\varphi'_t = c\varphi^2 + f\varphi + g, \quad (2)$$

$$\psi'_t = (a\lambda^2\varphi + 2c\varphi + f)\psi. \quad (3)$$

Equation (2) for  $\varphi = \varphi(t)$  is a Riccati equation; it can be reduced to a second-order linear equation. The books by Kamke (1977) and Zaitsev and Polyanin (1995) present many solutions of equation (2) for various  $f$  and  $g$ .

Given a solution of equation (2), the solution of equation (3) for  $\psi = \psi(t)$  is evaluated by

$$\psi(t) = C \exp \left[ \int (a\lambda^2\varphi + 2c\varphi + f) dt \right], \quad (4)$$

where  $C$  is an arbitrary constant.

2°. Exact solution ( $A$  is an arbitrary constant):

$$w(x, t) = \varphi(t) + \psi(t) \cosh(\lambda x + A), \quad \lambda = \left( \frac{-c}{a+b} \right)^{1/2}, \quad (5)$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following first-order ordinary differential equations with variable coefficients (the arguments of  $f$  and  $g$  are not specified):

$$\varphi'_t = c\varphi^2 - b\lambda^2\psi^2 + f\varphi + g, \quad (6)$$

$$\psi'_t = (a\lambda^2\varphi + 2c\varphi + f)\psi. \quad (7)$$

One can express  $\varphi$  from (7) in terms of  $\psi$  and substitute the resulting  $\varphi$  into (6). As a result, one arrives at a second-order nonlinear equation for  $\psi$  (if  $f$ ,  $g = \text{const}$ , this equation is autonomous and, hence, admits reduction of order).

3°. Exact solution ( $A$  is an arbitrary constant):

$$w(x, t) = \varphi(t) + \psi(t) \sinh(\lambda x + A), \quad \lambda = \left( \frac{-c}{a+b} \right)^{1/2},$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the system of first-order ordinary differential equations

$$\varphi'_t = c\varphi^2 + b\lambda^2\psi^2 + f\varphi + g,$$

$$\psi'_t = (a\lambda^2\varphi + 2c\varphi + f)\psi.$$

4°. Exact solution ( $A$  is an arbitrary constant):

$$w(x, t) = \varphi(t) + \psi(t) \cos(\lambda x + A), \quad \lambda = \left( \frac{c}{a+b} \right)^{1/2}, \quad (8)$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the system of first-order ordinary differential equations

$$\varphi'_t = c\varphi^2 + b\lambda^2\psi^2 + f\varphi + g, \quad (9)$$

$$\psi'_t = (-a\lambda^2\varphi + 2c\varphi + f)\psi. \quad (10)$$

One can express  $\varphi$  from (10) in terms of  $\psi$  and substitute the resulting  $\varphi$  into (9). Thus, one arrives at a second-order nonlinear equation for  $\psi$  (if  $f, g = \text{const}$ , this equation is autonomous and, hence, admits reduction of order).

⊙ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

$$31. \quad \frac{\partial w}{\partial t} = aw \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + [g_1(t)x + g_0(x)] \frac{\partial w}{\partial x} + h(t)w + p_2(t)x^2 + p_1(t)x + p_0(t).$$

The equation has an exact solution of the form

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by a system of first-order ordinary differential equations with variable coefficients (the system is not specified here).

|                                                                                                                                                                                                |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| B.5.1-5. Equations of the form $\frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left[ f(w) \frac{\partial w}{\partial x} \right] + f(x, t, w, \frac{\partial w}{\partial x})$ . |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

$$32. \quad \frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left( w^m \frac{\partial w}{\partial x} \right) + f(t)w^{1-m}.$$

The change of variable  $u = w^m$  leads to an equation of the form B.5.1.29,

$$\frac{\partial u}{\partial t} = au \frac{\partial^2 u}{\partial x^2} + \frac{a}{m} \left( \frac{\partial u}{\partial x} \right)^2 + mf(t),$$

which admits solutions with the form  $u = \varphi(t)x^2 + \psi(t)x + \chi(t)$ .

$$33. \quad \frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left( w^m \frac{\partial w}{\partial x} \right) + f(t)w + g(t)w^{1-m}.$$

The change of variable  $u = w^m$  leads to an equation of the form B.5.1.29,

$$\frac{\partial u}{\partial t} = au \frac{\partial^2 u}{\partial x^2} + \frac{a}{m} \left( \frac{\partial u}{\partial x} \right)^2 + mf(t)u + mg(t),$$

which admits solutions with the form  $u = \varphi(t)x^2 + \psi(t)x + \chi(t)$ .

⊙ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

$$34. \quad \frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left( w^m \frac{\partial w}{\partial x} \right) + bw^{1+m} + f(t)w + g(t)w^{1-m}.$$

For  $b = 0$ , see equation B.5.1.33.

The change of variable  $u = w^m$  leads to an equation of the form B.5.1.30,

$$\frac{\partial u}{\partial t} = au \frac{\partial^2 u}{\partial x^2} + \frac{a}{m} \left( \frac{\partial u}{\partial x} \right)^2 + bmu^2 + mf(t)u + mg(t),$$

which admits solutions with the forms

$$u(x, t) = \varphi(t) + \psi(t) \exp(\pm \lambda x),$$

$$u(x, t) = \varphi(t) + \psi(t) \cosh(\lambda x + C),$$

$$u(x, t) = \varphi(t) + \psi(t) \sinh(\lambda x + C),$$

$$u(x, t) = \varphi(t) + \psi(t) \cos(\lambda x + C),$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by a system of first-order ordinary differential equations; the parameter  $\lambda$  is a root of a quadratic equation and  $C$  is an arbitrary constant.

⊙ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

$$35. \quad \frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left( e^{\lambda w} \frac{\partial w}{\partial x} \right) + f(t) + g(t)e^{-\lambda w}.$$

The change of variable  $u = e^{\lambda w}$  leads to an equation of the form B.5.1.27,

$$\frac{\partial u}{\partial t} = au \frac{\partial^2 u}{\partial x^2} + \lambda f(t)u + \lambda g(t),$$

which admits solutions with the form  $u = \varphi(t)x^2 + \psi(t)x + \chi(t)$ .

$$36. \quad \frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left( e^{\lambda w} \frac{\partial w}{\partial x} \right) + f(x) + (bx + c)e^{-\lambda w}.$$

The change of variable  $u = e^{\lambda w}$  leads to an equation of the form B.5.1.26,

$$\frac{\partial u}{\partial t} = au \frac{\partial^2 u}{\partial x^2} + \lambda f(x)u + \lambda(bx + c),$$

which admits solutions with the form  $u = \lambda(bx + c)t + \varphi(x)$ .

$$37. \quad \frac{\partial w}{\partial t} = a \frac{\partial}{\partial x} \left( e^{\lambda w} \frac{\partial w}{\partial x} \right) + be^{\lambda w} + f(t) + g(t)e^{-\lambda w}.$$

For  $b = 0$ , see equation B.5.1.35.

The change of variable  $u = e^{\lambda w}$  leads to an equation of the form B.5.1.30,

$$\frac{\partial u}{\partial t} = au \frac{\partial^2 u}{\partial x^2} + bu^2 + \lambda f(t)u + \lambda g(t),$$

which admits solutions with the forms

$$u(x, t) = \varphi(t) + \psi(t) \exp(\pm \mu x),$$

$$u(x, t) = \varphi(t) + \psi(t) \cosh(\mu x + C),$$

$$u(x, t) = \varphi(t) + \psi(t) \sinh(\mu x + C),$$

$$u(x, t) = \varphi(t) + \psi(t) \cos(\mu x + C),$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by a system of first-order ordinary differential equations; the parameter  $\mu$  is a root of a quadratic equation and  $C$  is an arbitrary constant.

⊙ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

**B.5.1-6.** Equations of the form  $\frac{\partial w}{\partial t} = f(x, t, w) \frac{\partial^2 w}{\partial x^2} + g(x, t, w, \frac{\partial w}{\partial x})$ .

$$38. \quad \frac{\partial w}{\partial t} = \frac{f(x)}{aw + b} \frac{\partial^2 w}{\partial x^2}.$$

Exact solution:

$$w(x, t) = \frac{1}{a} [\varphi(x)t + \psi(x) - b],$$

where the functions  $\varphi(x)$  and  $\psi(x)$  are determined by the ordinary differential equations

$$f(x)\varphi''_{xx} - \varphi^2 = 0, \quad f(x)\psi''_{xx} - \varphi\psi = 0.$$

The first equation can be treated independently. The second equation has a particular solution  $\psi(x) = \varphi(x)$ , and hence, its general solution is given by

$$\psi(x) = C_1\varphi(x) + C_2\varphi(x) \int \frac{dx}{\varphi^2(x)},$$

where  $C_1$  and  $C_2$  are arbitrary constants.

$$39. \quad \frac{\partial w}{\partial t} = f(t) \frac{\partial}{\partial x} \left( w^m \frac{\partial w}{\partial x} \right) + g(t)w^{1-m}.$$

Exact solution:

$$w(x, t) = [\varphi(t)x^2 + \psi(t)]^{1/m},$$

where the functions  $\varphi = \varphi(x)$  and  $\psi = \psi(x)$  are determined by the first-order ordinary differential equations

$$\varphi'_t = \frac{2(m+2)}{m} f\varphi^2, \quad \psi'_t = 2f\varphi\psi + mg.$$

Integrating yields

$$\varphi = \frac{1}{F}, \quad \psi = F^{-\frac{m}{m+2}} \left( A + m \int gF^{\frac{m}{m+2}} dt \right), \quad F = B - \frac{2(m+2)}{m} \int f dt,$$

where  $A$  and  $B$  are arbitrary constants.

☉ *Reference:* V. F. Zaitsev, A. D. Polyanin (1996).

$$40. \quad \frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ f(x)e^{\beta w} \frac{\partial w}{\partial x} \right].$$

Exact solution in additive form:

$$w(x, t) = -\frac{1}{\beta} \ln(\beta t + C) + \frac{1}{\beta} \ln \left[ \int \frac{A - \beta x}{f(x)} dx + B \right],$$

where  $A$ ,  $B$ , and  $C$  are arbitrary constants.

**B.5.1-7.** Equations with three independent variables.

$$41. \quad \frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ f(x, y) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(x, y) \frac{\partial w}{\partial y} \right] + kw \ln w.$$

Exact solution in multiplicative form:

$$w(x, y, t) = \exp(Ae^{kt})\Theta(x, y),$$

where  $A$  is an arbitrary constant and the function  $\Theta(x, y)$  satisfies the stationary equation

$$\frac{\partial}{\partial x} \left[ f(x, y) \frac{\partial \Theta}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(x, y) \frac{\partial \Theta}{\partial y} \right] + k\Theta \ln \Theta = 0.$$

$$42. \quad \frac{\partial w}{\partial t} = \frac{\partial}{\partial x} \left[ f(x, t) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(y, t) \frac{\partial w}{\partial y} \right] + h(t) w \ln w.$$

Incomplete separable exact solution (the solution is separable in the space coordinates  $x$  and  $y$  but not in time  $t$ ):

$$w(x, y, t) = \varphi(x, t) \psi(y, t).$$

The functions  $\varphi(x, t)$  and  $\psi(y, t)$  are determined from the one-dimensional nonlinear parabolic differential equations

$$\begin{aligned} \frac{\partial \varphi}{\partial t} &= \frac{\partial}{\partial x} \left[ f(x, t) \frac{\partial \varphi}{\partial x} \right] + h(t) \varphi \ln \varphi + C(t) \varphi, \\ \frac{\partial \psi}{\partial t} &= \frac{\partial}{\partial y} \left[ g(y, t) \frac{\partial \psi}{\partial y} \right] + h(t) \psi \ln \psi - C(t) \psi, \end{aligned}$$

where  $C(t)$  is an arbitrary function.

## B.5.2. Hyperbolic Equations

|                                                                                                                                                        |
|--------------------------------------------------------------------------------------------------------------------------------------------------------|
| B.5.2-1. Equations of the form $\frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + f(x, t, w, \frac{\partial w}{\partial x})$ . |
|--------------------------------------------------------------------------------------------------------------------------------------------------------|

$$1. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + b w \ln w + [f(x) + g(t)] w.$$

Exact solution in multiplicative form:

$$w(x, t) = \varphi(t) \psi(x),$$

where the functions  $\varphi(t)$  and  $\psi(x)$  are determined by the second-order ordinary differential equations

$$\begin{aligned} \varphi''_{tt} - [b \ln \varphi + g(t) + C] \varphi &= 0, \\ a \psi''_{xx} + [b \ln \psi + f(x) - C] \psi &= 0, \end{aligned}$$

where  $C$  is an arbitrary constant.

$$2. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + c w + f(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(t) + \theta(\xi), \quad \xi = x + \lambda t,$$

where  $\lambda$  is an arbitrary constant and the functions  $\varphi = \varphi(t)$  and  $\theta = \theta(\xi)$  are determined by solving the second-order ordinary differential equations

$$\varphi''_{tt} - c \varphi - f(t) = 0, \tag{1}$$

$$(a - \lambda^2) \theta''_{\xi\xi} + b (\theta'_\xi)^2 + c \theta = 0. \tag{2}$$

The general solution of equation (1) is given by

$$\varphi(t) = C_1 \cosh(kt) + C_2 \sinh(kt) + \frac{1}{k} \int_0^t f(\tau) \sinh[k(t - \tau)] d\tau \quad \text{if } c = k^2 > 0,$$

$$\varphi(t) = C_1 \cos(kt) + C_2 \sin(kt) + \frac{1}{k} \int_0^t f(\tau) \sin[k(t - \tau)] d\tau \quad \text{if } c = -k^2 < 0,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

Equation (2) can be solved with the change of variable  $z(\theta) = (\theta'_\xi)^2$ , which leads to a first-order linear equation.

$$3. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw + f(x).$$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + \psi(t).$$

Here,

$$\psi(t) = C_1 \cosh(kt) + C_2 \sinh(kt) \quad \text{if } c = k^2 > 0,$$

$$\psi(t) = C_1 \cos(kt) + C_2 \sin(kt) \quad \text{if } c = -k^2 < 0,$$

where  $C_1$  and  $C_2$  are arbitrary constants, and the function  $\varphi(x)$  is determined by the ordinary differential equation

$$a\varphi''_{xx} + b(\varphi'_x)^2 + c\varphi + f(x) = 0.$$

$$4. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw \frac{\partial w}{\partial x} + kw^2 + f(t)w + g(t).$$

Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\lambda x),$$

where  $\lambda$  is a root of the quadratic equation  $b\lambda^2 + c\lambda + k = 0$  and the functions  $\varphi(t)$  and  $\psi(t)$  are determined by the following system of second-order ordinary differential equations:

$$\varphi''_{tt} = k\varphi^2 + f(t)\varphi + g(t), \quad (1)$$

$$\psi''_{tt} = [(c\lambda + 2k)\varphi + f(t) + a\lambda^2]\psi. \quad (2)$$

In the special case  $f(t) = \text{const}$  and  $g(t) = \text{const}$ , equation (1) is autonomous and has particular solutions of the form  $\varphi = \text{const}$  and, hence, can be integrated in quadrature. Equation (2) is linear in  $\psi$ , and consequently, with  $\varphi = \text{const}$ , its general solution is expressed in terms of exponentials or sine and cosine.

$$5. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + g(t)w + h(t).$$

Exact solution:

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t), \quad (1)$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by solving the following system of second-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\varphi''_{tt} = 4f\varphi^2 + g\varphi, \quad (2)$$

$$\psi''_{tt} = (4f\varphi + g)\psi, \quad (3)$$

$$\chi''_{tt} = g\chi + f\psi^2 + h + 2a\varphi. \quad (4)$$

Equation (2) has the trivial particular solution  $\varphi(t) \equiv 0$ ; the corresponding solution (1) is linear in the coordinate  $x$ .

Equation (3) has a particular solution  $\psi = \bar{\varphi}(t)$ , where  $\bar{\varphi}(t)$  is any nontrivial particular solution of equation (2). Hence, the general solution of equation (3) is given by

$$\psi(t) = C_1 \bar{\varphi}(t) + C_2 \bar{\varphi}(t) \int \frac{dt}{\bar{\varphi}^2(t)},$$

where  $C_1$  and  $C_2$  are arbitrary constants. If the functions  $f$  and  $g$  proportional, then  $\varphi = -\frac{1}{4}g/f$  ( $\varphi = \text{const}$ ) is a particular solution of equation (2).

Equation (4) linear in  $\chi = \chi(t)$ .



$$6. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + f(x) \left( \frac{\partial w}{\partial x} \right)^2 + g(x) + h(t).$$

Exact solution in additive form:

$$w(x, t) = \frac{1}{2} At^2 + Bt + C + \int_0^t (t - \tau) h(\tau) d\tau + \varphi(x).$$

Here,  $A$ ,  $B$ , and  $C$  are arbitrary constants, and the function  $\varphi(x)$  is determined by solving the second-order nonlinear ordinary differential equation

$$a\varphi''_{xx} + f(x)(\varphi'_x)^2 + g(x) - A = 0.$$

$$7. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + f(x) \left( \frac{\partial w}{\partial x} \right)^2 + bw + g(x) + h(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(t) + \psi(x).$$

Here, the functions  $\varphi(t)$  and  $\psi(x)$  are determined by solving the second-order ordinary differential equations

$$\begin{aligned} \varphi''_{tt} - b\varphi - h(t) &= 0, \\ a\psi''_{xx} + f(x)(\psi'_x)^2 + b\psi + g(x) &= 0. \end{aligned}$$

The general solution of the first equation is given by

$$\begin{aligned} \varphi(t) &= C_1 \cosh(kt) + C_2 \sinh(kt) + \frac{1}{k} \int_0^t h(\tau) \sinh[k(t - \tau)] d\tau \quad \text{if } b = k^2 > 0, \\ \varphi(t) &= C_1 \cos(kt) + C_2 \sin(kt) + \frac{1}{k} \int_0^t h(\tau) \sin[k(t - \tau)] d\tau \quad \text{if } b = -k^2 < 0, \end{aligned}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

$$8. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + bf(t)w^2 + g(t)w + h(t).$$

1°. Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\pm x\sqrt{-b}), \quad b < 0,$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following second-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\begin{aligned} \varphi''_{tt} &= bf\varphi^2 + g\varphi + h, \\ \psi''_{tt} &= (2bf\varphi + g - ab)\psi. \end{aligned}$$

2°. Exact solution of a more general form:

$$w(x, t) = \varphi(t) + \psi(t) [A \exp(x\sqrt{-b}) + B \exp(-x\sqrt{-b})], \quad b < 0,$$

where  $A$  and  $B$  are arbitrary constants and the functions  $\varphi(t)$  and  $\psi(t)$  are determined by the following system of second-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\begin{aligned} \varphi''_{tt} &= bf(\varphi^2 + 4AB\psi^2) + g\varphi + h, \\ \psi''_{tt} &= (2bf\varphi + g - ab)\psi. \end{aligned}$$

3°. Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \cos(x\sqrt{b} + C), \quad b > 0, \quad (9)$$

where  $C$  is an arbitrary constant and the functions  $\varphi(t)$  and  $\psi(t)$  are determined by the following system of second-order ordinary differential equations with variable coefficients:

$$\begin{aligned}\varphi''_{tt} &= bf(\varphi^2 + \psi^2) + g\varphi + h, \\ \psi''_{tt} &= (2bf\varphi + g - ab)\psi.\end{aligned}$$

9. 
$$\frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + f\left(x, \frac{\partial w}{\partial x}\right) + g(t).$$

Exact solution in additive form:

$$w(x, t) = \frac{1}{2}At^2 + Bt + C + \int_0^t (t - \tau)g(\tau) d\tau + \varphi(x).$$

Here,  $A$ ,  $B$ , and  $C$  are arbitrary constants and the function  $\varphi(x)$  is determined by solving the second-order nonlinear ordinary differential equation

$$a\varphi''_{xx} + f(x, \varphi'_x) - A = 0.$$

10. 
$$\frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + f\left(x, \frac{\partial w}{\partial x}\right) + bw + g(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(t) + \psi(x).$$

Here, the functions  $\varphi(t)$  and  $\psi(x)$  are determined by the second-order ordinary differential equations

$$\begin{aligned}\varphi''_{tt} - b\varphi - g(t) &= 0, \\ a\psi''_{xx} + f(x, \psi'_x) + b\psi &= 0.\end{aligned}$$

The general solution of the first equation is given by

$$\begin{aligned}\varphi(t) &= C_1 \cosh(kt) + C_2 \sinh(kt) + \frac{1}{k} \int_0^t g(\tau) \sinh[k(t - \tau)] d\tau \quad \text{if } b = k^2 > 0, \\ \varphi(t) &= C_1 \cos(kt) + C_2 \sin(kt) + \frac{1}{k} \int_0^t g(\tau) \sin[k(t - \tau)] d\tau \quad \text{if } b = -k^2 < 0,\end{aligned}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

11. 
$$\frac{\partial^2 w}{\partial t^2} = a \frac{\partial^2 w}{\partial x^2} + wf\left(t, \frac{1}{w} \frac{\partial w}{\partial x}\right).$$

Exact solution in multiplicative form:

$$w(x, t) = e^{\lambda x} \varphi(t),$$

where  $\lambda$  is an arbitrary constant and the function  $\varphi(t)$  is determined by the second-order linear ordinary differential equation

$$\varphi''_{tt} = [a\lambda^2 + f(t, \lambda)]\varphi.$$

$$12. \quad \frac{\partial^2 w}{\partial t^2} = a(x + \beta)^n \frac{\partial^2 w}{\partial x^2} + f(w), \quad a > 0.$$

Exact solution for  $n \neq 2$ :

$$w = w(z), \quad z = \left[ \frac{1}{4}a(2-n)^2(t+C)^2 - (x+\beta)^{2-n} \right]^{\frac{n}{2(2-n)}},$$

where  $C$  is an arbitrary constant and the function  $w = w(z)$  is determined by the generalized Emden–Fowler equation

$$w''_{zz} - \frac{4}{an^2} z^{\frac{4(1-n)}{n}} f(w) = 0. \quad (1)$$

A number of exact solutions to equation (1) for some specific  $f = f(w)$  can be found in Polyanin and Zaitsev (1995). In the special case  $n = 1$ , the general solution of equation (1) is given by

$$\int \left[ C_1 + \frac{8}{a} F(w) \right]^{-1/2} dw = \pm z + C_2, \quad F(w) = \int f(w) dw,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

$$13. \quad \frac{\partial^2 w}{\partial t^2} = ax^n \frac{\partial^2 w}{\partial x^2} + bx^{n-1} \frac{\partial w}{\partial x} + f(w), \quad a > 0.$$

Exact solution for  $n \neq 2$ :

$$w = w(\xi), \quad \xi = \frac{1}{4}a(2-n)^2(t+C)^2 - x^{2-n},$$

where  $C$  is an arbitrary constant and the function  $w = w(\xi)$  is determined by the ordinary differential equation

$$\xi w''_{\xi\xi} + Aw'_\xi - Bf(w) = 0, \quad \text{where} \quad A = \frac{a(4-3n)+2b}{2a(2-n)}, \quad B = \frac{1}{a(2-n)^2}.$$

For  $A \neq 1$ , the change of variable  $\xi = kz^{\frac{1}{1-A}}$  ( $k = \pm 1$ ) brings this equation to the generalized Emden–Fowler equation

$$w''_{zz} - \frac{kB}{(1-A)^2} z^{\frac{2A-1}{1-A}} f(w) = 0,$$

whose solvable cases are presented in Polyanin and Zaitsev (1995).

$$14. \quad \frac{\partial^2 w}{\partial t^2} = ax^n \frac{\partial^2 w}{\partial x^2} + x^{n-1} f(w) \frac{\partial w}{\partial x}.$$

Exact solution for  $n \neq 2$ :

$$w = w(z), \quad z = [ka(2-n)^2(t+C)^2 - 4kx^{2-n}]^{1/2}, \quad k = \pm 1,$$

where  $C$  is an arbitrary constant and the function  $w = w(z)$  is determined by the ordinary differential equation

$$w''_{zz} + \frac{2}{a(2-n)} [a(1-n) + f(w)] \frac{1}{z} w'_z = 0.$$

The change of variable  $u(w) = zw'_z$  leads to a first-order separable equation. Integrating this equation yields the general solution in implicit form:

$$\int \frac{dw}{anw - 2F(w) + C_1} = \frac{1}{a(2-n)} \ln |z| + C_2, \quad F(w) = \int f(w) dw,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

$$15. \quad \frac{\partial^2 w}{\partial t^2} = ax^n \frac{\partial^2 w}{\partial x^2} + x^{n-1} f(w) \frac{\partial w}{\partial x} + g(w).$$

Exact solution for  $n \neq 2$ :

$$w = w(z), \quad z = [ka(2-n)^2(t+C)^2 - 4kx^{2-n}]^{1/2}, \quad k = \pm 1,$$

where  $C$  is an arbitrary constant and the function  $w = w(z)$  is determined by the ordinary differential equation

$$w''_{zz} + \frac{2}{a(2-n)} [a(1-n) + f(w)] \frac{1}{z} w'_z - \frac{1}{ak(2-n)^2} g(w) = 0.$$

$$16. \quad \frac{\partial^2 w}{\partial t^2} = ae^{\lambda x} \frac{\partial^2 w}{\partial x^2} + be^{\lambda x} \frac{\partial w}{\partial x} + f(w), \quad a > 0.$$

Exact solution for  $\lambda \neq 0$ :

$$w = w(z), \quad z = [4ke^{-\lambda x} - ak\lambda^2(t+C)^2]^{1/2}, \quad k = \pm 1,$$

where  $C$  is an arbitrary constant and the function  $w = w(z)$  is determined by the ordinary differential equation

$$w''_{zz} + \frac{2(a\lambda - b)}{a\lambda} \frac{1}{z} w'_z + \frac{1}{ak\lambda^2} f(w) = 0. \quad (1)$$

For  $b = a\lambda$ , the solution of equation (1) is given by

$$\int \left[ C_1 - \frac{2}{ak\lambda^2} F(w) \right]^{-1/2} dw = \pm z + C_2, \quad F(w) = \int f(w) dw,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

For  $b \neq \frac{1}{2}a\lambda$ , the change of variable  $\xi = z \frac{2b-a\lambda}{a\lambda}$  brings (1) to the generalized Emden–Fowler equation

$$w''_{\xi\xi} + \frac{a}{k(2b-a\lambda)^2} \xi^{\frac{4(a\lambda-b)}{2b-a\lambda}} f(w) = 0,$$

whose solvable cases are presented in Polyanin and Zaitsev (1995).

$$17. \quad \frac{\partial^2 w}{\partial t^2} = ae^{\lambda x} \frac{\partial^2 w}{\partial x^2} + e^{\lambda x} f(w) \frac{\partial w}{\partial x}.$$

Exact solution for  $\lambda \neq 0$ :

$$w = w(z), \quad z = [4ke^{-\lambda x} - ak\lambda^2(t+A)^2]^{1/2}, \quad k = \pm 1.$$

Here,  $A$  is an arbitrary constant and the function  $w = w(z)$  is determined by the ordinary differential equation

$$w''_{zz} + \frac{2}{z} \left[ 1 - \frac{1}{a\lambda} f(w) \right] w'_z = 0,$$

which by the change of variable  $u(w) = zw'_z$  is reduced to a separable first-order equation. Integrating this equation yields the general solution in implicit form:

$$\int \frac{dw}{2F(w) - a\lambda w + C_1} = \frac{1}{a\lambda} \ln |z| + C_2, \quad F(w) = \int f(w) dw,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

$$18. \quad \frac{\partial^2 w}{\partial t^2} = ae^{\lambda x} \frac{\partial^2 w}{\partial x^2} + e^{\lambda x} f(w) \frac{\partial w}{\partial x} + g(w).$$

Exact solution for  $\lambda \neq 0$ :

$$w = w(z), \quad z = [4ke^{-\lambda x} - ak\lambda^2(t+C)^2]^{1/2}, \quad k = \pm 1,$$

where  $C$  is an arbitrary constant and the function  $w = w(z)$  is determined by the ordinary differential equation

$$w''_{zz} + \frac{2}{z} \left[ 1 - \frac{1}{a\lambda} f(w) \right] w'_z + \frac{1}{ak\lambda^2} g(w) = 0.$$

### B.5.2-3. Other equations.

$$19. \quad \frac{\partial^2 w}{\partial t^2} = f(t) \frac{\partial}{\partial x} \left( w \frac{\partial w}{\partial x} \right).$$

1°. Exact solutions:

$$w(x, t) = (C_1 t + C_2)(C_3 x + C_4)^{1/2},$$

$$w(x, t) = (C_1 t + C_2)x + \int_a^t (t - \tau)(C_1 \tau + C_2)^2 f(\tau) d\tau + C_3 t + C_4,$$

where  $C_1, C_2, C_3, C_4$ , and  $a$  are arbitrary constants.

2°. Exact solution:

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t),$$

where the functions  $\varphi = \varphi(t)$ ,  $\psi = \psi(t)$ , and  $\chi = \chi(t)$  are determined by the system of ordinary differential equations

$$\varphi''_{tt} = 6f(t)\varphi^2,$$

$$\psi''_{tt} = 6f(t)\varphi\psi,$$

$$\chi''_{tt} = 2f(t)\varphi\chi + f(t)\psi^2.$$

3°. Exact solution in multiplicative form:

$$w(x, t) = \Phi(t)\Psi(x),$$

where the functions  $\Phi = \Phi(t)$  and  $\Psi = \Psi(x)$  are determined by the ordinary differential equations ( $C$  is an arbitrary constant)

$$\Phi''_{tt} = C f(t)\Phi^2,$$

$$(\Psi\Psi'_x)'_x = C\Psi.$$

The latter equation is autonomous and has a particular solution  $\Psi = \frac{1}{6}Cx^2$  and, hence, is integrable in quadrature.

$$20. \quad \frac{\partial^2 w}{\partial t^2} = f(t) \frac{\partial}{\partial x} \left( w \frac{\partial w}{\partial x} \right) + g_2(t)x^2 + g_1(t)x + g_0(t).$$

Exact solution:

$$w(x, t) = \varphi(t)x^2 + \psi(t)x + \chi(t),$$

where the functions  $\varphi = \varphi(t)$ ,  $\psi = \psi(t)$ , and  $\chi = \chi(t)$  are determined by the system of ordinary differential equations

$$\varphi''_{tt} = 6f(t)\varphi^2 + g_2(t),$$

$$\psi''_{tt} = 6f(t)\varphi\psi + g_1(t),$$

$$\chi''_{tt} = 2f(t)\varphi\chi + f(t)\psi^2 + g_0(t).$$

### B.5.3. Elliptic Equations

B.5.3-1. Equations of the form  $a \frac{\partial^2 w}{\partial x^2} + b \frac{\partial^2 w}{\partial y^2} = f\left(x, y, w, \frac{\partial w}{\partial x}, \frac{\partial w}{\partial y}\right), \quad ab > 0.$

$$1. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = aw \ln w + [f(x) + g(y)]w.$$

Exact solution in multiplicative form:

$$w(x, y) = \varphi(x)\psi(y),$$

where the functions  $\varphi(x)$  and  $\psi(y)$  are determined by the ordinary differential equations

$$\varphi''_{xx} - [a \ln \varphi + f(x) + C] \varphi = 0,$$

$$\psi''_{yy} - [a \ln \psi + g(y) - C] \psi = 0,$$

where  $C$  is an arbitrary constant.

$$2. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = f(x)w \ln w + [af(x)y + g(x)]w.$$

Exact solution in multiplicative form:

$$w(x, y) = e^{-ay} \varphi(x),$$

where the function  $\varphi(x)$  is determined by the ordinary differential equation

$$\varphi''_{xx} = f(x)\varphi \ln \varphi + [g(x) - a^2]\varphi.$$

$$3. \quad \frac{\partial^2 w}{\partial x^2} + a \frac{\partial^2 w}{\partial y^2} = f(x) \left( \frac{\partial w}{\partial y} \right)^2 + g(x)w + h(x).$$

Exact solution:

$$w(x, y) = \varphi(x)y^2 + \psi(x)y + \chi(x). \quad (1)$$

Here, the functions  $\varphi(x)$ ,  $\psi(x)$ , and  $\chi(x)$  are determined by the following second-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\varphi''_{xx} = 4f\varphi^2 + g\varphi, \quad (2)$$

$$\psi''_{xx} = (4f\varphi + g)\psi, \quad (3)$$

$$\chi''_{xx} = g\chi + f\psi^2 + h - 2a\varphi. \quad (4)$$

If a solution  $\varphi = \varphi(x)$  of the nonlinear equation (2) is found, then the functions  $\psi = \psi(x)$  and  $\chi = \chi(x)$  can be determined successively from equations (3) and (4), which are linear in  $\psi$  and  $\chi$ .

By comparing equations (2) and (3), one can see that equation (3) has a particular solution  $\psi = \varphi(x)$ . Hence, the general solution of (3) is given by (see Polyanin and Zaitsev 1995)

$$\psi(x) = C_1\varphi(x) + C_2\varphi(x) \int \frac{dx}{\varphi^2(x)}, \quad \varphi \neq 0.$$

Note that equation (2) has the trivial particular solution  $\varphi(x) \equiv 0$ , to which there is a corresponding solution (1) linear in the coordinate  $y$ . If the functions  $f$  and  $g$  are proportional, then  $\varphi = -\frac{1}{4}g/f$  ( $\varphi = \text{const}$ ) is a particular solution of equation (2).

$$4. \quad \frac{\partial^2 w}{\partial x^2} + a \frac{\partial^2 w}{\partial y^2} = f(x) \left( \frac{\partial w}{\partial y} \right)^2 + bf(x)w^2 + g(x)w + h(x).$$

1°. Exact solution:

$$w(x, y) = \varphi(x) + \psi(x) \exp(\pm y\sqrt{-b}), \quad b < 0,$$

where the functions  $\varphi(x)$  and  $\psi(x)$  are determined by the following second-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\varphi''_{xx} = bf\varphi^2 + g\varphi + h,$$

$$\psi''_{xx} = (2bf\varphi + g + ab)\psi.$$

2°. Exact solution of a more general form:

$$w(x, y) = \varphi(x) + \psi(x) [A \exp(y\sqrt{-b}) + B \exp(-y\sqrt{-b})], \quad b < 0,$$

where the functions  $\varphi(x)$  and  $\psi(x)$  are determined by the following system of second-order ordinary differential equations with variable coefficients:

$$\varphi''_{xx} = bf(\varphi^2 + 4AB\psi^2) + g\varphi + h,$$

$$\psi''_{xx} = 2bf\varphi\psi + g\psi + ab\psi.$$

3°. Exact solution:

$$w(x, y) = \varphi(x) + \psi(x) \cos(y\sqrt{b} + C), \quad b > 0,$$

where  $C$  is an arbitrary constant and the functions  $\varphi(x)$  and  $\psi(x)$  are determined by the following system of second-order ordinary differential equations with variable coefficients:

$$\varphi''_{xx} = bf(\varphi^2 + \psi^2) + g\varphi + h,$$

$$\psi''_{xx} = 2bf\varphi\psi + g\psi + ab\psi.$$

⊙ Reference: V. F. Zaitsev, A. D. Polyanin (1996).

$$5. \quad a \frac{\partial^2 w}{\partial x^2} + b \frac{\partial^2 w}{\partial y^2} = f_1 \left( x, \frac{\partial w}{\partial x} \right) + f_2 \left( y, \frac{\partial w}{\partial y} \right) + kw.$$

Exact solution in additive form:

$$w(x, y) = \varphi(x) + \psi(y).$$

Here, the functions  $\varphi(x)$  and  $\psi(y)$  are determined by solving the second-order ordinary differential equations

$$\begin{aligned} a\varphi''_{xx} - f_1(x, \varphi'_x) - k\varphi &= C, \\ b\psi''_{yy} - f_2(y, \psi'_y) - k\psi &= -C, \end{aligned}$$

where  $C$  is an arbitrary constant.

**B.5.3-2.** Equations of the form  $\frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(y) \frac{\partial w}{\partial y} \right] = h(w).$

$$6. \quad \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( by^m \frac{\partial w}{\partial y} \right) = f(w).$$

1°. For  $n \neq 2$  and  $m \neq 2$ , there are exact solutions of the form

$$w = w(\xi), \quad \xi = [b(2-m)^2 x^{2-n} + a(2-n)^2 y^{2-m}]^{1/2}.$$

Here, the function  $w = w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{A}{\xi} w'_\xi = Bf(w), \quad (1)$$

where

$$A = \frac{4-nm}{(2-n)(2-m)}, \quad B = \frac{4}{ab(2-n)^2(2-m)^2}.$$

For  $m = 4/n$ , one obtains from (1) the following exact solution to the original equation with arbitrary  $f = f(w)$ :

$$\int \left[ C_1 + \frac{2n^2}{ab(2-n)^4} F(w) \right]^{-1/2} dw = C_2 \pm \xi, \quad F(w) = \int f(w) dw,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

2°. The change of variable  $\zeta = \xi^{1-A}$  brings (1) to the generalized Emden–Fowler equation

$$w''_{\zeta\zeta} = \frac{B}{(1-A)^2} \zeta^{\frac{2A}{1-A}} f(w). \quad (2)$$

A lot of exact solutions to equation (2) with various  $f = f(w)$  can be found in Polyanin and Zaitsev (1995).

⊙ *Reference:* V. F. Zaitsev and A. D. Polyanin (1996).

$$7. \quad \frac{\partial}{\partial x} \left( ae^{\beta x} \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( be^{\mu y} \frac{\partial w}{\partial y} \right) = f(w).$$

For  $\beta\mu \neq 0$ , there are exact solutions of the form

$$w = w(\xi), \quad \xi = (b\mu^2 e^{-\beta x} + a\beta^2 e^{-\mu y})^{1/2},$$

where the function  $w = w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} - \frac{1}{\xi} w'_\xi = Af(w), \quad A = \frac{4}{ab\beta^2\mu^2}. \quad (1)$$

The change of variable  $\zeta = \xi^2$  brings (1) to the generalized Emden–Fowler equation

$$w''_{\zeta\zeta} = \frac{1}{4} A \zeta^{-1} f(w),$$

whose solutions with  $f(w) = (kw + s)^{-1}$  and  $f(w) = (kw + s)^{-2}$  ( $k, s = \text{const}$ ) can be found in Polyanin and Zaitsev (1995).

⊙ *Reference:* V. F. Zaitsev and A. D. Polyanin (1996).

$$8. \quad \frac{\partial}{\partial x} \left( a x^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( b e^{\mu y} \frac{\partial w}{\partial y} \right) = f(w).$$

For  $n \neq 2$  and  $\mu \neq 0$ , there are exact solutions of the form

$$w = w(\xi), \quad \xi = [b\mu^2 x^{2-n} + a(2-n)^2 e^{-\mu y}]^{1/2},$$

where the function  $w = w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{n}{n-2} \frac{1}{\xi} w'_{\xi} = \frac{4}{ab\mu^2(2-n)^2} f(w).$$

$$9. \quad \frac{\partial}{\partial x} \left[ f(x) \frac{\partial w}{\partial x} \right] + \frac{\partial}{\partial y} \left[ g(y) \frac{\partial w}{\partial y} \right] = a w \ln w + [h_1(x) + h_2(y)]w.$$

Exact solution in multiplicative form:

$$w(x, y) = \varphi(x)\psi(y),$$

where  $\varphi = \varphi(x)$  and  $\psi = \psi(y)$  are determined by the ordinary differential equations

$$[f(x)\varphi'_x]'_x = [a \ln \varphi + h_1(x) + C]\varphi,$$

$$[g(y)\psi'_y]'_y = [a \ln \psi + h_2(y) - C]\psi,$$

where  $C$  is an arbitrary constant.

**B.5.3-3. Other equations with two independent variables.**

$$10. \quad \frac{\partial^2 w}{\partial x^2} + \frac{\partial}{\partial y} \left\{ [f(x)w + g(x)] \frac{\partial w}{\partial y} \right\} = 0.$$

1°. Exact solution:

$$w(x, y) = (Ax + B)y - \int_{x_0}^x (x-t)(At+B)^2 f(t) dt + C_1 x + C_2,$$

where  $A, B, C_1, C_2$ , and  $x_0$  are arbitrary constants.

2°. Exact solution:

$$w(x, y) = \varphi(x)y^2 + \psi(x)y + \chi(x),$$

where the functions  $\varphi = \varphi(x)$ ,  $\psi = \psi(x)$ , and  $\chi = \chi(x)$  are determined by the ordinary differential equations

$$\varphi''_{xx} + 6f\varphi^2 = 0, \tag{1}$$

$$\psi''_{xx} + 6f\varphi\psi = 0, \tag{2}$$

$$\chi''_{xx} + 2f\varphi\chi + 2\varphi g + f\psi^2 = 0. \tag{3}$$

The nonlinear equation (1) can be treated independently. For  $f \equiv \text{const}$ , its solution can be expressed in terms of elliptic integrals. For  $f = ae^{\lambda x}$ , a particular solution of (1) is  $\varphi = -\frac{\lambda^2}{6a}e^{-\lambda x}$ . Equations (2) and (3) can be solved successively (these are linear in the unknowns). Because  $\psi = \varphi(x)$  is a particular solution of equation (2), the general solution of (2) is given by (see Polyanin and Zaitsev 1995)

$$\psi(x) = C_1\varphi(x) + C_2\varphi(x) \int \frac{dx}{\varphi^2(x)},$$

where  $C_1$  and  $C_2$  are arbitrary constants.



$$11. \quad ax^n \frac{\partial^2 w}{\partial x^2} + by^m \frac{\partial^2 w}{\partial y^2} + kx^{n-1} \frac{\partial w}{\partial x} + sy^{m-1} \frac{\partial w}{\partial y} = f(w).$$

For  $n \neq 2$  and  $m \neq 2$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi = [b(2-m)^2 x^{2-n} + a(2-n)^2 y^{2-m}]^{1/2}.$$

Here, the function  $w = w(\xi)$  is determined by the ordinary differential equation

$$Aw''_{\xi\xi} + \frac{B}{\xi} w'_\xi = f(w),$$

where

$$A = \frac{1}{4}ab(2-n)^2(2-m)^2,$$

$$B = \frac{1}{4}(2-n)(2-m)[ab(3nm-4n-4m+4) + 2bk(2-m) + 2as(2-n)].$$

⊙ Reference: V. F. Zaitsev and A. D. Polyanin (1996).

$$12. \quad ax^n \frac{\partial^2 w}{\partial x^2} + by^m \frac{\partial^2 w}{\partial y^2} + kx^{n-1} f(w) \frac{\partial w}{\partial x} + sy^{m-1} f(w) \frac{\partial w}{\partial y} = g(w).$$

For  $n \neq 2$  and  $m \neq 2$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi = [b(2-m)^2 x^{2-n} + a(2-n)^2 y^{2-m}]^{1/2}.$$

$$13. \quad ae^{\beta x} \frac{\partial^2 w}{\partial x^2} + be^{\mu y} \frac{\partial^2 w}{\partial y^2} + ke^{\beta x} \frac{\partial w}{\partial x} + se^{\mu y} \frac{\partial w}{\partial y} = f(w).$$

For  $\beta\mu \neq 0$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi = (b\mu^2 e^{-\beta x} + a\beta^2 e^{-\mu y})^{1/2}.$$

Here, the function  $w = w(\xi)$  is determined by the ordinary differential equation

$$Aw''_{\xi\xi} + \frac{B}{\xi} w'_\xi = f(w),$$

where

$$A = \frac{1}{4}ab\beta^2\mu^2, \quad B = \frac{1}{4}\beta\mu(3ab\beta\mu - 2bk\mu - 2as\beta).$$

$$14. \quad ae^{\beta x} \frac{\partial^2 w}{\partial x^2} + be^{\mu y} \frac{\partial^2 w}{\partial y^2} + ke^{\beta x} f(w) \frac{\partial w}{\partial x} + se^{\mu y} f(w) \frac{\partial w}{\partial y} = g(w).$$

For  $\beta\mu \neq 0$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi = (b\mu^2 e^{-\beta x} + a\beta^2 e^{-\mu y})^{1/2}.$$

$$15. \quad ax^n \frac{\partial^2 w}{\partial x^2} + be^{\beta y} \frac{\partial^2 w}{\partial y^2} + kx^{n-1} \frac{\partial w}{\partial x} + se^{\beta y} \frac{\partial w}{\partial y} = f(w).$$

For  $\beta \neq 0$  and  $n \neq 2$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi = [b\beta^2 x^{2-n} + a(2-n)^2 e^{-\beta y}]^{1/2}.$$

Here, the function  $w = w(\xi)$  is determined by the ordinary differential equation

$$Aw''_{\xi\xi} + \frac{B}{\xi} w'_\xi = f(w),$$

where

$$A = \frac{1}{4}ab\beta^2(2-n)^2, \quad B = \frac{1}{4}\beta(2-n)[ab\beta(4-3n) + 2bk\beta - 2as(2-n)].$$

⊙ Reference: V. F. Zaitsev and A. D. Polyanin (1996).

$$16. \quad ax^n \frac{\partial^2 w}{\partial x^2} + be^{\beta y} \frac{\partial^2 w}{\partial y^2} + kx^{n-1} f(w) \frac{\partial w}{\partial x} + se^{\beta y} f(w) \frac{\partial w}{\partial y} = g(w).$$

For  $\beta \neq 0$  and  $n \neq 2$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi = [b\beta^2 x^{2-n} + a(2-n)^2 e^{-\beta y}]^{1/2}.$$

**B.5.3-4. Equations with three independent variables.**

$$17. \quad \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( by^m \frac{\partial w}{\partial y} \right) + \frac{\partial}{\partial z} \left( cz^l \frac{\partial w}{\partial z} \right) = f(w).$$

For  $n \neq 2$ ,  $m \neq 2$ , and  $l \neq 2$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi^2 = 4 \left[ \frac{x^{2-n}}{a(2-n)^2} + \frac{y^{2-m}}{b(2-m)^2} + \frac{z^{2-l}}{c(2-l)^2} \right],$$

where the function  $w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{A}{\xi} w'_\xi = f(w), \quad A = 2 \left( \frac{1}{2-n} + \frac{1}{2-m} + \frac{1}{2-l} \right) - 1.$$

⊙ *Reference:* A. D. Polyanin and A. I. Zhurov (1998).

$$18. \quad \frac{\partial}{\partial x} \left( ae^{\lambda x} \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( be^{\mu y} \frac{\partial w}{\partial y} \right) + \frac{\partial}{\partial z} \left( ce^{\nu z} \frac{\partial w}{\partial z} \right) = f(w).$$

For  $\lambda \neq 0$ ,  $\mu \neq 0$ , and  $\nu \neq 0$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi^2 = 4 \left( \frac{e^{-\lambda x}}{a\lambda^2} + \frac{e^{-\mu y}}{b\mu^2} + \frac{e^{-\nu z}}{c\nu^2} \right),$$

where the function  $w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} - \frac{1}{\xi} w'_\xi = f(w).$$

$$19. \quad \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( by^m \frac{\partial w}{\partial y} \right) + \frac{\partial}{\partial z} \left( ce^{\nu z} \frac{\partial w}{\partial z} \right) = f(w).$$

For  $n \neq 2$ ,  $m \neq 2$ , and  $\nu \neq 0$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi^2 = 4 \left[ \frac{x^{2-n}}{a(2-n)^2} + \frac{y^{2-m}}{b(2-m)^2} + \frac{e^{-\nu z}}{c\nu^2} \right],$$

where the function  $w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{A}{\xi} w'_\xi = f(w), \quad A = 2 \left( \frac{1}{2-n} + \frac{1}{2-m} \right) - 1.$$

$$20. \quad \frac{\partial}{\partial x} \left( ax^n \frac{\partial w}{\partial x} \right) + \frac{\partial}{\partial y} \left( be^{\mu y} \frac{\partial w}{\partial y} \right) + \frac{\partial}{\partial z} \left( ce^{\nu z} \frac{\partial w}{\partial z} \right) = f(w).$$

For  $n \neq 2$ ,  $\mu \neq 0$ , and  $\nu \neq 0$ , there is an exact solution of the form

$$w = w(\xi), \quad \xi^2 = 4 \left[ \frac{x^{2-n}}{a(2-n)^2} + \frac{e^{-\mu y}}{b\mu^2} + \frac{e^{-\nu z}}{c\nu^2} \right],$$

where the function  $w(\xi)$  is determined by the ordinary differential equation

$$w''_{\xi\xi} + \frac{n}{2-n} \frac{1}{\xi} w'_\xi = f(w).$$

⊙ *Reference:* A. D. Polyanin and A. I. Zhurov (1998).

## B.5.4. Equations Containing Mixed Derivatives

B.5.4-1. Monge–Ampère equations.

$$1. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x).$$

1°. Exact solutions:

$$w(x, y) = C_1 y^2 + C_2 x y + \frac{C_2^2}{4C_1} x^2 - \frac{1}{2C_1} \int_0^x (x-t)f(t) dt + C_3 y + C_4 x + C_5,$$

$$w(x, y) = \frac{1}{x + C_1} \left( C_2 y^2 + C_3 y + \frac{C_3^2}{4C_2} \right) - \frac{1}{2C_2} \int_0^x (x-t)(t + C_1)f(t) dt + C_4 y + C_5 x + C_6,$$

where  $C_1, C_2, C_3, C_4, C_5$ , and  $C_6$  are arbitrary constants.

2°. Exact solutions for  $f(x) > 0$ :

$$w(x, y) = \pm y \int \sqrt{f(x)} dx + \varphi(x) + C_1 y,$$

where  $\varphi(x)$  is an arbitrary function.

$$2. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)y.$$

1°. Exact solution:

$$w(x, y) = C_1 y^2 - y \int F(x) dx + \frac{1}{2C_1} \int_a^x (x-t)F^2(t) dt + C_2 x + C_3 y + C_4,$$

$$F(x) = \frac{1}{2C_1} \int f(x) dx + C_5,$$

where  $C_1, \dots, C_5$ , and  $a$  are arbitrary constants.

2°. Exact solution:

$$w(x, y) = \varphi(x)y^2 + \psi(x)y + \chi(x),$$

where

$$\varphi(x) = \frac{1}{C_1 x + C_2}, \quad \psi(x) = C_3 \varphi(x) + C_4 + \frac{\varphi(x)}{2C_1} \int \frac{f(x) dx}{[\varphi(x)]^3} - \frac{1}{2C_1} \int \frac{f(x) dx}{[\varphi(x)]^2},$$

$$\chi(x) = \frac{1}{2} \int_a^x (x-t) \frac{[\psi'_t(t)]^2}{\varphi(t)} dt + C_5 x + C_6,$$

3°. Exact solutions cubic in  $y$ :

$$w(x, y) = C_1 y^3 - \frac{1}{6C_1} \int_a^x (x-t)f(t) dt + C_2 x + C_3 y + C_4,$$

$$w(x, y) = \frac{y^3}{(C_1 x + C_2)^2} - \frac{1}{6} \int_a^x (x-t)(C_1 t + C_2)^2 f(t) dt + C_3 x + C_4 y + C_5,$$

where  $C_1, \dots, C_5$ , and  $a$  are arbitrary constants.

4°. See the solution of equation B.5.4.5 in Item 2° with  $k = 1$ .

$$3. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)y^2.$$

1°. Exact solution quadratic in  $y$ :

$$w(x, y) = \varphi(x)y^2 + \left[ C_1 \int \varphi^2(x) dx + C_2 \right] y + \frac{1}{2} C_1^2 \int_a^x (x-t)\varphi^3(t) dt + C_3 x + C_4.$$

The function  $\varphi = \varphi(x)$  is determined by the ordinary differential equation

$$\varphi \varphi_{xx}'' = 2(\varphi_x')^2 - \frac{1}{2} f(x).$$

2°. Exact solutions quartic in  $y$ :

$$w(x, y) = C_1 y^4 - \frac{1}{12 C_1} \int_a^x (x-t)f(t) dt + C_2 x + C_3 y + C_4,$$

$$w(x, y) = \frac{y^4}{(C_1 x + C_2)^3} - \frac{1}{12} \int_a^x (x-t)(C_1 t + C_2)^3 f(t) dt + C_3 x + C_4 y + C_5,$$

where  $C_1, \dots, C_5$ , and  $a$  are arbitrary constants.

3°. See the solution of equation B.5.4.5 in Item 2° with  $k = 2$ .

$$4. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)y^2 + g(x)y + h(x).$$

Exact solution:

$$w(x, y) = \varphi(x)y^2 + \psi(x)y + \chi(x),$$

where the functions  $\varphi = \varphi(x)$ ,  $\psi = \psi(x)$ , and  $\chi = \chi(x)$  are determined by the system of ordinary differential equations

$$\varphi \varphi_{xx}'' = 2(\varphi_x')^2 - \frac{1}{2} f(x),$$

$$\varphi \psi_{xx}'' = 2\varphi_x' \psi_x' - \frac{1}{2} g(x),$$

$$\varphi \chi_{xx}'' = \frac{1}{2} (\psi_x')^2 - \frac{1}{2} h(x).$$

$$5. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)y^k.$$

1°. Exact solutions:

$$w(x, y) = \frac{C_1 y^{k+2}}{(k+1)(k+2)} - \frac{1}{C_1} \int_a^x (x-t)f(t) dt + C_2 x + C_3 y + C_4,$$

$$w(x, y) = \frac{y^{k+2}}{(C_1 x + C_2)^{k+1}} - \frac{1}{(k+1)(k+2)} \int_a^x (x-t)(C_1 t + C_2)^{k+1} f(t) dt + C_3 x + C_4 y + C_5,$$

where  $C_1, \dots, C_5$ , and  $a$  are arbitrary constants.

2°. Exact solution:

$$w(x, y) = \varphi(x)y^{\frac{k+2}{2}}.$$

The function  $\varphi = \varphi(x)$  is determined by the ordinary differential equation

$$k(k+2)\varphi \varphi_{xx}'' - (k+2)^2(\varphi_x')^2 + 4f(x) = 0.$$

$$6. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)y^{2k+2} + g(x)y^k.$$

Exact solution:

$$w(x, y) = \varphi(x)y^{k+2} - \frac{1}{(k+1)(k+2)} \int_a^x (x-t) \frac{g(t)}{\varphi(t)} dt + C_1 x + C_2 y + C_3,$$

where  $\varphi = \varphi(x)$  is determined by the ordinary differential equation

$$(k+1)(k+2)\varphi\varphi''_{xx} - (k+2)^2(\varphi'_x)^2 + f(x) = 0.$$

$$7. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)e^{\lambda y}.$$

1°. Exact solutions:

$$w(x, y) = C_1 \int_a^x (x-t)f(t) dt + C_2 x - \frac{1}{C_1 \lambda^2} e^{\lambda y} + C_3 y + C_4,$$

$$w(x, y) = C_1 e^{\beta x + \lambda y} - \frac{1}{C_1 \lambda^2} \int_a^x (x-t)e^{-\beta t} f(t) dt + C_2 x + C_3 y + C_4,$$

where  $C_1, C_2, C_3, C_4, a$ , and  $\beta$  are arbitrary constants.

2°. Exact solution:

$$w(x, y) = \varphi(x) \exp\left(\frac{1}{2}\lambda y\right),$$

where  $\varphi = \varphi(x)$  is determined by the ordinary differential equation

$$\varphi\varphi''_{xx} - (\varphi'_x)^2 + 4\lambda^{-2}f(x) = 0.$$

$$8. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)e^{2\lambda y} + g(x)e^{\lambda y}.$$

Exact solution:

$$w(x, y) = \varphi(x)e^{\lambda y} - \frac{1}{\lambda^2} \int_a^x (x-t) \frac{g(t)}{\varphi(t)} dt + C_1 x + C_2 y + C_3,$$

where the function  $\varphi = \varphi(x)$  is determined by the ordinary differential equation

$$\varphi\varphi''_{xx} - (\varphi'_x)^2 + \lambda^{-2}f(x) = 0.$$

$$9. \quad \left( \frac{\partial^2 w}{\partial x \partial y} \right)^2 = \frac{\partial^2 w}{\partial x^2} \frac{\partial^2 w}{\partial y^2} + f(x)g(y).$$

Exact solution:

$$w(x, y) = C_1 \int_a^x (x-t)f(t) dt - \frac{1}{C_1} \int_b^y (y-\xi)g(\xi) d\xi + C_2 x + C_3 y + C_4,$$

where  $C_1, C_2, C_3, C_4, a$ , and  $b$  are arbitrary constants.

#### B.5.4-2. Other equations with quadratic nonlinearities.

$$10. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = f(x).$$

1°. Suppose  $w(x, y)$  is a solution of the equation. Then the functions

$$w_1 = \pm w(x, \pm y + \varphi(x)) + C,$$

where  $\varphi(x)$  is an arbitrary function and  $C$  is an arbitrary constant, are also solutions of the equation.

2°. Exact solutions:

$$w(x, y) = \pm y \left[ 2 \int f(x) dx + C_1 \right]^{1/2} + \varphi(x),$$

$$w(x, y) = C_1 y^2 + \varphi(x)y + \frac{1}{4C_1} \left[ \varphi^2(x) - 2 \int f(x) dx \right] + C_2,$$

where  $\varphi(x)$  is an arbitrary function and  $C_1$  and  $C_2$  are arbitrary constants.

3°. Exact solutions in implicit form:

$$\int \frac{dw}{\sqrt{2F(x) + \psi(w)}} = \pm y + \varphi(x),$$

where  $\varphi(x)$  and  $\psi(w)$  are arbitrary functions and  $F(x) = \int f(x) dx$ .

$$11. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} + f(y) \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = g(y)w + h(y)x + s(y).$$

Exact solution:

$$w = \varphi(y)x + \psi(y),$$

where the functions  $\varphi(y)$  and  $\psi(y)$  are determined by the system of ordinary differential equations

$$f\varphi\varphi''_{yy} + (\varphi'_y)^2 = g\varphi + h,$$

$$f\varphi\psi''_{yy} + \varphi'_y\psi'_y = g\psi + s.$$

### B.5.5. General Form Equations

|                                                                                                                                                           |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|
| B.5.5-1. Equations of the form $\frac{\partial w}{\partial t} = F\left(x, t, w, \frac{\partial w}{\partial x}, \frac{\partial^2 w}{\partial x^2}\right).$ |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|

$$1. \quad \frac{\partial w}{\partial t} = F\left(x, \frac{\partial^2 w}{\partial x^2}\right).$$

Exact solution:

$$w(x, t) = (Ax + B)t + \varphi(x),$$

where  $A$  and  $B$  are arbitrary constants and the function  $\varphi(x)$  is determined by the ordinary differential equation

$$F(x, \varphi''_{xx}) = Ax + B.$$

$$2. \quad \frac{\partial w}{\partial t} = F\left(\frac{\partial w}{\partial x}, \frac{\partial^2 w}{\partial x^2}\right).$$

Exact solution:

$$w(x, t) = At + B + \varphi(kx + \lambda t),$$

where  $A$ ,  $B$ ,  $k$ , and  $\lambda$  are arbitrary constants and the function  $\varphi(z)$  is determined by the ordinary differential equation

$$F(k\varphi'_z, k^2\varphi''_{zz}) - \lambda\varphi'_z - A = 0, \quad z = kx + \lambda t.$$

$$3. \quad \frac{\partial w}{\partial t} = wF\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right).$$

Exact solution in multiplicative form:

$$w(x, t) = (Ae^{\lambda x} + Be^{-\lambda x})E_1(t), \quad E_1(t) = \exp\left[\int F(t, \lambda^2) dt\right],$$

$$w(x, t) = [A \cos(\lambda x) + B \sin(\lambda x)]E_2(t), \quad E_2(t) = \exp\left[\int F(t, -\lambda^2) dt\right],$$

where  $A$ ,  $B$ , and  $\lambda$  are arbitrary constants.

$$4. \quad \frac{\partial w}{\partial t} = wF\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + f(t)e^{\lambda x} + g(t)e^{-\lambda x}.$$

Exact solution:

$$w(x, t) = e^{\lambda x} E(t) \left[ A + \int \frac{f(t)}{E(t)} dt \right] + e^{-\lambda x} E(t) \left[ B + \int \frac{g(t)}{E(t)} dt \right],$$

$$E(t) = \exp\left[\int F(t, \lambda^2) dt\right],$$

where  $A$ ,  $B$ , and  $\lambda$  are arbitrary constants.

$$5. \quad \frac{\partial w}{\partial t} = w F_1\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + e^{\lambda x} F_2\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + e^{-\lambda x} F_3\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right).$$

There are solutions of the form

$$w(x, t) = e^{\lambda x} \varphi(t) + e^{-\lambda x} \psi(t).$$

$$6. \quad \frac{\partial w}{\partial t} = w F\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + f(t) \cos(\lambda x) + g(t) \sin(\lambda x).$$

Exact solution:

$$w(x, t) = \cos(\lambda x) E(t) \left[ A + \int \frac{f(t)}{E(t)} dt \right] + \sin(\lambda x) E(t) \left[ B + \int \frac{g(t)}{E(t)} dt \right],$$

$$E(t) = \exp \left[ \int F(t, -\lambda^2) dt \right],$$

where  $A$ ,  $B$ , and  $\lambda$  are arbitrary constants.

$$7. \quad \frac{\partial w}{\partial t} = w F_1\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + \cos(\lambda x) F_2\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + \sin(\lambda x) F_3\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right).$$

There are solutions of the form

$$w(x, t) = \cos(\lambda x) \varphi(t) + \sin(\lambda x) \psi(t).$$

$$8. \quad \frac{\partial w}{\partial t} = w F\left(t, \frac{f(x)}{w} \frac{\partial^2 w}{\partial x^2}\right).$$

Exact solution in multiplicative form:

$$w(x, t) = \varphi(x) \exp \left[ \int F(t, \lambda) dt \right],$$

where the function  $\varphi = \varphi(x)$  satisfies the linear ordinary differential equation  $f(x) \varphi''_{xx} = \lambda \varphi$ .

$$9. \quad \frac{\partial w}{\partial t} = w F\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}, w \frac{\partial^2 w}{\partial x^2} - \left(\frac{\partial w}{\partial x}\right)^2\right).$$

1°. Exact solution in multiplicative form:

$$w(x, t) = C \exp \left[ \lambda x + \int F(t, \lambda^2, 0) dt \right],$$

where  $C$  is an arbitrary constant.

2°. Exact solution in multiplicative form:

$$w(x, t) = (A e^{\lambda x} + B e^{-\lambda x}) \varphi(t),$$

where  $A$  and  $B$  are arbitrary constants, and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation  $\varphi'_t = \varphi F(t, \lambda^2, 4AB\lambda^2\varphi^2)$ .

3°. Exact solution in multiplicative form:

$$w(x, t) = [A \sin(\lambda x) + B \cos(\lambda x)] \varphi(t),$$

where  $A$  and  $B$  are arbitrary constants, and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation  $\varphi'_t = \varphi F(t, -\lambda^2, -\lambda^2(A^2 + B^2)\varphi^2)$ .

⊙ *Reference:* Ph. W. Doyle (1996), the case  $\partial_t F \equiv 0$  was considered.

$$10. \quad \frac{\partial w}{\partial t} = w F\left(t, \frac{\partial^2 w}{\partial x^2}, \frac{\partial w}{\partial x} - x \frac{\partial^2 w}{\partial x^2}, 2w - 2x \frac{\partial w}{\partial x} + x^2 \frac{\partial^2 w}{\partial x^2}\right).$$

Exact solution in multiplicative form:

$$w(x, t) = (C_2 x^2 + C_1 x + C_0) \varphi(t),$$

where  $C_0$ ,  $C_1$ , and  $C_2$  are arbitrary constants, and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation  $\varphi'_t = \varphi F(t, 2C_2\varphi, C_1\varphi, 2C_0\varphi)$ .

⊙ *Reference:* Ph. W. Doyle (1996), the case  $\partial_t F \equiv 0$  was considered.

**B.5.5-2.** Equations of the form  $\frac{\partial^2 w}{\partial t^2} = F\left(x, t, w, \frac{\partial w}{\partial t}, \frac{\partial w}{\partial x}, \frac{\partial^2 w}{\partial x^2}\right)$ .

$$11. \quad \frac{\partial^2 w}{\partial t^2} = F\left(x, \frac{\partial w}{\partial x}, \frac{\partial^2 w}{\partial x^2}\right) + G\left(t, \frac{\partial w}{\partial t}\right) + bw.$$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + \psi(t),$$

where the functions  $\varphi(x)$  and  $\psi(t)$  are determined by solving the second-order nonlinear ordinary differential equations ( $C$  is an arbitrary constant)

$$F(x, \varphi'_x, \varphi''_{xx}) + b\varphi = C,$$

$$\psi''_{tt} - G(t, \psi'_t) - b\psi = C.$$

$$12. \quad \frac{\partial^2 w}{\partial t^2} = wF\left(t, \frac{1}{w} \frac{\partial w}{\partial t}, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}, w \frac{\partial^2 w}{\partial x^2} - \left(\frac{\partial w}{\partial x}\right)^2\right).$$

1°. Exact solution in multiplicative form:

$$w(x, t) = (Ae^{\lambda x} + Be^{-\lambda x})\varphi(t),$$

where  $A$  and  $B$  are arbitrary constants, and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation  $\varphi''_{tt} = \varphi F(t, \varphi'_t/\varphi, \lambda^2, 4AB\lambda^2\varphi^2)$ .

2°. Exact solution in multiplicative form:

$$w(x, t) = [A \sin(\lambda x) + B \cos(\lambda x)]\varphi(t),$$

where  $A$  and  $B$  are arbitrary constants, and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation  $\varphi''_{tt} = \varphi F(t, \varphi'_t/\varphi, -\lambda^2, -\lambda^2(A^2 + B^2)\varphi^2)$ .

$$13. \quad \frac{\partial^2 w}{\partial t^2} = wF\left(t, \frac{\partial^2 w}{\partial x^2}, \frac{\partial w}{\partial x} - x \frac{\partial^2 w}{\partial x^2}, 2w - 2x \frac{\partial w}{\partial x} + x^2 \frac{\partial^2 w}{\partial x^2}\right).$$

Exact solution in multiplicative form:

$$w(x, t) = (C_2 x^2 + C_1 x + C_0)\varphi(t),$$

where  $C_0$ ,  $C_1$ , and  $C_2$  are arbitrary constants, and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation  $\varphi''_{tt} = \varphi F(t, 2C_2\varphi, C_1\varphi, 2C_0\varphi)$ .

$$14. \quad \frac{\partial^2 w}{\partial t^2} = wF_1\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + e^{\lambda x} F_2\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + e^{-\lambda x} F_3\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right).$$

There are solutions of the form

$$w(x, t) = e^{\lambda x} \varphi(t) + e^{-\lambda x} \psi(t).$$

$$15. \quad \frac{\partial^2 w}{\partial t^2} = wF_1\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + \cos(\lambda x) F_2\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right) + \sin(\lambda x) F_3\left(t, \frac{1}{w} \frac{\partial^2 w}{\partial x^2}\right).$$

There are solutions of the form

$$w(x, t) = \cos(\lambda x)\varphi(t) + \sin(\lambda x)\psi(t).$$



## B.6. Third-Order Nonlinear Equations

### B.6.1. Stationary Hydrodynamic Boundary Layer Equations

$$1. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = \nu \frac{\partial^3 w}{\partial y^3}.$$

The system of equations of stationary laminar boundary layer on a flat plate (Schlichting 1981, Loitsyanskiy 1996),

$$\begin{aligned} u_1 \frac{\partial u_1}{\partial x} + u_2 \frac{\partial u_1}{\partial y} &= \nu \frac{\partial^2 u_1}{\partial y^2}, \\ \frac{\partial u_1}{\partial x} + \frac{\partial u_2}{\partial y} &= 0, \end{aligned}$$

can be reduced to this equation by introducing the stream function  $w$  in accordance with the relations  $u_1 = \frac{\partial w}{\partial y}$  and  $u_2 = -\frac{\partial w}{\partial x}$  ( $x$  and  $y$  are the longitudinal and transverse coordinates,  $u_1$  and  $u_2$  are the longitudinal and transverse components of the fluid velocity, and  $\nu$  is the kinematic fluid viscosity).

1°. Suppose  $w = w(x, y)$  is a solution of the stationary hydrodynamic boundary layer equation. Then the function

$$w_1 = C_1 w (C_2 x + C_3, C_1 C_2 y + \varphi(x)) + C_4,$$

where  $\varphi(x)$  is an arbitrary function and  $C_1, C_2, C_3$ , and  $C_4$  are arbitrary constants, is also a solution of the equation.

⊙ *Reference:* Yu. N. Pavlovskii (1961), L. V. Ovsiyannikov (1978).

2°. Exact solutions involving arbitrary functions:

$$\begin{aligned} w(x, y) &= C_1 y + \varphi(x), \\ w(x, y) &= C_1 y^2 + \varphi(x)y + \frac{1}{4C_1} \varphi^2(x) + C_2, \\ w(x, y) &= \frac{6\nu x + C_1}{y + \varphi(x)} + \frac{C_2}{[y + \varphi(x)]^2} + C_3, \\ w(x, y) &= \varphi(x) \exp(-C_1 y) + \nu C_1 x + C_2, \\ w(x, y) &= C_1 \exp[-C_2 y - C_2 \varphi(x)] + C_3 y + C_3 \varphi(x) + \nu C_2 x + C_4, \\ w(x, y) &= 6\nu C_1 x^{1/3} \tanh \xi + C_2, \quad \xi = C_1 y x^{-2/3} + \varphi(x), \\ w(x, y) &= -6\nu C_1 x^{1/3} \tan \xi + C_2, \quad \xi = C_1 y x^{-2/3} + \varphi(x), \end{aligned}$$

where  $C_1, C_2, C_3$ , and  $C_4$  are arbitrary constants and  $\varphi(x)$  is an arbitrary function. The second solution is specified in Zwillinger (1998) and the fourth and fifth were obtained by Ignatovich (1993).

3°. Exact solution:

$$w(x, y) = x f(y) + g(y), \quad (1)$$

where the functions  $f = f(y)$  and  $g = g(y)$  are determined by the system of ordinary differential equations

$$(f'_y)^2 - f f''_{yy} = \nu f'''_{yyy}, \quad (2)$$

$$f'_y g'_y - f g''_{yy} = \nu g'''_{yyy}. \quad (3)$$

The order of equation (2) can be reduced by two. Assume that a solution  $f = f(y)$  of equation (2) is known. Then equation (3), which is linear in  $g$ , has two linearly independent particular solutions

$$g_1 = 1, \quad g_2 = f(y)$$

The second particular solution is apparent from comparing equations (2) and (3). The general solution of equation (2) can be represented in the form (see Zaitsev and Polyanin 1995):

$$\begin{aligned} g(y) &= C_1 + C_2 f + C_3 \left( f \int \psi dy - \int f \psi dy \right), \\ f &= f(y), \quad \psi = \frac{1}{(f'_y)^2} \exp \left( -\frac{1}{\nu} \int f dy \right). \end{aligned} \quad (4)$$

It is not difficult to check that equation (2) has the particular solutions

$$\begin{aligned} f(y) &= 6\nu(y + C)^{-1}, \\ f(y) &= C e^{\lambda y} - \lambda \nu, \end{aligned} \quad (5)$$

where  $C$  and  $\lambda$  are arbitrary constants. With reference to (1) and (4), one can see that the first solution in (5) leads to the third solution in Item 2° with  $\varphi(x) = \text{const}$ . Substituting the second expression in (5) into (1) and (4) yields another solution.

⊙ *Reference:* A. D. Polyanin (2001b, 2001c).

$$2. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = \nu \frac{\partial^3 w}{\partial y^3} + f(x).$$

The equation of laminar boundary layer with pressure gradient.

1°. Suppose  $w = w(x, y)$  is a solution of the equation in question. Then the functions (Pavlovskii 1961)

$$w_1 = \pm w(x, \pm y + \varphi(x)) + C,$$

where  $\varphi(x)$  is an arbitrary function and  $C$  is an arbitrary constant, are also solutions of the equation.

2°. Nonviscous solutions (independent of the viscosity  $\nu$ ):

$$\begin{aligned} w(x, y) &= \pm y \left[ 2 \int f(x) dx + C_1 \right]^{1/2} + \varphi(x), \\ w(x, y) &= C_1 y^2 + \varphi(x) y + \frac{1}{4C_1} \left[ \varphi^2(x) - 2 \int f(x) dx \right] + C_2, \end{aligned}$$

where  $\varphi(x)$  is an arbitrary function and  $C_1$  and  $C_2$  are arbitrary constants.

3°. Exact solution for  $f(x) = ax + b$ :

$$w(x, y) = xF(y) + G(y),$$

where the functions  $F = F(y)$  and  $G = G(y)$  are determined by the system of ordinary differential equations

$$(F'_y)^2 - FF''_{yy} = \nu F'''_{yyy} + a, \quad (1)$$

$$F'_y G'_y - FG''_{yy} = \nu G'''_{yyy} + b. \quad (2)$$

The order of the autonomous equation (1) can be reduced by one. If a particular solution  $F(y)$  of equation (1) is known, the corresponding equation (2) can be reduced to a second-order linear equation by the change of variable  $H(y) = G'_y$ . If  $F(y) = \pm \sqrt{a}y + C$ , equation (2) can be integrated in quadrature, because its two particular solutions are known if  $b = 0$ , namely,  $G_1 = 1$  and  $G_2 = \pm \frac{1}{2} \sqrt{a} y^2 + Cy$ .

4°. Exact solution for  $f(x) = ae^{\beta x}$ :

$$w(x, y) = \varphi(x) e^{\lambda y} - \frac{a}{2\beta \lambda^2 \varphi(x)} e^{\beta x - \lambda y} - \nu \lambda x + \frac{2\nu \lambda^2}{\beta} y + \frac{2\nu \lambda}{\beta} \ln |\varphi(x)|,$$

where  $\varphi(x)$  is an arbitrary function and  $\lambda$  is an arbitrary constant.

⊙ *Reference:* A. D. Polyanin (2001b, 2001c).

$$3. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = \frac{\partial}{\partial y} \left[ f(y) \frac{\partial^2 w}{\partial y^2} \right] + g(y)x + h(y).$$

This equation can be used to model turbulent boundary layer.

Exact solution:

$$w = \varphi(y)x + \psi(y),$$

where the functions  $\varphi(y)$  and  $\psi(y)$  are determined by the system of ordinary differential equations

$$\begin{aligned} (f\varphi''_{yy})'_y + \varphi\varphi''_{yy} - (\varphi'_y)^2 + g &= 0, \\ (f\psi''_{yy})'_y + \varphi\psi''_{yy} - \varphi'_y\psi'_y + h &= 0. \end{aligned}$$

## B.6.2. Nonstationary Hydrodynamic Boundary Layer Equations

$$1. \quad \frac{\partial^2 w}{\partial t \partial y} + \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = \nu \frac{\partial^3 w}{\partial y^3}.$$

This is the equation of nonstationary laminar boundary layer on a flat plate;  $x$  and  $y$  are the longitudinal and transverse coordinates and  $w$  is the stream function (Schlichting 1981, Loitsyanskiy 1996).

1°. Suppose  $w = w(x, y, t)$  is a solution of the equation is question. Then the function (see Vereshchagina 1973)

$$w_1 = w(x, y + \varphi(x, t), t) + \frac{\partial}{\partial t} \int \varphi(x, t) dx + \psi(t),$$

where  $\varphi(x, t)$  and  $\psi(t)$  are arbitrary functions, is also a solution of the equation.

2°. Exact solutions:

$$w = C_1 y + \varphi(x, t),$$

$$w = C_1 y^2 + \varphi(x, t)y + \frac{1}{4C_1} \varphi^2(x, t) + \frac{\partial}{\partial t} \int \varphi(x, t) dx,$$

$$w = \frac{6\nu x + C_1}{y + \varphi(x, t)} + \frac{C_2}{[y + \varphi(x, t)]^2} + \frac{\partial}{\partial t} \int \varphi(x, t) dx,$$

$$w = C_1 \exp[-C_2 y - C_2 \varphi(x, t)] + C_3 y + C_3 \varphi(x, t) + \nu C_2 x + \frac{\partial}{\partial t} \int \varphi(x, t) dx,$$

$$w = 6\nu C_1 x^{1/3} \tanh \xi + \frac{\partial}{\partial t} \int \varphi(x, t) dx, \quad \xi = C_1 \frac{y + \varphi(x, t)}{x^{2/3}},$$

$$w = -6\nu C_1 x^{1/3} \tan \xi + \frac{\partial}{\partial t} \int \varphi(x, t) dx, \quad \xi = C_1 \frac{y + \varphi(x, t)}{x^{2/3}},$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments and  $C_1, C_2, C_3$ , and  $C_4$  are arbitrary constants.

3°. Exact solution:

$$w(x, y, t) = xF(y, t) + G(y, t), \quad (3)$$

where the functions  $F = F(y, t)$  and  $G = G(y, t)$  are determined by the simpler equations with two variables

$$\frac{\partial^2 F}{\partial t \partial y} + \left( \frac{\partial F}{\partial y} \right)^2 - F \frac{\partial^2 F}{\partial y^2} = \nu \frac{\partial^3 F}{\partial y^3}, \quad (4)$$

$$\frac{\partial^2 G}{\partial t \partial y} + \frac{\partial F}{\partial y} \frac{\partial G}{\partial y} - F \frac{\partial^2 G}{\partial y^2} = \nu \frac{\partial^3 G}{\partial y^3}. \quad (5)$$

TABLE B4  
Exact solutions of equation (4)

| No | Function $F = F(y, t)$<br>(or general form of solution)                                                                        | Remarks (or determining equation)                                                 |
|----|--------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------|
| 1  | $F = \psi(t)$                                                                                                                  | $\psi(t)$ is an arbitrary function                                                |
| 2  | $F = \frac{y}{t + C_1} + \psi(t)$                                                                                              | $\psi(t)$ is an arbitrary function,<br>$C_1$ is any number                        |
| 3  | $F = \frac{6\nu}{y + \psi(t)} + \psi'_t(t)$                                                                                    | $\psi(t)$ is an arbitrary function                                                |
| 4  | $F = C_1 \exp[-\lambda y + \lambda \psi(t)] - \psi'_t(t) + \nu \lambda$                                                        | $\psi(t)$ is an arbitrary function,<br>$C_1, \lambda$ are any numbers             |
| 5  | $F = \frac{C_1 \exp[-\lambda y + \lambda \psi(t)] + 1}{\lambda t + C_2} - \psi'_t(t) + \nu \lambda$                            | $\psi(t)$ is an arbitrary function,<br>$C_1, C_2, \lambda$ are any numbers        |
| 6  | $F = \frac{\beta + C_1 \exp[-\lambda y + \lambda \psi(t)]}{1 + C_2 \exp(-\lambda \beta t)} - \psi'_t(t) + \nu \lambda - \beta$ | $\psi(t)$ is an arbitrary function,<br>$C_1, C_2, \beta, \lambda$ are any numbers |
| 7  | $F = F(\xi), \quad \xi = y + \lambda t$                                                                                        | $\lambda F''_{\xi\xi} + (F'_\xi)^2 - F F''_{\xi\xi} = \nu F'''_{\xi\xi\xi}$       |
| 8  | $F = t^{-1/2} [H(\xi) - \frac{1}{2}\xi], \quad \xi = y t^{-1/2}$                                                               | $\frac{3}{4} - 2H'_\xi + (H'_\xi)^2 - H H''_{\xi\xi} = \nu H'''_{\xi\xi\xi}$      |

Equation (4) is independent of (5). If a particular solution  $F = F(y, t)$  of equation (4) is known, then the corresponding equation (5) can be reduced by the change of variable  $U = \frac{\partial G}{\partial y}$  to the second-order linear equation

$$\frac{\partial U}{\partial t} - F \frac{\partial U}{\partial y} = \nu \frac{\partial^2 U}{\partial y^2} - \frac{\partial F}{\partial y} U. \quad (6)$$

Exact solutions of equation (4) are listed in [Table B4](#). The ordinary differential equations in the last two rows are autonomous and, therefore, admit reduction of order.

[Table B5](#) presents solutions of equation (6) that correspond to the solutions of equation (4) specified in [Table B4](#). One can see that in the first three cases the solutions of equation (6) are expressed in terms of solutions to the classical heat equation with constant coefficients. There are other three cases where equation (6) is reduced to a separable equation.

4°. Exact solution:

$$\begin{aligned}
 w(x, y, t) &= [A(t)e^{k_1 x} + B(t)e^{k_2 x}] e^{\lambda y + \varphi(t)x + ay}, \\
 A(t) &= C_1 \exp \left[ (\nu \lambda^2 - a k_1) t + \lambda \int \varphi(t) dt \right], \\
 B(t) &= C_2 \exp \left[ (\nu \lambda^2 - a k_2) t + \lambda \int \varphi(t) dt \right],
 \end{aligned}$$

where  $\varphi(t)$  is an arbitrary function and  $C_1, C_2, a, k_1, k_2$ , and  $\lambda$  are arbitrary parameters.

TABLE B5

Transformations of equation (6) for the corresponding exact solutions of equation (4)  
[the number in the first column corresponds to the  
number of the exact solution  $F = F(y, t)$  in Table B4]

| No | Transformations of equation (6)                                                                                             | Resulting equation                                                                                                                                           |
|----|-----------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | $U = u(\zeta, t), \quad \zeta = y + \int \psi(t) dt$                                                                        | $\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial \zeta^2}$                                                                                  |
| 2  | $U = \frac{1}{t+C_1} u(z, \tau), \quad \tau = \frac{1}{3}(t+C_1)^3 + C_2,$<br>$z = (t+C_1)y + \int \psi(t)(t+C_1) dt + C_3$ | $\frac{\partial u}{\partial \tau} = \nu \frac{\partial^2 u}{\partial z^2}$                                                                                   |
| 3  | $U = \zeta^{-3} u(\zeta, t), \quad \zeta = y + \psi(t)$                                                                     | $\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial \zeta^2}$                                                                                  |
| 4  | $U = e^\eta Z(\eta, t), \quad \eta = -\lambda y + \lambda \psi(t)$                                                          | $\frac{\partial Z}{\partial t} = \nu \lambda^2 \frac{\partial^2 Z}{\partial \eta^2} + (\nu \lambda^2 - C_1 \lambda e^\eta) \frac{\partial Z}{\partial \eta}$ |
| 7  | $U = u(\xi, t), \quad \xi = y + \lambda t$                                                                                  | $\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial \xi^2} + [F(\xi) - \lambda] \frac{\partial u}{\partial \xi} - F'_\xi(\xi) u$               |
| 8  | $U = t^{-1/2} u(\xi, \tau), \quad \xi = y t^{-1/2}, \quad \tau = \ln t$                                                     | $\frac{\partial u}{\partial \tau} = \nu \frac{\partial^2 u}{\partial \xi^2} + H(\xi) \frac{\partial u}{\partial \xi} + [1 - H'_\xi(\xi)] u$                  |

5°. Exact solution:

$$w(x, y, t) = A(t) \exp(kx + \lambda y) + B(t) \exp(\beta kx + \beta \lambda y) + \varphi(t)x + ay,$$

$$A(t) = C_1 \exp \left[ (\nu \lambda^2 - ak)t + \lambda \int \varphi(t) dt \right],$$

$$B(t) = C_2 \exp \left[ (\nu \beta^2 \lambda^2 - ak\beta)t + \beta \lambda \int \varphi(t) dt \right],$$

where  $\varphi(t)$  is an arbitrary function and  $C_1, C_2, a, k, \beta$ , and  $\lambda$  are arbitrary parameters.

6°. Exact solution:

$$w(x, y, t) = \int u(z, t) dz + \varphi(t)y + \psi(t)x, \quad z = kx + \lambda y,$$

where  $\varphi(t)$  and  $\psi(t)$  are arbitrary functions,  $k$  and  $\lambda$  are arbitrary parameters, and  $u(z, t)$  is a function satisfying the second-order linear parabolic equation

$$\frac{\partial u}{\partial t} + [k\varphi(t) - \lambda\psi(t)] \frac{\partial u}{\partial z} = \nu \lambda^2 \frac{\partial^2 u}{\partial z^2} - \frac{1}{\lambda} \varphi'_t(t).$$

The transformation

$$u = U(\xi, t) - \frac{1}{\lambda} \varphi(t), \quad \xi = z - \int [k\varphi(t) - \lambda\psi(t)] dt$$

takes the last equation to the customary heat equation

$$\frac{\partial U}{\partial t} = \nu \lambda^2 \frac{\partial^2 U}{\partial \xi^2}.$$

7°. Exact solutions:

$$w = e^{\nu \lambda^2 t} (C_1 e^{\lambda z} + C_2 e^{-\lambda z}) + \frac{\partial}{\partial t} \int \varphi(x, t) dx, \quad z = y + \varphi(x, t),$$

$$w = e^{-\nu \lambda^2 t} [C_1 \sin(\lambda z) + C_2 \cos(\lambda z)] + \frac{\partial}{\partial t} \int \varphi(x, t) dx, \quad z = y + \varphi(x, t),$$

$$w = C_1 e^{-\nu \lambda^2 z} \sin(\lambda z - 2\nu \lambda^2 t + C_2) + \frac{\partial}{\partial t} \int \varphi(x, t) dx, \quad z = y + \varphi(x, t),$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments and  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants.

⊙ *Reference:* A. D. Polyinin (2001b).

$$2. \quad \frac{\partial^2 w}{\partial t \partial y} + \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = \nu \frac{\partial^3 w}{\partial y^3} + f(x, t).$$

The equation of nonstationary laminar boundary layer with pressure gradient.

1°. Suppose  $w(x, y, t)$  is a solution of the equation in question. Then the functions (see Vereshchagina 1973)

$$w_1 = \pm w(x, \pm y + \varphi(x, t), t) + \frac{\partial}{\partial t} \int \varphi(x, t) dx + \psi(t),$$

where  $\varphi(x, t)$  and  $\psi(t)$  are arbitrary functions, are also solutions of the equation.

2°. Nonviscous solution for any  $f(x, t)$  (independent of the viscosity  $\nu$ ):

$$w(x, y, t) = ay^2 + \varphi(x, t)y + \frac{1}{4a}\varphi^2(x, t) + \frac{1}{2a} \int \left[ \frac{\partial \varphi}{\partial t} - f(x, t) \right] dx + \psi(t).$$

where  $\varphi(x, t)$  and  $\psi(t)$  are arbitrary functions and  $a$  is an arbitrary constant.

Another nonviscous solution for any  $f(x, t)$ :

$$w(x, y, t) = \psi(x, t)y + \varphi(x, t),$$

where  $\varphi(x, t)$  is an arbitrary function, and the function  $\psi = \psi(x, t)$  is determined by the first-order equation

$$\frac{\partial \psi}{\partial t} + \psi \frac{\partial \psi}{\partial x} = f(x, t).$$

Nonviscous solutions for  $f(x, t) = f(x)$ :

$$w(x, y, t) = \pm y \left[ 2 \int f(x) dx + C_1 \right]^{1/2} + \varphi(x, t).$$

3°. Exact solutions for  $f(x, t) = f_1(t)x + f_2(t)$ :

$$w(x, y, t) = xF(y, t) + G(y, t),$$

where the functions  $F = F(y, t)$  and  $G = G(y, t)$  are determined from the simpler equations with two variables

$$\frac{\partial^2 F}{\partial t \partial y} + \left( \frac{\partial F}{\partial y} \right)^2 - F \frac{\partial^2 F}{\partial y^2} = \nu \frac{\partial^3 F}{\partial y^3} + f_1(t), \quad (1)$$

$$\frac{\partial^2 G}{\partial t \partial y} + \frac{\partial F}{\partial y} \frac{\partial G}{\partial y} - F \frac{\partial^2 G}{\partial y^2} = \nu \frac{\partial^3 G}{\partial y^3} + f_2(t). \quad (2)$$

Equation (1) is independent of (2). If  $F = F(y, t)$  is a solution of equation (1), then the function  $F_1 = F(y + \psi(t), t) + \psi'_t(t)$  with arbitrary  $\psi(t)$  is also a solution of equation (1). [Table B6](#) presents exact solutions of equation (1) for various  $f_1 = f_1(t)$ .

The change of variable  $U = \frac{\partial G}{\partial y}$  brings equation (2) to the second-order linear equation

$$\frac{\partial U}{\partial t} - F \frac{\partial U}{\partial y} = \nu \frac{\partial^2 U}{\partial y^2} - \frac{\partial F}{\partial y} U + f_2(t). \quad (3)$$

Let us dwell on the first solution of equation (1) in [Table B6](#):

$$F(y, t) = a(t)y + \psi(t), \quad \text{where} \quad a'_t + a^2 = f_1(t). \quad (4)$$

TABLE B6

Exact solutions of equation (1) for various  $f_1(t)$ ;  $\psi(t)$  is an arbitrary function

| Function<br>$f_1 = f_1(t)$                           | Function $F = F(y, t)$<br>(or general form of solution)                                                                                                          | Determining equation<br>(or determining coefficients)                           |
|------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------|
| Any                                                  | $F = a(t)y + \psi(t)$                                                                                                                                            | $a_t' + a^2 = f_1(t)$                                                           |
| $f_1(t) = Ae^{-\beta t}$ ,<br>$A > 0, \beta > 0$     | $F = Be^{-\frac{1}{2}\beta t} \sin[\lambda y + \lambda \psi(t)] + \psi_t'(t)$ ,<br>$F = Be^{-\frac{1}{2}\beta t} \cos[\lambda y + \lambda \psi(t)] + \psi_t'(t)$ | $B = \pm \sqrt{\frac{2A\nu}{\beta}}, \lambda = \sqrt{\frac{\beta}{2\nu}}$       |
| $f_1(t) = Ae^{\beta t}$ ,<br>$A > 0, \beta > 0$      | $F = Be^{\frac{1}{2}\beta t} \sinh[\lambda y + \lambda \psi(t)] + \psi_t'(t)$                                                                                    | $B = \pm \sqrt{\frac{2A\nu}{\beta}}, \lambda = \sqrt{\frac{\beta}{2\nu}}$       |
| $f_1(t) = Ae^{\beta t}$ ,<br>$A < 0, \beta > 0$      | $F = Be^{\frac{1}{2}\beta t} \cosh[\lambda y + \lambda \psi(t)] + \psi_t'(t)$                                                                                    | $B = \pm \sqrt{\frac{2 A \nu}{\beta}}, \lambda = \sqrt{\frac{\beta}{2\nu}}$     |
| $f_1(t) = Ae^{\beta t}$ ,<br>$A$ is any, $\beta > 0$ | $F = \psi(t)e^{\lambda y} - \frac{Ae^{\beta t - \lambda y}}{4\lambda^2\psi(t)} + \frac{\psi_t'(t)}{\lambda\psi(t)} - \nu\lambda$                                 | $\lambda = \pm \sqrt{\frac{\beta}{2\nu}}$                                       |
| $f_1(t) = At^{-2}$                                   | $F = t^{-1/2} [H(\xi) - \frac{1}{2}\xi], \xi = yt^{-1/2}$                                                                                                        | $\frac{3}{4} - A - 2H_\xi' + (H_\xi')^2 - HH_{\xi\xi}'' = \nu H_{\xi\xi\xi}'''$ |
| $f_1(t) = A$                                         | $F = F(\xi), \xi = y + \lambda t$                                                                                                                                | $-A + \lambda F_{\xi\xi}'' + (F_\xi')^2 - FF_{\xi\xi}'' = \nu F_{\xi\xi\xi}'''$ |

Exact solutions of the Riccati equation for  $a = a(t)$  with various  $f_1(t)$  can be found in Polyanin and Zaitsev (1995). The substitution  $a = h_t'/h$  brings this equation to a second-order linear equation for  $h(t)$ :  $h_{tt}'' - f_1(t)h = 0$ . In particular, if  $f_1(t) = \text{const}$ , we have

$$a(t) = k \frac{C_1 \cos(kt) - C_2 \sin(kt)}{C_1 \sin(kt) + C_2 \cos(kt)} \quad \text{for } f_1 = -k^2 < 0,$$

$$a(t) = k \frac{C_1 \cosh(kt) + C_2 \sinh(kt)}{C_1 \sinh(kt) + C_2 \cosh(kt)} \quad \text{for } f_1 = k^2 > 0.$$

On substituting solution (4) with arbitrary  $f_1(t)$  into equation (3), we obtain

$$\frac{\partial U}{\partial t} = \nu \frac{\partial^2 U}{\partial y^2} + [a(t)y + \psi(t)] \frac{\partial U}{\partial y} - a(t)U + f_2(t). \quad (5)$$

The transformation

$$U = \frac{1}{\Phi(t)} \left[ u(z, \tau) + \int f_2(t) \Phi(t) dt \right], \quad \tau = \int \Phi^2(t) dt + C_1,$$

$$z = y\Phi(t) + \int \psi(t)\Phi(t) dt + C_2, \quad \Phi(t) = \exp \left[ \int a(t) dt \right],$$

takes (5) to the classical constant coefficient heat equation

$$\frac{\partial u}{\partial \tau} = \nu \frac{\partial^2 u}{\partial z^2}.$$

*Remark 1.* The ordinary differential equations in the last two rows in Table B6 (see the last column) are autonomous and, hence, can be reduced in order.

*Remark 2.* Suppose  $w(x, y, t)$  is a solution of the nonstationary hydrodynamic boundary layer equation with  $f(x, t) = f_1(t)x + f_2(t)$ . Then the function

$$w_1 = w(x + h(t), y, t) - h_t'(t)y, \quad \text{where } h_{tt}'' + f_1(t)h = 0,$$

is also a solution of this equation.

4°. Exact solution for  $f(x, t) = g(x)e^{\beta t}$ ,  $\beta > 0$ :

$$w(x, y, t) = \varphi(x, t)e^{\lambda y} + \psi(x, t)e^{-\lambda y} + \frac{1}{\lambda} \frac{\partial}{\partial t} \int \ln |\varphi(x, t)| dx - \nu \lambda x,$$

$$\psi(x, t) = -\frac{e^{\beta t}}{2\lambda^2 \varphi(x, t)} \int g(x) dx, \quad \lambda = \pm \sqrt{\frac{\beta}{2\nu}},$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments.

5°. Exact solutions for  $f(x, t) = g(x)e^{\beta t}$ ,  $\beta > 0$ :

$$w(x, y, t) = \pm \frac{1}{\lambda} \exp\left(\frac{1}{2}\beta t\right) \sqrt{\psi(x)} \sinh[\lambda y + \varphi(x, t)] + \frac{\partial}{\partial t} \int \varphi(x, t) dx,$$

$$w(x, y, t) = \pm \frac{1}{\lambda} \exp\left(\frac{1}{2}\beta t\right) \sqrt{\psi(x)} \cosh[\lambda y + \varphi(x, t)] + \frac{\partial}{\partial t} \int \varphi(x, t) dx,$$

$$\psi(x) = 2 \int g(x) dx + C_1, \quad \lambda = \sqrt{\frac{\beta}{2\nu}},$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments.

6°. Exact solution for  $f(x, t) = g(x)e^{-\beta t}$ ,  $\beta < 0$ :

$$w(x, y, t) = \pm \frac{1}{\lambda} \exp\left(-\frac{1}{2}\beta t\right) \sqrt{\psi(x)} \sin[\lambda y + \varphi(x, t)] + \frac{\partial}{\partial t} \int \varphi(x, t) dx,$$

$$w(x, y, t) = \pm \frac{1}{\lambda} \exp\left(-\frac{1}{2}\beta t\right) \sqrt{\psi(x)} \cos[\lambda y + \varphi(x, t)] + \frac{\partial}{\partial t} \int \varphi(x, t) dx,$$

$$\psi(x) = 2 \int g(x) dx + C_1, \quad \lambda = \sqrt{\frac{\beta}{2\nu}}.$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments.

7°. Exact solution for  $f(x, t) = ae^{\beta x - \gamma t}$ :

$$w(x, y, t) = \varphi(x, t)e^{\lambda y} - \frac{a}{2\beta\lambda^2 \varphi(x, t)} e^{\beta x - \lambda y - \gamma t}$$

$$+ \frac{1}{\lambda} \frac{\partial}{\partial t} \int \ln |\varphi(x, t)| dx - \nu \lambda x + \frac{2\nu\lambda^2 + \gamma}{\beta} \left( y + \frac{1}{\lambda} \ln |\varphi(x, t)| \right),$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments and  $\lambda$  is an arbitrary constant.

8°. Exact solution for  $f(x, t) = f(t)$ :

$$w(x, y, t) = \int u(z, t) dz + \varphi(t)y + \psi(t)x, \quad z = kx + \lambda y,$$

where  $\varphi(t)$  and  $\psi(t)$  are arbitrary functions,  $k$  and  $\lambda$  are arbitrary parameters, and  $u(z, t)$  is a function satisfying the second-order linear parabolic equation

$$\frac{\partial u}{\partial t} + [k\varphi(t) - \lambda\psi(t)] \frac{\partial u}{\partial z} = \nu\lambda^2 \frac{\partial^2 u}{\partial z^2} - \frac{1}{\lambda} \varphi'_t(t) + \frac{1}{\lambda} f(t).$$

The transformation

$$u = U(\xi, t) - \frac{1}{\lambda} \varphi(t) + \frac{1}{\lambda} \int f(t) dt, \quad \xi = z - \int [k\varphi(t) - \lambda\psi(t)] dt$$

brings it to the customary heat equation

$$\frac{\partial U}{\partial t} = \nu\lambda^2 \frac{\partial^2 U}{\partial \xi^2}.$$

9°. Exact solution for  $f(x, t) = f(t)$ :

$$w(x, y, t) = C_1 e^{-\lambda y + \lambda \varphi(x, t)} - a(t) \varphi(x, t) - \frac{\partial}{\partial t} \int \varphi(x, t) dx + a(t)y + \nu \lambda x, \quad a(t) = \int f(t) dt + C_2,$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments and  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants.



10°. Exact solution for  $f(x, t) = f(t)$ :

$$w(x, y, t) = \varphi(x, t)e^{\lambda y} + \psi(x, t)e^{-\lambda y} + \chi(x, t) + a(t)y,$$

where  $\lambda$  is any number,  $\varphi(x, t)$  is an arbitrary function of two arguments, and the other functions are defined by

$$\begin{aligned}\psi(x, t) &= \frac{C\nu e^{2\nu\lambda^2 t}}{\varphi(x, t)} \left[ x - \int a(t) dt \right], \quad a(t) = \int f(t) dt + Ce^{2\nu\lambda^2 t}; \\ \chi(x, t) &= \frac{1}{\lambda} a(t) \ln |\varphi(x, t)| + \frac{1}{\lambda} \frac{\partial}{\partial t} \int \ln |\varphi(x, t)| dx - \nu \lambda x.\end{aligned}$$

11°. Exact solutions for  $f(x, t) = f(t)$ :

$$\begin{aligned}w &= e^{\nu\lambda^2 t} (C_1 e^{\lambda z} + C_2 e^{-\lambda z}) + \frac{\partial}{\partial t} \int \varphi(x, t) dx + z \int f(t) dt, \quad z = y + \varphi(x, t), \\ w &= e^{-\nu\lambda^2 t} [C_1 \sin(\lambda z) + C_2 \cos(\lambda z)] + \frac{\partial}{\partial t} \int \varphi(x, t) dx + z \int f(t) dt, \quad z = y + \varphi(x, t), \\ w &= C_1 e^{-\lambda z} \sin(\lambda z - 2\nu\lambda^2 t + C_2) + \frac{\partial}{\partial t} \int \varphi(x, t) dx + z \int f(t) dt, \quad z = y + \varphi(x, t),\end{aligned}$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments and  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary constants.

12°. Exact solutions for  $f(x, t) = A$ :

$$\begin{aligned}w &= -\frac{A}{6\nu} z^3 + C_2 z^2 + C_1 z + \frac{\partial}{\partial t} \int \varphi(x, t) dx, \quad z = y + \varphi(x, t), \\ w &= kx + C_1 \exp\left(-\frac{k}{\nu} z\right) - \frac{A}{2k} z^2 + C_2 z + \frac{\partial}{\partial t} \int \varphi(x, t) dx, \quad z = y + \varphi(x, t),\end{aligned}$$

where  $\varphi(x, t)$  is an arbitrary function of two arguments and  $C_1$ ,  $C_2$ , and  $k$  are arbitrary constants.

⊙ *Reference:* A. D. Polyanin (2001b).

$$3. \quad \frac{\partial^2 w}{\partial t \partial y} + \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = \frac{\partial}{\partial y} \left[ f \left( \frac{\partial^2 w}{\partial y^2} \right) \right] + g(x, t).$$

This equation describes the flow of a non-Newtonian fluid in a two-dimensional nonstationary boundary layer with a pressure gradient. Here,  $w$  is the stream function and the function  $f = f(u)$  depends on the rheological properties of the fluid. For power-law fluids,  $f = k|u|^{n-1}u$ .

1°. The assertion of Item 1°, equation 1 in Subsection B.6.2, remains valid for this equation.

2°. The equation admits the nonviscous solutions presented in Item 2°, equation 2 in Subsection B.6.2, where  $f(x, t)$  must be replaced by  $g(x, t)$ .

3°. Exact solution for  $g(x, t) = g(t)$ :

$$w(x, y, t) = a(t)x + \int U(y, t) dy,$$

where  $U = U(y, t)$  is a function satisfying the second-order equation

$$\frac{\partial U}{\partial t} - a(t) \frac{\partial U}{\partial y} = \frac{\partial}{\partial y} \left[ f \left( \frac{\partial U}{\partial y} \right) \right] + g(t). \quad (1)$$

The transformation

$$U = u(z, t) + \int g(t) dt, \quad z = y + \int a(t) dt$$

brings equation (1) to the simpler equation

$$\frac{\partial u}{\partial t} = \frac{\partial}{\partial z} \left[ f \left( \frac{\partial u}{\partial z} \right) \right]. \quad (2)$$

Equation (2) admits exact solutions with the forms [for any  $f = f(v)$ ]

$$\begin{aligned} u(z, t) &= H(\zeta), & \zeta &= kz + \lambda t & \implies & \text{equation} & \lambda H = k f(k H'_\zeta) + C; \\ u(z, t) &= az + H(\zeta), & \zeta &= kz + \lambda t & \implies & \text{equation} & \lambda H = k f(k H'_\zeta + a) + C; \\ u(z, t) &= \sqrt{t} H(\zeta), & \zeta &= z/\sqrt{t} & \implies & \text{equation} & \frac{1}{2} H - \frac{1}{2} \zeta H'_\zeta = [f(H'_\zeta)]'_\zeta, \end{aligned}$$

where  $a$ ,  $k$ ,  $C$ , and  $\lambda$  are arbitrary constants. The first two equations for  $H = H(\zeta)$  can be solved in parametric form.

4°. Exact solution for  $g(x, t) = g(t)$ :

$$w(x, y, t) = \int v(\eta, t) d\eta + \varphi(t)y + \psi(t)x, \quad \eta = y + kx,$$

where  $\varphi(t)$  and  $\psi(t)$  are arbitrary functions,  $k$  is an arbitrary parameter, and  $v(\eta, t)$  is a function satisfying the second-order nonlinear parabolic equation

$$\frac{\partial v}{\partial t} + [k\varphi(t) - \psi(t)] \frac{\partial v}{\partial \eta} = \frac{\partial}{\partial \eta} \left[ f \left( \frac{\partial v}{\partial \eta} \right) \right] - \varphi'_t(t) + g(t).$$

The transformation

$$v = R(\zeta, t) - \varphi(t) + \int g(t) dt, \quad \zeta = \eta - \int [k\varphi(t) - \psi(t)] dt$$

leads to the simpler equation

$$\frac{\partial R}{\partial t} = \frac{\partial}{\partial \zeta} \left[ f \left( \frac{\partial R}{\partial \zeta} \right) \right].$$

For exact solutions of this equation, see Item 3°.

5°. Exact solution for  $g(x, t) = s(t)x + h(t)$ :

$$w(x, y, t) = [a(t)y + \psi(t)]x + \int Q(y, t) dy,$$

where  $\psi(t)$  is an arbitrary function,  $a = a(t)$  is determined by the Riccati equation

$$a'_t + a^2 = s(t),$$

and  $Q = Q(y, t)$  satisfies the second-order equation

$$\frac{\partial Q}{\partial t} = \frac{\partial}{\partial y} \left[ f \left( \frac{\partial Q}{\partial y} \right) \right] + [a(t)y + \psi(t)] \frac{\partial Q}{\partial y} - a(t)Q + h(t).$$

The transformation

$$Q = \frac{1}{\Phi(t)} \left[ Z(\xi, \tau) + \int h(t)\Phi(t) dt \right], \quad \tau = \int \Phi^2(t) dt + A, \quad \xi = y\Phi(t) + \int \psi(t)\Phi(t) dt + B,$$

where  $\Phi(t) = \exp \left[ \int a(t) dt \right]$ , leads to the simpler equation

$$\frac{\partial Z}{\partial \tau} = \frac{\partial}{\partial \xi} \left[ f \left( \frac{\partial Z}{\partial \xi} \right) \right].$$

For exact solutions of this equation, see Item 3°.

## B.7. Fourth-Order Nonlinear Equations

### B.7.1. Stationary Hydrodynamic Equations (Navier–Stokes Equations)

$$1. \quad \frac{\partial w}{\partial y} \frac{\partial}{\partial x} (\Delta w) - \frac{\partial w}{\partial x} \frac{\partial}{\partial y} (\Delta w) = \nu \Delta \Delta w, \quad \Delta w = \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2}.$$

The two-dimensional equations of steady-state motion of a viscous incompressible fluid (stationary Navier–Stokes equations)

$$\begin{aligned} u_1 \frac{\partial u_1}{\partial x} + u_2 \frac{\partial u_1}{\partial y} &= -\frac{1}{\rho} \frac{\partial p}{\partial x} + \nu \Delta u_1, \\ u_1 \frac{\partial u_2}{\partial x} + u_2 \frac{\partial u_2}{\partial y} &= -\frac{1}{\rho} \frac{\partial p}{\partial y} + \nu \Delta u_2, \\ \frac{\partial u_1}{\partial x} + \frac{\partial u_2}{\partial y} &= 0 \end{aligned}$$

are reduced to the equation under consideration. To this end, one introduces the stream function  $w$  by the formulas  $u_1 = \frac{\partial w}{\partial y}$  and  $u_2 = -\frac{\partial w}{\partial x}$  and eliminates the pressure  $p$ , using cross differentiation, from the first two equations.

1°. Exact solutions in additive form:

$$\begin{aligned} w(x, y) &= C_1 y^3 + C_2 y^2 + C_3 y + C_4, \\ w(x, y) &= C_1 x^2 + C_2 x + C_3 y^2 + C_4 y + C_5, \\ w(x, y) &= C_1 \exp(-\lambda y) + C_2 y^2 + C_3 y + C_4 + \nu \lambda x, \\ w(x, y) &= C_1 \exp(\lambda x) - \nu \lambda x + C_2 \exp(\lambda y) + \nu \lambda y + C_3, \\ w(x, y) &= C_1 \exp(\lambda x) + \nu \lambda x + C_2 \exp(-\lambda y) + \nu \lambda y + C_3, \end{aligned}$$

where  $C_1, \dots, C_5$ , and  $\lambda$  are arbitrary constants.

2°. Exact solutions:

$$\begin{aligned} w(x, y) &= (Ax + B)e^{-\lambda y} + \nu \lambda x + C, \\ w(x, y) &= Ae^{-\lambda(y+kx)} + B(y+kx)^2 + C(y+kx) + \nu \lambda(k^2 + 1)x + D, \\ w(x, y) &= [A \sinh(\beta x) + B \cosh(\beta x)] e^{-\lambda y} + \frac{\nu}{\lambda}(\beta^2 + \lambda^2)x + C, \\ w(x, y) &= [A \sin(\beta x) + B \cos(\beta x)] e^{-\lambda y} + \frac{\nu}{\lambda}(\lambda^2 - \beta^2)x + C, \\ w(x, y) &= Ae^{\lambda y + \beta x} + Be^{\gamma x} + \nu \gamma y + \frac{\nu}{\lambda} \gamma(\beta - \gamma)x + C, \quad \gamma = \pm \sqrt{\lambda^2 + \beta^2}, \end{aligned}$$

where  $A, B, C, D, k, \beta$ , and  $\lambda$  are arbitrary constants.

3°. Exact solution:

$$w(x, y) = F(y)x + G(y),$$

where the functions  $F = F(y)$  and  $G = G(y)$  are determined by the system of fourth-order ordinary differential equations

$$F'_y F''_{yy} - F F'''_{yyy} = \nu F''''_{yyyy}, \quad (1)$$

$$G'_y F''_{yy} - F G'''_{yyy} = \nu G''''_{yyyy}. \quad (2)$$

Integrating yields the system of third-order equations

$$(F'_y)^2 - F F''_{yy} = \nu F'''_{yyy} + A, \quad (3)$$

$$G'_y F'_y - F G''_{yy} = \nu G'''_{yyy} + B, \quad (4)$$

where  $A$  and  $B$  are arbitrary constants. The order of the autonomous equation (3) can be reduced by one.

It is not difficult to verify that equation (1) has the particular solutions

$$F(y) = ay + b, \quad (5)$$

$$F(y) = 6\nu(y + a)^{-1}, \quad (6)$$

$$F(y) = ae^{-\lambda y} + \lambda\nu, \quad (7)$$

where  $a$ ,  $b$ , and  $\lambda$  are arbitrary constants.

In general, equation (4) can be reduced by the change of variable  $U = G'_y$  to the second-order nonhomogeneous linear equation

$$\nu U''_{yy} + FU'_y - F_y U + B = 0, \quad \text{where } U = G'_y. \quad (8)$$

The corresponding homogeneous equation (with  $B = 0$ ) has two linearly independent particular solutions

$$U_1 = \begin{cases} F''_{yy} & \text{if } F''_{yy} \neq 0, \\ F & \text{if } F''_{yy} = 0, \end{cases} \quad U_2 = U_1 \int \frac{\Phi dy}{U_1^2}, \quad \text{where } \Phi = \exp\left(-\frac{1}{\nu} \int F dy\right). \quad (9)$$

(The first solution is apparent from comparing equations (1) and (8) with  $B = 0$ .) The general solutions of equations (8) and (2) are given by

$$U = C_1 U_1 + C_2 U_2 + C_3 \left( U_2 \int \frac{U_1}{\Phi} dy - U_1 \int \frac{U_2}{\Phi} dy \right), \quad G = \int U dy + C_4, \quad C_3 = -\frac{B}{\nu}. \quad (10)$$

The general solution of equation (2) corresponding to the particular solution (6) is represented as

$$G(y) = \tilde{C}_1(y + a)^3 + \tilde{C}_2 + \tilde{C}_3(y + a)^{-1} + \tilde{C}_4(y + a)^{-2},$$

where  $\tilde{C}_1$ ,  $\tilde{C}_2$ ,  $\tilde{C}_3$ , and  $\tilde{C}_4$  are arbitrary constants (these are expressed in terms of  $C_1$ ,  $C_2$ ,  $C_3$ , and  $C_4$ ).

The general solutions of (2) corresponding to the particular solutions (5) and (7) are determined from (9) and (10).

4°. Exact solution of a more general form:

$$w(x, y) = F(z)x + G(z), \quad z = y + kx,$$

where the functions  $F = F(z)$  and  $G = G(z)$  are determined by the system of fourth-order ordinary differential equations

$$F'_z F''_{zz} - F F'''_{zzz} = \nu(k^2 + 1) F''''_{zzzz}, \quad (11)$$

$$G'_z F''_{zz} - F G'''_{zzz} = \nu(k^2 + 1) G''''_{zzzz} + 4k\nu F'''_{zzz} + \frac{2k}{(k^2 + 1)} F F''_{zz}. \quad (12)$$

Integrating yields the system of third-order equations

$$(F'_z)^2 - F F''_{zz} = \nu(k^2 + 1) F'''_{zzz} + A, \quad (13)$$

$$G'_z F'_z - F G''_{zz} = \nu(k^2 + 1) G'''_{zzz} + 4k\nu F''_{zz} + \frac{2k}{k^2 + 1} \int F F''_{zz} dz + B, \quad (14)$$

where  $A$  and  $B$  are arbitrary constants. The order of the autonomous equation (13) can be reduced by one.

It is not difficult to verify that equation (11) has the particular solutions

$$F(z) = az + b, \quad z = y + kx,$$

$$F(z) = 6\nu(k^2 + 1)(z + a)^{-1},$$

$$F(z) = ae^{-\lambda z} + \lambda\nu(k^2 + 1),$$

where  $a$ ,  $b$ , and  $\lambda$  are arbitrary constants.

In general, equation (14) can be reduced by the change of variable  $U = G'_z$  to a second-order nonhomogeneous linear equation.

© Reference: A. D. Polyanin (2001d).

$$2. \quad \frac{\partial w}{\partial y} \frac{\partial}{\partial x} (\Delta w) - \frac{\partial w}{\partial x} \frac{\partial}{\partial y} (\Delta w) = \nu \Delta \Delta w + f(y), \quad \Delta w = \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2}.$$

This equation describes plane flow of a viscous incompressible fluid under the action of a transverse force ( $w$  is the stream function). The case  $F(y) = a \sin(\lambda y)$  corresponds to A. N. Kolmogorov's model which is used to describe subcritical and transcritical (laminar-turbulent) modes of flow.

1°. Exact solution in additive form for arbitrary  $f(y)$ :

$$w(x, y) = -\frac{1}{2\nu} \int_0^y (y-z)^2 \Phi(z) dz + C_1 e^{-\lambda y} + C_2 y^2 + C_3 y + C_4 + \nu \lambda x, \quad \Phi(z) = e^{-\lambda z} \int e^{\lambda z} f(z) dz,$$

where  $C_1, C_2, C_3, C_4$ , and  $\lambda$  are arbitrary constants.

**Example.** In the case  $f(y) = a\beta \cos(\beta y)$ , which corresponds to  $F(y) = a \sin(\beta y)$ , it follows from the previous formula with  $C_1 = C_2 = C_4 = 0$  and  $B = -\nu\lambda$  that

$$w(x, y) = -\frac{a}{\beta^2(B^2 + \nu^2\beta^2)} [B \sin(\beta y) + \nu\beta \cos(\beta y)] + Cy - Bx,$$

where  $B$  and  $C$  are arbitrary constants. This solution was indicated by Belotserkovskii and Oparin (2000); it describes the flow with a periodic structure.

2°. Exact solution in additive form for  $f(y) = Ae^{\lambda y} + Be^{-\lambda y}$ :

$$w(x, y) = C_1 e^{-\lambda x} + C_2 x - \frac{A}{\lambda^3(C_2 + \nu\lambda)} e^{\lambda y} + \frac{B}{\lambda^3(C_2 - \nu\lambda)} e^{-\lambda y} - \nu \lambda y,$$

where  $C_1$  and  $C_2$  are arbitrary constants.

3°. Generalized separable solution for arbitrary  $f(y)$ :

$$w(x, y) = \varphi(y)x + \psi(y),$$

where the functions  $\varphi = \varphi(y)$  and  $\psi = \psi(y)$  are determined by the system of fourth-order ordinary differential equations

$$\varphi'_y \varphi''_{yy} - \varphi \varphi'''_{yyy} = \nu \varphi''''_{yyyy}, \quad (1)$$

$$\psi'_y \varphi''_{yy} - \varphi \psi'''_{yyy} = \nu \psi''''_{yyyy} + f(y). \quad (2)$$

Integrating yields the system of third-order equations

$$(\varphi'_y)^2 - \varphi \varphi''_{yy} = \nu \varphi'''_{yyy} + A, \quad (3)$$

$$\psi'_y \varphi'_y - \varphi \psi''_{yy} = \nu \psi'''_{yyy} + \int f(y) dy + B, \quad (4)$$

where  $A$  and  $B$  are arbitrary constants. The order of the autonomous equation (3) can be reduced by one.

It is not difficult to verify that equation (1) has the particular solutions

$$\varphi(y) = ay + b, \quad (5)$$

$$\varphi(y) = 6\nu(y+a)^{-1}, \quad (6)$$

$$\varphi(y) = ae^{-\lambda y} + \lambda\nu, \quad (7)$$

where  $a, b$ , and  $\lambda$  are arbitrary constants.

In general, equation (4) can be reduced by the change of variable  $U = \psi'_y$  to the second-order nonhomogeneous linear equation

$$\nu U''_{yy} + \varphi U'_y - \varphi'_y U + F = 0, \quad \text{where } U = \psi'_y, \quad F = \int f(y) dy + B. \quad (8)$$

The corresponding homogeneous equation (with  $F = 0$ ) has two linearly independent particular solutions:

$$U_1 = \begin{cases} \varphi''_{yy} & \text{for } \varphi \neq ay + b, \\ \varphi & \text{for } \varphi = ay + b, \end{cases} \quad U_2 = U_1 \int \frac{\Phi dy}{U_1^2}, \quad \text{where } \Phi = \exp\left(-\frac{1}{\nu} \int \varphi dy\right).$$

(The first solution is apparent from comparing equations (1) and (8) with  $F = 0$ .) The general solutions of equations (8) and (2) are given by

$$U = C_1 U_1 + C_2 U_2 + \frac{1}{\nu} U_1 \int U_2 \frac{F}{\Phi} dy - \frac{1}{\nu} U_2 \int U_1 \frac{F}{\Phi} dy, \quad \psi = \int U dy + C_4.$$

$$3. \quad \frac{1}{r} \frac{\partial w}{\partial \theta} \frac{\partial}{\partial r} (\Delta w) - \frac{1}{r} \frac{\partial w}{\partial r} \frac{\partial}{\partial \theta} (\Delta w) = \nu \Delta \Delta w, \quad \Delta w = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2}.$$

The equations of steady-state flow of a viscous incompressible fluid (stationary Navier–Stokes equations) written in polar coordinates ( $x = r \cos \theta$ ,  $y = r \sin \theta$ ) reduce to this equation. The radial and tangential components of the fluid velocity are expressed in terms of the stream function  $w$  by the formulas  $u_r = \frac{1}{r} \frac{\partial w}{\partial \theta}$  and  $u_\theta = -\frac{\partial w}{\partial r}$ .

1°. Exact solution in additive form:

$$w(r, \theta) = \nu C_1 \theta + C_2 r^{C_1+2} + C_3 r^2 + C_4 \ln r + C_5,$$

where  $C_1, \dots, C_5$  are arbitrary constants.

2°. Exact solution:

$$w(r, \theta) = f(r)\theta + g(r).$$

The functions  $f = f(r)$  and  $g = g(r)$  are determined by the system of ordinary differential equations

$$-f'_r \mathbf{L}(f) + f[\mathbf{L}(f)]'_r = \nu r \mathbf{L}^2(f), \quad (1)$$

$$-g'_r \mathbf{L}(f) + f[\mathbf{L}(g)]'_r = \nu r \mathbf{L}^2(g), \quad (2)$$

where  $\mathbf{L}(f) = r^{-1}(r f'_r)'_r$ .

Exact solution of system (1)–(2):

$$f(r) = C_1 \ln r + C_2, \quad g(r) = C_3 r^2 + C_4 \ln r + C_5 \int \left[ \int r Q(r) dr \right] \frac{dr}{r} + C_6,$$

$$Q(r) = \int r^{(C_2/\nu)-1} \exp\left(\frac{C_1}{2\nu} \ln^2 r\right) dr,$$

where  $C_1, \dots, C_6$  are arbitrary constants.

## B.7.2. Nonstationary Hydrodynamic Equations

$$1. \quad \frac{\partial}{\partial t} (\Delta w) + \frac{\partial w}{\partial y} \frac{\partial}{\partial x} (\Delta w) - \frac{\partial w}{\partial x} \frac{\partial}{\partial y} (\Delta w) = \nu \Delta \Delta w, \quad \Delta w = \frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2}.$$

The two-dimensional equations of steady-state flow of a viscous incompressible fluid (nonstationary Navier–Stokes equations) can be reduced to this equation by introducing a stream function  $w$ .

1°. Exact solution:

$$w(x, y, t) = F(y, t)x + G(y, t), \quad (1)$$

where the functions  $F(y, t)$  and  $G = G(y, t)$  are determined from the system of fourth-order one-dimensional equations

$$\frac{\partial^3 F}{\partial t \partial y^2} + \frac{\partial F}{\partial y} \frac{\partial^2 F}{\partial y^2} - F \frac{\partial^3 F}{\partial y^3} = \nu \frac{\partial^4 F}{\partial y^4}, \quad (2)$$

$$\frac{\partial^3 G}{\partial t \partial y^2} + \frac{\partial G}{\partial y} \frac{\partial^2 F}{\partial y^2} - F \frac{\partial^3 G}{\partial y^3} = \nu \frac{\partial^4 G}{\partial y^4}. \quad (3)$$

Equation (2) is independent of (3). Integrating (2) and (3) with respect to  $y$  yields

$$\frac{\partial^2 F}{\partial t \partial y} + \left( \frac{\partial F}{\partial y} \right)^2 - F \frac{\partial^2 F}{\partial y^2} = \nu \frac{\partial^3 F}{\partial y^3} + f_1(t), \quad (4)$$

$$\frac{\partial^2 G}{\partial t \partial y} + \frac{\partial F}{\partial y} \frac{\partial G}{\partial y} - F \frac{\partial^2 G}{\partial y^2} = \nu \frac{\partial^3 G}{\partial y^3} + f_2(t), \quad (5)$$

where  $f_1(t)$  and  $f_2(t)$  are arbitrary functions. Equation (5) is linear in  $G$ . The change of variable

$$G = \int U dy - hF + h'_t y, \quad \text{where } U = U(y, t), \quad F = F(y, t), \quad (6)$$

with  $h = h(t)$  satisfying the linear ordinary differential equation

$$h''_{tt} - f_1(t)h = f_2(t), \quad (7)$$

brings (5) to the second-order homogenous linear equation

$$\frac{\partial U}{\partial t} = \nu \frac{\partial^2 U}{\partial y^2} + F \frac{\partial U}{\partial y} - \frac{\partial F}{\partial y} U. \quad (8)$$

So, if a particular solution of equation (2) or (4) is known, then determining the function  $G$  reduces to solving the linear equations (7)–(8) followed by integrating in accordance with (6).

Table B7 lists exact solutions of equation (2). The ordinary differential equations in the last two rows, which determine a traveling wave solution and a self-similar solution, are autonomous and hence admit reduction of order.

The general solution of the nonhomogeneous equation (7) can be found with the aid of the fundamental system of solutions for the corresponding homogeneous equation (with  $f_2 \equiv 0$ ). The necessary formulas and fundamental solutions of the homogeneous equation (7) that correspond to all exact solutions of equation (2) listed in Table B7 can be found in the handbooks by Kamke (1977) and Polyanin and Zaitsev (1995).

Equation (8) for any function  $F = F(y, t)$  has the trivial solution,  $U = 0$ . The expressions in Table B7 and relation (6) with  $U = 0$  define some exact solutions of the form (1). By analyzing nontrivial solutions of equation (8), one can obtain a wider class of exact solutions.

Table B8 lists transformations that simplify equation (8) for some of the solutions of equation (2) [or (4)] given in Table B7. One can see that in the first two cases, solutions to equation (8) are expressed in terms of solutions to the classical constant coefficient heat equation. In the remaining three cases, the equation reduces to a separable equation.

2°. Exact solution of a more general form:

$$w(x, y, t) = F(\xi, t)x + G(\xi, t), \quad \xi = y + kx,$$

where the functions  $F(\xi, t)$  and  $G = G(\xi, t)$  are determined from the system of fourth-order one-dimensional equations

$$\frac{\partial^3 F}{\partial t \partial \xi^2} + \frac{\partial F}{\partial \xi} \frac{\partial^2 F}{\partial \xi^2} - F \frac{\partial^3 F}{\partial \xi^3} = \nu(k^2 + 1) \frac{\partial^4 F}{\partial \xi^4}, \quad (9)$$

$$\frac{\partial^3 G}{\partial t \partial \xi^2} + \frac{\partial G}{\partial \xi} \frac{\partial^2 F}{\partial \xi^2} - F \frac{\partial^3 G}{\partial \xi^3} = \nu(k^2 + 1) \frac{\partial^4 G}{\partial \xi^4} + 4\nu k \frac{\partial^3 F}{\partial \xi^3} + \frac{2k}{k^2 + 1} \left( F \frac{\partial^2 F}{\partial \xi^2} - \frac{\partial^2 F}{\partial t \partial \xi} \right). \quad (10)$$

Integrating equations (9) and (10) with respect to  $\xi$  yields

$$\frac{\partial^2 F}{\partial t \partial \xi} + \left( \frac{\partial F}{\partial \xi} \right)^2 - F \frac{\partial^2 F}{\partial \xi^2} = \nu(k^2 + 1) \frac{\partial^3 F}{\partial \xi^3} + f_1(t), \quad (11)$$

$$\frac{\partial^2 G}{\partial t \partial \xi} + \frac{\partial F}{\partial \xi} \frac{\partial G}{\partial \xi} - F \frac{\partial^2 G}{\partial \xi^2} = \nu(k^2 + 1) \frac{\partial^3 G}{\partial \xi^3} + Q(\xi, t), \quad (12)$$

TABLE B7

Exact solution of equations (2) and (4);  $\varphi(t)$ ,  $\psi(t)$  are arbitrary functions and  $A$ ,  $B$ ,  $\lambda$  are arbitrary constants

| No | Function $F = F(y, t)$<br>(or general form of solution)                                                                                | Function $f_1(t)$<br>in equation (4) | Determining coefficients<br>(or determining equation)                             |
|----|----------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------|-----------------------------------------------------------------------------------|
| 1  | $F = \varphi(t)y + \psi(t)$                                                                                                            | $f_1(t) = \varphi'_t + \varphi^2$    | N/A                                                                               |
| 2  | $F = \frac{6\nu}{y + \psi(t)} + \psi'_t(t)$                                                                                            | $f_1(t) = 0$                         | N/A                                                                               |
| 3  | $F = A \exp[-\lambda y - \lambda \psi(t)] + \psi'_t(t) + \nu \lambda$                                                                  | $f_1(t) = 0$                         | N/A                                                                               |
| 4  | $F = \frac{A \exp[-\lambda y + \lambda \psi(t)] + 1}{\lambda t + B} - \psi'_t(t) + \nu \lambda$                                        | $f_1(t) = 0$                         | N/A                                                                               |
| 5  | $F = \frac{\beta + A \exp[-\lambda y + \lambda \psi(t)]}{1 + B \exp(-\lambda \beta t)} - \psi'_t(t) + \nu \lambda - \beta$             | $f_1(t) = 0$                         | $\beta$ is an arbitrary constant                                                  |
| 6  | $F = A e^{-\beta t} \sin[\lambda y + \lambda \psi(t)] + \psi'_t(t)$                                                                    | $f_1(t) = B e^{-2\beta t}$           | $\beta = \nu \lambda^2$ , $B = A^2 \lambda^2 > 0$                                 |
| 7  | $F = A e^{-\beta t} \cos[\lambda y + \lambda \psi(t)] + \psi'_t(t)$                                                                    | $f_1(t) = B e^{-2\beta t}$           | $\beta = \nu \lambda^2$ , $B = A^2 \lambda^2 > 0$                                 |
| 8  | $F = A e^{\beta t} \sinh[\lambda y + \lambda \psi(t)] + \psi'_t(t)$                                                                    | $f_1(t) = B e^{2\beta t}$            | $\beta = \nu \lambda^2$ , $B = A^2 \lambda^2 > 0$                                 |
| 9  | $F = A e^{\beta t} \cosh[\lambda y + \lambda \psi(t)] + \psi'_t(t)$                                                                    | $f_1(t) = B e^{2\beta t}$            | $\beta = \nu \lambda^2$ , $B = -A^2 \lambda^2 < 0$                                |
| 10 | $F = \psi(t) e^{\lambda y} - \frac{A e^{\beta t - \lambda y}}{4 \lambda^2 \psi(t)} + \frac{\psi'_t(t)}{\lambda \psi(t)} - \nu \lambda$ | $f_1(t) = A e^{\beta t}$             | $\beta = 2 \nu \lambda^2$                                                         |
| 11 | $F = F(\xi)$ , $\xi = y + \lambda t$                                                                                                   | $f_1(t) = A$                         | $-A + \lambda F''_{\xi\xi} + (F'_\xi)^2 - F F''_{\xi\xi} = \nu F'''_{\xi\xi\xi}$  |
| 12 | $F = t^{-1/2} [H(\xi) - \frac{1}{2} \xi]$ , $\xi = y t^{-1/2}$                                                                         | $f_1(t) = A t^{-2}$                  | $\frac{3}{4} - A - 2 H'_\xi + (H'_\xi)^2 - H H''_{\xi\xi} = \nu H'''_{\xi\xi\xi}$ |

where  $f_1(t)$  is an arbitrary function and

$$Q(\xi, t) = 4\nu k \frac{\partial^2 F}{\partial \xi^2} - \frac{2k}{k^2 + 1} \frac{\partial F}{\partial t} + \frac{2k}{k^2 + 1} \int F \frac{\partial^2 F}{\partial \xi^2} d\xi + f_2(t) \quad [f_2(t) \text{ is any}].$$

Equation (12) is linear in  $G$ . The change of variable  $U = \frac{\partial G}{\partial \xi}$  takes it to the second-order linear equation

$$\frac{\partial U}{\partial t} = \nu(k^2 + 1) \frac{\partial^2 U}{\partial \xi^2} + F \frac{\partial U}{\partial \xi} - \frac{\partial F}{\partial \xi} U + Q(\xi, t). \quad (13)$$

So, if a particular solution of equation (9) or (11) is known, then determining the function  $G$  reduces to solving the linear equation (13). Scaling the independent variables by the formulas  $\xi = (k^2 + 1)\zeta$  and  $t = (k^2 + 1)\tau$ , one can reduce equation (9) to equation (2) in which  $y$  and  $t$  must be replaced by  $\zeta$  and  $\tau$ , respectively. Exact solutions of equation (2) are listed in [Table B7](#).



TABLE B8

Transformations of equation (8) for the corresponding exact solutions of equation (4)  
[the number in the first column corresponds to the  
number of the exact solution  $F = F(y, t)$  in Table B7]

| No | Transformations of equation (8)                                                                                                                                    | Resulting equation                                                                                                                                         |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 1  | $U = \frac{1}{\Phi(t)}u(z, \tau), \quad \tau = \int \Phi^2(t) dt,$<br>$z = y\Phi(t) + \int \psi(t)\Phi(t) dt, \quad \Phi(t) = \exp\left[\int \varphi(t) dt\right]$ | $\frac{\partial u}{\partial \tau} = \nu \frac{\partial^2 u}{\partial z^2}$                                                                                 |
| 2  | $U = \zeta^{-3}u(\zeta, t), \quad \zeta = y + \psi(t)$                                                                                                             | $\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial \zeta^2}$                                                                                |
| 3  | $U = e^\eta Z(\eta, t), \quad \eta = -\lambda y - \lambda \psi(t)$                                                                                                 | $\frac{\partial Z}{\partial t} = \nu \lambda^2 \frac{\partial^2 Z}{\partial \eta^2} + (\nu \lambda^2 - A \lambda e^\eta) \frac{\partial Z}{\partial \eta}$ |
| 11 | $U = u(\xi, t), \quad \xi = y + \lambda t$                                                                                                                         | $\frac{\partial u}{\partial t} = \nu \frac{\partial^2 u}{\partial \xi^2} + [F(\xi) - \lambda] \frac{\partial u}{\partial \xi} - F'_\xi(\xi)u$              |
| 12 | $U = t^{-1/2}u(\xi, \tau), \quad \xi = yt^{-1/2}, \quad \tau = \ln t$                                                                                              | $\frac{\partial u}{\partial \tau} = \nu \frac{\partial^2 u}{\partial \xi^2} + H(\xi) \frac{\partial u}{\partial \xi} + [1 - H'_\xi(\xi)]u$                 |

3°. Exact solution [special case of (1)]:

$$\begin{aligned}
 w(x, y, t) &= e^{-\lambda y} [f(t)x + g(t)] + \varphi(t)x + \psi(t)y + \chi(t), \\
 f(t) &= C_1 E(t), \quad E(t) = \exp \left[ \nu \lambda^2 t - \lambda \int \varphi(t) dt \right], \\
 g(t) &= C_2 E(t) - C_1 E(t) \int \psi(t) dt,
 \end{aligned}$$

where  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are arbitrary functions and  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary parameters.

4°. Exact solution:

$$\begin{aligned}
 w(x, y, t) &= e^{-\lambda y} [A(t)e^{\beta x} + B(t)e^{-\beta x}] + \varphi(t)x + \psi(t)y + \chi(t), \\
 A(t) &= C_1 \exp \left[ \nu(\lambda^2 + \beta^2)t - \beta \int \psi(t) dt - \lambda \int \varphi(t) dt \right], \\
 B(t) &= C_2 \exp \left[ \nu(\lambda^2 + \beta^2)t + \beta \int \psi(t) dt - \lambda \int \varphi(t) dt \right],
 \end{aligned}$$

where  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are arbitrary functions and  $C_1$ ,  $C_2$ ,  $\lambda$ , and  $\beta$  are arbitrary parameters.

5°. Exact solution:

$$w(x, y, t) = e^{-\lambda y} [A(t) \sin(\beta x) + B(t) \cos(\beta x)] + \varphi(t)x + \psi(t)y + \chi(t),$$

where  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are arbitrary functions,  $\lambda$  and  $\beta$  are arbitrary parameters, and the functions  $A(t)$  and  $B(t)$  are determined by the nonautonomous system of linear ordinary differential equations

$$\begin{aligned}
 A'_t &= [\nu(\lambda^2 - \beta^2) - \lambda\varphi(t)] A + \beta\psi(t)B, \\
 B'_t &= [\nu(\lambda^2 - \beta^2) - \lambda\varphi(t)] B - \beta\psi(t)A.
 \end{aligned} \tag{14}$$

The general solution of system (14) is given by

$$\begin{aligned}
 A(t) &= \exp \left[ \nu(\lambda^2 - \beta^2)t - \lambda \int \varphi dt \right] \left[ C_1 \sin \left( \beta \int \psi dt \right) + C_2 \cos \left( \beta \int \psi dt \right) \right], \\
 B(t) &= \exp \left[ \nu(\lambda^2 - \beta^2)t - \lambda \int \varphi dt \right] \left[ C_1 \cos \left( \beta \int \psi dt \right) - C_2 \sin \left( \beta \int \psi dt \right) \right],
 \end{aligned}$$

where  $\varphi = \varphi(t)$  and  $\psi = \psi(t)$ ;  $C_1$  and  $C_2$  are arbitrary constants. In particular, if  $\varphi = \frac{\nu}{\lambda}(\lambda^2 - \beta^2)$  and  $\psi = a$ , we obtain a periodic solution

$$\begin{aligned} A(t) &= C_1 \sin(a\beta t) + C_2 \cos(a\beta t), \\ B(t) &= C_1 \cos(a\beta t) - C_2 \sin(a\beta t). \end{aligned}$$

6°. Exact solutions:

$$w(x, y, t) = A(t) \exp(k_1 x + \lambda_1 y) + B(t) \exp(k_2 x + \lambda_2 y) + \varphi(t)x + \psi(t)y + \chi(t),$$

where  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are arbitrary functions and  $k_1$ ,  $\lambda_1$ ,  $k_2$ , and  $\lambda_2$  are arbitrary parameters that satisfy one of the two relations

$$\begin{aligned} k_1^2 + \lambda_1^2 &= k_2^2 + \lambda_2^2 \quad (\text{first family of solutions}), \\ k_1 \lambda_2 &= k_2 \lambda_1 \quad (\text{second family of solutions}), \end{aligned}$$

and the functions  $A(t)$  and  $B(t)$  are determined by the ordinary differential equations

$$\begin{aligned} A'_t &= [\nu(k_1^2 + \lambda_1^2) + \lambda_1 \varphi(t) - k_1 \psi(t)] A, \\ B'_t &= [\nu(k_2^2 + \lambda_2^2) + \lambda_2 \varphi(t) - k_2 \psi(t)] B. \end{aligned}$$

These equations are easy to integrate:

$$\begin{aligned} A(t) &= C_1 \exp \left[ \nu(k_1^2 + \lambda_1^2)t + \lambda_1 \int \varphi(t) dt - k_1 \int \psi(t) dt \right], \\ B(t) &= C_2 \exp \left[ \nu(k_2^2 + \lambda_2^2)t + \lambda_2 \int \varphi(t) dt - k_2 \int \psi(t) dt \right]. \end{aligned}$$

7°. Exact solution:

$$w(x, y, t) = [C_1 \sin(\lambda x) + C_2 \cos(\lambda x)] [A(t) \sin(\beta y) + B(t) \cos(\beta y)] + \varphi(t)x + \chi(t),$$

where  $\varphi(t)$  and  $\chi(t)$  are arbitrary functions,  $C_1$ ,  $C_2$ ,  $\lambda$ , and  $\beta$  are arbitrary parameters, and the functions  $A(t)$  and  $B(t)$  are determined by the nonautonomous system of linear ordinary differential equations

$$\begin{aligned} A'_t &= -\nu(\lambda^2 + \beta^2)A - \beta \varphi(t)B, \\ B'_t &= -\nu(\lambda^2 + \beta^2)B + \beta \varphi(t)A. \end{aligned} \tag{15}$$

The general solution of system (15) is given by

$$\begin{aligned} A(t) &= \exp[-\nu(\lambda^2 + \beta^2)t] \left[ C_3 \sin \left( \beta \int \varphi dt \right) + C_4 \cos \left( \beta \int \varphi dt \right) \right], \quad \varphi = \varphi(t), \\ B(t) &= \exp[-\nu(\lambda^2 + \beta^2)t] \left[ -C_3 \cos \left( \beta \int \varphi dt \right) + C_4 \sin \left( \beta \int \varphi dt \right) \right], \end{aligned}$$

where  $C_3$  and  $C_4$  are arbitrary constants.

8°. Exact solution:

$$w(x, y, t) = [C_1 \sinh(\lambda x) + C_2 \cosh(\lambda x)] [A(t) \sin(\beta y) + B(t) \cos(\beta y)] + \varphi(t)x + \chi(t),$$

where  $\varphi(t)$  and  $\chi(t)$  are arbitrary functions,  $C_1$ ,  $C_2$ ,  $\lambda$ , and  $\beta$  are arbitrary parameters, and the functions  $A(t)$  and  $B(t)$  are determined by the nonautonomous system of linear ordinary differential equations

$$\begin{aligned} A'_t &= \nu(\lambda^2 - \beta^2)A - \beta \varphi(t)B, \\ B'_t &= \nu(\lambda^2 - \beta^2)B + \beta \varphi(t)A. \end{aligned} \tag{16}$$

The general solution of system (16) is given by

$$\begin{aligned} A(t) &= \exp[\nu(\lambda^2 - \beta^2)t] \left[ C_3 \sin \left( \beta \int \varphi dt \right) + C_4 \cos \left( \beta \int \varphi dt \right) \right], \quad \varphi = \varphi(t), \\ B(t) &= \exp[\nu(\lambda^2 - \beta^2)t] \left[ -C_3 \cos \left( \beta \int \varphi dt \right) + C_4 \sin \left( \beta \int \varphi dt \right) \right], \end{aligned}$$

where  $C_3$  and  $C_4$  are arbitrary constants.

9°. Exact solution:

$$w(x, y, t) = u(z, t) + \varphi(t)x + \psi(t)y, \quad z = kx + \lambda y,$$

where  $\varphi(t)$  and  $\psi(t)$  are arbitrary functions,  $k$  and  $\lambda$  are arbitrary parameters, and the function  $u(z, t)$  is determined by the fourth-order linear equation

$$\frac{\partial^3 u}{\partial t \partial z^2} + [k\psi(t) - \lambda\varphi(t)] \frac{\partial^3 u}{\partial z^3} = \nu(k^2 + \lambda^2) \frac{\partial^4 u}{\partial z^4}.$$

The transformation

$$U(\xi, t) = \frac{\partial^2 u}{\partial z^2}, \quad \xi = z - \int [k\psi(t) - \lambda\varphi(t)] dt$$

brings it to the customary heat equation

$$\frac{\partial U}{\partial t} = \nu(k^2 + \lambda^2) \frac{\partial^2 U}{\partial \xi^2}.$$

⊙ Reference: A. D. Polyanin (2001d).

$$2. \quad \frac{\partial Q}{\partial t} + \frac{1}{r} \frac{\partial w}{\partial \theta} \frac{\partial Q}{\partial r} - \frac{1}{r} \frac{\partial w}{\partial r} \frac{\partial Q}{\partial \theta} = \nu \Delta Q, \quad Q = \Delta w = \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial w}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 w}{\partial \theta^2}.$$

The two-dimensional equations of steady-state flow of a viscous incompressible fluid written in polar coordinates are reduced to this equation ( $w$  is the stream function).

Exact solution:

$$w(r, \theta, t) = f(r, t)\theta + g(r, t).$$

The functions  $f = f(r, t)$  and  $g = g(r, t)$  satisfy the system of equations

$$\mathbf{L}(f_t) - r^{-1} f_r \mathbf{L}(f) + r^{-1} f [\mathbf{L}(f)]_r = \nu \mathbf{L}^2(f), \quad (1)$$

$$\mathbf{L}(g_t) - r^{-1} g_r \mathbf{L}(f) + r^{-1} f [\mathbf{L}(g)]_r = \nu \mathbf{L}^2(g), \quad (2)$$

where the subscripts  $r$  and  $t$  denote partial derivatives;  $\mathbf{L}(f) = r^{-1}(r f_r)_r$  and  $\mathbf{L}^2(f) = \mathbf{L}\mathbf{L}(f)$ .

For the particular solution  $f = \varphi(t) \ln r + \psi(t)$  of equation (1), with  $\varphi$  and  $\psi$  arbitrary, equation (2) can be reduced by the change of variable  $U = \mathbf{L}(g)$  to a second-order linear equation.

## B.8. Higher-Order Nonlinear Equations

### B.8.1. Equations of the Form $\frac{\partial w}{\partial t} = F\left(x, t, w, \frac{\partial w}{\partial x}, \dots, \frac{\partial^n w}{\partial x^n}\right)$

$$1. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + bw \ln w + f(t)w.$$

1°. Exact solution:

$$w(x, t) = \exp \left[ A e^{bt} x + B e^{bt} + \frac{a A^n}{b(n-1)} e^{nbt} + e^{bt} \int e^{-bt} f(t) dt \right],$$

where  $A$  and  $B$  are arbitrary constants.

2°. Exact solution:

$$w(x, t) = \exp \left[ A e^{bt} + e^{bt} \int e^{-bt} f(t) dt \right] \varphi(z), \quad z = x + \lambda t,$$

where  $A$  and  $\lambda$  are arbitrary constants and the function  $\varphi = \varphi(z)$  is determined from the autonomous ordinary differential equation

$$a\varphi_z^{(n)} - \lambda\varphi'_z + b\varphi \ln \varphi = 0,$$

whose order can be reduced by one.

$$2. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + bw \ln w + [f(x) + g(t)]w.$$

Exact solution in multiplicative form:

$$w(x, t) = \exp \left[ C e^{bt} + e^{bt} \int e^{-bt} g(t) dt \right] \varphi(x),$$

where  $C$  is an arbitrary constant and the function  $\varphi(t)$  is determined by the ordinary differential equation

$$a\varphi_x^{(n)} + b\varphi \ln \varphi + f(x)\varphi = 0.$$

$$3. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + f(t)w \ln w + g(t)w.$$

Exact solution:

$$w(x, t) = \exp[\varphi(t)x + \psi(t)].$$

The functions  $\varphi(t)$  and  $\psi(t)$  are determined from

$$\varphi(t) = Ae^F, \quad \psi(t) = Be^F + e^F \int e^{-F} (aA^n e^{nF} + g) dt, \quad F = \int f dt,$$

where  $A$  and  $B$  are arbitrary constants.

$$4. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + f(t)w \ln w + [g(t)x + h(t)]w.$$

Exact solution:

$$w(x, t) = \exp[\varphi(t)x + \psi(t)].$$

The functions  $\varphi(t)$  and  $\psi(t)$  are determined from

$$\begin{aligned} \varphi(t) &= Ae^F + e^F \int e^{-F} g dt, \quad F = \int f dt, \\ \psi(t) &= Be^F + e^F \int e^{-F} (a\varphi^n + h) dt, \end{aligned}$$

where  $A$  and  $B$  are arbitrary constants.

$$5. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw + f(t).$$

1°. Exact solution:

$$w(x, t) = \varphi_2(t)x^2 + \varphi_1(t)x + \varphi_0(t),$$

where the functions  $\varphi_k(t)$  satisfy an appropriate system of ordinary differential equations.

2°. Exact solution:

$$w(x, t) = Ae^{ct} + e^{ct} \int e^{-ct} f(t) dt + \Theta(\xi), \quad \xi = x + \lambda t,$$

where  $A$  and  $\lambda$  are arbitrary constants, and the function  $\Theta(\xi)$  is determined by solving the autonomous ordinary differential equation

$$a\Theta_\xi^{(n)} + b(\Theta'_\xi)^2 - \lambda\Theta'_\xi + c\Theta = 0.$$

$$6. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw \frac{\partial w}{\partial x} + kw^2 + f(t)w + g(t).$$

Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\lambda x),$$

where  $\lambda$  is a root of the quadratic equation  $b\lambda^2 + c\lambda + k = 0$ , and the functions  $\varphi(t)$  and  $\psi(t)$  are determined from the system of first-order ordinary differential equations

$$\varphi'_t = k\varphi^2 + f(t)\varphi + g(t), \quad (1)$$

$$\psi'_t = [(c\lambda + 2k)\varphi + f(t) + a\lambda^n]\psi. \quad (2)$$

For the Riccati equation (1), see Kamke (1977) and Polyanin and Zaitsev (1995). It is integrable in quadrature if, for example,

$$(a) \quad k = 0, \quad (b) \quad g(t) \equiv 0, \quad (c) \quad f(t) = \text{const}, \quad g(t) = \text{const}.$$

On solving equation (1), one can readily solve equation (2), which is linear in  $\psi$ .

$$7. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + b f(t) w^2 + g(t) w + h(t).$$

1°. Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\pm x\sqrt{-b}), \quad b < 0,$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following first-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\varphi'_t = b f \varphi^2 + g \varphi + h, \quad (1)$$

$$\psi'_t = [2b f \varphi + g + a(\pm\sqrt{-b})^n] \psi. \quad (2)$$

Equation (1) is a Riccati equation for  $\varphi = \varphi(t)$ ; it can be reduced to a second-order linear equation. A lot of exact solutions to equation (1) with various  $f$ ,  $g$ , and  $h$  can be found in Kamke (1977) and Polyanin and Zaitsev (1995).

Given a solution of equation (1), the corresponding solution of (2) is calculated by

$$\psi(t) = C \exp \left[ a(\pm\sqrt{-b})^n t + \int (2b f \varphi + g) dt \right],$$

where  $C$  is an arbitrary constant.

2°. Exact solution of a more general form:

$$w(x, t) = \varphi(t) + \psi(t) \exp(x\sqrt{-b}) + \chi(t) \exp(-x\sqrt{-b}), \quad b < 0, \quad (3)$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = b f \varphi^2 + g \varphi + h + 4b f \psi \chi, \quad (4)$$

$$\psi'_t = [2b f \varphi + g + a(\sqrt{-b})^n] \psi, \quad (5)$$

$$\chi'_t = [2b f \varphi + g + a(-\sqrt{-b})^n] \chi. \quad (6)$$

For equations of even order with  $n = 2m$  ( $m = 1, 2, \dots$ ), it follows from (5) and (6) that the functions  $\psi(t)$  and  $\chi(t)$  are proportional. Setting  $\psi(t) = A\theta(t)$  and  $\chi(t) = B\theta(t)$ , one can rewrite solution (3) as

$$w(x, t) = \varphi(t) + \theta(t) [A \exp(x\sqrt{-b}) + B \exp(-x\sqrt{-b})], \quad b < 0,$$

where the functions  $\varphi(t)$  and  $\theta(t)$  are determined by the system of ordinary differential equations

$$\varphi'_t = b f (\varphi^2 + 4AB\theta^2) + g \varphi + h, \quad (7)$$

$$\theta'_t = [2b f \varphi + g + (-1)^m a b^m] \theta. \quad (8)$$

On expressing  $\varphi$  from (8) in terms of  $\theta$  and substituting the result into (7), one arrives at a second-order nonlinear equation for  $\theta$ ; if  $f$ ,  $g$ ,  $h = \text{const}$ , this equation is autonomous and hence admits reduction of order.

3°. Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \cos(x\sqrt{b}) + \chi(t) \sin(x\sqrt{b}), \quad b > 0,$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\chi(t)$  are determined by a system of ordinary differential equations (not specified here).

For equations of even order with  $n = 2m$  ( $m = 1, 2, \dots$ ), there are exact solutions with the following form ( $c$  is any number):

$$w(x, t) = \varphi(t) + \theta(t) \cos(x\sqrt{b} + c), \quad b > 0.$$

The functions  $\varphi(t)$  and  $\theta(t)$  are determined by the following system of first-order ordinary differential equations with variable coefficients:

$$\varphi'_t = b f(\varphi^2 + \theta^2) + g\varphi + h, \quad (9)$$

$$\theta'_t = [2b f\varphi + g + (-1)^m ab^m] \theta. \quad (10)$$

On expressing  $\varphi$  from (10) in terms of  $\theta$  and substituting the result into (9), one arrives at a second-order nonlinear equation for  $\theta$ ; if  $f, g, h = \text{const}$ , this equation is autonomous and hence admits reduction of order.

$$8. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + f\left(x, \frac{\partial w}{\partial x}\right) + g(t).$$

Exact solution in additive form:

$$w(x, t) = At + B + \int g(t) dt + \varphi(x).$$

Here,  $A$  and  $B$  are arbitrary constants and the function  $\varphi(x)$  is determined from the nonlinear ordinary differential equation

$$a\varphi_x^{(n)} + f(x, \varphi'_x) - A = 0.$$

$$9. \quad \frac{\partial w}{\partial t} = a \frac{\partial^n w}{\partial x^n} + f\left(x, \frac{\partial w}{\partial x}\right) + bw + g(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(x) + Ae^{bt} + e^{bt} \int e^{-bt} g(t) dt.$$

Here,  $A$  is an arbitrary constant and the function  $\varphi(x)$  is determined from the nonlinear ordinary differential equation

$$a\varphi_x^{(n)} + f(x, \varphi'_x) + b\varphi = 0.$$

$$10. \quad \frac{\partial w}{\partial t} = aw \frac{\partial^n w}{\partial x^n} + f(t)w + g(t).$$

1°. Exact solution:

$$w(x, t) = F(t)(A_{n-1}x^{n-1} + \dots + A_1x + A_0) + F(t) \int \frac{g(t)}{F(t)} dt, \quad F(t) = \exp \left[ \int f(t) dt \right],$$

where  $A_0, A_1, \dots, A_{n-1}$  are arbitrary constants.

2°. Exact solution:

$$w(x, t) = \varphi(t)(x^n + A_{n-1}x^{n-1} + \dots + A_1x + A_0) + \varphi(t) \int \frac{g(t)}{\varphi(t)} dt,$$

$$\varphi(t) = F(t) \left[ C - an! \int F(t) dt \right]^{-1}, \quad F(t) = \exp \left[ \int f(t) dt \right],$$

where  $A_0, A_1, \dots, A_{n-1}$ , and  $C$  are arbitrary constants.

$$11. \quad \frac{\partial w}{\partial t} = aw \frac{\partial^n w}{\partial x^n} + bw^2 + f(t)w + g(t).$$

Exact solution:

$$w(x, t) = \varphi(t)\Theta(x) + \psi(t),$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined from the following system of first-order ordinary differential equations ( $C$  is any number):

$$\begin{aligned}\varphi'_t &= C\varphi^2 + b\varphi\psi + f(t)\varphi, \\ \psi'_t &= C\varphi\psi + b\psi^2 + f(t)\psi + g(t),\end{aligned}$$

and the function  $\Theta(x)$  is determined by the  $n$ th-order linear ordinary differential equation

$$a\Theta_x^{(n)} + b\Theta = C.$$

$$12. \quad \frac{\partial w}{\partial t} = aw \frac{\partial^n w}{\partial x^n} + f(x)w \frac{\partial w}{\partial x} + g(t)w + h(t).$$

Exact solution:

$$w(x, t) = \varphi(t)\Theta(x) + \psi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\Theta(x)$  are determined by the ordinary differential equations

$$\begin{aligned}\varphi'_t &= C\varphi^2 + g(t)\varphi, \\ \psi'_t &= [C\varphi + g(t)]\psi + h(t), \\ a\Theta_x^{(n)} + f(x)\Theta'_x &= C,\end{aligned}$$

where  $C$  is an arbitrary constant. Integrating successively, for  $\varphi(t)$  and  $\psi(t)$  we obtain

$$\begin{aligned}\varphi(t) &= G(t) \left[ A - C \int G(t) dt \right]^{-1}, \quad G(t) = \exp \left[ \int g(t) dt \right], \\ \psi(t) &= B\varphi(t) + \varphi(t) \int \frac{h(t)}{\varphi(t)} dt,\end{aligned}$$

where  $A$  and  $B$  are arbitrary constants.

### B.8.2. Equations of the Form $\frac{\partial^2 w}{\partial t^2} = F\left(x, t, w, \frac{\partial w}{\partial x}, \dots, \frac{\partial^n w}{\partial x^n}\right)$

$$1. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + f(x) \frac{\partial w}{\partial x} + bw \ln w + [g(x) + h(t)]w.$$

Exact solution in multiplicative form:

$$w(x, t) = \varphi(t)\psi(x).$$

The functions  $\varphi(t)$  and  $\psi(x)$  are determined by the ordinary differential equations

$$\begin{aligned}\varphi''_{tt} - [b \ln \varphi + h(t) + C]\varphi &= 0, \\ a\psi_x^{(n)} + f(x)\psi'_x + [b \ln \psi + g(x) - C]\psi &= 0,\end{aligned}$$

where  $C$  is an arbitrary constant.

$$2. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw + f(t).$$

1°. Exact solution:

$$w(x, t) = \varphi_2(t)x^2 + \varphi_1(t)x + \varphi_0(t),$$

where the functions  $\varphi_k(t)$  satisfy an appropriate system of ordinary differential equations.

2°. Exact solution:

$$w(x, t) = \psi(t) + \Theta(\xi), \quad \xi = x + \lambda t.$$

The functions  $\psi(t)$  and  $\Theta(\xi)$  are determined from the ordinary differential equations

$$\psi''_{tt} - c\psi - f(t) = A,$$

$$a\Theta^{(n)}_{\xi} - \lambda^2 \Theta''_{\xi\xi} + b(\Theta'_{\xi})^2 + c\Theta = A,$$

where  $\lambda$  and  $A$  are arbitrary constants.

$$3. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + b \left( \frac{\partial w}{\partial x} \right)^2 + cw \frac{\partial w}{\partial x} + kw^2 + f(t)w + g(t).$$

Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\lambda x),$$

where  $\lambda$  is a root of the quadratic equation  $b\lambda^2 + c\lambda + k = 0$ , and the functions  $\varphi(t)$  and  $\psi(t)$  are determined from the system of second-order ordinary differential equations

$$\varphi''_{tt} = k\varphi^2 + f(t)\varphi + g(t), \quad (1)$$

$$\psi''_{tt} = [(c\lambda + 2k)\varphi + f(t) + a\lambda^n] \psi. \quad (2)$$

In the special case  $f(t) = \text{const}$  and  $g(t) = \text{const}$ , equation (1) is autonomous and has particular solutions of the form  $\varphi = \text{const}$  and, hence, can be integrated in quadrature. Equation (2) is linear in  $\psi$ ; therefore, for  $\varphi = \text{const}$ , its general solution is expressed in terms of exponentials or sine and cosine.

$$4. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + f(x) \left( \frac{\partial w}{\partial x} \right)^2 + g(x) + h(t).$$

Exact solution in additive form:

$$w(x, t) = \frac{1}{2}At^2 + Bt + C + \int_0^t (t - \tau)h(\tau) d\tau + \varphi(x).$$

Here,  $A$ ,  $B$ , and  $C$  are arbitrary constants, and the function  $\varphi(x)$  is determined by solving the nonlinear ordinary differential equation

$$a\varphi^{(n)}_x + f(x)(\varphi'_x)^2 + g(x) - A = 0.$$

$$5. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + f(x) \left( \frac{\partial w}{\partial x} \right)^2 + bw + g(x) + h(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(t) + \psi(x).$$

The functions  $\varphi(t)$  and  $\psi(x)$  are determined by solving the nonlinear ordinary differential equations

$$\varphi''_{tt} - b\varphi - h(t) = 0,$$

$$a\psi^{(n)}_x + f(x)(\psi'_x)^2 + b\psi + g(x) = 0.$$

The general solution of the first equation is given by

$$\varphi(t) = C_1 \cosh(kx) + C_2 \sinh(kx) + \frac{1}{k} \int_0^t h(\tau) \sinh[k(t - \tau)] d\tau \quad \text{for } b = k^2 > 0,$$

$$\varphi(t) = C_1 \cos(kx) + C_2 \sin(kx) + \frac{1}{k} \int_0^t h(\tau) \sin[k(t - \tau)] d\tau \quad \text{for } b = -k^2 < 0,$$

where  $C_1$  and  $C_2$  are arbitrary constants.



$$6. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^{2n} w}{\partial x^{2n}} + f(t) \left( \frac{\partial w}{\partial x} \right)^2 + b f(t) w^2 + g(t) w + h(t).$$

1°. Exact solution:

$$w(x, t) = \varphi(t) + \psi(t) \exp(\pm x \sqrt{-b}), \quad b < 0,$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following first-order ordinary differential equations with variable coefficients (the arguments of  $f$ ,  $g$ , and  $h$  are not specified):

$$\varphi''_{tt} = b f \varphi^2 + g \varphi + h, \quad (1)$$

$$\psi''_{tt} = [2b f \varphi + g + (-1)^n a b^n] \psi. \quad (2)$$

In the special case where  $f$ ,  $g$ ,  $h$  are constant, equation (1) has particular solutions of the form  $\varphi = \text{const}$ . Here, the general solution of equation (2) is expressed in terms of exponentials or sine and cosine.

2°. Exact solution of a more general form:

$$w(x, t) = \varphi(t) + \psi(t) [A \exp(x \sqrt{-b}) + B \exp(-x \sqrt{-b})], \quad b < 0,$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of second-order ordinary differential equations with variable coefficients:

$$\varphi''_{tt} = b f (\varphi^2 + 4AB\psi^2) + g \varphi + h, \quad (3)$$

$$\psi''_{tt} = [2b f \varphi + g + (-1)^n a b^n] \psi. \quad (4)$$

On expressing  $\varphi$  from (4) in terms of  $\psi$  and substituting the result into (3), one arrives at a fourth-order nonlinear equation for  $\psi$ ; if  $f$ ,  $g$ ,  $h = \text{const}$ , this equation is autonomous and hence admits reduction of order.

3°. Exact solution ( $c$  is an arbitrary constant):

$$w(x, t) = \varphi(t) + \psi(t) \cos(x \sqrt{b} + c), \quad b > 0,$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by solving the following system of second-order ordinary differential equations with variable coefficients:

$$\varphi''_{tt} = b f (\varphi^2 + \psi^2) + g \varphi + h,$$

$$\psi''_{tt} = [2b f \varphi + g + (-1)^n a b^n] \psi.$$

$$7. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + f \left( x, \frac{\partial w}{\partial x} \right) + g(t).$$

Exact solution in additive form:

$$w(x, t) = \frac{1}{2} A t^2 + B t + C + \int_0^t (t - \tau) g(\tau) d\tau + \varphi(x).$$

Here,  $A$ ,  $B$ , and  $C$  are arbitrary constants and the function  $\varphi(x)$  is determined by the nonlinear ordinary differential equation

$$a \varphi_x^{(n)} + f(x, \varphi'_x) - A = 0.$$

$$8. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + f\left(x, \frac{\partial w}{\partial x}\right) + bw + g(t).$$

Exact solution in additive form:

$$w(x, t) = \varphi(t) + \psi(x).$$

The functions  $\varphi(t)$  and  $\psi(x)$  are determined by the ordinary differential equations

$$\begin{aligned}\varphi''_{tt} - b\varphi - g(t) &= 0, \\ a\psi^{(n)}_x + f(x, \psi'_x) + b\psi &= 0.\end{aligned}$$

The general solution of the first equation is given by

$$\begin{aligned}\varphi(t) &= C_1 \cosh(kx) + C_2 \sinh(kx) + \frac{1}{k} \int_0^t g(\tau) \sinh[k(t - \tau)] d\tau \quad \text{for } b = k^2 > 0, \\ \varphi(t) &= C_1 \cos(kx) + C_2 \sin(kx) + \frac{1}{k} \int_0^t g(\tau) \sin[k(t - \tau)] d\tau \quad \text{for } b = -k^2 < 0,\end{aligned}$$

where  $C_1$  and  $C_2$  are arbitrary constants.

$$9. \quad \frac{\partial^2 w}{\partial t^2} = a \frac{\partial^n w}{\partial x^n} + w f\left(t, \frac{1}{w} \frac{\partial w}{\partial x}\right).$$

Exact solution in multiplicative form:

$$w(x, t) = e^{\lambda x} \varphi(t),$$

where  $\lambda$  is an arbitrary constant and the function  $\varphi(t)$  is determined from the second-order linear ordinary differential equation

$$\varphi''_{tt} = [a\lambda^n + f(t, \lambda)]\varphi.$$

$$10. \quad \frac{\partial^2 w}{\partial t^2} = aw \frac{\partial^n w}{\partial x^n} + f(t)w + g(t).$$

Exact solution:

$$w(x, t) = \varphi(t)(A_n x^n + \cdots + A_1 x) + \psi(t),$$

where  $A_1, \dots, A_n$  are arbitrary constants and the functions  $\varphi(t)$  and  $\psi(t)$  are determined from the second-order ordinary differential equations

$$\begin{aligned}\varphi''_{tt} &= A_n a n! \varphi^2 + f(t)\varphi, \\ \psi''_{tt} &= A_n a n! \varphi \psi + f(t)\psi + g(t).\end{aligned}$$

$$11. \quad \frac{\partial^2 w}{\partial t^2} = aw \frac{\partial^n w}{\partial x^n} + bw^2 + f(t)w + g(t).$$

Exact solution:

$$w(x, t) = \varphi(t)\Theta(x) + \psi(t),$$

where the functions  $\varphi(t)$  and  $\psi(t)$  are determined by the following system of second-order ordinary differential equations ( $C$  is an arbitrary constant):

$$\begin{aligned}\varphi''_{tt} &= C\varphi^2 + b\varphi\psi + f(t)\varphi, \\ \psi''_{tt} &= C\varphi\psi + b\psi^2 + f(t)\psi + g(t).\end{aligned}$$

The function  $\Theta(x)$  satisfies the  $n$ th-order linear ordinary differential equation

$$a\Theta^{(n)}_x + b\Theta = C.$$

$$12. \quad \frac{\partial^2 w}{\partial t^2} = aw \frac{\partial^n w}{\partial x^n} + f(x)w \frac{\partial w}{\partial x} + g(t)w + h(t).$$

Exact solution:

$$w(x, t) = \varphi(t)\Theta(x) + \psi(t),$$

where the functions  $\varphi(t)$ ,  $\psi(t)$ , and  $\Theta(x)$  are determined by the ordinary differential equations

$$\begin{aligned}\varphi''_{tt} &= C\varphi^2 + g(t)\varphi, \\ \psi''_{tt} &= [C\varphi + g(t)]\psi + h(t), \\ a\Theta_x^{(n)} + f(x)\Theta'_x &= C,\end{aligned}$$

where  $C$  is an arbitrary constant.

### B.8.3. Other Equations

$$1. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = f(x) \frac{\partial^n w}{\partial y^n}.$$

This is a special case of equation B.8.3.3.

Exact solution:

$$w(x, y) = \varphi(x)e^{\lambda y} - \lambda^{n-2} \int f(x) dx + C,$$

where  $\varphi(x)$  is an arbitrary function;  $C$  and  $\lambda$  are arbitrary constants.

$$2. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = f(x) \frac{\partial^{2n} w}{\partial y^{2n}} + g(x).$$

This is a special case of equation B.8.3.3.

Exact solution:

$$w(x, y) = \varphi(x)e^{\lambda y} - \frac{1}{2\lambda^2\varphi(x)} \left[ \int g(x) dx + C_1 \right] e^{-\lambda y} - \lambda^{2n-2} \int f(x) dx + C_2,$$

where  $\varphi(x)$  is an arbitrary function and  $C_1$ ,  $C_2$ , and  $\lambda$  are arbitrary parameters.

$$3. \quad \frac{\partial w}{\partial y} \frac{\partial^2 w}{\partial x \partial y} - \frac{\partial w}{\partial x} \frac{\partial^2 w}{\partial y^2} = F\left(x, w, \frac{\partial w}{\partial y}, \dots, \frac{\partial^n w}{\partial y^n}\right).$$

1°. If  $w(x, y)$  is a solution of the equation in question, then the function

$$w_1(x, y) = w(x, y + \varphi(x)),$$

where  $\varphi(x)$  is an arbitrary function, is also a solution of the equation.

2°. Let the right-hand side of the equation be independent of  $x$  explicitly. Then there are exact solutions of the form

$$w = w(z), \quad z = y + \varphi(x),$$

where  $\varphi(x)$  is an arbitrary function and  $w(z)$  is a solution of the ordinary differential equation  $F(w, w'_z, \dots, w_z^{(n)}) = 0$ .

3°. Let the right-hand side of the equation be independent of  $x$  and  $w$  explicitly. Then there are exact solutions of the form

$$w = Cx + g(z), \quad z = y + \varphi(x),$$

where  $\varphi(x)$  is an arbitrary function,  $C$  is an arbitrary constant, and  $g(z)$  is a solution of the ordinary differential equation  $F(g'_z, \dots, g_z^{(n)}) + Cg''_{zz} = 0$ .

$$4. \quad \frac{\partial w}{\partial t} = wF(t, \zeta_0, \zeta_1, \dots, \zeta_n), \quad \zeta_k = \sum_{i=k}^n \frac{(-1)^{i+k}}{k!(i-k)!} x^{i-k} \frac{\partial^i w}{\partial x^i}, \quad k = 0, 1, \dots, n.$$

Exact solution in multiplicative form:

$$w(x, t) = (C_0 + C_1 x + \dots + C_n x^n) \varphi(t),$$

where  $C_0, C_1, \dots, C_n$  are arbitrary constants and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation

$$\varphi'_t = \varphi F(t, C_0 \varphi, C_1 \varphi, \dots, C_n \varphi).$$

⊙ *Reference:* Ph. W. Doyle (1996), the case  $\partial_t F \equiv 0$  was considered.

$$5. \quad \frac{\partial^2 w}{\partial t^2} = wF(t, \zeta_0, \zeta_1, \dots, \zeta_n), \quad \zeta_k = \sum_{i=k}^n \frac{(-1)^{i+k}}{k!(i-k)!} x^{i-k} \frac{\partial^i w}{\partial x^i}, \quad k = 0, 1, \dots, n.$$

Exact solution in multiplicative form:

$$w(x, t) = (C_0 + C_1 x + \dots + C_n x^n) \varphi(t),$$

where  $C_0, C_1, \dots, C_n$  are arbitrary constants and the function  $\varphi = \varphi(t)$  satisfies the ordinary differential equation

$$\varphi''_{tt} = \varphi F(t, C_0 \varphi, C_1 \varphi, \dots, C_n \varphi).$$

$$6. \quad F\left(x, \frac{1}{w} \frac{\partial w}{\partial x}, \dots, \frac{1}{w} \frac{\partial^n w}{\partial x^n}; \frac{1}{w} \frac{\partial w}{\partial y}, \dots, \frac{1}{w} \frac{\partial^m w}{\partial y^m}\right) = 0.$$

Exact solution in multiplicative form:

$$w(x, y) = A e^{\lambda y} \varphi(x),$$

where  $A$  and  $\lambda$  are arbitrary constants and the function  $\varphi(x)$  is determined from the  $n$ th-order ordinary differential equation

$$F(x, \varphi'_x/\varphi, \dots, \varphi^{(n)}_x/\varphi; \lambda, \dots, \lambda^m) = 0.$$

$$7. \quad F\left(x, \frac{1}{w} \frac{\partial w}{\partial x}, \dots, \frac{1}{w} \frac{\partial^n w}{\partial x^n}; \frac{1}{w} \frac{\partial^2 w}{\partial y^2}, \dots, \frac{1}{w} \frac{\partial^{2m} w}{\partial y^{2m}}\right) = 0.$$

1°. Exact solution:

$$w(x, y) = [A \cosh(\lambda y) + B \sinh(\lambda y)] \varphi(x),$$

where  $A, B$ , and  $\lambda$  are arbitrary constants and the function  $\varphi(x)$  is determined by solving the  $n$ th-order ordinary differential equation

$$F(x, \varphi'_x/\varphi, \dots, \varphi^{(n)}_x/\varphi; \lambda^2, \dots, \lambda^{2m}) = 0.$$

2°. Exact solution:

$$w(x, y) = [A \cos(\lambda y) + B \sin(\lambda y)] \varphi(x),$$

where  $A, B$ , and  $\lambda$  are arbitrary constants and the function  $\varphi(x)$  is determined by solving the  $n$ th-order ordinary differential equation

$$F(x, \varphi'_x/\varphi, \dots, \varphi^{(n)}_x/\varphi; -\lambda^2, \dots, (-1)^m \lambda^{2m}) = 0.$$

$$8. \quad f_1\left(x, \frac{\partial w}{\partial x}, \dots, \frac{\partial^n w}{\partial x^n}\right) + f_2\left(y, \frac{\partial w}{\partial y}, \dots, \frac{\partial^m w}{\partial y^m}\right) = k w.$$

Exact solution in additive form:

$$w(x, y) = \varphi(x) + \psi(y).$$

The functions  $\varphi(x)$  and  $\psi(y)$  are determined by the ordinary differential equations

$$f_1(x, \varphi'_x, \dots, \varphi^{(n)}_x) - k\varphi = C,$$

$$f_2(y, \psi'_y, \dots, \psi^{(m)}_y) - k\psi = -C,$$

where  $C$  is an arbitrary constant.

$$9. \quad f_1\left(x, \frac{1}{w} \frac{\partial w}{\partial x}, \dots, \frac{1}{w} \frac{\partial^n w}{\partial x^n}\right) + w^k f_2\left(y, \frac{1}{w} \frac{\partial w}{\partial y}, \dots, \frac{1}{w} \frac{\partial^m w}{\partial y^m}\right) = 0.$$

Exact solution in multiplicative form:

$$w(x, y) = \varphi(x)\psi(y).$$

The functions  $\varphi(x)$  and  $\psi(y)$  are determined by the ordinary differential equations

$$\begin{aligned} \varphi^{-k} f_1\left(x, \varphi'_x/\varphi, \dots, \varphi_x^{(n)}/\varphi\right) &= C, \\ \psi^k f_2\left(y, \psi'_y/\psi, \dots, \psi_y^{(m)}/\psi\right) &= -C, \end{aligned}$$

where  $C$  is an arbitrary constant.

$$10. \quad f_1\left(x, \frac{\partial w}{\partial x}, \dots, \frac{\partial^n w}{\partial x^n}\right) + e^{\lambda w} f_2\left(y, \frac{\partial w}{\partial y}, \dots, \frac{\partial^m w}{\partial y^m}\right) = 0.$$

Exact solution in additive form:

$$w(x, y) = \varphi(x) + \psi(y).$$

The functions  $\varphi(x)$  and  $\psi(y)$  are determined by the ordinary differential equations

$$\begin{aligned} e^{-\lambda \varphi} f_1\left(x, \varphi'_x, \dots, \varphi_x^{(n)}\right) &= C, \\ e^{\lambda \psi} f_2\left(y, \psi'_y, \dots, \psi_y^{(m)}\right) &= -C, \end{aligned}$$

where  $C$  is an arbitrary constant.

$$11. \quad f_1\left(x, \frac{1}{w} \frac{\partial w}{\partial x}, \dots, \frac{1}{w} \frac{\partial^n w}{\partial x^n}\right) + f_2\left(y, \frac{1}{w} \frac{\partial w}{\partial y}, \dots, \frac{1}{w} \frac{\partial^m w}{\partial y^m}\right) = k \ln w.$$

Exact solution in multiplicative form:

$$w(x, y) = \varphi(x)\psi(y).$$

The functions  $\varphi(x)$  and  $\psi(y)$  are determined by the ordinary differential equations

$$\begin{aligned} f_1\left(x, \varphi'_x/\varphi, \dots, \varphi_x^{(n)}/\varphi\right) - k \ln \varphi &= C, \\ f_2\left(y, \psi'_y/\psi, \dots, \psi_y^{(m)}/\psi\right) - k \ln \psi &= -C, \end{aligned}$$

where  $C$  is an arbitrary constant.